

Supplementary Material

Use of grit supplements by waterbirds: an experimental assessment of strategies to reduce lead poisoning

Monica Martinez-Haro^{§, #}, *Andy J. Green*[#], *Pelayo Acevedo*[‡], *Rafael Mateo*^{§, *}

[§] Instituto de Investigación en Recursos Cinegéticos, IREC (CSIC-UCLM-JCCM), Ronda de Toledo s/n, 13071 Ciudad Real, Spain

[#] Department of Wetland Ecology, Estación Biológica de Doñana, EBD (CSIC), Américo Vespucio s/n, 41092 Sevilla, Spain

[‡] Biogeography, Diversity, and Conservation Research Team, Department of Animal Biology, Faculty of Science, University of Málaga, 29071 Málaga, Spain.

* Corresponding author:

tlf: +34 926 29 54 50

fax: +34 926 24 54 51

e-mail: rafael.mateo@uclm.es

No. Pages – 2

No. Tables – 1

Table S1. Mass of grit taken by non-waterbirds species from the control treatments of different experiments (2-4) carried out in the Concerted Nature Reserve of Cañada de los Pájaros and Medina Lagoon (Andalusia, southern Spain).

Controls	Treatment	N	Grit taken (g)	
			Mean	SE
Experiment 2 (Semi-captive)	Grit-food-water	11	2.0	1.0
	Grit-food-dry	12	3.0	1.0
	Grit-water	3	0.0	0.0
	Grit-dry	3	2.0	1.0
Experiment 3 (dry medium) (Semi-captive)	Si grit	6	0.5	0.2
	Ca grit	6	0.7	0.4
	Si grit-food	9	3.1	0.9
	Ca grit-food	9	5.6	1.9
Experiment 3 (adding water) (Semi-captive)	Si grit	6	0.5	0.3
	Ca grit	6	0.8	0.2
	Si grit-food	9	3.8	0.6
	Ca grit-food	9	2.6	0.7
Experiment 4 (Wild)	Shore	7	0.0	0.0
	Lagoon	7	0.3	0.2
	Platform	7	0.1	0.1
	Shore-food	7	3.6	0.3
	Lagoon-food	7	1.3	0.5
	Platform-food	7	1.1	0.3