

1 **Comparative mapping of the Oregon Wolfe Barley using doubled haploid lines derived**
2 **from female and male gametes**

3 L. Cistué^{1, 2}, S. Chao³, B. Echávarri², Y. Chutimanitsakun¹, A. Corey¹, A. Cuesta-Marcos¹, T.
4 Filichkina¹, N. Garcia-Mariño^{1, 2}, I. Romagosa⁴ and P.M. Hayes¹

5

6 1. Department of Crop and Soil Sciences. Oregon State University. Corvallis, OR 97331. USA

7 2. Departamento de Genética y Producción Vegetal. Estación Experimental de Aula Dei.
8 Consejo Superior de Investigaciones Científicas. Apartado 202. 50059 Zaragoza, Spain.

9 3. USDA-ARS Biosciences Research Lab. 1605 Albrecht Blvd. Fargo, North Dakota. USA

10 4. Centre UdL-IRTA. Universitat de Lleida. Avda. Alcalde Rovira Roure 91, 25198 Lleida,
11 Spain.

12

13

14 Corresponding author: lcistue@eead.csic.es

15

16 Key words: *Hordeum vulgare*, *Hordeum bulbosum*, anther culture, doubled haploids, segregation
17 distortion, quantitative trait loci, Oregon Wolfe Barley.

18

19

20

21

1 **Abstract**

2 The Oregon Wolfe Barley mapping population is a resource for genetics research and instruction.
3 Prior reports are based on a population of doubled haploid (DH) lines developed by the *Hordeum*
4 *bulbosum* (H.b.) method, which samples female gametes. We developed new DH lines from the
5 same cross using anther culture (A.C.), which samples male gametes. Linkage maps were
6 generated in each of the two subpopulations using the same 1,328 single nucleotide
7 polymorphism (SNP) markers. The linkage maps based on DH lines derived from the products of
8 megasporogenesis and microsporogenesis revealed minor differences in terms of estimated
9 recombination rates. There were no differences in locus ordering. There was greater segregation
10 distortion in the A.C.-derived subpopulation than in the H.b.-derived subpopulation, but in the
11 region showing the greatest distortion, the cause was more likely allelic variation at the *ZEO1*
12 plant height locus rather than to DH production method. The effects of segregation distortion and
13 pleiotropy had greater impacts on estimates of QTL effect than population size for reproductive
14 fitness traits assayed under greenhouse conditions. The OWB population and data are
15 community resources. Seed is available from three distribution centers located in North America,
16 Europe, and Asia. Details on ordering seed sets, as well as complete genotype and phenotype
17 data files, are available at (<http://wheat.pw.usda.gov/ggpages/maps/OWB/>).

1 **Introduction**

2 Doubled haploid (DH) techniques, by accelerating the approach to homozygosity, are a useful
3 tool for conventional and molecular plant breeding (Thomas 2003; Forster et al. 2007). In the
4 case of barley, DH populations have been widely used for constructing the linkage maps that
5 underlie Quantitative Trait Locus (QTL) mapping and marker assisted selection (MAS)
6 (Karakousis et al. 2003; Wenzl et al. 2006; Stein et al. 2007; Hearnden et al. 2007; Varshney et al.
7 2007; Szücs et al. 2009; Close et al. 2009; and <http://wheat.pw.usda.gov/GG2/index.shtml>).

8 Barley DH populations can be produced using female or male gametes. The former involves the
9 interspecific crossing of an F₁ (as the female) with *Hordeum bulbosum*, followed by embryo
10 rescue, plant regeneration and artificial chromosome doubling (Kasha and Kao 1970). The latter
11 involves using the F₁ (as the male) followed by anther or microspore culture (Maluszynski et al.
12 2003). The potential efficiencies of androgenetic systems are much greater than those of
13 gynogenetic systems because each inflorescence produces more pollen than eggs.

14 The principal issues related to the use of DH populations for linkage map construction relate to
15 segregation and recombination. Segregation distortion - the deviation of observed genotypic
16 frequencies from their expected values - complicates the application of genetic theory and
17 analysis (Lu et al. 2002). The allele transmission and gamete survival frequencies that cause
18 segregation distortion can be caused by exogenous factors such as temperature (Xu et al. 1997)
19 and *in vitro* culture conditions (Graner et al. 1991; Foisset and Delourne 1996; Manninen 2000).
20 Distortion can also be caused by genetic factors (Lambrides et al. 2004; Törjék et al. 2006) and
21 may be more prevalent in some species than in others (Lu et al. 2002; Marshall et al. 2007). The
22 implications of segregation distortion for genetic analyses and breeding are reviewed by Xian-
23 Liang et al. (2006). To generalize - in the case of androgenetic systems in wheat, barley and rice
24 - it appears that selection for genes favoring microspore growth and development in culture
25 media may not have negative effects on agronomic traits in the derived populations (Ma et al.
26 1999; Guzy-Wrobelska and Szarejo 2003; Cistué et al. 2005; Lapitan et al. 2009).

27 Representative linkage maps require “normal” rates of recombination between homologous
28 chromosomes (Lenormand and Dutheil 2005). The levels of recombination observed in DH
29 populations derived by the *Hordeum bulbosum* technique (hereafter referred to as H.b.) and

1 anther culture (hereafter referred to as A.C.) reveal the crossover frequencies in the megaspore
2 and microspore mother cells. For barley, Devaux et al. (1995) found 1.047 and 0.912
3 recombination events per chromosome in A.C.- and H.b.- derived DH populations, respectively.
4 Recent studies have demonstrated the effects of various factors on rates of recombination and the
5 distribution of recombination breakpoints in plants (Li et al. 2007). Genome regions where
6 recombination rates are significantly higher or lower than the genome average are termed
7 recombination hot- and cold-spots, respectively (Mezard 2006) and in barley the relationship
8 between genetic and physical maps was explored in depth by Kuenzel et al. (2000).

9 Of the many barley linkage mapping populations available (summarized in GrainGenes;
10 <http://wheat.pw.usda.gov/GG2/index.shtml>), one of the most widely-used is the Oregon Wolfe
11 Barley (OWB). This population of DH lines was developed by the H.b. technique from the F₁ of
12 the cross between the dominant and recessive morphological marker stocks developed by R.
13 Wolfe (Wolfe 1972). The alternative alleles at the morphological traits loci determining the
14 major germplasm groups of barley are represented in this population (Costa et al. 2001) and the
15 sequential addition of new generations of markers (e.g. restriction fragment length
16 polymorphisms (RFLP), amplified length fragment polymorphisms (AFLPs), simple sequence
17 repeats (SSRs), diversity array technologies (DArTs), single nucleotide polymorphisms (SNPs),
18 and restriction site associated DNAs (RADs) has allowed the population to serve as a resource
19 for linkage map and QTL integration. Szücs et al. (2009) integrated prior marker data with 1,472
20 SNPs represented in the three barley PilotOPAs (Oligonucleotide Pooled Assay) (POPA1, 2 and
21 3) and Chutimanitsakun et al. (submitted) added over 450 RAD loci to the Szücs et al. (2009)
22 map. Qualitative and quantitative loci determining morphological, phenological, and disease
23 resistance traits of importance to barley improvement have been mapped in the OWB (Börner et
24 al. 2002; Costa et al. 2001; Jafary et al. 2008; Rostocks et al. 2005; Stein et al. 2007). However,
25 as empirically demonstrated by Vales et al. (2005), limited population size can lead to
26 underestimation of QTL number, overestimation of QTL effects, and failure to quantify QTL
27 interactions.

28 The goal of this project was to increase the size of the OWB mapping population in order to
29 improve it as a resource for genetic mapping and QTL detection. Since the A.C. technique was
30 used to develop the new DH plants, we were able to compare linkage maps based on A.C. and

1 H.b.-derived subpopulations genotyped with the same high-throughput SNP assays. This allowed
2 for direct comparisons of segregation distortion and linkage distance with the each of the two
3 linkage maps and empirical assessment of improvements in QTL detection afforded by doubling
4 the size of the mapping population.

5

6 **Materials and methods**

7

8 **Plant materials**

9 Complete information on the “Oregon Wolfe Barley” population can be found at
10 <http://barleyworld.org/oregonwolfe.php>. Briefly, the mapping of the original set of 94 H.b.-
11 derived plants was described by Costa et al. (2001), a population of 93 was mapped by Szücs et
12 al. (2009) due to incomplete data on one line, and 82 were mapped by Chutimanitsakun et al.
13 (submitted) after the discovery of nine sets of identical DH lines. The reduction in population
14 size has not affected estimates of locus order, distance, nor estimate of QTL number and effect
15 (see “OWB population size” at (<http://wheat.pw.usda.gov/ggpages/maps/OWB/>)). The discovery
16 of the identical sets of lines was not made until after the research described in this paper was
17 undertaken; as a consequence, this report was intended to compare maps based on 93 H.b.-
18 derived lines and 93 A.C.-derived lines. However, in order not to confound comparisons by the
19 possible effects of identical genotypes, in this report we compare the H.b. (n = 82), A.C. (n = 93)
20 subpopulations and combined populations (n = 175). The 93 A.C.-derived lines were produced
21 from the F₁ of the cross between the recessive (OWB-R) and dominant (OWB-D) morphological
22 marker spring barley genetic stocks (Wolfe 1972; Wolfe and Franckowiak 1991) as described by
23 Cistué et al. (2003). This method includes a stress pretreatment of the anthers with 0.7 M
24 mannitol for four days at 24°C followed by 25 days on a modified FHG culture medium (Hunter
25 1988). Embryos were transplanted to FHG regeneration medium. DH lines were obtained by
26 spontaneous doubling of the chromosome number of each haploid plant.

27 For measuring reproductive fitness traits (see “Phenotypes”, below) the two sets of DH lines
28 (H.b.-derived and A.C.-derived) were grown in separate greenhouse experiments at Oregon State

1 University, Corvallis, Oregon, USA. For both experiments, greenhouse temperatures were
2 $18\pm 1.5^{\circ}\text{C}$ (day and night) with a photoperiod of 16 h day/8 h night provided by supplemental
3 illumination from Sylvania Lumalux-Eco ET18 400 w lights suspended 1.5 meters over the
4 bench surface. Each DH line was replicated twice. The parents were replicated four times.

5 **Markers**

6 The process of developing the high confidence SNP markers has been described in detail by
7 Close et al. (2009). Briefly, SNPs observed in ESTs and sequenced amplicons were used to
8 design three Illumina 1536-plex pilot Oligonucleotide Pool Assays (pilot OPAs; POPA1,
9 POPA2, POPA3). Based on the technical performance and other criteria, 3,072 SNPs were
10 selected from three POPAs to generate two production barley OPAs (BOPA1 and BOPA2). The
11 H.b.-derived OWB population was genotyped with three POPAs, whereas the A.C.-derived
12 population was genotyped with two BOPAs. The common SNPs mapped on both H.b- and A.C-
13 derived populations were used for analysis in this study. The SNP genotyping assays were
14 conducted using the Illumina GoldenGate BeadArray SNP detection platform. The POPA assays
15 were conducted at the Southern California Genotyping Consortium at the University of
16 California, Los Angeles. Genomic DNA extractions for A.C.-derived population and BOPA SNP
17 assays were performed at USDA-ARS Small Grains Genotyping Center in Fargo, ND. The
18 naming convention for SNP loci appearing on the maps shown in this report is by the POPA
19 numbers (e.g. 1_1311), where 1 = the POPA number (POPA1 in this case) and the subsequent
20 four digits correspond to the SNP order in the corresponding POPA.

21 **Mapping**

22 JoinMap 4 (Van Ooijen 2006) was used for map construction. SNP-only maps were constructed
23 for the 82 H.b.-derived DH lines and the 93 A.C.-derived DH lines. The two data sets were then
24 merged and a combined map made based on 175 DH lines. For each of the data sets, at linkage
25 LOD score 5, the 1,328 polymorphic markers formed seven linkage groups. The Monte Carlo
26 maximum likelihood (ML) mapping algorithm was used to determine the marker order in each of
27 the seven linkage groups. Recombination frequencies were converted to centiMorgans (cM)
28 using Haldane's mapping function. The positions of ten morphological loci (*VRS1*, *ZEO1*, *WST1*,

1 *ALM, HSH, SRH, ROB, GBSSI (WX), NUD, and LKS2*) are shown on the A.C.-derived and
2 combined map based on the flanking SNPs reported by Szücs et al. (2009).

3 **Segregation distortion and recombination events**

4 A full genome assessment of segregation distortion involves a series of chi-square tests for
5 equality of allelic frequencies for every individual locus. This requires a multiple test correction
6 of the level for assessing significance of each individual test. A very conservative Bonferroni
7 correction would simply take the significance level for individual markers as the genome-wide
8 level divided by the total number of polymorphic markers, (1,328 in this study) Thus, for 0.05,
9 the required significance level for assessing significance of each test would be
10 $\alpha_{\text{BON}}=0.05/1328\approx 0.00004$. However this approach wrongly assumes independence of markers.
11 Less conservative corrections make an effort to consider dependence between markers, as will be
12 the case with markers within a linkage group. One such method estimates the effective number
13 of independent tests across the genome by dividing the total map length in cM by 20-30 cM,
14 which is an arbitrarily taken distance between any two marker pairs for relative independence
15 (van Eeuwijk, *personal communication*). In this study the number of effective independent test
16 would be approximately equal to 50, that is the ratio between 1,250 cM, reported roughly map
17 length of barley, and 25 cM arbitrarily taken for relative independence. The significance level for
18 individual tests would be equal to the genome-wide significance level divided by 50, the number
19 of putative independent tests. This criterion would lead to an, $\alpha'_{0.05}=0.001$, $\alpha'_{0.01}=0.0002$ and
20 $\alpha'_{0.001} = 0.00002$. Allelic frequencies giving rise to these significance levels are shown in Fig. 3
21 by differential dotted lines.

22 The number of apparent single crossovers (CO) was counted in each linkage group of each of the
23 175 DH. These values underestimate the actual number due to undetected CO in regions of
24 monomorphic markers. An analysis of variance was performed, using GenStat (Payne 2006), to
25 test for significant differences between DH production method and between chromosomes.

26 **Phenotypes**

27 Phenology and reproductive fitness (yield component) traits were measured on each DH line as
28 follows. Heading date (HD) was recorded as the number of days from seedling emergence until
29 the first appearance of awns (or hoods). The number of fertile tillers per plant (spike number;

1 SN) was counted and the spikes were harvested from each fertile tiller. Three fertile spikes were
2 selected at random and used to determine: spike length (SL) in cm; number of florets per spike
3 (floret number; FN); number of grains per spike (grain number; GN); hundred grain weight
4 (HGW) in g; and plant height (PH) in cm.

5 **QTL mapping**

6 QTL analyses for each of the phenotypic characters were conducted for each of the data sets (82
7 DH-H.b.; 93 DH-A.C.; 175 DH-H.b.+A.C.) using the Composite Interval Mapping (CIM)
8 procedure (Zeng 1994) implemented in Windows QTL Cartographer 2.5 (Wang et al. 2001–
9 2003). Skeleton maps - constructed from a set of 622 evenly-distributed and non-co-segregating
10 markers were used for QTL analysis (Supplemental Fig. 2). A forward-selection backward-
11 elimination stepwise regression procedure was used to identify co-factors for CIM. The
12 maximum number of cofactors used was seven. A 30-cM scan window was used for all analyses.
13 Experiment-wise significance likelihood ratio (LR) test statistic thresholds ($P < 0.05$) for QTL
14 identification were determined with 1,000 permutations and expressed as LOD (LOD =
15 $0.217LR$). For every significant QTL, we calculated individual R^2 (proportion of phenotypic
16 variance explained by the individual QTL) and additive effect (expressed as one half of the
17 difference between the two allelic classes). Negative values indicate that the parent line
18 contributing the allele with the highest value was OWB-R. Epistatic interactions between QTL
19 were tested by means of Multiple Interval Mapping (MIM, Kao et al. 1999) using Windows QTL
20 Cartographer 2.5 and a score statistic test with significance level of 0.05. Using MIM we also
21 calculated for each trait the R^2 of the multilocus model that included the QTL detected in the
22 CIM analysis (main and interaction effects). All genotype and phenotype data are available at
23 <http://wheat.pw.usda.gov/ggpages/maps/OWB/>.

24

25 **Results**

26 The numbers of SNPs mapping to each chromosome are shown in Table 1. Of the 3,072 SNPs
27 represented on BOPA1 and BOPA2, a total of 1,328 (43%) were mapped in the H.b.- and A.C.-
28 derived sub-populations. The lengths (in Haldane cM) for five of the seven chromosomes (for
29 the sake of brevity, the term “chromosome” will be considered synonymous with “linkage

1 group”) were longer for the A.C.-derived than they were for the H.b-derived population (Figure
2 2 and Supplemental Figure 1). However, these differences were only significant for 5H (Figure
3 1). For this chromosome, the H.b. map is significantly longer. In the map based on the combined
4 data from the H.b.- and A.C.-derived populations (hereafter referred to as the combined
5 population and/or combined map) cM values are intermediate between those for the two sub-
6 populations. The average number of apparent crossovers (hereafter referred to as crossovers) per
7 chromosome for each of the two subpopulations are shown graphically in Fig. 1; the significant
8 differences between chromosomes (averaged across methods) are apparent, as are the significant
9 interactions between method and chromosome (Supplemental Table 1).

10 Marker orders are consistent between the A.C. and the H.b. maps, although markers co-
11 segregating in one map sometimes showed recombination in the other map. On average, there is
12 one SNP/cM. However, there are regions of monomorphism (e.g. gaps) in both the H.b. and A.C.
13 maps. The largest such gap is on the short arm of chromosome 2H (22.8 and 22.7 cM,
14 respectively) and large gaps were also observed on 6HS and 7HL. These three gaps persist in the
15 combined map. For the purposes of illustration, the chromosome 2H maps for the H.b., A.C., and
16 combined populations are shown in Fig. 2. Linkage maps for the other six chromosomes are
17 shown in Supplemental Fig. 1.

18 The long arm of chromosome 2H was associated with significant segregation distortion in the
19 A.C. population and in the combined population (Fig. 3). The *ZEO-1* locus is associated with the
20 most significant distortion in each case. For example, in the A.C. subpopulation there were 25
21 and 68 plants in each of the allele classes. The region of significant distortion extends from *VRS-*
22 *1* to *WST-1*. Genome-wide allele frequencies for OWB-D (dominant parent) alleles in the H.b.-
23 and A.C.-derived populations are shown in the Figs. 3A and 3B. For the H.b. population, based
24 on the significance threshold calculated by the modified Bonferroni system, there is significant
25 distortion only on chromosome 3H. The distortion was in favor of the OWB-R (recessive parent)
26 alleles. For the A.C. population, there was significant segregation distortion on 2H, 3H, 5H, 6H,
27 and 7H. For 2H, 3H, 6H and 7H, the distortion was in favor of the OWB-R alleles and on 5H it
28 was in favor of the OWB-D allele. In the combined population (Fig. 3C) there was significant
29 distortion on 2H, 3H, and 5H.

1 Considering all seven traits, 22 QTLs were detected in the H.b. subpopulation, 19 in the A.C.
2 subpopulation and 22 in the combined data set (Table 2 and Supplemental Tables 2 and 3).
3 QTLs for all traits, except for HD, were detected on 2H in the two subpopulations and in the
4 combined population (Figs. 4A – C). Based on all three datasets, QTLs for SL and PH were
5 coincident with *ZEO-1*. QTLs for GN, FN, HGW, and SN were coincident with *VRS-1* in the
6 three populations. A QTL for HGW (coincident with *NUD*) on 7H was detected in the three
7 populations. Six QTL (three for number of seeds on 1H, 2H and 6H, one for number of florets on
8 1H and two for heading date on 4H and 6H) were detected in the H.b. and the combined
9 populations but not in the A.C. population. Two QTL (one for number of florets on 3H and one
10 plant height on 4H) were detected in the A.C. and combined populations but not in the H.b.
11 population (Table 2, Supplemental Tables 2 and 3). Four QTL (one for number of seeds on 4H,
12 two for number of florets on 4H and 6H and one for hundred grain weight on 6H) were detected
13 only in the H.b. population (Supplemental Table 2). Five QTL (one for spike length on 7H, one
14 for hundred grain weight on 4H one for plant height on 7H and two for heading date on 3H and
15 7H) were detected only in the A.C. population (Supplemental Table 3). Two QTL for spike
16 length (6H and 7H) were detected only in the combined population (Table 2). Twelve QTL,
17 usually those with the highest effects, were detected for the seven traits in the three populations.
18 For these, LOD scores were always highest in the combined population and lowest in the H.b.
19 population (Table 2 and Supplemental Tables 2 and 3). Based on the score test, no significant
20 epistatic interaction was detected at a significance level of 0.05.

21

22 Discussion

23 The Oregon Wolfe Barley population provides a highly polymorphic and connective mapping
24 resource (Costa et al. 2001; Rostocks et al. 2005; Stein et al. 2007; Szücs et al. 2009; Close et al.
25 2010; Chutimanitsakun et al. submitted) and a unique genetic background for mapping
26 determinants of certain phenotypes (Börner et al. 2002; Jafary et al. 2008). In order to further
27 improve this resource for the genetics community, we increased the size of the mapping
28 population. Larger populations allow for higher resolution linkage maps and better estimates of
29 QTL number, location, effect, and interaction (Melchinger et al. 1998; Vales et al. 2005). One
30 hundred seventy five DH lines (82 previously developed by the *Hordeum bulbosum* technique

1 and 93 by anther culture) are now available. The 175 lines are mapped with the same 1,328
2 SNPs.

3 The development of 93 lines by anther culture afforded an opportunity to empirically assess the
4 effects of DH derivation method on estimates of recombination and linkage map quality.
5 Devaux et al. (1995) reported recombination rates of 1.05 and 0.91 events per chromosome for
6 anther culture and *Hordeum bulbosum*-derived DH lines. Our estimates were 1.78 and 1.72,
7 respectively, for the two techniques. The differences between the two reports may be due to the
8 germplasm (Steptoe x Morex vs. Wolfe dominant marker stock x Wolfe recessive marker stock)
9 and marker density (much higher in our case). In these two barley mapping populations, there are
10 only slightly higher average rates of recombination in barley microsporogenesis vs.
11 megasporogenesis. In contrast, Guzy-Wrobelska et al. (2007) compared recombination
12 frequencies in wheat DH populations obtained via maize pollination and anther culture and
13 reported that there was significantly higher recombination in pollen mother cells. Interestingly,
14 we did find significant differences in recombination rates between chromosomes and an
15 interaction of chromosome and method (Supplemental Table 1). Even if map lengths differ, key
16 issues in linkage map quality are locus order and segregation distortion.

17 There were no differences in locus orders between the A.C. and H.b. maps. We did observe more
18 cases of significant segregation distortion in the A.C.-derived lines. The key question is whether
19 this distortion is due to method or to germplasm. A lack of common markers between
20 populations, and differences in anther/microspore culture protocols between labs, complicates
21 the identification of genome regions associated with doubled haploid production capacity. The
22 region of distortion we observed on 3H is coincident with a QTL for *in vitro* shoot regeneration
23 on 3H (Manninen 2000) and the distorted region on 5H is coincident with a QTL for green plant
24 regeneration (Muñoz-Amatriain et al. 2008). The distortion observed in the A.C.-derived
25 subpopulation may be due to these QTLs. However, the region with the greatest segregation
26 distortion (on chromosome 2H) was observed in the A.C.- derived and the combined populations
27 and this is most likely due to the specific alleles segregating at the Zeocriton 1 (*ZEO1*) locus in
28 the OWB. The dominant X-ray induced mutant allele at this locus causes extreme dwarfism
29 (Lundqvist and Lundqvist 1998). We hypothesize that the *Zeo1* (dominant dwarfing) allele has a
30 negative pleiotropic effect on *in vitro* growth and regeneration. A similar situation, related to

1 negative effects on shoot regeneration in barley, was reported for the dwarfing allele at the *UZU*
2 locus (Rikiishi et al. 2008). Androgenetic doubled haploid production systems are more efficient
3 than gynogenetic methods (Maluszynski et al. 2003; Forster et al. 2007) and our results support
4 that localized segregation distortion is a small price to pay for doubled haploid efficiency.
5 Continued progress in techniques has reduced the incidence of albinism (Torp and Andersen
6 2009; Jacquard et al. 2009), led to the direct regeneration of well-developed embryos (Supena et
7 al. 2008; Cistué et al. 2009), and the isolation of plants from the embryogenic phase rather than
8 the callus phase (Maluszynski et al. 2003). Continued improvements in technique should further
9 reduce the incidence of segregation distortion in A.C.-derived populations.

10 One of the objectives of developing linkage maps is to locate genes determining qualitative and
11 quantitative phenotypes. One of the unique attributes of the OWB is that many of the genes
12 determining the principal germplasm groups of barley are segregating in a single population. On
13 the one hand, this allows for simultaneous mapping of these genes and multiple marker loci. On
14 the other hand, some of these genes are likely to have pleiotropic effects on other phenotypes.
15 This was certainly the case for the reproductive fitness traits measured under greenhouse
16 conditions. Clearly, controlled environment conditions cannot reflect the complexity of
17 conditions encountered under field conditions and it is not appropriate to equate yield component
18 QTLs detected in an exotic cross such as the OWB under greenhouse conditions to
19 agronomically relevant germplasm assayed under field conditions. This is certainly the case for
20 *ZEO1*, which, as shown in Table 2, was coincident with highly significant QTLs for spike length,
21 grain number and plant height (with LODs of 93.4, 5.6 and 53.4, respectively). Dwarfing genes,
22 such as the *sdw1/denso* gene in barley (Jia et al. 2009) and the *Rht* genes of wheat (Febrer et al.
23 2009) are of tremendous agronomic value. In its current background, the dwarfing allele at *ZEO1*
24 has too extreme an effect on plant height and negative pleiotropic effects on spike length and
25 grain number to be of immediate agronomic interest. Furthermore, the pleiotropic effects of the
26 *ZEO1* locus altered the expected pattern of favorable allele effects at the *VRS1* locus, which is
27 the principal determinant of the two-row and six-row germplasm groups of barley.

28 Two-row and six-row refer to the number of fertile florets per rachis node. Although most barley
29 breeders prefer to work within and inflorescence group type, crosses between groups are not
30 uncommon and several important biparental QTL mapping populations have been derived from

1 two-row x six-row crosses (e.g. Cali sib x Bowman (Chen et al. 1994), Gobernadora x CMB643
2 (Zhu et al. 1999), Harrington x Morex (Marquez-Cedillo et al. 2000), and Morex x Barke (Kota
3 et al. 2001). Typically, two-row genotypes have fewer kernels/spike (lower grain number),
4 higher kernel weight and more spikes per plant. In the case of the OWBs, the pleiotropic effects
5 of the *ZEO1* locus on spike length reversed this pattern for grain number. However, the effects in
6 this exotic cross, under greenhouse conditions, were consistent with those reported from
7 agronomically relevant crosses assessed under field conditions for hundred grain weight and
8 spike number (Table 2). Although QTL mapping of reproductive fitness traits in this exotic cross
9 under greenhouse conditions may be of limited practical utility, there is value in the OWB
10 population as a model for QTL analysis and instruction.

11 For example, the full set of 175 lines should be useful for obtaining better estimates of QTL
12 number, effect, and interaction than either of the smaller subpopulations. In the case of the traits
13 reported herein, however, pleiotropy and segregation distortion – attributed to *ZEO1* – may be
14 more responsible for the higher LODs observed for most traits on chromosome 2H in the n= 175
15 population vs. the original population of n= 82 (Fig. 4). The same figure provides an illustrative
16 example of two large-effect QTLs, both with candidate genes, which are on the same
17 chromosome determining the same traits. We expected to optimize estimates of epistatic
18 interaction with the larger population. However, no significant epistasis was detected.

19 In conclusion, the comparisons of linkage maps derived from DH population derived from the
20 products of megasporogenesis and microsporogenesis revealed minor differences in terms of
21 estimated recombination rates and were made possible by the very high quality and repeatability
22 of the SNP data generated by the Illumina Golden Gate assay. For example, the OWB H.b.
23 subpopulation was genotyped with three Pilot OPAS in one laboratory (Close et al. 2009) and
24 the A.C. subpopulation was genotyped in another laboratory using Barley OPAS 1 and 2 (this
25 report) and yet the two data sets were integrated seamlessly. There was greater segregation
26 distortion in the A.C.-derived subpopulation than in the H.b.-derived population, but in the
27 region showing the greatest distortion, the cause was more likely a unique allele at a plant height
28 locus rather than an effect of the DH production method. The effects of segregation distortion
29 and pleiotropy had greater impacts on estimates of QTL effect than population size for the traits
30 studied. The OWB population and data are community resources. Seed is available from three

1 distribution centers located in North America, Europe, and Asia. Details on ordering seed sets, as
2 well as complete genotype and phenotype data files, are available at
3 (<http://wheat.pw.usda.gov/ggpages/maps/OWB/>).

1 **Acknowledgements**

2 L. Cistué was recipient of a Senior Research Fellowship from the Spanish Ministry of Science
3 and Innovation during his time with the Oregon State University Barley Project and his research
4 was financed by the Spanish Ministry of Science and Innovation through the National Plan
5 Projects AGL2005-07195-C02-01 and AGL2008-05541-C02-01. Alfonso Cuesta-Marcos was
6 supported by a postdoctoral fellowship from the Spanish Ministerio de Ciencia e Innovación
7 (MICINN). The BOPA SNPs were developed under the auspices of USDA-CSREES-NRI Grant
8 No 2006- 55606-16722 “Barley Coordinated Agricultural Project: Leveraging Genomics,
9 Genetics, and Breeding for Gene Discovery and Barley Improvement”. This paper is dedicated to
10 Dr. Bob Wolfe, whose vision and persistence led to the development of the Wolfe dominant and
11 recessive marker stocks.

1 **References**

- 2
- 3 Börner A, Buck-Sorlin GH, Hayes PM, Malyshev S, Korzun V (2002) Molecular mapping of
4 major genes and quantitative trait loci determining flowering time in response to photoperiod in
5 barley. *Plant Breed* 121:129-132
- 6 Chen FQ, Prehn D, Hayes PM, Mulrooney D, Corey A, Vivar H (1994) Mapping genes for
7 resistance to barley stripe rust (*Puccinia striiformis* f. sp. *hordei*). *Theor Appl Genet* 88:215-219
- 8 Cistué L, Echávarri B, Batlle F, Soriano M, Castillo A, Vallés MP, Romagosa I (2005)
9 Segregation distortion for agronomic traits in doubled haploid lines of barley. *Plant Breed*
10 124:546-550
- 11 Cistué L, Romagosa I, Batlle F, Echávarri B (2009) Improvements in the production of doubled
12 haploids in durum wheat (*Triticum turgidum* L.) through isolated microspore culture. *Plant Cell*
13 *Rep* 28:727-735
- 14 Cistué L, Vallés MP, Echávarri B, Sanz JM, Castillo AM (2003) Barley anther culture. In:
15 Maluszynski M, Kasha KJ, Forster BP, Szarejko I (2003) Doubled haploid production in crop
16 plants: a manual. Kluwer Academic Publishers, Dordrecht, The Netherlands. pp 29-35.
- 17 Close TJ, Bhat PR, Lonardi S, Wu Y, Rostoks N, Ramsay L, Druka A, Stein N, Svensson JT,
18 Wanamaker S, Bozdag S, Roose ML, Moscou M, Chao S, Varshney RK, Sücs P, Sato K, Hayes
19 PM, Matthews DE, Kleinhofs A, Muehlbauer G, DeYoung J, Marshall D, Madishetty K, Fenton
20 RD, Condamine P, Graner A, Waugh R (2009) Development and implementation of high-
21 throughput SNP genotyping in barley. *BMC Genomics* 10:582
- 22 Costa JM, Corey A, Hayes PM, Jobet C, Kleinhofs A, Kopisch-Obusch A, Kramer SF, Kudrna
23 D, Li M, Riera-Lizarazu O, Sato K, Szücs P, Toojinda T, Vales MI, Wolfe RI (2001) Molecular
24 mapping of the Oregon Wolfe Barleys: a phenotypically polymorphic doubled-haploid
25 population. *Theor Appl Genet* 103:415–424
- 26 Devaux P, Kilian A, Kleinhofs A (1995) Comparative mapping of the barley genome with male
27 and female recombination-derived, doubled haploid populations. *Mol Gen Genet* 249:600–608

- 1 Febrer M, Wilhelm E, Al-Kaff N, Wright J, Powell W, Bevan MW, Boulton MI (2009) Rapid
2 identification of the three homoeologues of the wheat dwarfing gene *Rht* using a novel PCR-
3 based screen of three-dimensional BAC pools. *Genome* 52:993-1000.
- 4 Foisset N, Delourne R (1996) Segregation distortion in androgenic plants. In: Jain SM, Spory
5 SK, and Veilleux RE (eds), *In Vitro Haploid Production in Higher Plants*, Kluwer Academic
6 Publishers, Dordrecht 2:189—201
- 7 Forster BP, Heberle-Bors E, Kasha KJ, Touraev A (2007) The resurgence of haploids in higher
8 plants. *Trends in Plant Sci* 12:368-375
- 9 Graner A, Jahoor A, Schondelmaier J, Siedler H, Pillen K, Fischbeck G, Wenzel G, Herrmann
10 RG (1991) Construction of an RFLP map of barley. *Theor Appl Genet* 83:250-256
- 11 Guzy–Wrobelska J, Szarejko I (2003) Molecular and agronomic evaluation of wheat doubled
12 haploid lines obtained through maize pollination and anther culture methods. *Plant Breed*
13 122:305–313
- 14 Hearnden PR, Eckermann PJ, McMichael GL, Hayden MJ, Eglinton JK, Chalmers KJ (2007) A
15 genetic map of 1,000 SSR and DArT markers in a wide barley cross. *Theor Appl Genet* 115:383-
16 391
- 17 Hunter CP (1988) Plant regeneration from microspores of barley, *Hordeum vulgare*. PhD Thesis,
18 Wye College, University of London, London.
- 19 Jacquard C, Nolin F, Hécart C, Grauda D, Rashal I, Dhondt-Cordelier S, Sangwan RS, Devaux P,
20 Mazeyrat-Gourbeyre F, Clément C (2009) Microspore embryogenesis and programmed cell
21 death in barley: effects of copper on albinism in recalcitrant cultivars. *Plant Cell Rep* 28:1329-
22 1339
- 23 Jafary H, Albertazzi G, Marcel TC, Niks R (2008) High diversity of genes for nonhost resistance
24 of barley to heterologous rust fungi. *Genetics* 178:2327-2339
- 25 Jia QJ, JJ Zhang, Westcott S, Zhang XQ, Bellgard M, Lance R, Li CD (2009) GA-20 oxidase as
26 a candidate for the semidwarf gene *sdw1/denso* in barley. *Funct Integr Genomics* 9: 255-262.

- 1 Kao CH, Zeng ZB, Teasdale RD (1999) Multiple interval mapping for quantitative trait loci.
2 Genetics 152:1203-1216
- 3 Kasha KJ, Kao KN (1970) High frequency haploid production in barley (*Hordeum vulgare* L.).
4 Nature 225:874-876
- 5 Karakousis A, Gustafson JP, Chalmers KJ, Barr AR, Langridge P (2003) A consensus map of
6 barley integrating SSR, RFLP, and AFLP markers. Aust J Agric Res 54:1173-1185
- 7 Kota R, Varshney RK, Thiel T, Dehmer KJ, Graner A (2001) Generation and comparison of
8 EST-derived SSRs and SNPs in barley (*Hordeum vulgare* L.). J Hered 135:145-151
- 9 Kuenzel G, Korzun L, Meister A (2000) Cytologically integrated physical restriction fragment
10 length polymorphism maps for the barley genome based on translocation breakpoints. Genetics
11 154: 397-412
- 12 Lambrides CJ, Godwin ID, Lawn RJ, Imrie BC (2004) Segregation distortion for seed testa color
13 in mungbean (*Vigna radiata* L. Wilcek). J Hered 95(6):532-535
- 14 Lapitan VC, Redoña ED, Abe T, Brar DS (2009) Molecular characterization and agronomic
15 performance of DH lines from the F1 of *indica* and *japonica* cultivar of rice (*Oryza sativa* L.).
16 Field Crops Res 112:222-228
- 17 Lenormand T, Dutheil L (2005) Recombination difference between sexes: a role for haploid
18 selection. PLOS Biol 3:396-403
- 19 Li J, Hsia A, Schnable PS (2007) Recent advances in plant recombination. Curr Opin Plant Biol
20 10:131-135
- 21 Lu H, Romero-Severson J, Bernardo R (2002) Chromosomal regions associated with segregation
22 distortion in maize. Theor Appl Genet 105:622–628
- 23 Lundqvist U, Lundqvist A (1998) Intermedium mutants in barley (*Hordeum vulgare* L.):
24 diversity, interactions and plant breeding value. J Appl Genet 39:85-96

- 1 Ma H, Busch RH, Riera-Lizarazu O, Rines HW, Dill-Macky R (1999) Agronomic performance
2 of lines derived from anther culture, maize pollination and single-seed descent in a spring wheat
3 cross. *Theor Appl Genet* 99:432–436
- 4 Maluszynski M, Kasha KJ, Forster BP, Szarejko I (2003) Doubled haploid production in crop
5 plants: a manual. Kluwer Academic Publishers, Dordrecht, The Netherlands pp 21-52
- 6 Manninen OM (2000) Associations between anther-culture response and molecular markers on
7 chromosomes 2H, 3H and 4H of barley (*Hordeum vulgare* L.). *Theor Appl Genet* 100:57-62
- 8 Marquez-Cedillo LA, Hayes PM, Kleinhofs A, Legge WG, Rossnagel BG, Sato K, Ullrich SE,
9 Wesenberg DM (2001) QTL analysis of agronomic traits in barley based on the doubled haploid
10 progeny of two elite North American varieties representing different germplasm groups. *Theor*
11 *Appl Genet* 103:625–637
- 12 Marshall DL, Marieken G. M. Shaner MGM, Oliva JP (2007) Effects of pollen load size on seed
13 paternity in wild radish: the roles of pollen competition and mate choice. *Evolution* 61:1925-
14 1937
- 15 Melchinger AE, Utz HF, Schon CC (1998) Quantitative Trait Locus (QTL) mapping using
16 different testers and independent population samples in maize reveals low power of QTL
17 detection and large bias in estimates of QTL effects. *Genetics* 149:383-403
18
- 19 Mezard C (2006) Meiotic recombination hotspots in plants. *Biochem Soc Trans* 34:531-534
- 20 Muñoz-Amatriain M, Castillo AM, Chen XW, Cistue L, Valles MP (2008) Identification and
21 validation of QTLs for green plant percentage in barley (*Hordeum vulgare* L.) anther culture.
22 *Mol Breed* 22:119-129
- 23 Payne RW (2006) *The Guide to GenStat Release 9*. VSN International, Hertfordshire.
- 24 Rikiishi K, Saisho D, Takeda K (2008) *Uzu*, a barley semi-dwarf gene, suppresses plant
25 regeneration in calli derived from immature embryos. *Breed Sci* 58:149-155
- 26 Rostoks N, Mudie S, Cardle L, Russell J, Ramsay L, Booth A, Svensson JT, Wanamaker SI,
27 Walia H, Rodriguez EM, Hedley PE, Liu H, Morris J, Close TJ, Marshall DF, Waugh R (2005)

- 1 Genome-wide SNP discovery and linkage analysis in barley based on genes responsive to abiotic
2 stress. *Mol Genet Genomics* 274:515-527
- 3 Stein N, Prasad M, Scholz U, Thiel T, Zhang H, Wolf M, Kota R, Varshney RK, Perovic D,
4 Grosse I, Graner A (2007) A 1,000-loci transcript map of the barley genome: new anchoring
5 points for integrative grass genomics *Theor Appl Genet* 114:823–839
- 6 Supena EDJ, Winarto B, Riksen T, Dubas E. van Lammeren A, Offringa R, Boutilier K, Custers
7 J (2008) Regeneration of zygotic-like microspore-derived embryos suggests an important role for
8 the suspensor in early embryo patterning. *J Exp Bot* 59: 803-814
- 9 Szücs P, Blake VC, Bhat PR, Chao S, Close TJ, Cuesta-Marcos A, Muehlbauer GJ, Ramsay L,
10 Waugh R, Hayes PM (2009) An integrated resource for barley linkage map and malting quality
11 QTL alignment. *The Plant Genome* 2:1-7
- 12 Thomas WTB, Forster BP, Gertsson B (2003) Doubled haploids in breeding. In: Maluszynski M,
13 Kasha KJ, Forster BP, Szarejko I (eds) *Doubled haploid production in crop plants: a manual*.
14 Kluwer Acad Publ, Dordrecht, pp 337–350
- 15 Törjék O, Witucka-Wall H, Meyer RC, von Korff M, Kusterer B, Rautengarten C, Altmann T
16 (2006) Segregation distortion in *Arabidopsis* C24/Col-0 and Col-0/C24 recombinant inbred line
17 populations is due to reduced fertility caused by epistatic interaction of two loci. *Theor Appl*
18 *Genet* 113:1551-1561
- 19 Torp AM, Andersen (2009) Albinism in Microspore Culture. A. Touraev et al. (eds) *Advances in*
20 *haploid production in higher Plants*. Chapter 12 pp.155-160. Springer Science.
- 21 Vales MI, Schón CC, Capettini F, Chen XM, Corey AE, Mather DE, Mundt CC, Richardson KL,
22 Sandoval-Islas JS, Utz HF, Hayes PM (2005) Effect of population size on the estimation of QTL:
23 a test using resistance to barley stripe rust. *Theor Appl Genet* 111:1260-1270
- 24 Van Ooijen JW (2006) JoinMap^R 4.0, Software for the calculation of genetic linkage maps in
25 experimental populations. Kyazma V.V., Wageningen, Netherlands.

- 1 Varshney RK, Marcel TC, Ramsay L, Russell J, Röder MS, Stein N, Waugh R, Langridge P,
2 Niks RE, Graner A (2007) A high density barley microsatellite consensus map with 775 SSR
3 loci. *Theor Appl Genet* 114:1091-1103
- 4 Wang S, Basten CJ, Zeng Z-B (2001–2003) Windows QTL cartographer 2.5. Department of
5 Statistics, North Carolina State University, Raleigh, NC, USA
- 6 Wenzl P, Li H, Carling J, Zhou M, Raman H, Paul E, Hearnden P, Maier C, Xia L, Caig V,
7 Ovesná J, Cakir M, Poulsen D, Wang J, Raman R, Smith KP, Muehlbauer GJ, Chalmers KJ,
8 Kleinhofs A, Huttner E, Kilian A (2006) A high-density consensus map of barley linking DArT
9 markers to SSR, RFLP and STS loci and agricultural traits. *BMC Genomics* 7:206
- 10 Wolfe RI (1972) A multiple stock in Brandon, Canada. *Barley Genet Newsl* 2:170
- 11 Wolfe RI, Franckowiak JD (1991) Multiple dominant and recessive genetic marker stocks in
12 spring barley. *Barley Genet Newsl* 20:117-121
- 13 Xian-Liang S, Xue-Zhen S, Tian-Zhen Z (2006) Segregation distortion and its effect on genetic
14 mapping in plants. *Chin J Agric Biotech* 3:163-169
- 15 Xu Y, Zhu L, Xiao J, Huang N, McCouch SR (1997) Chromosomal regions associated with
16 segregation distortion of molecular markers in F₂, backcross, doubled haploid, and recombinant
17 inbred populations in rice (*Oryza sativa* L.). *Mol Gen Genet* 253:535–545
- 18 Zeng ZB (1994) Precision mapping of quantitative trait loci. *Genetics* 136:1457-1468
- 19 Zhu H, Gilchrist L, Hayes P, Kleinhofs A, Kudrna D, Liu Z, Prom L, Steffenson B, Toojinda T,
20 Vivar H. (1999) Does function follow form? Principal QTLs for Fusarium head blight (FHB)
21 resistance are coincident with QTLs for inflorescence traits and plant height in a doubled-haploid
22 population of barley. *Theor Appl Genet* 99:1221-1232

Table 1. Number of SNP loci used for mapping each of the seven chromosomes of barley in *Hordeum bulbosum*-derived (H.b.) and anther culture-derived (A.C.) doubled haploid (DH) populations of barley, as well as in the combined (H.b.+A.C.) population and the total length of each linkage group, and all linkage groups, in Haldane cM.

Chrom.	Number of SNP	H.b. 82 DH	A.C. 93 DH	H.b.+A.C. 175 DH
1H	148	157.3	186.2	171.2
2H	199	188.8	193.8	190.4
3H	211	205.8	208.7	205.4
4H	187	127.4	156.9	141.9
5H	236	240.2	200.4	216.7
6H	182	159.1	146.3	151.0
7H	173	206.6	225.8	215.6
Total	1,328	1,285.2	1,318.1	1,292.2

Table 2. Summary of reproductive fitness trait QTL detected in the Oregon Wolfe Barley mapping population (175 DH lines).

Trait	QTL No.	Chrom.	Peak Position (2-LOD conf. Interval)	Closest Marker	LOD	R ²	Additive effect	LOD Threshold	MIM R ²
Spike Length	1	1H	170.7 (166.1-171.3)	2_0840	8.2	0.02	-0.48	3.0	0.90
	2	2H	161.8 (160.6-163.3)	3_0396	93.4	0.83	-3.26		
	3	6H	92.2 (89.9-93.3)	3_0573	3.9	0.01	0.32		
	4	7H	124.6 (108.1-135.1)	2_1201	3.4	0.01	-0.32		
Grain Number	1	1H	170.7 (103.4-106.3)	2_0840	4.0	0.03	-3.70	2.9	0.69
	2	2H	104.8 (103.4-106.3)	1_1100	37.8	0.52	-14.61		
	3	2H	161.8 (154.0-168.8)	3_0396	5.6	0.05	-4.83		
	4	6H	82.8 (80.5-87.5)	2_0468	4.1	0.03	3.74		
Floret Number	1	1H	169.0 (163.2-170.6)	1_0041	5.8	0.02	-3.24	3.0	0.89
	2	2H	105.8 (104.1-106.8)	3_0897	87.3	0.81	-21.81		
	3	3H	62.4 (57.7-67.1)	3_0721	4.0	0.01	-2.64		
Hundred Grain Weight	1	2H	104.8 (102.9-106.9)	1_1100	38.0	0.54	0.61	3.0	0.67
	2	7H	117.6 (113.3-127.3)	2_0685	10.5	0.09	0.25		
Plant Height	1	1H	79.3 (78.7-88.9)	2_0696	5.4	0.03	-4.25	3.1	0.75
	2	2H	160.8 (159.8-162.8)	3_0396	53.4	0.64	-20.84		
	3	3H	57.7 (48.6-68.1)	2_1189	5.3	0.03	4.23		
	4	4H	138.1 (134.8-140.1)	3_1422	5.6	0.04	4.50		
	5	6H	86.9 (81.7-91)	2_0673	5.7	0.03	4.39		
Spike Number	1	2H	105.8 (99.0-109.6)	3_0897	12.9	0.24	3.03	3.0	0.29
Heading Date	1	1H	170.7 (163.1-171.3)	2_0840	4.9	0.08	-2.41	2.9	0.30
	2	4H	134.1 (129.4-140.1)	2_0272	5.4	0.10	2.64		
	3	6H	88.5 (82.8-91)	2_0577	4.6	0.08	2.41		

Figure 1. Number of apparent crossovers for each of the seven chromosomes of barley in *Hordeum bulbosum*-derived (*H.b.*) and anther culture-derived (A.C.) doubled haploid populations of the Oregon Wolfe Barley. Numbers on the X axis below chromosome numbers represent *p*-values for the statistical contrasts between doubled haploid production methods, using square root-transformed data.






Fig. 2. Chromosome 2H linkage maps from two subpopulations of doubled haploid lines (H.b. = *Hordeum bulbosum*-derived and A.C. = anther culture-derived) and the combined population (H.b. + A.C.) All maps were constructed using the same 199 SNPs. The map positions of three loci determining key morphological traits (*VRS1*, *ZEO1*, and *WST1*) are shown in large font. The single locus p -values of the χ^2 test for segregation distortion are denoted by * $p < 0.05$; ** $p < 0.01$; and *** $p < 0.001$

H.b.



A.C.



H.b.+A.C.




Fig. 3. Allelic frequencies for the OWB-Dominant parental alleles across 1328 loci sorted by map position. Genome-wide segregation distortion thresholds are calculated according to a modified Bonferroni correction, considering 50 effective independent tests. Panel A shows results based on 82 DH lines derived by the *Hordeum bulbosum* (H.b.) technique. Panel B shows results based on 93 DH lines derived by anther culture (A.C.). Panel C shows results based on 175 DH lines (82 H.b.-derived lines and 93 A.C.-derived lines)




Fig. 4. LOD plots for reproductive fitness trait QTL on chromosome 2H mapped in the doubled haploid (DH) Oregon Wolfe Barley population. Panel A shows results based on 82 DH lines derived by the *Hordeum bulbosum* (H.b.) technique. Panel B shows results based on 93 DH lines derived by anther culture (A.C.). Panel C shows results based on 175 DH lines (82 H.b.-derived lines and 93 A.C.-derived lines). Positions of two morphological trait loci – *VRS1* and *ZEO1* are shown.

