

Aquatic Invasion Records

First record of *Percnon gibbesi* (H. Milne Edwards, 1853) (Crustacea: Decapoda: Percnidae) from Egyptian waters

Ernesto Azzurro^{1,2*}, Marco Milazzo³, Francesc Maynou¹, Pere Abelló¹ and Tarek Temraz⁴

¹Institut de Ciències del Mar (CSIC) Passeig Marítim de la Barceloneta, 37-49, E-08003 Barcelona, Spain

²High Institute for Environmental Protection and Research (ISPRA), Laboratory of Milazzo, Via dei Mille 44, 98057 Milazzo, Messina, Italy

³Dipartimento di Ecologia, Università di Palermo, via Archirafi 28, I-90123, Palermo, Italy

⁴Marine Science Department, Suez Canal University, Egypt

E-mail: eazzurr@tin.it (EA), marmilazzo@iol.it (MM), maynouf@icm.csic.es (FM), pabello@icm.csic.es (PA), ttemraz@yahoo.com (TT)

*Corresponding author

Received: 12 November 2010 / Accepted: 30 November 2010 / Published online: 22 December 2010

Abstract

On July 2010, the invasive crab *Percnon gibbesi* was photographed and captured along the coast of Alexandria (Egypt, Eastern Mediterranean Sea). This represents the first observation of this species in Egyptian waters and the easternmost record for the southern rim of the Mediterranean.

Key words: *Percnon gibbesi*, Percnidae, Egypt, Mediterranean, invasive species

Introduction

The invasive crab *Percnon gibbesi* (H. Milne Edwards, 1853) is one the most recent and successful invaders in the Mediterranean Sea. This species was formerly included within the family Plagusiidae Dana, 1851 but a recent study based on molecular and morphological analyses (Schubart and Cuesta 2010) assigned this genus to the family Percnidae Števčić, 2005. It inhabits the infralittoral rocky shore and its native range extends over a wide latitudinal and temperature range along the eastern Pacific, from California to Chile, western Atlantic, from Florida to Brazil, and eastern Atlantic, from the Gulf of Guinea to Portugal, including Madeira, Azores, and Cape Verde Islands (Nobre 1931; Manning and Holthuis 1981; Udekom d'Acoz 1999).

In the Mediterranean Sea, the species distribution has been recently updated by Katsanevakis et al. (2010) and by Galil et al. (2009). It was firstly recorded in 1999 at Linosa, Pelagie Islands (Relini et al. 2000) where an established population was sampled soon afterwards (Puccio et al. 2006) and at the Balearic Islands (Garcia and Reviriego 2000;

Mueller 2001). Subsequent records were from Sicilian shores (Pipitone et al. 2001; Mori and Vacchi 2002), Malta (Borg and Attard-Montalto 2002), Tyrrhenian and Ionian shores of Italy (Russo and Villani 2005; Faccia and Bianchi 2007), Catalan Sea (Abelló et al. 2003), Greece (Cannicci et al. 2006; Thessalou-Legaki et al. 2006; Katsanevakis and Tsiamis 2009; Katsanevakis et al. 2010) and Tunisia (MedMPA 2004). Recently the species was observed along the southern Turkish coast (Yokes and Galil 2006) and in Libya (Elkrwe et al. 2008).

Results and discussion

On July 2nd 2010, several specimens of *Percnon gibbesi* were observed at Miami beach (Alexandria, Egypt: 31°16'20.81"N, 29°59'27.43"E) (Figure 1) while snorkeling. They occurred on artificial rocky boulders and natural rocky bottoms in the narrow subtidal zone between 0.5 and 2.5 m depth. The crabs were hiding under the blocks and in narrow crevices as observed in other Mediterranean regions (Pipitone et al. 2001; Russo and Villani 2005; Deudero et al. 2005).

Figure 1. Record location of *Percnon gibbesi* at Miami beach, Alexandria, Egypt. Crab drawing by Lucio Valdez.




Figure 2. *Percnon gibbesi* photographed underwater at Miami beach, Alexandria, Egypt, on July 2th 2010. Photograph by E. Azzurro.



A small male (17.5 mm carapace length and 17.0 mm carapace width) was photographed (Figure 2), captured, preserved in ethanol and deposited in the Biological Reference Collections of the Institut de Ciències del Mar - CSIC of Barcelona (accession number ICMD_201011-01).

This record establishes the presence of *Percnon gibbesi* in Egypt and represents the easternmost record along the southern rim of the Mediterranean Sea, almost 800 km east of previous record (Haniyah, Libya - 32°50.205'N, 21°30.803'E) (Elkrwe et al. 2008). This would confirm the rapid expansion of the crab in the Mediterranean basin and its recent spread

eastwards. It is possible that this species has been present along the Egyptian coasts for some time, as the population observed seemed well established. Keeping track of alien species is often a difficult task and regular faunistic monitoring programmes are essential for information on these biodiversity changes.

Acknowledgements

We acknowledge the Euromediterranean Center for Climatic Changes and the Italian Ministry for the Environment and the Territory (Project: The impacts of biological invasions and climate change on the biodiversity of the Mediterranean Sea). The suggestions of two anonymous referees improved the quality of this paper.

References

- Abelló P, Visauta E, Buci A, Demestre M (2003) Noves dades sobre l'expansió del cranc *Percnon gibbesi* (Brachyura: Grapsidae: Plagusiinae) a la Mediterrània occidental. *Bulleti de la Societat d'Historia Natural de les Balears* 46: 73–77
- Borg JJ, Attard-Montalto J (2002) The grapsid crab *Percnon gibbesi* (Milne Edwards, 1853) (Crustacea, Decapoda, Brachyura), a new addition to the marine fauna of Malta. *Central Mediterranean Naturalist* 3: 159–160
- Cannicci S, Garcia L, Galil BS (2006) Racing across the Mediterranean – first record of *Percnon gibbesi* (Crustacea: Decapoda: Grapsidae) in Greece. *JMBA2 - Biodiversity Records*, doi: 10.1017/S1755267206003009
- Deudero S, Frau A, Cerdá M, Hampel H (2005) Distribution and densities of the decapod crab *Percnon gibbesi*, an invasive Grapsidae, in western Mediterranean waters. *Marine Ecology Progress Series* 285: 151–158, doi:10.3354/meps285151
- Elkrwe HM, Elhawaj HM, Galil BS, Ben Abdallah A (2008) The first record of *Percnon gibbesi* (H. Milne Edwards, 1853) (Crustacea: Decapoda: Plagusiidae) from the southern rim of the Mediterranean. *Aquatic Invasions* 3: 243–245, doi:10.3391/ai.2008.3.2.16
- Faccia I, Bianchi CN (2007) Prima segnalazione del granchio *Percnon gibbesi* (H. Milne-Edwards, 1853) nella Calabria ionica. *Thalassia Salentina* 30: 39–43
- Galil B, Froglio C, Noël O (2009) CIESM Atlas of exotic species in the Mediterranean Sea: Online publication, CIESM Monaco. <http://www.ciesm.org/atlas> (Accessed 11 November 2010)
- Garcia L, Reviriego B (2000) Presència del cranc subtropical *Percnon gibbesi* (H. Milne Edwards, 1853) (Crustacea, Brachyura, Grapsidae) a les Illes Balears. Primera cita a la Mediterrània occidental. *Bulleti de la Societat d'Història Natural de les Balears* 43: 81–89
- Katsanevakis S, Poursanidis D, Issaris Y, Tsiamis K, Salomidi M, Maroulakis M, Kyritsou E, Thessalou-Legaki M, Zenetos A (2010) The invasive crab *Percnon gibbesi* is spreading in the Aegean and Ionian Seas. *Marine Biodiversity Records* 3: 1–5, doi:10.1017/S1755267210000163
- Katsanevakis S, Tsiamis K (2009) Records of alien marine species in the shallow coastal waters of Chios Island. *Mediterranean Marine Science* 10(2): 99–107
- Manning R, Holthuis BL (1981) West African Brachyuran crabs (Crustacea: Decapoda). *Smithsonian Contributions to Zoology* 306: 1–379
- MedMPA (2004) Elaboration du plan de gestion de la partie marine du parc national de Zembra et Zembretta. Projet Ré'gional pour le dé'veloppe'ment d'aires proté'gées marines et co^tie'res dans la ré'gion Mé'diterrané'enne (Projet MedMPA), Rapport global des travaux de prospection marine, 35 pp
- Mori M, Vacchi M (2002) On a new occurrence of the alien flat crab, *Percnon gibbesi* (H. Milne Edwards), in the southern Sicily (Central Mediterranean Sea) (Crustacea, Brachyura, Grapsidae). *Annali del Museo Civico di Storia Naturale "Giacomo Doria"* 114: 295–302
- Mueller C (2001) First record of *Percnon gibbesi* (Crustacea: Brachyura: Grapsidae) for the Balearic Islands. *Senckenbergiana Maritima* 31: 83–89
- Nobre A (1931) Crustáceos Decápodes e Stomatópodes marinhos de Portugal. Imprensa Portuguesa, Porto, 307 pp
- Pipitone C, Badalamenti F, Sparrow A (2001) Contribution to the knowledge of *Percnon gibbesi* (Decapoda, Grapsidae), an exotic species spreading rapidly in Sicilian waters. *Crustaceana* 74(10): 1009–1017, doi:10.1163/15685400152691061
- Puccio V, Relini M, Azzurro E, Orsi Relini L (2006) Feeding habits of *Percnon gibbesi* (H. Milne Edwards, 1853) in the Sicily Strait. *Hydrobiologia* 557(1): 79–84, doi:10.1007/s10750-005-1310-2
- Relini M, Orsi L, Puccio V, Azzurro E (2000) The exotic crab *Percnon gibbesi* (H. Milne Edwards, 1853) (Decapoda, Grapsidae) in the central Mediterranean. *Scientia Marina* 64(3): 337–340
- Russo GF, Villani G (2005) Diffusione nel Tirreno centrale della specie alloctona *Percnon gibbesi* (Decapoda, Grapsidae). *Biologia Marina Mediterranea* 12: 329–330
- Schubart CD, Cuesta JA (2010) Phylogenetic relationships of the Plagusiidae Dana, 1851 (Brachyura), with description of a new genus and recognition of Percnidae Števčić, 2005 as an independent family. *Studies on Brachyura*: 279–299
- Thessalou-Legaki M, Zenetos A, Kambouroglou V, Corsini-Foka M, Kouraklis P, Dounas C, Nicolaïdou A (2006) The establishment of the invasive crab *Percnon gibbesi* (H. Milne Edwards, 1853) (Crustacea: Decapoda: Grapsidae) in Greek waters. *Aquatic Invasions* 1: 133–136, doi:10.3391/ai.2006.1.3.6
- Udekem d'Acoz CD (1999) Inventaire et distribution des crustacés décapodes de l'Atlantique nord-oriental, de la Méditerranée et des eaux continentales adjacentes au nord de 25°N. *Collection Patrimoines Naturels* 40: 383 pp
- Yokes B, Galil BS (2006) Touchdown – first record of *Percnon gibbesi* (H. Milne Edwards, 1853) (Crustacea: Decapoda: Grapsidae) along the Levant coast. *Aquatic Invasions* 1: 130–132, doi:10.3391/ai.2006.1.3.5