

DCPT/08/172
IPPP/08/86
BONN-TH-2008-17

HiggsBounds: Confronting Arbitrary Higgs Sectors with Exclusion Bounds from LEP and the Tevatron

P. Bechtle^a, O. Brein^b, S. Heinemeyer^c, G. Weiglein^b
and K.E. Williams^d.

^a*DESY, Notkestrasse 85, 22607 Hamburg, Germany*

^b*Institute for Particle Physics Phenomenology, Durham University,
Durham, DH1 3LE, United Kingdom*

^c*Instituto de Fisica de Cantabria (CSIC-UC), Santander, Spain*

^d*Bethe Center for Theoretical Physics, Physikalisches Institut der Universität
Bonn, Nussallee 12, 53115 Bonn, Germany*

Abstract

HiggsBounds is a computer code that tests theoretical predictions of models with arbitrary Higgs sectors against the exclusion bounds obtained from the Higgs searches at LEP and the Tevatron. The included experimental information comprises exclusion bounds at 95% C.L. on topological cross sections. In order to determine which search topology has the highest exclusion power, the program also includes, for each topology, information from the experiments on the expected exclusion bound, which would have been observed in case of a pure background distribution. Using the predictions of the desired model provided by the user as input, **HiggsBounds** determines the most sensitive channel and tests whether the considered parameter point is excluded at the 95% C.L. **HiggsBounds** is available as a Fortran 77 and Fortran 90 code. The code can be invoked as a command line version, a subroutine version and an online version. Examples of exclusion bounds obtained with **HiggsBounds** are discussed for the Standard Model, for a model with a fourth generation of quarks and leptons and for the Minimal Supersymmetric Standard Model with and without \mathcal{CP} -violation. The experimental information on the exclusion bounds currently implemented in **HiggsBounds** will be updated as new results from the Higgs searches become available.

Key words: Higgs bosons, Higgs search, LEP, Tevatron, Beyond the Standard Model

PROGRAM SUMMARY

Manuscript Title: HiggsBounds: Confronting Arbitrary Higgs Sectors with Exclusion Bounds from LEP and the Tevatron

Authors: P. Bechtle, O. Brein, S. Heinemeyer, G. Weiglein, K.E. Williams

Program Title: HiggsBounds

Journal Reference:

Catalogue identifier:

Licensing provisions:

Program requirements: HiggsBounds can be built with any compatible Fortran 77 or Fortran 90 compiler. The program has been tested on x86 CPUs running under Linux (Ubuntu 8.04) and with the following compilers: The Portland Group Inc. Fortran compilers (pgf77, pgf90), the GNU project Fortran compilers (g77, gfortran).

Programming language: Fortran 77, Fortran 90 (two code versions are offered)

RAM: minimum of about 6000 kbytes (dependent on the code version)

Keywords: Elementary Particle Physics; General, High Energy Physics and Computing; Higgs bosons; Higgs search; LEP; Tevatron; Beyond the Standard Model.

PACS: 14.80.Bn; 14.80.Cp; 12.60.Fr.

Classification:

External routines/libraries: HiggsBounds requires no external routines/libraries. Some sample programs in the distribution require the programs FeynHiggs 2.6.x or CPsuperH2 to be installed.

No. of lines in distributed program, including test data, etc.: 61645

No. of bytes in distributed program, including test data, etc.: 13721912

Distribution format: tar.gz

Nature of problem: Determine whether a parameter point of a given model is excluded or allowed by LEP and Tevatron Higgs-boson search results.

Solution method: The most sensitive channel from LEP and Tevatron searches is determined and subsequently applied to test this parameter point. The test requires as input model predictions for the Higgs-boson masses, branching ratios and ratios of production cross sections with respect to reference values.

Restrictions: In the current version, results from decay-mode independent Higgs searches and results of searches for charged Higgs bosons are not taken into account.

Running time: about 0.01 seconds (or less) for one parameter point using one processor of an Intel Core 2 Quad Q6600 CPU at 2.40GHz for sample model scenarios with three Higgs bosons. It depends on the complexity of the Higgs sector (e.g. the number of Higgs bosons and the number of open decay channels) and on the code version.

LONG WRITE-UP

1 Introduction

A major goal of the particle physics programme at the high energy frontier, currently being pursued at the Fermilab Tevatron collider and soon to be taken up by the CERN Large Hadron Collider (LHC), is to unravel the nature of electroweak symmetry breaking. While the existence of the massive electroweak gauge bosons (W^\pm, Z), together with the successful description of their behaviour by non-abelian gauge theory, requires some form of electroweak symmetry breaking to be present in nature, the underlying dynamics is not known yet. An appealing theoretical suggestion for such dynamics is the Higgs mechanism [1], which implies the existence of one or more Higgs bosons (depending on the specific model considered). Therefore, the search for Higgs bosons is a major cornerstone in the physics programmes of past, present and future high energy colliders.

Many theoretical models employing the Higgs mechanism in order to account for electroweak symmetry breaking have been studied in the literature, of which the most popular ones are the Standard Model (SM) [2] and the Minimal Supersymmetric Standard Model (MSSM) [3]. Within the SM, the Higgs boson is the last undiscovered particle, whereas the MSSM has a richer Higgs sector, containing three neutral and two charged Higgs bosons. Among alternative theoretical models beyond the SM which lead to the prediction of Higgs bosons, the most prominent are the Two Higgs Doublet Model (THDM) [4], non-minimal supersymmetric extensions of the SM (e.g. extensions of the MSSM by an extra singlet superfield [5]), little Higgs models [6] and models with more than three spatial dimensions [7]. Furthermore, several models have been proposed which mainly extend the Higgs sector of the SM, e.g. extensions of the SM by electroweak singlet scalar fields [8], or the Private Higgs Model [9] which introduces one Higgs doublet for each fermion generation, or models with Higgs-dependent Yukawa couplings [10].

LEP has searched for the SM Higgs boson [11], the MSSM Higgs bosons [12], and other, more “exotic”, manifestations of the Higgs mechanism [13]. After the termination of LEP, the search has been taken up again by the Tevatron. So far, no signals of Higgs bosons have been found, and LEP and Tevatron turned the non-observation of Higgs signals into cross section constraints. The constraints are provided by experiments in the form of limits on cross sections of individual signal topologies (such as $e^+e^- \rightarrow h_i Z \rightarrow b\bar{b}Z$ or $p\bar{p} \rightarrow h_i Z \rightarrow b\bar{b}l^+l^-$) or in the form of combined limits for a specific model, such as the SM. In the latter case, the individual topological cross sections have been combined using the proportions of the individual contributions as predicted

by the model.

An important part of the phenomenological exploration of physics models beyond the SM is the confrontation of such models with existing experimental constraints. In particular, checking the predictions of the model's Higgs sector with bounds from experimental searches for a Higgs boson is vital. In order to test whether the parameter space of a certain model is compatible with the limits from the Higgs searches at LEP and the Tevatron, it is, in general, not possible to employ the experimental results that have been obtained for the Higgs search in the SM (or any other specific model). This is because, in a general model of new physics, individual Higgs signal topologies will contribute in different proportions than, for instance, in the SM. Thus, for models for which no explicit experimental analysis has been carried out, or, like in the case of the MSSM, for model parameters that differ from the benchmark values chosen in the experimental analyses, one needs to resort to the limits on the individual topological cross sections provided by the experimental collaborations. These limits exist for various search channels from LEP and the Tevatron (the latter ones are frequently updated as new data becomes available). Comparing the predictions of a particular model with the existing experimental bounds on the various search topologies can be quite a tedious task as it involves the implementation of experimental results that are distributed over many different publications and combining these results requires a procedure to ensure the correct statistical interpretation of the exclusion bounds obtained on the parameter space of the model.

We present here the program `HiggsBounds`, which is a tool designed to facilitate the above task so that wide classes of models can easily be checked against the state-of-the-art results from Higgs searches. This should be useful for applications in Higgs phenomenology and model building. `HiggsBounds` takes theoretical Higgs sector predictions, e.g. for a particular parameter scenario of a model beyond the SM, as input and determines which Higgs search analysis has the highest exclusion power according to a list of expected exclusion limits from LEP and the Tevatron (an expected exclusion limit corresponds to the bound that one would obtain in the hypothetical case of an observed distribution that agrees precisely with the background expectation). In order to ensure the correct statistical interpretation of the obtained exclusion bound as a 95% C.L., the comparison of the model with the experimental limits has to be restricted to the single channel that possesses the highest statistical sensitivity for setting an exclusion limit. For this channel, the program then compares the theoretical prediction for the Higgs production cross section times decay branching ratio with the actual experimental limit and determines whether or not the considered parameter point of the model is excluded at 95% C.L. So far, no bounds on charged Higgs bosons are taken into account. This will be included in a later version. `HiggsBounds` is model-independent, in that it can be used for models with arbitrary Higgs sectors. The input needed is the

number of neutral Higgs bosons in the model and masses, decay branching ratios and ratios of production cross sections versus corresponding reference (usually SM) values for all neutral Higgs bosons in the model. The code has both a Fortran 77 and Fortran 90 version. It can be operated in a command line mode that can process input files in a variety of formats, as a subroutine suitable for inclusion in user applications, and as an online version, available at the URL: <http://www.ippp.dur.ac.uk/HiggsBounds>. `HiggsBounds` includes sample programs which demonstrate its usage. Sample programs which demonstrate how `HiggsBounds` can be used in conjunction with the widely used programs `FeynHiggs` [14,15,16,17] and `CPsuperH` [18] for Higgs-sector predictions in the MSSM are provided by default.

The rest of the paper is organised as follows. In Section 2 we describe our general approach for the implementation of experimental Higgs constraints followed by specific details on the implementation of LEP and Tevatron results. In Section 3, the general operating instructions for `HiggsBounds` are specified and accompanied by small sample programs, while in Section 4 some examples of use for `HiggsBounds` are described. Section 5 contains a summary and outlook and is followed by an appendix which contains details on the definition and statistical interpretation of the confidence limits.

2 Implementation of experimental Higgs constraints

2.1 General approach

Tevatron and LEP turn(ed) the non-observation of Higgs signals into 95% C.L. upper limits on cross sections for individual signal topologies. These limits are given as functions of the anticipated mass of the Higgs boson. By a certain Higgs signal topology X , we mean a specific combination of a single (double) Higgs boson production process, $P(h)$ ($P(h_1, h_2)$), and decay final state(s) F of the Higgs boson(s) h (h_1, h_2). In the limit of a narrow-width Higgs boson the cross section $\sigma(X)$ of the signal topology X factorises into the on-shell Higgs production cross sections $\sigma(P)$ times the appropriate Higgs decay branching ratio(s) $\text{BR}(h \rightarrow F)$ ($\text{BR}(h_1, h_2 \rightarrow F)$).

The Tevatron and LEP cross section limits are understood to be applicable to models which do not change the signature of the background processes considerably¹ which do not significantly change the kinematical distributions of

¹ This is not such a strong extra restriction on new physics models as it may seem, because viable models are required to agree with present experimental results which are well described by the SM, such as the LEP and SLD EW precision data [19] and

the signal topology X (e.g. η , p_T distributions of the final state particles) from what has been assumed in the corresponding analysis. This is an unavoidable limitation of the standard approach adopted in the literature of expressing search results in terms of limits on total cross sections times branching ratios for certain signal topologies. Detection efficiencies, for instance the b-jet tagging efficiency, usually depend on the kinematical configuration of the signal event (particularly the η and p_T of the final state particles). Therefore, a dedicated experimental analysis for a model giving rise to kinematical distributions that strongly differ from the ones assumed in the analysis on which the published cross section limit is based would lead to a somewhat modified cross section limit. For instance, applying the cross section limit of a Tevatron analysis which relies on b-jet tagging to a model where the distribution of the signal b-jets differs strongly from the one assumed in the Tevatron analysis would lead to a slightly too weak/strong limit if a higher/lower fraction of the signal b-jets than assumed in the analysis is radiated into the central part of the detector where the b-jet tagging efficiency is higher than in regions closer in angle to the beam axis. `HiggsBounds` provides three options for passing model predictions to the program: (`hadr`, `part` and `effC`). The extent to which the kinematical distributions may vary from those assumed in the analyses depends on which of these input options is chosen. We will discuss this issue in further detail in the sections describing the input options.

The Tevatron and LEP cross section limits are usually given for a narrow-width Higgs boson. From the narrow-width limits, some experimental analyses extract limits for Higgs bosons with a substantial decay width by representing the signal of a “large-width” Higgs boson as a weighted sum of mock narrow-width Higgs signals. Examples of this procedure can be found in [21,22,23,24]. In particular, [24] shows that the effect of a non-vanishing width on the extracted limit can be parametrised by a function of the Higgs mass m_h and the ratio $\Gamma_{\text{tot}}(h)/m_h$. If model-independent information of this kind also became available (for a fine grid of points) for other analyses, this would greatly facilitate the task to implement width-dependent Higgs exclusion limits.

At present, we constrain the implementation of `HiggsBounds` to the case of narrow-width Higgs boson exclusion limits. Neglecting the width of a Higgs boson usually leads to tighter bounds on the topological cross sections than those obtained through taking the width into account. The user should be aware of this fact when considering exclusion bounds on parameter regions that give rise to very large Higgs-boson widths. In a future version of `HiggsBounds` we

the precision measurements of gauge boson production at LEP and the Tevatron [20]. These analyses are much more sensitive to changes in the background to Higgs searches than the Higgs searches themselves. Therefore, the models with new physics in the Higgs sector which have been considered in the literature usually do not show strong deviations from the SM in the background processes.

plan to incorporate a proper width treatment for cases where width-dependent results are provided by the experimental collaborations. As mentioned above, **HiggsBounds** is not meant to be applied to models for which the background rates to the considered signal processes differ drastically to the SM case.

For some specific models (the SM and some benchmark scenarios of the MSSM) the results of several Higgs signal topologies have been combined by the experimental collaborations according to the relative contributions to the total cross section. In the combination, a detailed knowledge of the overlap between the individual experimental searches is used, which is not available using the limits on topological cross sections only. Hence, a combined limit, for instance for the SM, is more sensitive than a combination of individual topological limits. The use of detailed SM dependent information during the calculation of the combined limit prohibits its application to those Higgs bosons of new physics models which are not predicted to have all relevant couplings proportional to the SM values. However, such models can still contain one (or some) Higgs boson(s) which are SM-like. For those Higgs bosons, the SM combined limit can be applied.

The basic flow of **HiggsBounds** is as follows.

- (1) The user provides (in a form to be specified below) for all neutral Higgs bosons h_i ($i = 1, \dots, n_{\text{Higgs}}$) of the considered model the predictions for: Higgs masses, total decay widths, the relevant branching ratios for the decay of Higgs bosons h_i into known ordinary particles (OP, i.e. SM fermions and gauge bosons) and into pairs of other Higgs bosons h_j , and the relevant cross sections for Higgs production processes P normalised to reference (usually SM) values:

$$m_{h_i}, \Gamma_{\text{tot}}(h_i), \text{BR}(h_i \rightarrow \text{OP}), \text{BR}(h_i \rightarrow h_j h_j), \quad (1)$$

$$\frac{\sigma_{\text{model}}(P(h_i))}{\sigma_{\text{ref}}(P(H), m_H)} \Big|_{m_H=m_{h_i}}, \frac{\sigma_{\text{model}}(P(h_i, h_j))}{\sigma_{\text{ref}}(P(H, H'), m_H, m_{H'})} \Big|_{m_H, H'=m_{h_i, j}}.$$

The precise definitions of the reference cross sections are given in the LEP- and Tevatron-specific subsections below. The user can choose to specify all the above quantities directly, or, in the simplest form, one just needs to provide $m_{h_i}, \Gamma_{\text{tot}}(h_i), \text{BR}(h_i \rightarrow h_j h_j)$ and effective couplings of the Higgs bosons to ordinary particles. The remaining branching ratios and cross section ratios listed above can then be calculated from this input in the effective coupling approximation. For the preparation of normalised input, the user may apply the SM predictions for Higgs boson production cross sections and decay branching ratios which **HiggsBounds** provides as external routines (see Table 12 in Section 3.3).

- (2) With the above model input, the program calculates the model predictions for the cross section (normalised to a reference value in some cases)

for each Higgs signal topology $Q_{\text{model}}(X)$. Depending on whether the exclusion result for a particular search topology has been given by the experimental collaborations as a relative or absolute limit, the program evaluates

$$Q_{\text{model}} = \frac{[\sigma \times \text{BR}]_{\text{model}}}{[\sigma \times \text{BR}]_{\text{ref}}} \text{ or } [\sigma \times \text{BR}]_{\text{model}},$$

where σ and BR denote the production process and decay branching ratio of the Higgs boson, respectively.

- (3) From the experimental results, we read off the values $Q_{\text{obs}}(X)$ and $Q_{\text{expec}}(X)$ corresponding to the observed and expected 95% C.L. exclusion bound on the quantity $Q(X)$, respectively. To that end, we have implemented many of the available observed and expected exclusion limits (preliminary and final results) from LEP and the Tevatron as data tables which are read in by the program during start-up. In order to provide values for $Q_{\text{obs}}(X)$ and $Q_{\text{expec}}(X)$ for continuous Higgs mass values, the program interpolates Q -values linearly between neighbouring Higgs mass points given by the experimental collaborations. Details on the definition of 95% C.L. limits can be found in Appendix A.
- (4) The program determines the search topology X_0 with the highest ratio of Q_{model} to expected Q_{expec} , i.e. out of all topologies X , it finds the channel X_0 where

$$\frac{Q_{\text{model}}(X)}{Q_{\text{expec}}(X)}$$

is maximal. This topology has the highest statistical sensitivity for excluding the model prediction.

- (5) We then look at the ratio of Q_{model} to the observed Q_{obs} value for this channel X_0 and, if

$$\frac{Q_{\text{model}}(X_0)}{Q_{\text{obs}}(X_0)} > 1, \tag{2}$$

this particular parameter point is excluded by the corresponding experimental analysis for the search channel X_0 at 95 % C.L. In order to ensure the correct statistical interpretation of the obtained exclusion bound as a 95% C.L., it is crucial to perform the test of Eq. (2) only for exactly one search channel (namely the one with the highest statistical sensitivity), and not for all available search channels. In the latter case the derived constraint would in general not correspond to a constraint at 95 % C.L., due to the maximally 5 % probability of each individual comparison of Q_{model} and Q_{obs} to yield a false exclusion.

As additional information, the web version of the program **HiggsBounds** also informs the user which topologies had the second and third highest statistical sensitivity and whether or not the ratio of Q_{model} to observed 95% C.L. limit

Q_{obs} was greater than one for those topologies. However, this information does not contribute to the decision about whether or not this parameter point is excluded.

There are a few more general implementation details which should be mentioned here.

Using SM combined limits, SM-likeness

In order to use the best available exclusion limit, it makes sense to use the results of the SM combined analyses for any Higgs boson predicted to behave in a similar way to the SM Higgs boson (in terms of its production and decay). A SM analysis, combining several signal topologies assumes that the relative contribution to the event rate by each included search topology is according to the proportions in the SM. For a given Higgs mass m_H , the only free parameter left in such analyses is one overall scale factor, which multiplies all Higgs signal cross sections and which is varied in order to determine the value for which 95% C.L. exclusion occurs.

Therefore, the results of a SM analysis, combining M signal topologies, are exactly applicable to a Higgs boson h of an alternative model if for all of the M search topologies X_n the ratio of topological cross sections (model versus SM) is independent of X_n , i.e.

$$\frac{\sigma_{\text{model}}(X_n)}{\sigma_{\text{SM}}(X_n)} = \text{const.} =: \alpha, \quad n = 1, \dots, M, \quad (3)$$

with a proportionality constant α , or equivalently

$$\frac{\sigma_{\text{model}}(X_n)}{\sigma_{\text{model}}(X_k)} = \frac{\sigma_{\text{SM}}(X_n)}{\sigma_{\text{SM}}(X_k)} \quad \text{for all } n, k \in \{1, \dots, M\}. \quad (4)$$

For a model which passes the conditions (3) or (4) to sufficient accuracy for a set of M Higgs search topologies, the cross section prediction for a combination of those topologies, normalised to the SM reference value, fulfils

$$Q_{\text{model}}(\{X_1, \dots, X_M\}) := \frac{\sum_{n=1}^M \sigma_{\text{model}}(X_n)}{\sum_{n=1}^M \sigma_{\text{SM}}(X_n)} = \alpha. \quad (5)$$

Here and in the following, it is always understood that the mass of the SM Higgs boson m_H is set to m_h . In the narrow-width approximation, which we assume to hold throughout (see the discussion above), each cross section $\sigma(X_n)$ of the Higgs signal topologies X_n factorises into a production cross section $\sigma(P_i(h))$ and decay branching ratio $\text{BR}(h \rightarrow F_k)$, where i and k are functions of n , and $n = 1, \dots, M$. Note that the index n labels distinct signal

topologies, i.e. specific combinations of one Higgs production process with one decay final state. SM-likeness is fulfilled if

$$\frac{\sigma(P_i(h))_{\text{model}} \text{BR}_{\text{model}}(h \rightarrow F_k)}{\sigma(P_i(H))_{\text{SM}} \text{BR}_{\text{SM}}(H \rightarrow F_k)} = \text{const.} \quad (6)$$

holds to sufficient accuracy for all M topologies. If this is the case for a particular Higgs boson of a certain model, a SM analysis combining precisely those M channels also applies to that Higgs boson.

In order to decide whether or not a Higgs boson of a certain model is sufficiently SM-like such that the corresponding SM combined analysis can be applied, we use in `HiggsBounds` a pragmatic definition of SM-likeness. For each of the N_{CS} distinct production cross sections $\sigma_{\text{model}}(P_i(h))$ and the N_{BR} distinct decay branching ratios $\text{BR}(h \rightarrow F_k)$ which appear in the list of M signal topologies, we determine the normalised mean value and the deviation from the mean:

$$\bar{s} = \frac{1}{N_{\text{CS}}} \sum_{i=1}^{N_{\text{CS}}} s_i, \quad \bar{b} = \frac{1}{N_{\text{BR}}} \sum_{k=1}^{N_{\text{BR}}} b_k, \quad (7)$$

$$\delta s_i = s_i - \bar{s}, \quad \delta b_k = b_k - \bar{b}, \quad (8)$$

$$\text{with } s_i = \frac{\sigma_{\text{model}}(P_i(h))}{\sigma_{\text{SM}}(P_i(H))}, \quad b_k = \frac{\text{BR}_{\text{model}}(h \rightarrow F_k)}{\text{BR}_{\text{SM}}(H \rightarrow F_k)}. \quad (9)$$

The approximate SM-likeness of the given model is then ensured if the deviations from a constant expression according to Eq. (6) are small. For a given combination (i, k) of a production and decay process, the left hand side of Eq. (6) can be expressed as:

$$\frac{\sigma_{\text{model}}(P_i(h)) \text{BR}_{\text{model}}(h \rightarrow F_k)}{\sigma_{\text{SM}}(P_i(H)) \text{BR}_{\text{SM}}(H \rightarrow F_k)} = s_i b_k = \bar{s} \bar{b} \left(1 + \frac{\delta s_i}{\bar{s}} + \frac{\delta b_k}{\bar{b}} + \frac{\delta s_i \delta b_k}{\bar{s} \bar{b}} \right). \quad (10)$$

In the program, we consider a corresponding SM analysis applicable if the maximum relative deviation from the mean value $\bar{s} \bar{b}$ satisfies

$$\max_{i,k} \left(\frac{\delta s_i}{\bar{s}} + \frac{\delta b_k}{\bar{b}} + \frac{\delta s_i \delta b_k}{\bar{s} \bar{b}} \right) < \epsilon, \quad (11)$$

where we have chosen $\epsilon = 2\%$, i.e. the predictions for the different topological cross sections, normalised to the SM values, are required to be equal to each other with at least 2% accuracy. This restriction is quite stringent and may cause a SM combined analysis to be not applied to some models where it may actually be justifiable. The user can remedy this situation by changing the setting of the corresponding variable `eps` in the internal subroutine `check_SM_likeness` (Fortran 77) or `SMlikeness` (Fortran 90) in the code.

So far, **HiggsBounds** uses analyses with SM combinations of topologies only from the Tevatron experiments. This is the most important case, as CDF and DØ have not done combinations of their results for individual search topologies so far, while the LEP results we are using represent the combination of results from all four LEP experiments for each individual channel.

A more elaborate criterion for determining whether SM combined analyses are applicable to Higgs bosons of models beyond the SM can be sketched as follows. The variation of the ratios of signal topology cross sections (model versus SM) due to non-exact SM-likeness of the model prediction should stay within a fraction of the 1σ band of the sensitivity of the observed limit of the analysis with respect to signal-like fluctuations. The fixed maximum deviation of 2%, described above, lies well within the typical expected statistical fluctuation of the limits in the SM combined analyses. We plan to implement such a more elaborate criterion in future updates of the program.

Adding of signal channels

In models with more than one Higgs boson it can happen that some of the Higgs bosons, h_i , have masses which are quite close to each other, say

$$|m_{h_i} - m_{h_j}| \leq \Delta m_h \quad (12)$$

for certain i and j . This is the case for instance in the parameter space of the (\mathcal{CP} -conserving) MSSM for the heavy Higgs bosons H^0 and A^0 for $M_A \gg M_Z$. If the masses are so close to each other that their signals appear within a certain mass window, it may happen that the two signals are not experimentally distinguishable anymore, or, at least, that it makes sense to combine their signal contribution into one signal to be compared with the data.

In the narrow-width approximation (which is used in **HiggsBounds**, see the discussion above) it is assumed that the total widths of the Higgs bosons $\Gamma_{\text{tot}}(h_i)$ are all (much) smaller than their mutual mass differences,

$$\Gamma_{\text{tot}}(h_i) \ll |m_{h_i} - m_{h_k}| \quad \text{for all } i \text{ and } k \neq i . \quad (13)$$

In this case the signal channels factorise into production cross section and decay branching ratio, so that the addition of different signal cross sections may formally always be carried out where it is necessary. In parameter regions where the narrow-width approximation does not hold, however, one may worry about the possible effects of interference terms arising from squaring the amplitude of the full process involving both production and decay. It should be noted that even in cases where the total widths of two Higgs bosons are significantly larger than their mass difference it can happen that interference terms are small. An example for such a situation is the above-mentioned case of the MSSM with real soft-breaking parameters for $M_A \gg M_Z$. While for large $\tan \beta$ the widths of the two Higgs bosons can be very large, the different

quantum numbers of the \mathcal{CP} -even and the \mathcal{CP} -odd state nevertheless forbid an interference contribution.

In determining the cross sections $\sigma(X)$ for a given signal topology X , **HiggsBounds** does the following. Suppose there are n_{Higgs} neutral Higgs bosons h_i in a given model and let the numbering be according to a mass order, i.e.

$$m_{h_1} \leq m_{h_2} \leq \dots \leq m_{h_{n_{\text{Higgs}}}} . \quad (14)$$

Then, **HiggsBounds** works through a loop from $i = 1$ to n_{Higgs} and adds up all² production cross sections of Higgs bosons h_j ($j \geq i$) which satisfy

$$m_{h_j} - m_{h_i} \leq \Delta m_h(X) . \quad (15)$$

In the implementation, the X -dependence of Δm_h is limited to two different values: one for LEP and one for Tevatron search topologies, represented by the variables `delta_Mh_LEP` and `delta_Mh_TEV`, respectively. Adding of Higgs signal cross sections is mainly relevant for Tevatron Higgs search analyses, where the invariant mass distributions of the potential Higgs boson decay products usually use bin sizes of the order of 10 GeV. For the LEP analyses, the Higgs mass resolution is typically around 2–3 GeV [11]. For the LEP results the addition of Higgs signal cross sections is implemented in **HiggsBounds** only for the Higgsstrahlung process where the Higgs boson decays directly into SM particles. As default values for `delta_Mh_LEP` and `delta_Mh_TEV` we have chosen (in units of GeV):

$$\begin{aligned} \text{delta_Mh_TEV} &= 10 , \\ \text{delta_Mh_LEP} &= 2 . \end{aligned}$$

The default setting of `delta_Mh_TEV` for the Tevatron search channels has also been tested against known MSSM Higgs search results [23,24,25] which we could reproduce successfully with **HiggsBounds** within the limitations of the narrow-width approach.

In the calculation of normalised quantities, the reference cross section and branching ratio are evaluated at the average mass of the Higgs bosons contributing to a signal topology.

Provided SM normalisation

In the evaluation of the model predictions $Q_{\text{model}}(X)$ and $Q_{\text{model}}(\{X_1, \dots, X_M\})$ for single and multiple search topologies, respectively, to be compared with the expected and observed limits Q_{obs} and Q_{exp} , **HiggsBounds** uses internally SM predictions for Higgs production cross sections and decay branching

² If the cross section for a SM combination of signal topologies is calculated, production cross sections of those Higgs bosons are only included in the sum if they also fulfil the SM-likeness criterion.

ratios. Out of the set of implemented Higgs search results, the need for using such predictions internally arises only for Tevatron analyses. The provided SM predictions for Higgs boson production cross sections and decay branching ratios use results of HDECAY 3.303 [26] and the TEV4LHC Higgs Working Group [27] (see Table 12 below for references to the original publications). In addition, for the bH production cross section we need several predictions with different kinematic cuts on the b jet which are not available from [27]. We used HJET 1.1 [28] to calculate those, but checked that the uncut cross section closely resembles the result reported in [27] (see Section 3.3 for details). If the model input is given in terms of ratios of partonic cross sections, the calculation of hadronic Higgs production cross sections is facilitated by using internally SM cross section ratios which we have calculated (see Section 2.3.2 for details). If the model input is given in terms of effective couplings, we make use of results from VBFNLO [29] in order to facilitate the calculation of the Higgs production cross section via vector boson fusion (see Section 2.3.3 for details).

The rationale behind the choice of SM normalisation is that virtually all Tevatron analyses implemented in this program use these predictions when normalising their cross section limits to SM quantities. Thus, describing deviations from the SM of a new model by using the SM normalisation the experimental analyses have chosen allows for the most accurate interpretation of the limits. By using SM predictions for Higgs production cross sections which deviate from the internally used ones by a certain percentage (be it because of a different loop order, different numerical values of input parameters, renormalisation scheme or choice of parton distribution functions), a deviation of the same relative size will be caused in the quantities Q_{model} . The user should bear this in mind when interpreting the output of `HiggsBounds`.

2.2 LEP limits

The main Higgs boson production processes at LEP have been Higgsstrahlung and double Higgs production:

$$e^+e^- \rightarrow h_k Z, \quad e^+e^- \rightarrow h_k h_i. \quad (16)$$

Both processes are assumed to be mediated via s -channel Z boson exchange in leading order of perturbation theory.

Currently, `HiggsBounds` uses the exclusion limits on topological cross sections obtained from the combination of all LEP2 and some LEP1 Higgs boson search results from Ref. [12] which considers the Higgs decay final states $b\bar{b}$ and $\tau^+\tau^-$. Apart from SM search topologies, this analysis also considers double Higgs production topologies, and almost all possibilities for the decay of an

search topology X	reference
$e^+e^- \rightarrow (h_k)Z \rightarrow (b\bar{b})Z$	[12,30] table 14b
$e^+e^- \rightarrow (h_k)Z \rightarrow (\tau^+\tau^-)Z$	[12,30] table 14c
$e^+e^- \rightarrow (h_k \rightarrow h_i h_i)Z \rightarrow (b\bar{b}b\bar{b})Z$	[12,31] table 15
$e^+e^- \rightarrow (h_k \rightarrow h_i h_i)Z \rightarrow (\tau^+\tau^-\tau^+\tau^-)Z$	[12,31] table 16
$e^+e^- \rightarrow (h_k h_i) \rightarrow (b\bar{b}b\bar{b})$	[12,31] table 18
$e^+e^- \rightarrow (h_k h_i) \rightarrow (\tau^+\tau^-\tau^+\tau^-)$	[12,31] table 19
$e^+e^- \rightarrow (h_k \rightarrow h_i h_i)h_i \rightarrow (b\bar{b}b\bar{b})b\bar{b}$	[12,31] table 20
$e^+e^- \rightarrow (h_k \rightarrow h_i h_i)h_i \rightarrow (\tau^+\tau^-\tau^+\tau^-)\tau^+\tau^-$	[12,31] table 21
$e^+e^- \rightarrow (h_k \rightarrow h_i h_i)Z \rightarrow (b\bar{b})(\tau^+\tau^-)Z$	[31]
$e^+e^- \rightarrow (h_k \rightarrow b\bar{b})(h_i \rightarrow \tau^+\tau^-)$	[31]
$e^+e^- \rightarrow (h_k \rightarrow \tau^+\tau^-)(h_i \rightarrow b\bar{b})$	[31]

Table 1

LEP search topologies used by HiggsBounds. For each Higgs search topology which has been studied in [12], we indicate the corresponding table number. It is assumed in all tables that $m_{h_k} > m_{h_i}$. In the program, we use more fine-grained results [30,31] from the analysis [12] than have been displayed as tables in [12]. We also use equally fine-grained results for the expected limits [30,31], which have only been displayed in graphs in [12], and for the observed and expected limits of a few channels, which were not published [31].

on-shell Higgs boson into two lighter ones³ and their subsequent decay into matter particles. Because of the limitation in [12] to $b\bar{b}$ and $\tau^+\tau^-$ final states, Higgs bosons which decay predominantly invisibly cannot be well constrained by HiggsBounds at present. We will include the appropriate LEP results in future updates of the code. The search topologies included in HiggsBounds up to now are shown in Table 1.

The limits on topological cross sections in [12] have been given in the normalisation where all Higgs decay branching ratios are equal to 1. For given Higgs mass(es), limits on the production cross sections of the individual search topologies X have been derived for this normalisation in the form of scaling

³ The only possibilities not analysed in [12] are the ones where more than two distinct Higgs bosons appear. For example, this occurs if a primarily produced Higgs boson h_k decays into two different Higgs bosons h_j and h_i , or if two Higgs bosons $h_k h_j$ are primarily produced and one of these, say h_k , decays into two Higgs bosons h_i (i, j, k all different). We are not aware of any studies by the LEP experiments of those more general topologies.

factors $S_{95}(X)$, defined as

$$S_{95}(X) = \sigma_{\max}(X)/\sigma_{\text{ref}}(X), \quad (17)$$

where σ_{\max} is the largest cross section compatible with the data at 95 % C.L. and σ_{ref} is a reference cross section for the Higgs production process. The quantities $S_{95}(X)$ of [12] correspond to the quantities $Q(X)$ mentioned in the description of the program flow in Section 2.1.

For Higgsstrahlung processes, $e^+e^- \rightarrow h_k Z \rightarrow (\text{final state})Z$, the reference cross section σ_{ref} is the SM cross section

$$\sigma_{\text{ref}}(\text{Higgsstrahlung}) = \sigma(e^+e^- \rightarrow HZ)_{\text{SM}} \quad (18)$$

with the SM Higgs mass m_H chosen as m_{h_k} .

For double Higgs production processes, $e^+e^- \rightarrow H'H \rightarrow (\text{final state})$, there is no direct SM reference process. However, as in [12], we choose a reference cross section for a fictitious production process of two scalar particles (H' , H) with masses $m_{H'} = m_{h_k}$ and $m_H = m_{h_i}$ via a virtual Z exchange with a standardised coupling constant

$$g_{H'HZ}^{\text{ref}} = \frac{e}{2 s_w c_w}, \quad (19)$$

where e denotes the electromagnetic coupling constant, and s_w and c_w the sine and cosine of the electroweak mixing angle, respectively. The cross section of this process in leading order is then completely determined by the Higgs masses and SM input and related to the SM Higgsstrahlung cross section via a simple phase space factor:

$$\begin{aligned} \sigma_{\text{ref}}(H' H \text{ production}) &= \bar{\lambda}(m_{H'}, m_H, s) \sigma_{HZ}^{\text{SM}}(m_H), \quad (20) \\ \bar{\lambda}(m_{H'}, m_H, s) &= \frac{\lambda_{H'H}^{3/2}(s)}{\lambda_{HZ}^{1/2}(s) \left(\lambda_{HZ}(s) + 12 \frac{m_Z^2}{s} \right)}, \\ \lambda_{ab}(s) &= \left[1 - \frac{(m_a + m_b)^2}{s} \right] \left[1 - \frac{(m_a - m_b)^2}{s} \right]. \end{aligned}$$

This reference cross section coincides with the MSSM or THDM tree-level cross section for the process $e^+e^- \rightarrow h^0 A^0$ if the Higgs mixing-angle dependent factor $\cos(\beta - \alpha)$ is divided out and m_{A^0} and m_{h^0} are chosen as $m_{H'}$ and m_H .

In order to apply the LEP Higgs constraints implemented in `HiggsBounds` to a model, the user has to provide model information which allows model predictions to be calculated for all Higgs search topologies X listed in Table

1,

$$Q_{\text{model}}(X) = \frac{\sigma_{\text{model}}(P(X))\text{BR}_{\text{model}}(F(X))}{\sigma_{\text{ref}}(P(X))}, \quad (21)$$

where the symbol $P(X)$ denotes the corresponding Higgs production process and $F(X)$ the corresponding Higgs decay chain, e.g. $h_k \rightarrow b\bar{b}$ or $h_k \rightarrow h_i h_i \rightarrow b\bar{b}\tau^+\tau^-$. The quantity $Q_{\text{model}}(X)$ is compared with the experimental limits $Q_{\text{exp}}(X)$ and $Q_{\text{obs}}(X)$ according to the description in Section 2.1. Currently, **HiggsBounds** allows model information to be provided in three ways.

2.2.1 LEP input (*HiggsBounds Input Option part and hadr*): cross section and branching ratios

In order to be able to apply the implemented LEP limits to the model under study, the user is asked to provide model predictions for the Higgs boson masses m_{h_k} (in units of GeV), for the branching ratios

$$\text{BR}_{\text{model}}(h_i \rightarrow b\bar{b}), \quad \text{BR}_{\text{model}}(h_i \rightarrow \tau^+\tau^-), \quad \text{BR}_{\text{model}}(h_k \rightarrow h_i h_i), \quad (22)$$

and for the ratios of cross sections

$$\frac{\sigma_{\text{model}}(e^+e^- \rightarrow h_k Z)}{\sigma_{\text{ref}}(e^+e^- \rightarrow HZ)}, \quad \frac{\sigma_{\text{model}}(e^+e^- \rightarrow h_k h_i)}{\sigma_{\text{ref}}(e^+e^- \rightarrow H'H)}, \quad (23)$$

for $k, i \in \{1, \dots, n_{\text{Higgs}}\}$.

With this information, **HiggsBounds** obtains model predictions $Q_{\text{model}}(X)$ for all topologies X specified in Table 1. For example, the Q_{model} value for the channel $e^+e^- \rightarrow h_1 Z \rightarrow (b\bar{b})Z$ is given by (suppressing the initial state in the notation):

$$Q_{\text{model}}\left((h_1)Z \rightarrow (b\bar{b})Z\right) = \frac{\sigma_{\text{model}}(h_1 Z)}{\sigma_{\text{ref}}(HZ)} \text{BR}_{\text{model}}(h_1 \rightarrow b\bar{b}),$$

with $m_H = m_{h_1}$, and similarly for the channel $e^+e^- \rightarrow (h_2 \rightarrow h_1 h_1)Z \rightarrow (b\bar{b}b\bar{b})Z$ it is:

$$Q_{\text{model}}\left((h_2 \rightarrow h_1 h_1)Z \rightarrow (b\bar{b}b\bar{b})Z\right) = \frac{\sigma_{\text{model}}(h_2 Z)}{\sigma_{\text{ref}}(HZ)} \text{BR}_{\text{model}}(h_2 \rightarrow h_1 h_1) \left(\text{BR}_{\text{model}}(h_1 \rightarrow b\bar{b})\right)^2,$$

with $m_H = m_{h_2}$.

The **HiggsBounds Input Option hadr** amounts to exactly the same LEP related input as described above. It differs for the Tevatron related input (see Section 2.3).

If the model under study predicts kinematical distributions which differ significantly from the distributions of the reference processes (18) or (20), the exclusion result, returned by `HiggsBounds`, can only be considered an estimate.⁴

2.2.2 LEP input (`HiggsBounds Input Option effC`): effective couplings

The user is asked to provide model predictions for:

$$m_{h_k}, \Gamma_{\text{tot}}(h_k), \left(\frac{g_{h_k ZZ}^{\text{model}}}{g_{HZZ}^{\text{SM}}}\right)^2, \left(\frac{g_{h_k h_i Z}^{\text{model}}}{g_{H'HZ}^{\text{ref}}}\right)^2, \left(\frac{g_{h_k f \bar{f}, \text{eff}}^{\text{model}}}{g_{Hf\bar{f}}^{\text{SM}}}\right)^2, \\ \text{BR}_{\text{model}}(h_k \rightarrow h_i h_i), \quad (24)$$

for $k, i \in \{1, \dots, n_{\text{Higgs}}\}$ and $f \in \{b, \tau\}$. The normalised effective squared coupling of the Higgs boson fermion interaction is defined as an approximation for the ratio of partial decay widths,

$$\left(\frac{g_{h_k f \bar{f}, \text{eff}}^{\text{model}}}{g_{Hf\bar{f}}^{\text{SM}}}\right)^2 := \frac{\Gamma_{h_k \rightarrow f \bar{f}}^{\text{model}}(m_{h_k})}{\Gamma_{H \rightarrow f \bar{f}}^{\text{SM}}(m_H)} \Big|_{m_H = m_{h_k}}, \quad (25)$$

and the reference coupling constants are defined by Eq. (19) and by

$$(g_{HZZ}^{\text{SM}})^2 = \left(\frac{e m_W}{s_w c_w^2}\right)^2, \quad (26)$$

$$(g_{Hf\bar{f}}^{\text{SM}})^2 = \left(\frac{1}{2} \frac{e m_f}{s_w m_W}\right)^2, \quad (27)$$

where m_W and m_f denote the masses of the W boson and fermion f , respectively.

For the purpose of running `HiggsBounds`, only the ratio of partial widths corresponding to the right-hand-side of Eq. (25) is needed. In order to completely specify effective couplings for neutral Higgs boson interactions with fermions, in general a scalar and a pseudoscalar part of the coupling is needed. The Feynman rule for the coupling of a generic neutral Higgs boson h to fermions can be written as:

$$G(h f \bar{f}) = i(g_s \mathbf{1} + i g_p \gamma_5), \quad (28)$$

⁴ In such a case, the general cross section limits given by the experimental collaborations would have to be replaced by a dedicated experimental analysis of a particular model giving rise to these kinematical distributions, taking into account detailed information about the detector response to the signal events.

where g_s and g_p are real-valued coupling constants, and $\mathbf{1}$ and γ_5 are the usual matrices in Dirac space. A scalar particle, like the SM Higgs boson, has $g_p = 0$ and a pseudoscalar particle has $g_s = 0$. If the user has the couplings g_s and g_p available, the user may calculate the normalised effective squared couplings, specified in Eq. (25), by using the relation:

$$\left(\frac{g_{h_k f \bar{f}, \text{eff}}^{\text{model}}}{g_{H f \bar{f}}^{\text{SM}}}\right)^2 = \left(\frac{g_{s, h_k f \bar{f}}^{\text{model}}}{g_{H f \bar{f}}^{\text{SM}}}\right)^2 + \left(\frac{g_{p, h_k f \bar{f}}^{\text{model}}}{g_{H f \bar{f}}^{\text{SM}}}\right)^2 \frac{1}{\beta_f^2(m_{h_k})}, \quad (29)$$

with

$$\beta_f(m_{h_k}) = \sqrt{1 - \frac{4m_f^2}{m_{h_k}^2}}.$$

The extra factor of β_f^{-2} in Eq. (29) appears, because the partial width for a scalar particle decaying into $f\bar{f}$ is proportional to β_f^3 , while for a pseudoscalar particle it is proportional to β_f .

The user may also avoid the explicit use of the reference coupling, defined in Eq. (26), by using the relation

$$\left(\frac{g_{h_k Z Z}^{\text{model}}}{g_{H Z Z}^{\text{SM}}}\right)^2 = \frac{\Gamma_{h_k \rightarrow Z^{(*)} Z^{(*)}}^{\text{model}}(m_{h_k})}{\Gamma_{H \rightarrow Z^{(*)} Z^{(*)}}^{\text{SM}}(m_H)} \Bigg|_{m_H = m_{h_k}}, \quad (30)$$

where the asterisk in brackets indicates the possible off-shellness of the decay products. Knowledge of the off-shell decay matrix elements allows to use relation (30) for masses m_{h_k} smaller than $2M_Z$.

From the information specified in Eq. (24), **HiggsBounds** calculates the LEP part of the cross section and branching ratios required in Input Option **part** in the effective coupling approximation:

$$\frac{\sigma_{\text{model}}(e^+e^- \rightarrow h_k Z)}{\sigma_{\text{ref}}(e^+e^- \rightarrow H Z)} = \left(\frac{g_{h_k Z Z}^{\text{model}}}{g_{H Z Z}^{\text{SM}}}\right)^2, \quad (31)$$

$$\frac{\sigma_{\text{model}}(e^+e^- \rightarrow h_k h_i)}{\sigma_{\text{ref}}(e^+e^- \rightarrow h_k h_i)} = \left(\frac{g_{H' H Z}^{\text{model}}}{g_{H' H Z}^{\text{ref}}}\right)^2, \quad (32)$$

$$\begin{aligned} \text{BR}_{\text{model}}(h_k \rightarrow f \bar{f}) &= \text{BR}_{\text{SM}}(H \rightarrow f \bar{f})(m_H) \frac{\Gamma_{\text{tot}}^{\text{SM}}(m_H)}{\Gamma_{\text{tot}}(h_k)} \Bigg|_{m_H = m_{h_k}} \times \\ &\times \left(\frac{g_{h_k f \bar{f}, \text{eff}}^{\text{model}}}{g_{H f \bar{f}}^{\text{SM}}}\right)^2, \end{aligned} \quad (33)$$

for $f \in \{b, \tau\}$.

From this point, the evaluation of the quantities $Q_{\text{model}}(X)$ can proceed exactly as exemplified in Section 2.2.1.

As far as the implemented LEP constraints are concerned, effective couplings just rescale all kinematical distributions of the reference processes (18) or (20) by common factors. Thus, for the class of models where the Higgs sector can be faithfully parametrised in this way, the kinematical distributions correspond to those of the reference processes. Consequently, limits on topological cross sections can exactly be applied.

2.3 Tevatron limits

The main SM Higgs boson production processes at the proton–anti-proton collider Tevatron at Fermilab are:

- Gluon fusion: $p\bar{p} \rightarrow H + \dots$ via $gg \rightarrow H$,
- Higgsstrahlung: $p\bar{p} \rightarrow HV + \dots$ via $q\bar{q}' \rightarrow VH$ ($V = Z, W^\pm$),
- Vector boson fusion (VBF): $p\bar{p} \rightarrow Hjj + \dots$ via $qq' \rightarrow V^*V'^*q''q''' \rightarrow Hq''q'''$,
- Higgs production associated with heavy quarks: $p\bar{p} \rightarrow Ht\bar{t}$ ($Hb\bar{b}$) + \dots ,

where the ellipses indicate the hadronic remainder of the reaction which we will suppress in our notation below.

Currently, the Tevatron experiments CDF and DØ provide separate limits on cross sections of individual Higgs search topologies X and limits on combinations of search topologies according to SM proportions. In particular, combinations of Higgs search results from *both* experiments are, so far, only available for a SM-like Higgs boson. The results implemented in `HiggsBounds` correspond to the Higgs search topologies that have been analysed by CDF and DØ, using Tevatron Run II data, listed in Tables 2 and 3. New results provided by CDF and DØ will be implemented once they appear, thus keeping `HiggsBounds` up-to-date.

In the absence of a Higgs signal, the Tevatron experiments obtain cross section limits from their measurements: $\sigma_{\text{max}}(X)$ is the highest cross section for a given assumed Higgs mass m_H which is compatible with the only-background hypothesis at 95% C.L. Therefore, if a particular model of the Higgs sector predicts a cross section $\sigma_{\text{model}}(X)$ for a Higgs boson with mass m_H , which is higher than the limit, i.e.

$$\sigma_{\text{model}}(X) > \sigma_{\text{max}}(X), \quad (34)$$

then the probability that such a prediction is consistent with observation is less than 5% and the model parameter point will be called excluded with (at

search topology X (analysis)	reference
$p\bar{p} \rightarrow ZH \rightarrow l^+l^-b\bar{b}$ (CDF with 1.0 fb^{-1})	[32]*
$p\bar{p} \rightarrow ZH \rightarrow l^+l^-b\bar{b}$ (CDF with 2.4 fb^{-1})	[33]
$p\bar{p} \rightarrow ZH \rightarrow l^+l^-b\bar{b}$ (DØ with 2.3 fb^{-1})	[34]
$p\bar{p} \rightarrow WH \rightarrow l\nu b\bar{b}$ (DØ with 1.7 fb^{-1})	[35]
$p\bar{p} \rightarrow WH \rightarrow l\nu b\bar{b}$ (CDF with 2.7 fb^{-1})	[36]
$p\bar{p} \rightarrow WH \rightarrow W^+W^-W^\pm$ (DØ with 1.0 fb^{-1})	[37]
$p\bar{p} \rightarrow WH \rightarrow W^+W^-W^\pm$ (CDF with 1.9 fb^{-1})	[38]
$p\bar{p} \rightarrow H \rightarrow W^+W^- \rightarrow l^+l'^-$ (DØ with 3.0 fb^{-1})	[39]
$p\bar{p} \rightarrow H \rightarrow W^+W^- \rightarrow l^+l'^-$ (CDF with 3.0 fb^{-1})	[40]*
$p\bar{p} \rightarrow H \rightarrow \gamma\gamma$ (DØ with 1.1 fb^{-1})	[41]*
$p\bar{p} \rightarrow H \rightarrow \gamma\gamma$ (DØ with 2.68 fb^{-1})	[42]
$p\bar{p} \rightarrow H \rightarrow \tau^+\tau^-$ (DØ with 1.0 fb^{-1})	[24]*
$p\bar{p} \rightarrow H \rightarrow \tau^+\tau^-$ (CDF with 1.8 fb^{-1})	[25]
$p\bar{p} \rightarrow bH, H \rightarrow b\bar{b}$ (CDF with 1.9 fb^{-1})	[22]
$p\bar{p} \rightarrow bH, H \rightarrow b\bar{b}$ (DØ with 1.0 fb^{-1})	[23]*
$p\bar{p} \rightarrow bH, H \rightarrow b\bar{b}$ (DØ with 2.6 fb^{-1})	[43]

Table 2

Tevatron analyses for single search topologies, the results of which are used by HiggsBounds. The leptons l and l' can be an electron or a muon. References marked by an asterisk (\star) refer to analyses which have been published or submitted for publication.

least) 95% C.L. While some Tevatron Higgs search results are presented in the form of absolute cross section limits, $\sigma_{\max}(X)$, most of the results are presented as limits on signal cross sections normalised to the corresponding SM cross section, $\sigma_{\text{SM}}(X)$.⁵ In the latter case, the exclusion condition (34) becomes:

$$\frac{\sigma_{\text{model}}(X)}{\sigma_{\text{SM}}(X)} > \frac{\sigma_{\max}(X)}{\sigma_{\text{SM}}(X)}. \quad (35)$$

In order to test the exclusion of a given model by CDF or DØ results according to the procedure outlined in Section 2.1, the user needs to provide model

⁵ So far, no Tevatron Higgs search results have appeared which would require to use a normalisation to non-SM cross sections.

search topologies $\{X\}$ (analysis)	reference
$p\bar{p} \rightarrow WH/ZH \rightarrow b\bar{b} + E_T^{\text{miss.}}$ (CDF with 2.3 fb^{-1})	[44]
$p\bar{p} \rightarrow WH/ZH \rightarrow b\bar{b} + E_T^{\text{miss.}}$ (DØ with 2.1 fb^{-1})	[45]
$p\bar{p} \rightarrow H/HW/HZ/H$ via VBF, $H \rightarrow \tau^+\tau^-$ (CDF with 2.0 fb^{-1})	[46]
Combined SM analysis (CDF & DØ with $0.9 - 1.9 \text{ fb}^{-1}$)	[47]
Combined SM analysis (CDF & DØ with $1.0 - 2.4 \text{ fb}^{-1}$)	[48]
Combined SM analysis (CDF & DØ with 3.0 fb^{-1})	[49]

Table 3

CDF and DØ analyses which are contained in `HiggsBounds` and which combine search topologies according to SM proportions.

predictions which allow, for each relevant Higgs boson search topology X , either absolute cross sections,

$$Q_{\text{model}}(X) = \sigma_{\text{model}}(P(X))\text{BR}_{\text{model}}(F(X)), \quad (36)$$

or cross section ratios,

$$Q_{\text{model}}(X) = \frac{\sigma_{\text{model}}(P(X))\text{BR}_{\text{model}}(F(X))}{\sigma_{\text{SM}}(P(X))\text{BR}_{\text{SM}}(F(X))}, \quad (37)$$

of hadronic processes to be calculated, where the symbol $P(X)$ denotes the corresponding Higgs production process and $F(X)$ the corresponding Higgs decay chain. The program allows both variants to be calculated from the same input, as the SM predictions for Higgs production processes and decay branching ratios are internally available. In Eqs. (36) and (37), we have already used the implicit assumption that Higgs bosons have narrow width, for which the CDF and DØ limits we employ are valid. We should also stress again that the limits are based on analyses which assume that the backgrounds to the signal process under study are similar to the backgrounds predicted by the SM. For that reason, the branching ratios for the decays of the W^\pm or the Z for a particular individual Higgs search topology X are assumed to cancel in the ratio (37).

For the calculation of signal cross section predictions where several Higgs search topologies $\{X_1, \dots, X_M\}$ are combined (all cases listed in Table 3), the definition in Eq. (5) applies. In the narrow-width approximation, this definition reads:

$$Q_{\text{model}}(\{X_1, \dots, X_M\}) = \frac{\sum_{n=1}^M \sigma_{\text{model}}(P(X_n))\text{BR}_{\text{model}}(F(X_n))}{\sum_{n=1}^M \sigma_{\text{SM}}(P(X_n))\text{BR}_{\text{SM}}(F(X_n))}. \quad (38)$$

The quantities Q_{model} are to be compared with the experimental limits Q_{expec} and Q_{obs} according to the description in Section 2.1. There are three options, which allow the user to provide **HiggsBounds** with the necessary model information in order to calculate the quantities Q_{model} . These are described in the following.

2.3.1 Tevatron input (*HiggsBounds Input Option `hadr`*): hadronic cross section and branching ratios

In order to make use of all the implemented Tevatron limits, the user is asked to provide model predictions for the Higgs boson masses m_{h_k} (in units of GeV), for the branching ratios

$$\begin{aligned} \text{BR}_{\text{model}}(h_k \rightarrow b\bar{b}), & & \text{BR}_{\text{model}}(h_k \rightarrow \tau^+\tau^-), \\ \text{BR}_{\text{model}}(h_k \rightarrow W^+W^-), & & \text{BR}_{\text{model}}(h_k \rightarrow \gamma\gamma), \end{aligned} \quad (39)$$

and for the ratios of the hadronic cross sections

$$\begin{aligned} \frac{\sigma_{\text{model}}(p\bar{p} \rightarrow h_k Z)}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H Z)}, & \quad \frac{\sigma_{\text{model}}(p\bar{p} \rightarrow h_k W^\pm)}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H W^\pm)}, & \quad \frac{\sigma_{\text{model}}(p\bar{p} \rightarrow h_k \text{ via VBF})}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via VBF})}, \\ \frac{\sigma_{\text{model}}(p\bar{p} \rightarrow h_k)}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H)}, & \quad \frac{\sigma_{\text{model}}(p\bar{p} \rightarrow h_k b)}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H b)}, & \end{aligned} \quad (40)$$

for $k \in \{1, \dots, n_{\text{Higgs}}\}$.

With this information and the internally available SM predictions for Higgs production cross sections and decay branching ratios, **HiggsBounds** calculates all values Q_{model} , according to Eqs. (36), (37) or (38), which are needed in order to test the chosen model against the implemented Tevatron results. Note that the branching ratios $\text{BR}_{\text{model}}(h_k \rightarrow h_i h_i)$ are currently not used in comparisons with Tevatron results, because, so far, no Tevatron analyses have appeared which consider such decays.

While **hadr** is the most widely applicable input option, it can also be the one that is most difficult for the user to provide, as this option requires the user to provide predictions for *hadronic* cross sections, which involve a convolution with parton distribution functions. Therefore, the program offers the Input Option **part**, which requires only ratios of *partonic* cross sections. This should in general be more convenient for the user. However, this input option is applicable to a narrower class of models than Input Option **hadr**.

If the model under study predicts kinematical distributions which differ significantly from the distributions of the reference SM processes, the exclusion result, returned by **HiggsBounds**, can only be considered an estimate (as discussed at the end of Section 2.2.1).

2.3.2 *Tevatron input (HiggsBounds Input Option part): partonic cross section and branching ratios*

The user is asked to provide model predictions for the Higgs boson masses m_{h_k} (in units of GeV), for the branching ratios

$$\begin{aligned} \text{BR}_{\text{model}}(h_k \rightarrow b\bar{b}), & & \text{BR}_{\text{model}}(h_k \rightarrow \tau^+\tau^-), \\ \text{BR}_{\text{model}}(h_k \rightarrow W^+W^-), & & \text{BR}_{\text{model}}(h_k \rightarrow \gamma\gamma), \end{aligned}$$

for the following ratios of partonic cross sections

$$\begin{aligned} \frac{\hat{\sigma}_{\text{model}}(gg \rightarrow h_k)}{\hat{\sigma}_{\text{SM}}(gg \rightarrow H)}, & \quad \frac{\hat{\sigma}_{\text{model}}(b\bar{b} \rightarrow h_k)}{\hat{\sigma}_{\text{SM}}(b\bar{b} \rightarrow H)}, & \quad \frac{\hat{\sigma}_{\text{model}}(bg \rightarrow h_k b)}{\hat{\sigma}_{\text{SM}}(bg \rightarrow Hb)}, \\ \frac{\hat{\sigma}_{\text{model}}(q\bar{q}' \rightarrow h_k W^+)}{\hat{\sigma}_{\text{SM}}(q\bar{q}' \rightarrow HW^+)}, & \quad \frac{\hat{\sigma}_{\text{model}}(q'\bar{q} \rightarrow h_k W^-)}{\hat{\sigma}_{\text{SM}}(q'\bar{q} \rightarrow HW^-)}, & \quad (q, q') \in \{(u, d), (c, s)\}, \\ \frac{\hat{\sigma}_{\text{model}}(q\bar{q} \rightarrow h_k Z)}{\hat{\sigma}_{\text{SM}}(q\bar{q} \rightarrow HZ)}, & \quad q \in \{u, d, c, s, b\}, & \end{aligned}$$

and for the hadronic cross section ratio

$$\frac{\sigma_{\text{model}}(p\bar{p} \rightarrow h_k \text{ via VBF})}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via VBF})},$$

for $k \in \{1, \dots, n_{\text{Higgs}}\}$.

With this information and some internally provided functions (described below) `HiggsBounds` can calculate the same input quantities as required for Input Option `hadr` (subject to some approximations, as described below). It is assumed that the model cross sections for the processes $\bar{b}g \rightarrow h_k \bar{b}$ and $bg \rightarrow h_k b$ are equal to each other.

Hadronic Higgs production cross sections can be written as a sum of cross sections of partonic processes $\hat{\sigma}_{nm \rightarrow H+y}(\hat{s}, m_H)$ convoluted by parton luminosity functions:

$$\begin{aligned} \sigma(p\bar{p} \rightarrow H + y, m_H) &= \int_{\tau_0}^1 d\tau \sum_{\{n,m\}} \frac{d\mathcal{L}_{nm}^{p\bar{p}}}{d\tau} \hat{\sigma}_{nm \rightarrow H+y}(\hat{s} = \tau S, m_H), & (41) \\ &=: \sum_{\{n,m\}} \sigma(p\bar{p} \rightarrow nm \rightarrow H + y, m_H). \end{aligned}$$

Here, the sum includes all distinct parton pairs only once. The centre-of-mass energy of the colliding proton (p) and anti-proton (\bar{p}) is assumed to be $\sqrt{S} = 1.96$ TeV. The symbol y indicates possible further particles in the hard partonic process, like the W^\pm or Z . The parton luminosity functions are

defined as

$$\frac{d\mathcal{L}_{nm}^{AB}}{d\tau} = \int_{\tau}^1 \frac{dx}{x} \frac{1}{1 + \delta_{nm}} \left[f_{n/A}(x, \mu_F) f_{m/B}\left(\frac{\tau}{x}, \mu_F\right) + f_{m/A}(x, \mu_F) f_{n/B}\left(\frac{\tau}{x}, \mu_F\right) \right], \quad (42)$$

where $f_{n/A}(x, \mu_F)$ denotes the density of partons of type n in the hadron A carrying a fraction x of the hadron's momentum at the scale μ_F and $(A, B) = (p, \bar{p})$ for the Tevatron. The lower bound of the τ -integration (τ_0) is determined by the minimal invariant mass of the parton system, $\hat{s}_0 = \tau_0 S$. The ratio $Q_{\text{model}}(X)$ of Eq. (37) factorises into a ratio of branching ratio predictions and a ratio of hadronic production cross section predictions. The calculation of the ratio of branching ratios is straightforward. In the following, we show an approximate relation (that in special cases is exact), which allows to calculate the ratios of hadronic cross sections for the Higgs production process P ,

$$R_{\sigma}(P) := \frac{\sigma_{\text{model}}(P)}{\sigma_{\text{SM}}(P)}, \quad (43)$$

from the knowledge of ratios of partonic cross sections. Under the assumption that the ratio of the partonic cross sections,

$$R_{nm}^{H+y}(\hat{s}, m_H) := \frac{\hat{\sigma}_{nm \rightarrow H+y}^{\text{model}}(\hat{s}, m_H)}{\hat{\sigma}_{nm \rightarrow H+y}^{\text{SM}}(\hat{s}, m_H)}, \quad (44)$$

has, at most, a mild dependence on the parton-system centre-of-mass energy squared, \hat{s} , we can write Eq. (43) as

$$R_{\sigma}(P) \approx \sum_{\{n,m\}} R_{nm}^{H+y}(\hat{s}_0, m_H) \frac{\sigma_{\text{SM}}(p\bar{p} \rightarrow nm \rightarrow H+y, m_H)}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H+y, m_H)}, \quad (45)$$

with \hat{s}_0 denoting the partonic production threshold, $\hat{s}_0 = (m_H + m_y)^2$, with $m_y = 0$ in the case of single Higgs boson production. In this formula, only ratios of SM hadronic cross sections are needed, which we provide in our program.

In order to provide the cross section ratios in Eq. (45) with state-of-the-art theoretical accuracy, we pursue the following strategy. The K-factors (NNLO QCD + NLO EW) for the WH and ZH production cross sections at the Tevatron are, to a very good approximation, independent of the generation of the initial state quarks and thus drop out in the ratios. The only non-trivial case is the single Higgs production which has contributions from gluon fusion and $b\bar{b}$ annihilation. For the calculation of the gluon fusion cross section, we assume a constant NNLO QCD K-factor of 3.6 [51] in the relevant range of $70 \text{ GeV} < m_H < 300 \text{ GeV}$. In this range, our assumption is consistent with the m_H -dependent K-factor reported in [51] within the uncertainty estimate also given in [51]. In order to obtain an approximate NNLO QCD cross section




Fig. 1. Relative contributions (in %) of different SM partonic processes to cross sections of hadronic processes: (a) single Higgs production, (b) HW^\pm production and (c) HZ production.

prediction for the $b\bar{b}$ annihilation process, we used the fact that this cross section is known to be well approximated by the LO prediction in the $\overline{\text{MS}}$ scheme with the factorisation and renormalisation scale choice of $m_H/4$ [52]. The SM cross section ratios from Eq. (45) are then obtained by normalising the cross sections for one parton configuration to the appropriate sum of cross sections. As a cross-check, we confirmed that our results for Higgs production cross sections are in good agreement with the compilation of results by the TEV4LHC working group [27]. Fig. 1 shows our result: the relative contributions from different parton configurations to the total hadronic cross section for single Higgs production, HW^\pm production and HZ production in the range relevant for the implemented Tevatron analyses.

Relation (45) becomes an exact identity for all models which differ from the SM only by having different values for coupling constants, i.e. where the ratios R_{nm}^{H+y} become \hat{s} -independent. Moreover, for models which give \hat{s} -dependent ratios R_{nm}^{H+y} , Eq. (45) can still be a good approximation because the luminosity functions $\frac{d\mathcal{L}_{nm}^{p\bar{p}}}{d\tau}$ are steeply falling functions of τ and therefore strongly favour the threshold regions of the convolution integrals. However, in this case, the user should make sure that Eq. (45) is indeed a good approximation for the model under study before using it. In case of doubt, one can always resort to Input Option `hadr`.

As an example, we show here how the normalised model cross section $Q_{\text{model}}(X)$ is obtained on the basis of Eq. (45) for the search topology $p\bar{p} \rightarrow H \rightarrow$

$W^+W^- \rightarrow l^+l'^-$ of Table 2 for a model Higgs boson h_k :

$$Q_{\text{model}}(X) = \left\{ \left(\frac{\hat{\sigma}_{gg \rightarrow h_k}^{\text{model}}(\hat{s}_0, m_{h_k})}{\hat{\sigma}_{gg \rightarrow h_k}^{\text{SM}}(\hat{s}_0, m_H)} \frac{\sigma_{\text{SM}}(p\bar{p} \rightarrow gg \rightarrow H, m_H)}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H, m_H)} + \frac{\hat{\sigma}_{b\bar{b} \rightarrow h_k}^{\text{model}}(\hat{s}_0, m_{h_k})}{\hat{\sigma}_{b\bar{b} \rightarrow h_k}^{\text{SM}}(\hat{s}_0, m_H)} \frac{\sigma_{\text{SM}}(p\bar{p} \rightarrow b\bar{b} \rightarrow H, m_H)}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H, m_H)} \right) \times \frac{\text{BR}_{h_k \rightarrow W^+W^-}^{\text{model}}(m_{h_k})}{\text{BR}_{H \rightarrow W^+W^-}^{\text{SM}}(m_H)} \right\} \Big|_{m_H=m_{h_k}}.$$

If the model under study predicts kinematical distributions which differ significantly from the distributions of the reference SM processes, the exclusion result, returned by `HiggsBounds`, can only be considered an estimate. Further comments regarding this limitation from the end of Section 2.2.1 apply here too.

2.3.3 Tevatron input (`HiggsBounds Input Option effC`): effective couplings

The user is asked to provide model predictions for the Higgs boson masses m_{h_k} and total decay widths $\Gamma_{\text{tot}}(h_k)$ (in units of GeV), and for the following normalised effective couplings squared:

$$\left(\frac{g_{h_k gg}^{\text{model}}}{g_{H gg}^{\text{SM}}} \right)^2, \quad \left(\frac{g_{h_k \gamma\gamma}^{\text{model}}}{g_{H \gamma\gamma}^{\text{SM}}} \right)^2, \quad \left(\frac{g_{h_k ZZ}^{\text{model}}}{g_{H ZZ}^{\text{SM}}} \right)^2, \quad \left(\frac{g_{h_k WW}^{\text{model}}}{g_{H WW}^{\text{SM}}} \right)^2, \quad \left(\frac{g_{h_k ff, \text{eff}}^{\text{model}}}{g_{H ff}^{\text{SM}}} \right)^2, \quad (46)$$

with $k \in \{1, \dots, n_{\text{Higgs}}\}$ and $f \in \{b, \tau\}$. The reference coupling constants $g_{H ZZ}^{\text{SM}}$ and $g_{H ff}^{\text{SM}}$ are defined in Eqs. (26) and (27), respectively, and $g_{H WW}^{\text{SM}}$ by

$$(g_{H WW}^{\text{SM}})^2 = \left(\frac{e}{s_w} m_W \right)^2. \quad (47)$$

A convenient way to evaluate the ratios of the loop-induced effective couplings might be to use the ratio of partial decay widths:

$$\left(\frac{g_{h_k vv'}^{\text{model}}}{g_{H vv'}^{\text{SM}}} \right)^2 = \frac{\Gamma_{h_k \rightarrow vv'}^{\text{model}}(m_{h_k})}{\Gamma_{H \rightarrow vv'}^{\text{SM}}(m_H)} \Big|_{m_H=m_{h_k}}, \quad (48)$$

with $vv' \in \{gg, \gamma\gamma\}$. Of course, the user may apply Eq. (48) also for $vv' \in \{ZZ, WW\}$ if m_H is large enough or the Γ s for off-shell vector bosons are used. If the user has only the absolute partial decay widths, $\Gamma_{h_k \rightarrow F}^{\text{model}}(m_{h_k})$, available, he or she can use the built-in functions for the calculation of SM quantities (see Table 12 below for details) in order to calculate the right-hand side of Eq. (48).

From the input specified in Eqs. (24) and (46) `HiggsBounds` evaluates all partonic cross section ratios, as required in the Tevatron part of Input Option `part`, using the following formulae which are valid in the effective coupling approximation:

$$\frac{\hat{\sigma}_{\text{model}}(gg \rightarrow h_k)}{\hat{\sigma}_{\text{SM}}(gg \rightarrow H)} = \left(\frac{g_{h_k gg}^{\text{model}}}{g_{H gg}^{\text{SM}}} \right)^2, \quad (49)$$

$$\frac{\hat{\sigma}_{\text{model}}(b\bar{b} \rightarrow h_k)}{\hat{\sigma}_{\text{SM}}(b\bar{b} \rightarrow H)} = \frac{\hat{\sigma}_{\text{model}}(bg \rightarrow h_k b)}{\hat{\sigma}_{\text{SM}}(bg \rightarrow Hb)} = \left(\frac{g_{h_k b\bar{b}, \text{eff}}^{\text{model}}}{g_{H b\bar{b}}^{\text{SM}}} \right)^2, \quad (50)$$

$$\frac{\hat{\sigma}_{\text{model}}(q\bar{q}' \rightarrow h_k W^+)}{\hat{\sigma}_{\text{SM}}(q\bar{q}' \rightarrow HW^+)} = \frac{\hat{\sigma}_{\text{model}}(q'\bar{q} \rightarrow h_k W^-)}{\hat{\sigma}_{\text{SM}}(q'\bar{q} \rightarrow HW^-)} = \left(\frac{g_{h_k WW}^{\text{model}}}{g_{HW}^{\text{SM}}} \right)^2, \quad (51)$$

$$\frac{\hat{\sigma}_{\text{model}}(q''\bar{q}'' \rightarrow h_k Z)}{\hat{\sigma}_{\text{SM}}(q''\bar{q}'' \rightarrow HZ)} = \left(\frac{g_{h_k ZZ}^{\text{model}}}{g_{HZ}^{\text{SM}}} \right)^2, \quad (52)$$

for $(q, q') \in \{(u, d), (c, s)\}$ and $q'' \in \{u, d, c, s, b\}$.

In order to facilitate the calculation of the ratios of the hadronic cross sections for h_k production via VBF, we obtained the following SM cross section ratio for $p\bar{p}$ collisions with 1.96 TeV centre-of-mass energy using VBFNLO [29]:

$$\frac{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via } WW \text{ fusion})}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via } ZZ \text{ fusion})} \approx 3.35. \quad (53)$$

The right-hand side is a mean value. The actual value varies slightly with m_H ($\pm 3.6\%$ in the relevant mass range of $70 \text{ GeV} < m_H < 300 \text{ GeV}$). Using this mean and neglecting interference effects, which affect the result below 1% (cf. [53]), we can calculate the proportions of WW - and ZZ -fusion contributions to the SM VBF Higgs production cross section for $p\bar{p}$ collisions with 1.96 TeV centre-of-mass energy:

$$R_{\text{VBF}}^{WW} := \frac{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via } WW \text{ fusion})}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via VBF})} = 77\%, \quad (54)$$

$$R_{\text{VBF}}^{ZZ} := \frac{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via } ZZ \text{ fusion})}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via VBF})} = 23\%. \quad (55)$$

The slight variation of the right-hand side of Eq. (53) with m_H causes a variation of the above proportions by less than 1%.

Using the above SM predictions, we can calculate the model predictions for the ratios of the hadronic cross sections for h_k production via VBF quite accurately, as follows:

$$\frac{\sigma_{\text{model}}(p\bar{p} \rightarrow h_k \text{ via VBF})}{\sigma_{\text{SM}}(p\bar{p} \rightarrow H \text{ via VBF})} = R_{\text{VBF}}^{WW} \left(\frac{g_{h_k WW}^{\text{model}}}{g_{HW}^{\text{SM}}} \right)^2 + R_{\text{VBF}}^{ZZ} \left(\frac{g_{h_k ZZ}^{\text{model}}}{g_{HZ}^{\text{SM}}} \right)^2. \quad (56)$$

Branching ratios, additional to the ones we have already described when discussing the LEP limits in Section 2.2.2, are evaluated along the lines of Eq. (33):

$$\begin{aligned} \text{BR}_{\text{model}}(h_k \rightarrow \gamma\gamma) &= \text{BR}_{\text{SM}}(H \rightarrow \gamma\gamma)(m_H) \frac{\Gamma_{\text{tot}}^{\text{SM}}(m_H)}{\Gamma_{\text{tot}}(h_k)} \Big|_{m_H=m_{h_k}} \times \\ &\times \left(\frac{g_{h_k\gamma\gamma}^{\text{model}}}{g_{H\gamma\gamma}^{\text{SM}}} \right)^2, \end{aligned} \quad (57)$$

$$\begin{aligned} \text{BR}_{\text{model}}(h_k \rightarrow WW) &= \text{BR}_{\text{SM}}(H \rightarrow WW)(m_H) \frac{\Gamma_{\text{tot}}^{\text{SM}}(m_H)}{\Gamma_{\text{tot}}(h_k)} \Big|_{m_H=m_{h_k}} \times \\ &\times \left(\frac{g_{h_k WW}^{\text{model}}}{g_{HWW}^{\text{SM}}} \right)^2. \end{aligned} \quad (58)$$

At this point, all quantities which are required as Tevatron input by Input Option `part` are evaluated using the effective coupling approximation. The evaluation of the model predictions $Q_{\text{model}}(X)$ can then proceed as exemplified in Section 2.3.2 above.

For search topologies, of which the amplitude is proportional to one monomial of Higgs coupling constants, effective couplings just rescale all kinematical distributions of the reference process by a common factor without changing their shape. Consequently, in such cases, limits on topological cross sections can be exactly applied. For the application of limits obtained from SM analyses, which combine a set of search topologies, the SM-likeness check (see Section 2.1) ensures that the kinematical distributions arising from models that pass this check closely resemble the distributions of the SM. Thus, also in this case, limits on topological cross sections can be exactly applied.

Two slightly less straight-forward cases require further discussion.

a) For the single Higgs production topology, we add the gluon-fusion and $b\bar{b}$ annihilation contribution together. Firstly, one might expect that the boost distribution of the produced Higgs boson is different for the two partonic processes and thus the kinematical distributions of the Higgs decay products would depend on the relative contributions of the two processes to the signal. However, it turns out that the gluon and bottom parton distributions (using MSTW 2008 PDFs [54]) in the relevant region (lower x values) are, to a very good approximation, proportional to each other, i.e. the two parton processes do not give rise to different boost distributions of the produced Higgs boson. Secondly, it is known that the event selection cuts used in single Higgs analyses select Higgs bosons with a certain p_T , which is known to lead to different distributions for processes arising from gluon-gluon and $b\bar{b}$ initial states. This change in the kinematical distribution becomes relevant if the coupling of the Higgs to a bottom quark is strongly enhanced such that the $b\bar{b}$ annihilation

competes with gluon-fusion contribution, where the latter dominates in the SM. Indeed, it turns out that, for such a scenario, out of the implemented set of analyses, the most sensitive are those that involve the search topology $p\bar{p} \rightarrow H \rightarrow \tau\tau$ [24,25]. However, these analyses reject events with extra jets of large transverse momentum, where the deviation of the p_T -distributions becomes substantial [55]. Therefore, we do not expect a strong influence of the moderately changed kinematical distributions on the cross section limits. This expectation is supported by the fact that we reproduce the MSSM exclusion plots in [24,25] reasonably well.

b) For Higgs production via vector boson fusion, a deviation from the SM proportion of WW and ZZ initiated processes caused by different effective couplings $g_{h_k ZZ}^2$ and $g_{h_k WW}^2$, does not lead to significant changes in the kinematical distributions compared to the SM reference process.

In summary, for the implemented set of analyses, we expect the limits on topological cross sections to be applicable with high accuracy for models where the Higgs sector can be faithfully parametrised with effective couplings.

3 HiggsBounds Operating Instructions

There are three formats in which the program `HiggsBounds` can be used:

- Library of subroutines
- Command-line version
- Online version

The most widely applicable format of `HiggsBounds` is the command-line version, since this reads all the model data from text files and thus this model data can be generated using any package the user wishes. The library of `HiggsBounds` subroutines allows `HiggsBounds` to be called within other programs. If the user just wishes to check a few parameter points, the online version provides quick access to all the functionality of `HiggsBounds`, without the need to install the code.

The `HiggsBounds` code, the online version and documentation can all be found at the URL www.ippp.dur.ac.uk/HiggsBounds .

The `HiggsBounds` code is provided in either Fortran 77 or Fortran 90. Both codes provide exactly the same functionality and have exactly the same operating instructions. In fact, the Fortran 77 and Fortran 90 versions of the `HiggsBounds` subroutines can even be called within codes written in Fortran 90 and Fortran 77, respectively. Both codes have also been tested with a variety of Fortran compilers, including the free gnu compilers which accompany

most Linux distributions. Therefore, the user may download either code and the difference will only be apparent if the user wishes to examine the structure of the code⁶.

The library of subroutines, the command-line version and the online version share a common set of features, which we will describe first. We will then give operating instructions for each of these three `HiggsBounds` formats individually.

3.1 Common features: Input

`HiggsBounds` requires four types of input:

- the number of neutral Higgs bosons in the model under study (`nH`)
- the set of experimental analyses which should be considered (`whichexpt`)
- the theoretical predictions of the model under study (a set of input arrays)
- the format of these theoretical predictions (`whichinput`)

Table 4 contains further information on the variable `nH`, and the possible values of `whichexpt` are described in Table 5. Note that the option `whichexpt='singH'` should only be used if neither processes involving Higgs pair production at LEP nor processes involving the $h_j \rightarrow h_i h_i$ decay are relevant. However, if these conditions are met, this option can save significantly on computing time.

`HiggsBounds` expects the theoretical input to be in one of three formats, labelled by the variable `whichinput`. These formats were described in detail in Section 2 and are briefly summarised in Table 6. In Table 7 and Table 8 we assign names to all of the possible input arrays (each array is defined in terms of the notation used in Section 2). These names will prove useful when we describe the input requirements of each version of `HiggsBounds` individually.

<code>nH</code>	(integer)
1-9	By default, <code>HiggsBounds</code> can be applied to models with between 1 and 9 neutral Higgs bosons. (This range can easily be extended by the user if required.)

Table 4

The possible values of the variable `nH`, which labels the number of neutral Higgs bosons in the model under study.

⁶ The maintenance of two separate codes is primarily intended to provide an efficient way for the authors to confirm that each update is free from implementation errors.

<code>whichexpt</code>	(<code>character(LEN=5)</code>)
<code>LandT</code>	both LEP and Tevatron analyses
<code>onlyL</code>	only LEP analyses
<code>onlyT</code>	only Tevatron analyses
<code>singH</code>	only analyses for processes involving one Higgs boson

Table 5

The possible values of the variable `whichexpt`, which indicates which subset of experimental analyses will be considered by `HiggsBounds`.

<code>whichinput</code>	(<code>character(LEN=4)</code>)
<code>effC</code>	Masses, total decay widths, ratios of effective couplings squared, $\text{BR}(h_j \rightarrow h_i h_i)$.
<code>part</code>	Masses, total decay widths, ratios of LEP cross sections, mainly ratios of partonic Tevatron cross sections, branching ratios.
<code>hadr</code>	Masses, total decay widths, ratios of LEP cross sections, ratios of hadronic Tevatron cross sections, branching ratios.

Table 6

The possible values of the variable `whichinput`, which indicates the format of the theoretical predictions provided by the user. (See Section 2 for a more detailed description of each of these settings).

input arrays	(double precision)	
Mh(nH)	m_{h_i}	in GeV
GammaTotal(nH)	$\Gamma_{\text{tot}}(h_i)$	in GeV
g2hjbb(nH)	$\left(\frac{g_{h_j(\text{OP})}^{\text{model}}}{g_{H(\text{OP})}^{\text{SM}}}\right)^2$,	OP = $b\bar{b}$
g2hjtautau(nH)		$\tau^+\tau^-$
g2hjWW(nH)		WW
g2hjZZ(nH)		ZZ
g2hjgaga(nH)		$\gamma\gamma$
g2hjgg(nH)		gg
g2hjhiZ(nH,nH)	$\left(\frac{g_{h_j h_i Z}^{\text{model}}}{g_{HH'Z}^{\text{ref}}}\right)^2$	
BR_hjbb(nH)	BR($h_j \rightarrow \text{OP}$),	OP = $b\bar{b}$
BR_hjtautau(nH)		$\tau^+\tau^-$
BR_hjWW(nH)		WW
BR_hjgaga(nH)		$\gamma\gamma$
BR_hjhihi(nH,nH)	BR($h_j \rightarrow h_i h_i$)	

Table 7

Input arrays for model predictions for effective normalised squared couplings and branching ratios recognised by HiggsBounds. The size of each array is given in brackets in the first column. See Section 2 for the description of the notation used in the second column. The elements of BR_hjhihi are ordered such that $\text{BR_hjhihi}(j,i) = \text{BR}(h_j \rightarrow h_i h_i)$.

input arrays cont.	(double precision)	
CS_lep_hjZ_ratio(nH)	$R_\sigma(P)$,	$P = e^+e^- \rightarrow h_j Z$
CS_lep_hjhi_ratio(nH,nH)		$e^+e^- \rightarrow h_j h_i$
CS_tev_pp_hj_ratio(nH)		$p\bar{p} \rightarrow h_j$
CS_tev_pp_hjb_ratio(nH)		$p\bar{p} \rightarrow b h_j$
CS_tev_pp_hjW_ratio(nH)		$p\bar{p} \rightarrow h_j W$
CS_tev_pp_hjZ_ratio(nH)		$p\bar{p} \rightarrow h_j Z$
CS_tev_pp_vbf_ratio(nH)		$p\bar{p} \rightarrow h_j$ via VBF
CS_tev_gg_hj_ratio(nH)	$R_{nm}^{h_j}$,	$nm = gg$
CS_tev_bb_hj_ratio(nH)		$b\bar{b}$
CS_tev_ud_hjWp_ratio(nH)	$R_{nm}^{h_j+W^+}$,	$nm = u\bar{d}$
CS_tev_cs_hjWp_ratio(nH)		$c\bar{s}$
CS_tev_ud_hjWm_ratio(nH)	$R_{nm}^{h_j+W^-}$	$nm = d\bar{u}$
CS_tev_cs_hjWm_ratio(nH)		$s\bar{c}$
CS_tev_dd_hjZ_ratio(nH)	$R_{nm}^{h_j+Z}$	$nm = d\bar{d}$
CS_tev_uu_hjZ_ratio(nH)		$u\bar{u}$
CS_tev_ss_hjZ_ratio(nH)		$s\bar{s}$
CS_tev_cc_hjZ_ratio(nH)		$c\bar{c}$
CS_tev_bb_hjZ_ratio(nH)		$b\bar{b}$
CS_tev_bg_hjb_ratio(nH)	$R_{nm}^{h_j+b}$	$nm = bg, \bar{b}g$

Table 8

Input arrays for model predictions for cross section ratios recognised by HiggsBounds. The size of each array is given in brackets in the first column. The LEP or hadronic Tevatron cross section ratios $R_\sigma(P)$ are defined in Eq. (43) and the partonic Tevatron cross section ratios $R_{nm}^{h_j+y}$ are defined in Eq. (44).

3.2 Common features: Output

HiggsBounds provides the user with four types of output:

- whether the parameter point is excluded at the 95% C.L. or not (**HBresult**)
- the identifying number of the process with the highest statistical sensitivity (**chan**).
- the number of Higgs bosons which have contributed to the theoretical rate for this process (**ncombined**)
- the ratio of the theoretical rate Q_{model} to the observed limit Q_{obs} for this process (**obsratio**).

Table 9 shows the possible values of **HBresult** and **obsratio**, which are complementary. Tables 10 and 11 discuss **chan** and **ncombined** respectively. If the library of subroutines or the command-line versions are used, the key to the process numbers is written in the file `Key.dat`. In the online version, this information appears on the screen.

HBresult	obsratio	
(integer)	(double precision)	
0	≥ 1.0	parameter point is excluded
1	< 1.0	parameter point is not excluded
-1	≤ 0.0	invalid parameter set

Table 9

*The possible values of the output variables **HBresult** and **obsratio**, which indicate whether a parameter point has been excluded at the 95% C.L. by the experimental results under consideration.*

chan	(integer)
1-[# of considered analyses]	See the file <code>Key.dat</code> for the definition of each process number. <code>Key.dat</code> is automatically generated when either the command line or the subroutine version of HiggsBounds are used.

Table 10

*Further information about the output variable **chan**, which stores the reference number of the process with the highest statistical sensitivity.*

<code>ncombined</code>	(integer)
<code>1-n_H</code>	Number of Higgs bosons which have contributed to the theoretical rate for this process. The number depends on <code>delta_Mh_LEP</code> or <code>delta_Mh_TEV</code> .

Table 11

Further information about the output variable `ncombined`.

3.3 Library of subroutines

Installation

The `HiggsBounds` code can be compiled to form a library of subroutines using the following commands:

```
./configure
make libHB
```

A program which wishes to use the `HiggsBounds` subroutines can be compiled and linked to the library by adding `-L<HBpath> -lHB` to the command line, for example,

```
gfortran myprog.f90 -o myprog -L<HBpath> -lHB
```

where `<HBpath>` is the location of the `HiggsBounds` library.

Subroutine `initialize_HiggsBounds`

The subroutine `initialize_HiggsBounds` must be called before any other `HiggsBounds` subroutine. It performs some preparatory operations such as reading in the tables of data. It is called as:

```
call initialize_HiggsBounds(nH, whichexpt)
```

When using the `HiggsBounds` subroutines in another code, the subroutine `initialize_HiggsBounds` must be called only once, before any other `HiggsBounds` subroutine is called.

Subroutines `run_HiggsBounds_effC`, `run_HiggsBounds_part` and `run_HiggsBounds_hadr`

The subroutines `run_HiggsBounds_effC`, `run_HiggsBounds_part` and `run_HiggsBounds_hadr` perform the main part of the `HiggsBounds` calculations. They set the value of `whichinput` to be `effC`, `part` and `hadr` respectively and therefore require different arguments. These subroutines are called

as:

```
call run_HiggsBounds_effC(nH,Mh,GammaTotal,  
  &      g2hjbb,g2hjtautau,g2hjWW,g2hjZZ,  
  &      g2hjgaga,g2hjgg,g2hjhiZ,  
  &      BR_hjhihi,  
  &      HBresult,chan,  
  &      obsratio, ncombined )
```

```
call run_HiggsBounds_part(nH,Mh,  
  &      CS_lep_hjZ_ratio,   CS_lep_hjhi_ratio,  
  &      CS_tev_gg_hj_ratio, CS_tev_bb_hj_ratio,  
  &      CS_tev_bg_hjb_ratio,  
  &      CS_tev_ud_hjWp_ratio,CS_tev_cs_hjWp_ratio,  
  &      CS_tev_ud_hjWm_ratio,CS_tev_cs_hjWm_ratio,  
  &      CS_tev_dd_hjZ_ratio, CS_tev_uu_hjZ_ratio,  
  &      CS_tev_ss_hjZ_ratio, CS_tev_cc_hjZ_ratio,  
  &      CS_tev_bb_hjZ_ratio,  
  &      CS_tev_pp_vbf_ratio,  
  &      BR_hjbb,BR_hjtautau,  
  &      BR_hjWW,BR_hjgaga,  
  &      BR_hjhihi,  
  &      HBresult,chan,  
  &      obsratio, ncombined )
```

```
call run_HiggsBounds_hadr(nH,Mh,  
  &      CS_lep_hjZ_ratio,   CS_lep_hjhi_ratio,  
  &      CS_tev_pp_hj_ratio,  CS_tev_pp_hjb_ratio,  
  &      CS_tev_pp_hjW_ratio, CS_tev_pp_hjZ_ratio,  
  &      CS_tev_pp_vbf_ratio,  
  &      BR_hjbb,BR_hjtautau,  
  &      BR_hjWW,BR_hjgaga,  
  &      BR_hjhihi,  
  &      HBresult,chan,  
  &      obsratio, ncombined )
```

Each of these arguments must be supplied. However, if a branching ratio, effective coupling or cross section is believed to be irrelevant, the corresponding array may be filled with zeros. This will ensure that the value of Q_{model} for processes involving this quantity will also be zero. For example, in the MSSM, processes involving the decay $h_j \rightarrow \gamma\gamma$ will rarely be the process with the highest statistical sensitivity of $Q_{\text{model}}/Q_{\text{expec}}$ and, consequently, it may be convenient to set the arrays `g2hjgaga` and `BR_hjgaga` to zero for simplicity.

Also, depending on the value given for `whichexpt`, some of the input arrays will be ignored within `HiggsBounds`. For example, if `whichexpt='onlyT'`, the branching ratio for the Higgs cascade decay $h_j \rightarrow h_i h_i$ will not be relevant. Therefore, setting this array to zero in this case will not affect the `HiggsBounds` results in this case.

Subroutine finish_HiggsBounds

The subroutine `finish_HiggsBounds` should be called once at the end of the program, after all other `HiggsBounds` subroutines⁷. It is called as:

```
call finish_HiggsBounds
```

Functions for Standard Model branching ratios, total decay width and cross sections

The `HiggsBounds` library also allows users access to the Standard Model Higgs branching ratios, total decay width and production cross sections, which are used internally by `HiggsBounds`. We use Standard Model Higgs branching ratios and total decay width from the program HDECAY 3.303 [26]. The SM hadronic cross sections have been obtained from the TEV4LHC Higgs Working Group [27] (see Table 12 for references to the original works) with the exception of the $\sigma^{\text{SM}}(p\bar{p} \rightarrow b\bar{g} \rightarrow bH)$ cross section. The latter cross section has been calculated with the program HJET 1.1 [28] for a set of different cuts on the transverse momentum and pseudo-rapidity of the b -quark, which are needed internally in order to apply correctly the results of some Tevatron analyses and which were not available from [27]. From this set, only the cross section without cuts is externally provided.

Examples

We have provided three example programs which demonstrate the use of the `HiggsBounds` subroutines. The first example relates to the Fourth Generation Model and is contained in the file `example-SM_vs_4thGen.F` (see Fig. 4 below for a code listing). This program uses the `HiggsBounds` functions for the SM branching ratios and SM total decay width to calculate the Higgs decay width and the effective normalised squared couplings in the SM and a simple Fourth Generation Model. This information is then used as input for the subroutine `run_HiggsBounds_effC`, which is called once with SM input and once with Fourth Generation Model input. The example will be described in more detail in Section 4.2. Once the `HiggsBounds` library has been compiled (using `./configure ; make libHB` as described previously), the code

⁷ In the Fortran 90 version of the code, the subroutine `finish_HiggsBounds` is used to deallocate the allocatable arrays used within `HiggsBounds`.

function	(double precision)		
SMGamma_h(Mh)	$\Gamma_{\text{tot}}^{\text{SM}}(h_i)$		[26]
SMBR_Hbb(Mh)	$\text{BR}^{\text{SM}}(H \rightarrow \text{OP}),$	OP= $b\bar{b}$	[26]
SMBR_Htautau(Mh)		$\tau^- \tau^+$	[26]
SMBR_HWW(Mh)		WW	[26]
SMBR_Hgamgam(Mh)		$\gamma\gamma$	[26]
SMBR_Hgg(Mh)		gg	[26]
SMBR_HZgam(Mh)		$Z\gamma$	[26]
SMBR_HZZ(Mh)		ZZ	[26]
SMBR_Httop(Mh)		$t\bar{t}$	[26]
SMBR_Hcc(Mh)		$c\bar{c}$	[26]
SMBR_Hss(Mh)		$s\bar{s}$	[26]
SMBR_Hmumu(Mh)		$\mu^- \mu^+$	[26]
SMCS_tev_pp_qq_HW(Mh)	$\sigma^{\text{SM}}(P),$	P= $p\bar{p} \rightarrow q\bar{q} \rightarrow HW$	[56]
SMCS_tev_pp_qq_HZ(Mh)		$p\bar{p} \rightarrow q\bar{q} \rightarrow HZ$	[56]
SMCS_tev_pp_gg_H(Mh)		$p\bar{p} \rightarrow gg \rightarrow H$	[50,51]
SMCS_tev_pp_bb_H(Mh)		$p\bar{p} \rightarrow b\bar{b} \rightarrow H$	[52]
SMCS_tev_pp_vbf_H(Mh)		$p\bar{p} \rightarrow H$ via VBF	[57]
SMCS_tev_pp_bg_Hb(Mh)		$p\bar{p} \rightarrow bg \rightarrow Hb$	[28]

Table 12

Standard Model branching ratios, total decay widths in units of GeV and hadronic Tevatron cross sections in units of pikobarn provided as functions by HiggsBounds, together with references. Each function takes a Higgs mass Mh (double precision) as its argument.

example-SM_vs_4thGen.F can be compiled and run with the commands:

```
gfortran example-SM_vs_4thGen.F -o HBfourthgen -L<HBpath> -lHB
./HBfourthgen
```

where <HBpath> is the location of the HiggsBounds library.

The files HBwithFH.F and HBwithCPsuperH.f demonstrate the use of HiggsBounds subroutines with the publically available programs FeynHiggs [14] and CPsuperH [18], respectively. We refer the reader to the extensive comments contained within these example files for further details.

3.4 Command line version

Installation

In order to be able to call `HiggsBounds` from the command line, it should be compiled using the commands

```
./configure  
make
```

Command line and input file format

In the command-line usage of `HiggsBounds`, the arrays containing the theoretical model predictions are read from text files. The other options are specified in the command line, which is of the form:

```
./HiggsBounds <whichexpt> <whichinput> <nH> <prefix>
```

The variable `<prefix>` is a string which is added to the front of input and output file names and may include directory names or other identifying information.

Table 13 describes the contents of each input file. Note that each input file should start with a line number. The input files should not contain any comments or blank lines. The line number identifies the predictions which belong to the same model parameter point in different files.

Care should be taken with the order of the array elements in the files. This is best illustrated by an example, where we will use $n_H = 3$. The one dimensional arrays, e.g. `Mh`, should be given in the order

`Mh(1), Mh(2), Mh(3)`

However, not all of the elements of the two dimensional arrays are required. Only the lower left triangle (including the diagonal) is required from the arrays `g2hjhiZ` and `lepCS_hjhi_ratio`, since they are symmetric, e.g.

$$\begin{pmatrix} g2hjhiZ(1,1) & g2hjhiZ(1,2) & g2hjhiZ(1,3) \\ g2hjhiZ(2,1) & g2hjhiZ(2,2) & g2hjhiZ(2,3) \\ g2hjhiZ(3,1) & g2hjhiZ(3,2) & g2hjhiZ(3,3) \end{pmatrix}$$

i.e. the elements in the input file should be written in the order

`g2hjhiZ(1,1), g2hjhiZ(2,1), g2hjhiZ(2,2), g2hjhiZ(3,1),
g2hjhiZ(3,2), g2hjhiZ(3,3)` .

file name	data format
MH_GammaTot.dat	k, Mh, GammaTotal
effC.dat	k, g2hjbb, g2hjtautau, g2hjWW, g2hjZZ, g2hjgaga, g2hjgg, some elements of g2hjhiZ (lower left triangle - see example)
LEP_HZ_CS_ratios.dat	k, CS_lep_hjZ_ratio
LEP_2H_CS_ratios.dat	some elements of CS_lep_hjhi_ratio (lower left triangle - see example)
TEV_H_0jet_partCS_ratios.dat	k, CS_tev_gg_hj_ratio, CS_tev_bb_hj_ratio
TEV_H_1jet_partCS_ratios.dat	k CS_tev_bb_hjb_ratio
TEV_HW_partCS_ratios.dat	k, CS_tev_ud_hjWp_ratio, CS_tev_cs_hjWp_ratio, CS_tev_ud_hjWm_ratio, CS_tev_cs_hjWm_ratio
TEV_HZ_partCS_ratios.dat	k, CS_tev_dd_hjZ_ratio, CS_tev_uu_hjZ_ratio, CS_tev_ss_hjZ_ratio, CS_tev_cc_hjZ_ratio, CS_tev_bb_hjZ_ratio
TEV_H_vbf_hadCS_ratios.dat	k, CS_tev_pp_vbf_ratio
TEV_1H_hadCS_ratios.dat	k, CS_tev_pp_hj_ratio, CS_tev_pp_hjb_ratio, CS_tev_pp_hjW_ratio, CS_tev_pp_hjZ_ratio, CS_tev_pp_vbf_ratio
BR_1H.dat	k, BR_hjbb, BR_hjtautau, BR_hjWW, BR_hjgaga
BR_2H.dat	k, some elements of BR_hjhihi (row by row, without diagonal - see example)
additional.dat(optional)	k, ...

Table 13

Names and data format of all HiggsBounds input files. The right column shows the order of the input data arrays within one line of the input file. For the order within the arrays, see example. k is the line number. Note that the array CS_tev_pp_vbf_ratio appears in two different input files. However, these files will never be required by HiggsBounds simultaneously.

For the array `BR_hjhihi`, only the off-diagonal components are required

$$\begin{pmatrix} \text{BR_hjhihi}(1, 1) & \text{BR_hjhihi}(1, 2) & \text{BR_hjhihi}(1, 3) \\ \text{BR_hjhihi}(2, 1) & \text{BR_hjhihi}(2, 2) & \text{BR_hjhihi}(2, 3) \\ \text{BR_hjhihi}(3, 1) & \text{BR_hjhihi}(3, 2) & \text{BR_hjhihi}(3, 3) \end{pmatrix}$$

since the diagonal elements are not physical quantities. Therefore, the elements should be written in the order

```
BR_hjhihi(1,2), BR_hjhihi(1,3), BR_hjhihi(2,1), BR_hjhihi(2,3),
BR_hjhihi(3,1), BR_hjhihi(3,2)
```

in the input file.

The file `additional.dat` is optional. If it is included, it can have any number of columns greater than 1 (as for the previous files, the first entry on each line should be the line number). It is envisaged that this input file will be particularly useful when parameter scans are performed over a variable which is not required by `HiggsBounds` but helpful plotting the results. For example, in the case of the complex MSSM, `additional.dat` could be used to store the values of $\tan\beta$ and the charged Higgs mass.

As in the subroutine version, the command line version of `HiggsBounds` expects a subset of the total list of input arrays, which depends on the chosen setting of `whichinput`. The maximal list of files used for each value of `whichinput` is given in Table 5.

As discussed for the subroutine version, some of the arrays will not be relevant for some of the choices for `whichexpt`. The command line version of `HiggsBounds` will consider the list of input files appropriate to the setting `whichinput` and then only attempt to read any of these input files if the value chosen for `whichexpt` means that at least one of the arrays it contains will be directly used. Table 15 contains a list of which input files are actually relevant to each value of `whichexpt`. For example, if `whichinput = 'hadr'`, and `whichexpt = 'LandT'`, then `HiggsBounds` requires the input files:

```
MH_GammaTot.dat, BR_2H.dat, BR_1H.dat, LEP_HZ_CS_ratios.dat,
LEP_2H_CS_ratios.dat, TEV_1H_hadCS_ratios.dat .
```

However, if `whichinput = 'hadr'` and `whichexpt = 'onlyL'`, `HiggsBounds` requires the input files:

```
MH_GammaTot.dat, BR_2H.dat, BR_1H.dat, LEP_HZ_CS_ratios.dat,
LEP_2H_CS_ratios.dat .
```

whichinput = 'part',	'hadr',	'effC'
MH_GammaTot.dat	MH_GammaTot.dat	MH_GammaTot.dat
BR_2H.dat	BR_2H.dat	effC.dat
BR_1H.dat	BR_1H.dat	BR_2H.dat
LEP_HZ_CS_ratios.dat	LEP_HZ_CS_ratios.dat	additional.dat
LEP_2H_CS_ratios.dat	LEP_2H_CS_ratios.dat	
TEV_H_0jet_partCS_ratios.dat	TEV_1H_hadCS_ratios.dat	
TEV_H_1jet_partCS_ratios.dat	additional.dat	
TEV_HW_partCS_ratios.dat		
TEV_HZ_partCS_ratios.dat		
TEV_H_vbf_hadCS_ratios.dat		
additional.dat		

Table 14

The list of possible input files for each value of whichinput. Note that some input files may not be relevant, depending on the value of whichexpt. In this case, they are not required. See Table 15 for more details.

As a third example, if whichinput = 'hadr' and whichexpt = 'onlyT', HiggsBounds requires the input files:

MH_GammaTot.dat, BR_1H.dat, TEV_1H_hadCS_ratios.dat .

In each of these three examples, HiggsBounds will also read the file

additional.dat

if it exists.

As for the subroutine version, if the user does not require processes involving a particular branching ratio or cross section ratio to be checked by HiggsBounds, that particular array can be filled with zeros.

Output file format

When the command line version of HiggsBounds is used, the output is written to the file <prefix>HiggsBounds_results.dat. A sample of the output is shown in Fig. 2. The key to the process numbering is written to <prefix>Key.dat.

name of input file	values of <code>whichexpt</code>			
	which this file is relevant to			
	LandT	onlyL	onlyT	singH
MH_GammaTot.dat	y	y	y	y
effC.dat	y	y	y	y
LEP_HZ_CS_ratios.dat	y	y		y
LEP_2H_CS_ratios.dat	y	y		
TEV_H_0jet_partCS_ratios.dat	y		y	y
TEV_H_1jet_partCS_ratios.dat	y		y	y
TEV_HW_partCS_ratios.dat	y		y	y
TEV_HZ_partCS_ratios.dat	y		y	y
TEV_H_vbf_hadCS_ratios.dat	y		y	y
TEV_1H_hadCS_ratios.dat	y		y	y
BR_1H.dat	y	y	y	y
BR_2H.dat	y	y		
additional.dat (optional)	y	y	y	y

Table 15

List of input files, specifying which values of `whichexpt` each input file is relevant to (marked by 'y').

Examples

The `HiggsBounds` package includes a full set of sample input files for the case $n_H = 3$, contained in the folder `example`. Each filename is prefixed with `nH3_`. To run the command-line version of `HiggsBounds` with these files as input, use, for example,

```
./configure
make
./HiggsBounds LandT effC 3 'example/nH3_'
```

where the values of `whichexpt` and `whichinput` can be varied as desired.

3.5 Website version

The website allows the user to select the required number of neutral Higgs bosons and then generates a `html` form accordingly. The values of `whichinput`

```

# generated with HiggsBounds on 31.10.2008 at 11:18
# settings: LandT, effC
#
# column abbreviations
# n          : line id of input
# Mh(i)      : Higgs boson masses
# HBresult   : scenario allowed flag (1: allowed, 0: excluded, -1: unphysical)
# chan       : most sensitive channel (see below). chan=0 if no channel applies
# obsratio   : ratio [sig x BR]_model/[sig x BR]_limit (<1: allowed, >1: excluded)
# ncomb      : number of Higgs bosons combined in most sensitive channel
# additional  : optional additional data stored in <prefix>additional.dat (e.g. tan beta)
#
# channel numbers used in this file
# 3 : (ee)->(h3)Z->(b b)Z (LEP table 14b)
# 4 : (ee)->(h1)Z->(tau tau)Z (LEP table 14c)
# 124 : (pp)->W(h1)->l nu (b b) (CDF Note 9463)
# 134 : (pp)->h2->tau tau (\protect\vrule width0pt\protect\href{http://arxiv.org/abs/0805.2491}{arXiv:0805.2491})
# 157 : (pp)->h1+... where h1 is SM-like (\protect\vrule width0pt\protect\href{http://arxiv.org/abs/0804.3423}{arXiv:0804.3423})
# (for full list of processes, see Key.dat)
#
#cols: n      Mh(1)    Mh(2)    Mh(3)    HBresult  chan   obsratio   ncomb  additional(1)
#
1      359.121  271.963  134.929     1    134   0.212206E-03   1    0.246862
2      75.0123  92.8677  71.9716     1     4    0.306172E-01   1    0.714964
3      136.293  345.483  330.026     1    124   0.640713E-01   1    0.434594
4      111.377  220.765  51.7469     1     3    0.162811       1    0.727173
5      186.131  355.002  146.448     0    157   15.2354        1    0.230522

```

Fig. 2. Sample output file (written to `<prefix>HiggsBounds_Results.dat`)

and `whichexpt` can be chosen and the appropriate theoretical input entered. `HiggsBounds` will then be called with these settings and the result outputted to screen. The website version contains the additional feature that it notifies the user about the processes with the second and third highest statistical sensitivities and the values of `obsratio` for these processes. This is designed to give guidance to the user who, for example, wishes to find an excluded region iteratively by adjusting the input quantities.

The website also contains a selection of pre-filled `html` forms as examples, including entries for the Standard Model, the fermiophobic Higgs model and the MSSM with real and complex parameters.

4 Examples of use

4.1 Re-evaluation of SM exclusion limit with an improved SM prediction

Recently, an improved prediction for the Higgs-boson production via gluon fusion appeared [58], which incorporates previously unknown mixed QCD–electroweak radiative corrections to this process as well as other known relevant corrections [59]. It turns out that those corrections are relevant, increasing the SM cross section prediction by 7–10% compared to the cross section prediction currently used by the Tevatron collaborations, when the MRST 2006




Fig. 3. Ratio $Q_{\text{obs}}/Q_{\text{SM}}$ for the most sensitive channel as a function of the Higgs mass m_H : (a) The current best available SM combined analyses of CDF and DØ (solid line) is compared to a re-evaluation using the recently improved SM prediction for the gluon fusion process with MRST 2006 PDFs (dashed line) and MSTW 2008 PDFs (dotted line) [58]; (b) The impact of the PDF uncertainty on the exclusion limit using the improved SM prediction [58] is shown.

NNLO parton distribution functions (PDFs) [60] are used. However, very recently, an updated set of PDFs, MSTW 2008 [54], appeared which show a decrease in the low- x gluon distribution. In an updated version of Ref. [58] the combined effect of using the improved prediction for the SM cross section and the latest set of MSTW 2008 NNLO PDFs has been found to result in a downward shift of the SM cross section prediction in the region where the Tevatron searches have the highest sensitivity by 4–6% with respect to the cross section prediction used by the Tevatron collaborations.

With `HiggsBounds` the impact of such changes in the SM cross section prediction on the SM exclusion limit can easily be assessed. Fig. 3 shows the result of a re-evaluation of the Tevatron exclusion limits. In Fig. 3(a), the limit obtained using the cross section predictions of [58] with MRST 2006 NNLO PDFs and MSTW 2008 NNLO PDFs are compared with the published limits from the CDF and DØ SM combined analyses ([48] for $m_H < 155$ GeV and [49] for $m_H > 155$ GeV) which do not include this improved prediction. While in the published Tevatron result the cross section limit just touches the SM prediction for $m_H \approx 170$ GeV, the updated SM cross section prediction exceeds the cross section limit within an interval of about 5 GeV if MRST 2006 PDFs are used. However, when the most up-to-date MSTW 2008 PDFs are used, the cross section prediction is lower than the one obtained by the Tevatron collaborations for m_H above about 160 GeV, resulting in a reduction of the exclusion power. With the updated prediction, no value of the SM Higgs boson mass is excluded at the 95% C.L. Fig. 3(b) shows how the comparison between the SM cross section prediction and the exclusion limits from the Tevatron is affected by PDF uncertainties. The solid line shows the result based on the updated SM cross section prediction using MSTW 2008 PDFs, while the dashed lines indicate the PDF uncertainties as given in Ref. [58].

It can be clearly seen that currently no firm exclusion of any SM Higgs mass values at the 95% C.L. can be established from the exclusion bounds obtained at the Tevatron once PDF uncertainties on the SM cross section prediction are taken into account.

4.2 Fourth Generation Model

A very simple example of physics beyond the SM is a model which extends the SM by a fourth generation of heavy fermions [61]. In particular, the masses of the 4th generation quarks and leptons are assumed to be (much) heavier than the mass of the top-quark. In this case, the effective coupling of the Higgs boson to two gluons is three times larger than in the SM. No other coupling, relevant to LEP and Tevatron searches, changes significantly⁸. Essentially, only the partial decay width $\Gamma(H \rightarrow gg)$ changes by a factor of 9 and, with it, the total Higgs width and therefore all the decay branching ratios [62].

We can easily calculate the new total decay width and the relevant decay branching ratios using the above information and the built-in functions to calculate SM quantities using the relations:

$$\begin{aligned}\Gamma_{\text{SM}}(H \rightarrow gg) &= \text{BR}_{\text{SM}}(H \rightarrow gg) \Gamma_{\text{tot}}^{\text{SM}}(H), \\ \Gamma_{\text{model}}(H \rightarrow gg) &= 9 \Gamma_{\text{SM}}(H \rightarrow gg), \\ \Gamma_{\text{tot}}^{\text{model}}(H) &= \Gamma_{\text{tot}}^{\text{SM}}(H) - \Gamma_{\text{SM}}(H \rightarrow gg) + \Gamma_{\text{model}}(H \rightarrow gg).\end{aligned}$$

It is then very simple to test the model with `HiggsBounds` using the effective coupling input.

In Fig. 4 we show a Fortran 77 code example which uses `HiggsBounds` for the calculation of the Higgs mass dependence of the ratio $Q_{\text{model}}/Q_{\text{obs}}$ for the SM and our simple Fourth Generation Model scenario. From the results returned by the subroutine `run_HiggsBounds_effC`, the set (`mh`, `HBresult`, `chan`, `obsratio`) is stored in a file, separately for the two models.

The results obtained by this sample program are presented in Fig. 5, which shows the ratio $(Q_{\text{model}}/Q_{\text{obs}})^{-1} = [\sigma \times \text{BR}]_{\text{obs. limit}}/[\sigma \times \text{BR}]_{\text{model}}$ as a function of the Higgs mass m_H for the SM (solid lines) and the Fourth Generation Model (dashed lines). For the SM cross section we use here the prediction employed by the Tevatron collaborations, see the discussion above concerning

⁸ The effective Higgs-photon-photon coupling is suppressed by a factor of about $\frac{1}{9} \Gamma_{\text{tot}}^{\text{model}}(H)/\Gamma_{\text{tot}}^{\text{SM}}(H)$ for a light Higgs H [62]. The exact suppression factor is irrelevant here, as the Higgs decay into $\gamma\gamma$ can only lead to relevant search topologies at the Tevatron if it is enhanced by more than an order of magnitude compared to the SM.

```

program SM_vs_fourth_generation_model
implicit none
integer nH,HBresult,chan,ncombined
real*8 obsratio,SMGammaTotal,GammaTotal,SMgamma_hgg,
&   Gamma_hgg,SMGamma_h,SMBR_Hgg,Mh,g2hjbb,g2hjtautau,
&   g2hjWW,g2hjZZ,g2hjgaga,g2hjgg,g2hjhiZ,BR_hjhihi
nH=1
call initialize_HiggsBounds(nH,'LandT')
open(10,file='example-SM-results.dat')
open(20,file='example-4thGen-results.dat')
do Mh=90,200,5
  SMGammaTotal=SMGamma_h(Mh)
  SMgamma_hgg=SMBR_Hgg(Mh)*SMGammaTotal
  Gamma_hgg=9d0*SMBR_Hgg(Mh)*SMGammaTotal
  GammaTotal=SMGammaTotal-SMgamma_hgg+Gamma_hgg
  g2hjbb=1d0
  g2hjtautau=1d0
  g2hjWW=1d0
  g2hjZZ=1d0
  g2hjgaga=1d0
  g2hjgg=1d0
  g2hjhiZ=0d0
  BR_hjhihi=0d0
  call run_HiggsBounds_effC(nH,Mh,SMGammaTotal,
&   g2hjbb,g2hjtautau,g2hjWW,g2hjZZ,g2hjgaga,g2hjgg,g2hjhiZ,
&   BR_hjhihi,HBresult,chan,obsratio,ncombined)
  write(10,*) Mh,HBresult,chan,obsratio
  g2hjgaga=1d0/9d0*GammaTotal/SMGammaTotal
  g2hjgg=9d0
  call run_HiggsBounds_effC(nH,Mh,GammaTotal,
&   g2hjbb,g2hjtautau,g2hjWW,g2hjZZ,g2hjgaga,g2hjgg,g2hjhiZ,
&   BR_hjhihi,HBresult,chan,obsratio,ncombined)
  write(20,*) Mh,HBresult,chan,obsratio
enddo
close(10)
close(20)
call finish_HiggsBounds
end

```

Fig. 4. Fortran77 code `example-SM_vs_4thGen.F` which calculates the ratio Q_{model}/Q_{obs} (`obsratio`) for the most sensitive Higgs search topology for the SM and a simple Fourth Generation Model as a function of the Higgs mass (M_h). This program has been used to produce the plot shown in Fig. 5.




Fig. 5. Ratio Q_{obs}/Q_{model} for the most sensitive channel as a function of the Higgs mass m_H : 4th Generation Model versus SM. The colours indicate whether the most sensitive search channel is from *LEP* (lighter grey) or the *Tevatron* (darker grey).

an update of this prediction. Exclusion at 95% C.L. is given if the value of $(Q_{model}/Q_{obs})^{-1}$ drops below 1. Both curves have sections in red colour (lighter grey), where the LEP results provide the most sensitive search topology, and in blue colour (darker grey), where the most sensitive search topology is provided by the Tevatron results. As expected, the LEP results have the highest sensitivity in the low-mass region up to the kinematical limit of about 115 GeV, while for higher mass values the Tevatron results take over.

The SM results show the familiar exclusion of Higgs masses below ≈ 115 GeV from LEP and the small excluded region at about $m_H = 170$ GeV from the latest Tevatron SM combined analysis [49] using the results of the slightly more conservative results obtained by the Bayesian method (see the discussion above for an update of the cross-section prediction employed here). For the Fourth Generation Model the LEP exclusion is a bit weaker than for the SM, because at low Higgs mass the larger branching ratio for Higgs decays into gluon pairs diminishes the $b\bar{b}$ and $\tau^+\tau^-$ branching ratios. In the region above 115 GeV, the most sensitive search topology for the Fourth Generation Model Higgs is $p\bar{p} \rightarrow H \rightarrow W^+W^- \rightarrow l^+l'^-$ throughout the displayed range. Because of the enhancement of the signal cross section due to the enhanced coupling of the Higgs boson to gluons, the region excluded by the Tevatron results extends from about 145 GeV to the end of the displayed range. Because of the SM-likeness criterion, the single topology analyses [39,40] apply, instead of the

SM combined analysis, for the Fourth Generation Model. Disregarding this criterion would change the $(Q_{\text{model}}/Q_{\text{obs}})^{-1}$ values in the Tevatron exclusion region to almost exactly 1/9 times the SM values.

4.3 MSSM

Theoretical background

The MSSM Higgs sector consists of two scalar weak isospin doublets and possesses therefore five physical Higgs bosons. At tree-level these are the \mathcal{CP} -even h and H (with $m_h < m_H$), the \mathcal{CP} -odd A and the charged H^\pm . The Higgs sector of the MSSM can be expressed at lowest order in terms of M_Z , M_A (or M_{H^\pm}) and $\tan\beta \equiv v_2/v_1$, the ratio of the vacuum expectation values of the two doublets. All other masses and mixing angles can therefore be predicted. The tree-level masses and couplings are strongly affected by higher-order corrections [63]. In the real MSSM (rMSSM), i.e. the model without \mathcal{CP} -violation, the higher-order corrected masses are denoted as M_h , M_H and M_A . In the MSSM with complex phases (cMSSM) the three neutral Higgs bosons can mix and the masses are denoted as M_{h_1} , M_{h_2} , M_{h_3} (with $M_{h_1} < M_{h_2} < M_{h_3}$).

In order to ensure that the external Higgs bosons have the correct on-shell properties, the tree-level Higgs states have to be supplemented with Higgs-propagator corrections (which are often the dominant corrections). This is done with the help of the \mathbf{Z} matrix [17,64], a non-unitary matrix, which mixes between the Higgs tree-level mass eigenstates (in the order h, H, A) and the loop-corrected mass eigenstates

$$\begin{pmatrix} h_1 \\ h_2 \\ h_3 \end{pmatrix} = \mathbf{Z} \cdot \begin{pmatrix} h \\ H \\ A \end{pmatrix}. \quad (59)$$

In the rMSSM, setting $h_1 = h$, $h_2 = H$, $h_3 = A$, one has

$$Z_{31} = Z_{32} = Z_{13} = Z_{23} = 0, \quad Z_{33} = 1. \quad (60)$$

In the following we give the formulas valid for the more general cMSSM, where the special case of the rMSSM can be derived with the help of Eq. (60). The normalised cross sections $\sigma_{h_i Z}^{\text{norm}}$, $\sigma_{h_j h_i}^{\text{norm}}$, defined through Eqs. (1), (18) and (20) as part of the input needed by `HiggsBounds`, can then be found using the elements of the matrix \mathbf{Z}

$$\sigma_{h_i Z}^{\text{norm}} = |Z_{i1} \sin(\beta - \alpha) + Z_{i2} \cos(\beta - \alpha)|^2$$

$$\sigma_{h_j h_i}^{\text{norm}} = |Z_{i3}(Z_{j1} \cos(\beta - \alpha) - Z_{j2} \sin(\beta - \alpha))$$
(61)

$$- Z_{j3}(Z_{i1} \cos(\beta - \alpha) - Z_{i2} \sin(\beta - \alpha))|^2,$$
(62)

where α is the angle that diagonalises the \mathcal{CP} -even Higgs sector at tree-level. It should be noted that for two identical Higgs bosons this leads to $\sigma_{h_i h_i}^{\text{norm}} = 0$ and that the following approximate relations hold:

$$\sigma_{h_1 Z}^{\text{norm}} + \sigma_{h_2 Z}^{\text{norm}} + \sigma_{h_3 Z}^{\text{norm}} \simeq 1,$$
(63)

$$\sigma_{h_k Z}^{\text{norm}} \simeq \sigma_{h_j h_i}^{\text{norm}},$$
(64)

where i, j, k are all different⁹.

Limits for the real MSSM

In Ref. [12] the final results for the LEP Higgs searches in the MSSM were published. In the case of the rMSSM they had been obtained in four benchmark scenarios: the m_h^{max} , no-mixing, small α_{eff} and gluophobic Higgs scenarios (including also some variations), see Ref. [65] for a motivation and detailed definition of these scenarios. The computer code used at that time was `FeynHiggs 2.0`, and in most cases the top-quark mass was fixed to $m_t = 174.3$ GeV. The limits had been obtained by a combination of all available channels.

Here we present the limits obtained by `HiggsBounds` using the latest version (2.6.4) of `FeynHiggs` to calculate the required model input and choosing different values for m_t . Note that the limits given here are in the narrow-width approximation, as discussed in Section 2.1.

In the left column of Fig. 6 we show the LEP and Tevatron exclusion bounds as obtained with `HiggsBounds` in the m_h^{max} (a), no-mixing (b), small α_{eff} (c) and gluophobic Higgs scenario (d) in the M_A - $\tan \beta$ plane for the current top-quark mass value, $m_t = 172.4$ GeV [66]. In the right column of Fig. 6 we show the most sensitive channels for each point in the M_A - $\tan \beta$ plane for the same four scenarios (e)–(h) (the colour code is given in the caption).

The four scenarios show similar features in the excluded regions, as presented in the left-hand column. Very low M_A values are excluded by LEP searches for all $\tan \beta$ values. This is due to the channel $e^+e^- \rightarrow hA \rightarrow b\bar{b}b\bar{b}$ as can be seen in the right column of Fig. 6. In this region the hZZ coupling is suppressed. Only at very low $\tan \beta$ values the channel $e^+e^- \rightarrow hZ, h \rightarrow b\bar{b}$ has

⁹ Note that these relations become equalities if a unitary Higgs mixing matrix is used. See, e.g., Ref. [17].




Fig. 6. Left column [(a),(b),(c),(d)]: 95% C.L. level excluded regions in the M_A - $\tan\beta$ plane for the the four rMSSM benchmark scenarios. The colours indicate exclusion from LEP (■) or the Tevatron (■). Right column [(e),(f),(g),(h)]: channel of highest statistical sensitivity: yellow (■): $e^+e^- \rightarrow hZ, h \rightarrow b\bar{b}$; cyan (■): $e^+e^- \rightarrow HZ, H \rightarrow b\bar{b}$; cyan (□): $e^+e^- \rightarrow hZ/HZ, h/H \rightarrow b\bar{b}$; orange (■/□): $e^+e^- \rightarrow hA \rightarrow b\bar{b}b\bar{b}/b\bar{b}\tau^+\tau^-$; green (■): $p\bar{p} \rightarrow h/A \rightarrow \tau^+\tau^-$ [25]; blue (■): $p\bar{p} \rightarrow h/H/A \rightarrow \tau^+\tau^-$ [25]; magenta (■): $p\bar{p} \rightarrow H/A \rightarrow \tau^+\tau^-$ [25]; magenta (□): $p\bar{p} \rightarrow A \rightarrow \tau^+\tau^-$ [25]; red (■): $p\bar{p} \rightarrow hW \rightarrow b\bar{b}l\nu$ [36]; black (■): $p\bar{p} \rightarrow H/A \rightarrow \tau^+\tau^-$ [24]; blue (□): $p\bar{p} \rightarrow H \rightarrow W^+W^-$ [39]. The contour borders the excluded regions.

the highest statistical sensitivity, but still with a somewhat suppressed hZZ coupling. At around $M_A \approx 100$ GeV the kinematical limit for the hA channel is reached and other channels take over. There the exclusion drops to a relatively low excluded $\tan\beta$ value, slightly different in the four scenarios. At around $M_A \approx 115$ GeV the exclusion in the $e^+e^- \rightarrow hZ, h \rightarrow b\bar{b}$ channel becomes SM-like, and Higgs bosons up to the kinematical limit can be excluded by LEP searches, reaching $\tan\beta = 8, 30, 10, 50$ in the m_h^{\max} , no-mixing, small α_{eff} and gluophobic Higgs scenario, respectively. In the latter one also larger values could be excluded, but the Tevatron channels become more important, see below. For larger M_A values the LEP exclusion drops to a lower $\tan\beta$ value between 3 and 11, depending on the scenario. These $\tan\beta$ values correspond to $M_h \approx 208$ GeV $- M_Z$, i.e. the kinematical limit and are excluded for all M_A values above ~ 115 GeV.

The Tevatron searches exclude a parameter space, depending on the scenario, between $M_A = 110$ GeV and 170 GeV, and between $\tan\beta = 39$ and 60 (where we stopped our scan, but in principle larger values can be excluded). The relevant channels are $p\bar{p} \rightarrow h/H/A + X, h/H/A \rightarrow \tau^+\tau^-$, see the right-hand column of Fig. 6. The variation in the colour here only indicates which Higgs bosons contribute together to the signal. At low M_A (green) one has $M_h \approx M_A$, at intermediate M_A (dark blue) we find $M_h \approx M_A \approx M_H$, and at higher M_A values one generically has $M_A \approx M_H$. Consequently, two or three neutral Higgs bosons can contribute to the signal. Two other Tevatron channels have the largest exclusion potential in some part of the M_A - $\tan\beta$ parameter space. Within the m_h^{\max} scenario $p\bar{p} \rightarrow hW \rightarrow b\bar{b}l\nu$ is the most sensitive channel for $M_A \gtrsim 130$ GeV and intermediate $\tan\beta$, see plot (e) in Fig. 6. In the small α_{eff} scenario $p\bar{p} \rightarrow H \rightarrow W^+W^-$ with the W s decaying leptonically is the most sensitive channel for low M_A and intermediate $\tan\beta$, see plot (g) of Fig. 6.

As mentioned before, the original exclusion bounds in Ref. [12] had been obtained for $m_t = 174.3$ GeV. The most prominent effect of increasing the top-quark mass from $m_t = 172.4$ GeV to $m_t = 174.3$ GeV is an increase in M_h with $\Delta M_h / \Delta m_t \lesssim 1$ [67]. This reduces the reach in the M_A - $\tan\beta$ plane at LEP for the $e^+e^- \rightarrow hZ$ channels. This effect can be observed to some extent in all four scenarios. In Fig. 7 we show the exclusion bounds as obtained with `HiggsBounds` with $m_t = 174.3$ GeV in the m_h^{\max} scenario in the M_h - $\tan\beta$ plane. This plot can be compared with Fig. 9(b) in the original LEP publication [12], which, however, terminated at $\tan\beta = 40$, whereas the plot shown here extends up to $\tan\beta = 60$. The values of M_h reached in Fig. 7 are smaller by ~ 2 GeV as compared to the original LEP figure, which is due to the newer version of `FeynHiggs` employed here. Apart from this, the LEP exclusion areas are almost identical in the two figures. This shows how well in this case our procedure (based on the LEP data for single topological cross sections) reproduces the full LEP analysis where different search channels were combined. In addition, we also find a region excluded by the Tevatron Higgs searches at




Fig. 7. Parameter region in the M_h - $\tan \beta$ plane for the m_h^{\max} scenario of the rMSSM which is excluded at the 95% C.L. The mass of the top quark has been set to 174.3 GeV. The colours indicate whether the most sensitive search channel which leads to exclusion is from **LEP** (lighter grey) or the **Tevatron** (darker grey). The solid line shows the theoretical upper bound on M_h in the m_h^{\max} scenario.

$\tan \beta \gtrsim 40$ and $M_h \gtrsim 100$ GeV. These results from the Tevatron were not yet available when Ref. [12] was published.

LEP limits for the complex MSSM in the CPX scenario

Ref. [12] also published the results of the LEP Higgs searches in the CPX scenario, which is a \mathcal{CP} -violating MSSM benchmark scenario [68]. This analysis incorporated Higgs sector predictions from the programs CPH [68,69] and FeynHiggs 2.0 [14,15,16,17].

More recently, in Ref. [70], a preliminary version of HiggsBounds was used in conjunction with full 1-loop vertex corrections for the $h_j \rightarrow h_i h_i$ and $h_j \rightarrow f \bar{f}$ decays to examine the effect of these corrections on the LEP exclusions in the CPX scenario. In Ref. [70], Higgs masses and the Higgs propagator corrections (as contained in the \mathbf{Z} matrix) were calculated using renormalised Higgs self-energies from FeynHiggs 2.6.4.




Fig. 8. Coverage of the LEP Higgs searches in the M_{H_1} - $\tan\beta$ plane of the CPX scenario. Plot (a) shows the channels that are predicted to have the highest statistical sensitivity for setting an exclusion limit. The colour codings are: red (■) = $(h_1 Z) \rightarrow (b\bar{b}Z)$, blue (■) = $(h_2 Z) \rightarrow (b\bar{b}Z)$, white (□) = $(h_2 Z) \rightarrow (h_1 h_1 Z) \rightarrow (b\bar{b}b\bar{b}Z)$, cyan (■) = $(h_2 h_1) \rightarrow (b\bar{b}b\bar{b})$, yellow (■) = $(h_2 h_1) \rightarrow (h_1 h_1 h_1) \rightarrow (b\bar{b}b\bar{b}b\bar{b})$, purple (■) = other channels. Plot (b) shows the parameter regions excluded at the 95% C.L. by the topological cross section limits obtained at LEP [11,12]. The colour codings are: green (■) = excluded, white (□) = unexcluded. In both plots, the grey (■) peripheral areas are theoretically inaccessible using the input parameters associated with the CPX scenario.

The results shown in Fig. 8 have been calculated using the conventions¹⁰ and method described in Ref. [70]. However, the results shown here include two significant updates: we use $m_t = 172.6$ GeV rather than $m_t = 170.9$ GeV and we use a newer version of `FeynHiggs` to calculate the renormalised Higgs self-energies, which contains the correct combination of the $\tan\beta$ -enhanced contributions to the Higgs-boson self-energies in the complex MSSM with the one-loop corrections for all options concerning the two-loop contributions.

In Fig. 8 (a), we show the channels with the highest statistical sensitivity for each point in CPX parameter space. The process $(h_1 Z) \rightarrow (b\bar{b}Z)$ (red)¹¹ dominates in regions where the $h_1 Z Z$ effective coupling is significant. Adjacent to this region is an area of parameter space with a relatively large $h_2 Z Z$ effective coupling and therefore the processes $(h_2 Z) \rightarrow (b\bar{b}Z)$ (blue) and $(h_2 Z) \rightarrow (h_1 h_1 Z) \rightarrow (b\bar{b}b\bar{b}Z)$ (white) have the highest statistical sensitivity in this region. The processes $(h_2 h_1) \rightarrow (b\bar{b}b\bar{b})$ (cyan) and $(h_2 h_1) \rightarrow (h_1 h_1 h_1) \rightarrow (b\bar{b}b\bar{b}b\bar{b})$

¹⁰Note that Ref. [12] and Ref. [70] use slightly different definitions of the CPX scenario (see Ref. [70] for a full description of these differences).

¹¹All processes discussed in this subsection are LEP processes. Therefore, for brevity, we do not include the initial state explicitly in the process descriptions.

(yellow) dominate in the region where the h_3ZZ effective coupling is relatively large (and therefore also the h_1h_2Z effective coupling, as predicted by Eq. (64)).

In the region of Fig. 8 (a) where $M_{h_1} \lesssim 45$ GeV, $\tan\beta \gtrsim 4$, the total decay width of h_2 is dominated by the partial decay width $h_2 \rightarrow h_1h_1$. This has a very significant effect on the exclusion prospects in this region. As Fig. 8 (a) shows, this region is partly covered by the $(h_2Z) \rightarrow (h_1h_1Z) \rightarrow (b\bar{b}b\bar{b}Z)$ (white) and $(h_2h_1) \rightarrow (h_1h_1h_1) \rightarrow (b\bar{b}b\bar{b}b\bar{b})$ (yellow) processes, which directly involve the $h_2 \rightarrow h_1h_1$ decay. However, searching for these topologies is very challenging experimentally as, for example, the $(h_2h_1) \rightarrow (h_1h_1h_1) \rightarrow (b\bar{b}b\bar{b}b\bar{b})$ process gives rise to complex final states involving six jets. Additionally, processes directly involving the $h_2 \rightarrow b\bar{b}$ decay (such as $(h_2Z) \rightarrow (b\bar{b}Z)$ (blue)) are more difficult to detect in this region because of the suppression of the $h_2 \rightarrow b\bar{b}$ branching ratio due to the large $h_2 \rightarrow h_1h_1$ partial decay width.

Fig. 8 (b) shows the areas of CPX parameter space which have been excluded by the LEP Higgs searches at the 95 % C.L. (green). There are two sizable unexcluded regions¹² (white), at $M_{h_1} \gtrsim 50$ GeV, $\tan\beta \gtrsim 8$ and a “hole” at $M_{h_1} \approx 45$ GeV, $\tan\beta \approx 8$. Both of these regions occur in parts of parameter space where processes involving the topology $(h_iZ) \rightarrow (b\bar{b}Z)$ predominantly have the highest statistical sensitivity, and the second of these regions occurs in the part of the CPX parameter space which, as discussed above, was expected to prove challenging experimentally due to a suppressed $h_2 \rightarrow b\bar{b}$ branching ratio. In addition, the two unexcluded regions are very sensitive to the slight excess in the LEP data for this topology at 89.6 GeV $\lesssim M_{h_i} \lesssim 107$ GeV, where the observed limit is more than 1 sigma above the predicted limit based on simulations with no signal, see Ref. [12].

Comparison of Fig. 8 (a) with Fig. 8 (b) also shows that there are some unexcluded parameter points along boundaries between areas where different processes have the highest statistical sensitivity, for example, at $M_{h_1} \approx 100$ GeV, $\tan\beta \approx 5$. Since in our analysis only the single channel having the highest statistical sensitivity can be used to set the 95% C.L. limit, the statistical information from all other decay and production channels in this region does not contribute. In these narrow regions, an analysis which can utilise more than one experimentally measured cross section limit for each parameter point (such as that performed in Ref. [12]) will have a higher exclusion power, and thus possibly be able to reduce the size of the unexcluded region.

¹²Note that we do not consider unexcluded regions to be significant if they only occur in a very small Higgs mass range and are situated on the borders of regions where different channels have the highest statistical sensitivity. As discussed in Section 2.1, we expect HiggsBounds to be less sensitive at these borders than a dedicated experimental analysis.

Using `HiggsBounds`, it has therefore been possible to investigate the effects of recent improvements in the theoretical Higgs sector predictions on the LEP exclusions in the CPX scenario. In particular, since it provides information about the process with the highest statistical sensitivity, `HiggsBounds` should be helpful for the user to obtain some guidance on which effects are important for the exclusion or non-exclusion of certain parts of parameter space. `HiggsBounds` also provides a convenient way to investigate the dependence of the calculation on the model input parameters, such as the effect of varying the top-quark mass.

5 Summary

We have presented the code `HiggsBounds`, which is a tool to test whether the predictions arising from the Higgs sector of arbitrary types of theoretical models are in accordance with the existing exclusion bounds from the Higgs searches at LEP and the Tevatron. To this end, the experimental bounds on topological cross sections made available by the LEP and Tevatron collaborations have been incorporated as look-up tables into the program. In order to utilise this information to set exclusion bounds on the parameter space of a certain model one first needs to determine the search channel with the highest statistical sensitivity for setting an exclusion limit. This is done by comparing the expected limits (which have also been provided by the experimental collaborations), i.e. the exclusion bounds that one would obtain in the hypothetical case of an observed distribution with a pure background shape, with the model predictions. In order to ensure the correct statistical interpretation of the obtained exclusion bound as a 95% C.L., the analysis has to be restricted to the single channel that possesses the highest statistical sensitivity for setting an exclusion limit. For this channel the program determines whether or not a certain parameter point is excluded at the 95% C.L.

The experimental information incorporated into `HiggsBounds` consists primarily of limits on single topological cross sections, either from a single collaboration or combinations of the results from different collaborations. Some experimental combinations have been done by the collaborations only for the special case of a SM-type Higgs boson. These have the advantage of an enhanced statistical power. In order to determine whether experimental results on the searches for the SM Higgs can be used for constraining a certain parameter region of a new physics scenario, the “SM likeness” of the Higgs boson(s) of the model under study is evaluated. Most experimental results that are currently available have been obtained under the assumption that a narrow width approximation is valid for the Higgs boson under study. If this assumption is not valid the experimental exclusion bounds are modified. The current version of `HiggsBounds` is based on exclusion limits obtained in the narrow-

width approximation. The proper treatment of experimental results, taking into account a non-negligible Higgs-boson width, will be incorporated in a forthcoming version of `HiggsBounds`. The exclusion bounds included in the `HiggsBounds` package will be updated as new results from the Higgs searches become available.

The predictions of the desired model, needed as input for `HiggsBounds`, have to be provided by the user. This can be done at various levels of sophistication, ranging from effective couplings to complete information on cross sections and branching ratios. Links to the widely used programs `FeynHiggs` and `CPsuperH` for Higgs-sector predictions in the MSSM are provided by default.

We have discussed various examples for running `HiggsBounds` and have given sample files for its input and output. In particular, we have considered the cases of the Standard Model, a model with a fourth generation of quarks and leptons and for the MSSM with and without \mathcal{CP} -violation.

The `HiggsBounds` code exists both in a Fortran 77 and Fortran 90 version. It can be operated in a command line mode, as a subroutine that can easily be linked to other codes, and as an online version, see www.ippp.dur.ac.uk/HiggsBounds.

Acknowledgements

We thank the LEP collaborations ALEPH, DELPHI, L3 and OPAL and the Tevatron collaborations CDF and DØ for providing us with detailed information about their Higgs search results. In particular, we are grateful to Alexander Read and Peter Igo-Kemenes (LEP Higgs Working Group), Mark Owen (DØ), Tom Junk and Matthew Herndon (CDF) for very valuable discussions on their experiment's results. We thank Terrance Figy for providing us with SM predictions for the Higgs production cross section via VBF at the Tevatron separated in W - and Z -fusion contribution. We would like to thank Philip Roffe and David Ambrose-Griffith for valuable computing support, and the IPPP (Durham, UK) for hosting the web version of `HiggsBounds`. Thanks also go to Lydia Heck, Sophy Palmer and Alison Fowler for their assistance in testing the code and to Valery Khoze for interesting discussions. KW was supported by the Helmholtz Alliance HA-101 'Physics at the Terascale'. This work has been supported in part by the European Community's Marie-Curie Research Training Network under contract MRTN-CT-2006-035505 'Tools and Precision Calculations for Physics Discoveries at Colliders' (HEPTOOLS) and MRTN-CT-2006-035657 'Understanding the Electroweak Symmetry Breaking and the Origin of Mass using the First Data of ATLAS' (ARTEMIS), by the Helmholtz Young Investigator Grant VH-NG-303, and by the DFG Collabo-

rative Research Grant SFB 676 'Particles, Strings and the Early Universe'.

A Determination of experimental S_{95} values

A.1 The CL_s method

In order to calculate the experimental S_{95} values, the modified frequentist method (or CL_s method) is used. As a first step, using the experimental data binned in i bins with d_i observed data events, s_i expected signal events for a given hypothesis of the signal, and b_i background events, a test statistics

$$Q = \prod_{i=1}^j \frac{e^{-(s_i+b_i)}(s_i + b_i)^{d_i}}{d_i!} \bigg/ \frac{e^{-(b_i)}(b_i)^{d_i}}{d_i!}$$

is defined, which orders the expected outcomes of test experiments according to their ‘‘signal-likeness’’. Then, confidence levels for the given hypothesis are calculated. The confidence level for the background hypothesis, CL_b , is defined as the probability to obtain values of a test statistics Q no larger than the observed value Q_{obs} , given a large number of hypothetical experiments with background processes only,

$$CL_b = P(Q \leq Q_{\text{obs}} | \text{background}).$$

Similarly, the confidence level for the signal+background hypothesis, CL_{s+b} , is defined as the probability to obtain values of Q smaller than observed, given a large number of hypothetical experiments with signal and background processes,

$$CL_{s+b} = P(Q \leq Q_{\text{obs}} | \text{signal} + \text{background}).$$

In principle, CL_{s+b} can be used to exclude the signal+background hypothesis, given a model for Higgs boson production. However, this procedure may lead to the undesired possibility that a downward fluctuation of the background would allow a hypotheses to be excluded for which the experiment has no sensitivity due to the small expected signal rate. This problem is avoided by introducing the ratio

$$CL_s = CL_{s+b}/CL_b.$$

Since CL_b is a positive number less than one, CL_s will always be greater than CL_{s+b} and the limit obtained in this way will therefore be conservative. This quantity is adopted for setting exclusion limits and a hypothesis is considered to be excluded at the 95 % confidence level if the corresponding value of CL_s is less than 0.05. This method is also called ‘‘modified frequentist approach’’, since not the frequentistically determined confidence levels CL_b and CL_{s+b} but their ratio is used.

A.2 Expected and observed confidence levels

The expected confidence levels are obtained by replacing the observed data configuration by a large number of simulated event configurations for the two hypotheses background only or signal+background. These can be used to estimate the expected sensitivity of a search and to compare the observed exclusion with the one expected with no signal present. The expected CL_s is then the median of the outcome of the large number of simulated experiments with background only event configuration.

Once the experimentally observed and the expected values of CL_s are derived as described above using the hypothesis of a Higgs production in the channel under study with SM strength (i.e. all BR and cross section scaling factors set to 1), the observed and expected values of S_{95} can be numerically determined by varying the production rate of the channel by a scaling factor s until

$$CL_s = 0.05$$

is reached. The scaling factor s is then interpreted as observed or expected S_{95} , depending of whether the observed or expected confidence level CL_s has been used.

The numerical calculation at LEP has been done with tools described in [71] and [72], while the Tevatron results use different methods described in the individual references.

References

- [1] P. W. Higgs, *Phys. Lett.* **12** (1964) 132; *Phys. Rev. Lett.* **13** (1964) 508; *Phys. Rev.* **145** (1966) 1156;
F. Englert and R. Brout, *Phys. Rev. Lett.* **13** (1964) 321;
G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, *Phys. Rev. Lett.* **13** (1964) 585.
- [2] S.L. Glashow, *Nucl. Phys.* **B 22** (1961) 579;
S. Weinberg, *Phys. Rev. Lett.* **19** (1967) 19;
A. Salam, in: *Proceedings of the 8th Nobel Symposium*, Editor N. Svartholm, Stockholm, 1968.
- [3] H. Nilles, *Phys. Rept.* **110** (1984) 1;
H. Haber and G. Kane, *Phys. Rept.* **117** (1985) 75;
R. Barbieri, *Riv. Nuovo Cim.* **11** (1988) 1.
- [4] S. Weinberg, *Phys. Rev. Lett.* **37** (1976) 657;
J. Gunion, H. Haber, G. Kane and S. Dawson, *The Higgs Hunter's Guide* (Perseus Publishing, Cambridge, MA, 1990), and references therein.
- [5] P. Fayet, *Nucl. Phys.* **B 90** (1975) 104; *Phys. Lett.* **B 64** (1976) 159; *Phys. Lett.* **B 69** (1977) 489; *Phys. Lett.* **B 84** (1979) 416;
H.P. Nilles, M. Srednicki and D. Wyler, *Phys. Lett.* **B 120** (1983) 346;
J.M. Frere, D.R. Jones and S. Raby, *Nucl. Phys.* **B 222** (1983) 11;
J.P. Derendinger and C.A. Savoy, *Nucl. Phys.* **B 237** (1984) 307;
J. Ellis, J. Gunion, H. Haber, L. Roszkowski and F. Zwirner, *Phys. Rev.* **D 39** (1989) 844;
M. Drees, *Int. J. Mod. Phys.* **A 4** (1989) 3635;
U. Ellwanger, M. Rausch de Traubenberg and C.A. Savoy, *Phys. Lett.* **B 315** (1993) 331 [arXiv:hep-ph/9307322]; *Z. Phys.* **C 67** (1995) 665 [arXiv:hep-ph/9502206]; *Nucl. Phys.* **B 492** (1997) 307 [arXiv:hep-ph/9611251];
T. Elliott, S.F. King and P. White, *Phys. Lett.* **B 351** (1995) 213 [arXiv:hep-ph/9406303];
S.F. King and P. White, *Phys. Rev.* **D 52** (1995) 4183 [arXiv:hep-ph/9505326];
C. Panagiotakopoulos and K. Tamvakis, *Phys. Lett.* **B 469** (1999) 145 [arXiv:hep-ph/9908351];
C. Panagiotakopoulos and A. Pilaftsis, *Phys. Rev.* **D 63** (2001) 055003 [arXiv:hep-ph/0008268];
A. Dedes, C. Hugonie, S. Moretti and K. Tamvakis, *Phys. Rev.* **D 63** (2001) 055009 [arXiv:hep-ph/0009125].
- [6] N. Arkani-Hamed, A. Cohen and H. Georgi, *Phys. Lett.* **B 513** (2001) 232 [arXiv:hep-ph/0105239];
N. Arkani-Hamed, A. Cohen, T. Gregoire and J. Wacker, *JHEP* **0208** (2002) 020 [arXiv:hep-ph/0202089].
- [7] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett.* **B 429** (1998) 263 [arXiv:hep-ph/9803315]; *Phys. Lett.* **B 436** (1998) 257 [arXiv:hep-ph/9804398];

- I. Antoniadis, *Phys. Lett.* **B 246** (1990) 377;
 J. Lykken, *Phys. Rev.* **D 54** (1996) 3693 [arXiv:hep-th/9603133];
 I. Antoniadis and M. Quiros, *Phys. Lett.* **B 392** (1997) 61
 [arXiv:hep-th/9609209];
 L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83** (1999) 3370
 [arXiv:hep-ph/9905221];
 D. Cremades, L. E. Ibanez and F. Marchesano, *Nucl. Phys.* **B 643** (2002) 93
 [arXiv:hep-th/0205074];
 C. Kokorelis, *Nucl. Phys.* **B 677** (2004) 115 [arXiv:hep-th/0207234].
- [8] T. Binoth and J. J. van der Bij, *Z. Phys.* **C 75** (1997) 17
 [arXiv:hep-ph/9608245];
 D. G. Cerdeno, A. Dedes and T. E. J. Underwood, *JHEP* **0609** (2006) 067
 [arXiv:hep-ph/0607157];
 D. O’Connell, M. J. Ramsey-Musolf and M. B. Wise, *Phys. Rev.* **D 75** (2007)
 037701 [arXiv:hep-ph/0611014].
- [9] R. A. Porto and A. Zee, arXiv:0712.0448 [hep-ph].
- [10] G. F. Giudice and O. Lebedev, *Phys. Lett.* **B 665** (2008) 79 [arXiv:0804.1753
 [hep-ph]].
- [11] R. Barate et al., *Phys. Lett.* **B 565** (2003) 61 [hep-ex/0306033].
- [12] S. Schael et al., *Eur. Phys. J.* **C 47** (2006) 547 [hep-ex/0602042].
- [13] [LEP Higgs Working Group for Higgs boson searches] arXiv: hep-ex/0107031;
 arXiv: hep-ex/0107032; arXiv: hep-ex/0107033; arXiv: hep-ex/0107034; arXiv:
 hep-ex/0107035.
- [14] S. Heinemeyer, W. Hollik and G. Weiglein, *Comput. Phys. Commun.* **124** (2000)
 76, [arXiv:hep-ph/9812320]; see: www.feynhiggs.de .
- [15] S. Heinemeyer, W. Hollik and G. Weiglein, *Eur. Phys. J.* **C 9** (1999) 343
 [arXiv:hep-ph/9812472].
- [16] G. Degrossi, S. Heinemeyer, W. Hollik, P. Slavich and G. Weiglein, *Eur. Phys.*
J. **C 28** (2003) 133 [arXiv:hep-ph/0212020].
- [17] M. Frank, T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak and G. Weiglein,
JHEP **0702** (2007) 047 [arXiv:hep-ph/0611326].
- [18] J. S. Lee, A. Pilaftsis, M. S. Carena, S. Y. Choi, M. Drees, J. R. Ellis
 and C. E. M. Wagner, *Comput. Phys. Commun.* **156** (2004) 283
 [arXiv:hep-ph/0307377];
 J. S. Lee, M. Carena, J. Ellis, A. Pilaftsis and C. E. M. Wagner, arXiv:0712.2360
 [hep-ph].
- [19] The ALEPH Collaboration, The DELPHI Collaboration, The L3 Collaboration,
 The OPAL Collaboration, The SLD Collaboration, The LEP Electroweak
 Working Group and The SLD Electroweak and Heavy Flavour Groups, *Phys.*
Rept. **427** (2006) 257. [arXiv:hep-ex/0509008].

- [20] J. Alcaraz et al. [ALEPH Collaboration and DELPHI Collaboration and L3 Collaboration and], arXiv:hep-ex/0612034.
- [21] The CDF Collaboration, CDF note 8954.
- [22] The CDF Collaboration, CDF note 9284.
- [23] The DØ Collaboration, Report FERMILAB-PUB-08-142-E, arXiv:0805.3556, submitted to Phys. Rev. Lett.
- [24] The DØ Collaboration, *Phys. Rev. Lett.* **101** (2008) 071804, arXiv:0805.2491 [hep-ex].
- [25] The CDF Collaboration, CDF note 9071.
- [26] A. Djouadi, J. Kalinowski and M. Spira, *Comput. Phys. Commun.* **108** (1998) 56 [arXiv:hep-ph/9704448].
- [27] T. Hahn, S. Heinemeyer, F. Maltoni, G. Weiglein and S. Willenbrock, arXiv:hep-ph/0607308;
U. Aglietti et al., arXiv:hep-ph/0612172;
See also: <http://maltoni.web.cern.ch/maltoni/TeV4LHC/>.
- [28] O. Brein and W. Hollik, *Phys. Rev. D* **68** (2003) 095006 [arXiv:hep-ph/0305321]; *Phys. Rev. D* **76** (2007) 035002 [arXiv:0705.2744 [hep-ph]].
- [29] T. Figy, C. Oleari and D. Zeppenfeld, *Phys. Rev. D* **68** (2003) 073005 [arXiv:hep-ph/0306109].
- [30] Alexander L. Read, private communication.
- [31] LEP Higgs Working Group, unpublished.
- [32] The CDF Collaboration, arXiv:0807.4493 [hep-ex].
- [33] The CDF Collaboration, CDF note 9475.
- [34] The DØ Collaboration, DØ note 5570.
- [35] The DØ Collaboration, DØ note 5472.
- [36] The CDF Collaboration, CDF note 9463.
- [37] The DØ Collaboration, DØ note 5485.
- [38] The CDF Collaboration, CDF note 7307.
- [39] The DØ Collaboration, DØ note 5757.
- [40] T. Aaltonen et al. [CDF Collaboration], arXiv:0809.3930 [hep-ex].
- [41] The DØ Collaboration, *Phys. Rev. Lett.* **101** (2008) 051801 [arXiv:0803.1514 [hep-ex]].
- [42] The DØ Collaboration, DØ note 5737.

- [43] The DØ Collaboration, DØ note 5726.
- [44] The CDF Collaboration, CDF note 9483.
- [45] The DØ Collaboration, DØ note 5586.
- [46] The CDF Collaboration, CDF note 9248.
- [47] TEVNPB Working Group Collaboration for the CDF Collaboration and D0 Collaboration, Report FERMILAB-PUB-07-656-E, CDF note 8961, DØ note 5536, arXiv:0712.2383 [hep-ex].
- [48] TEVNPB Working Group Collaboration for the CDF Collaboration and D0 Collaboration, Report FERMILAB-PUB-08-069-E, CDF note 9290, DØ note 5645, arXiv:0804.3423 [hep-ex].
- [49] TEVNPB Working Group Collaboration for the CDF Collaboration and D0 Collaboration, Report FERMILAB-PUB-08-270-E, CDF note 9465, DØ note 5754, arXiv:0808.0534 [hep-ex].
- [50] D. Graudenz, M. Spira and P. Zerwas, *Phys. Rev. Lett.* **70** (1993) 1372;
M. Spira, A. Djouadi, D. Graudenz and P. Zerwas, *Nucl. Phys. B* **453** (1995) 17 [arXiv:hep-ph/9504378];
R. Harlander and W. Kilgore, *Phys. Rev. Lett.* **88** (2002) 201801 [arXiv:hep-ph/0201206];
V. Ravindran, J. Smith and W. van Neerven, *Nucl. Phys. B* **665** (2003) 325 [arXiv:hep-ph/0302135];
C. Anastasiou and K. Melnikov, *Nucl. Phys. B* **646** (2002) 220 [arXiv:hep-ph/0207004];
S. Moch and A. Vogt, *Phys. Lett. B* **631** (2005) 48 [arXiv:hep-ph/0508265];
E. Laenen and L. Magnea, *Phys. Lett. B* **632** (2006) 270 [arXiv:hep-ph/0508284].
- [51] S. Catani, D. de Florian, M. Grazzini and P. Nason, *JHEP* **0307** (2003) 028 [arXiv:hep-ph/0306211];
- [52] R. V. Harlander and W. B. Kilgore, *Phys. Rev. D* **68** (2003) 013001 [arXiv:hep-ph/0304035].
- [53] A. Djouadi, *Phys. Rept.* **457** (2008) 1 [arXiv:hep-ph/0503172].
- [54] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, arXiv:0901.0002 [hep-ph].
- [55] U. Langenegger, M. Spira, A. Starodumov and P. Trueb, *JHEP* **0606** (2006) 035 [arXiv:hep-ph/0604156].
- [56] O. Brein, A. Djouadi and R. Harlander, *Phys. Lett. B* **579** (2004) 149 [arXiv:hep-ph/0307206];
M. L. Ciccolini, S. Dittmaier and M. Kramer, *Phys. Rev. D* **68** (2003) 073003 [arXiv:hep-ph/0306234];
K. A. Assamagan et al. [Higgs Working Group Collaboration], arXiv:hep-ph/0406152.

- [57] T. Han, G. Valencia and S. Willenbrock, *Phys. Rev. Lett.* **69** (1992) 3274 [arXiv:hep-ph/9206246];
 J. M. Campbell and R. K. Ellis, *Phys. Rev. D* **60** (1999) 113006 [arXiv:hep-ph/9905386];
 T. Figy, C. Oleari and D. Zeppenfeld, *Phys. Rev. D* **68** (2003) 073005 [arXiv:hep-ph/0306109];
 E. Berger and J. Campbell, *Phys. Rev. D* **70** (2004) 073011 [arXiv:hep-ph/0403194].
- [58] C. Anastasiou, R. Boughezal and F. Petriello, arXiv:0811.3458 [hep-ph].
- [59] U. Aglietti, R. Bonciani, G. Degrossi and A. Vicini, *Phys. Lett. B* **595** (2004) 432 [arXiv:hep-ph/0404071], arXiv:hep-ph/0610033;
 S. Actis, G. Passarino, C. Sturm and S. Uccirati, *Phys. Lett. B* **670** (2008) 12 [arXiv:0809.1301 [hep-ph]];
 C. Anastasiou, S. Beerli, S. Bucherer, A. Daleo and Z. Kunszt, *JHEP* **0701** (2007) 082 [arXiv:hep-ph/0611236].
- [60] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, *Phys. Lett.* **652** (2007) 292 [arXiv:0706.0459 [hep-ph]].
- [61] P. H. Frampton, P. Q. Hung and M. Sher, *Phys. Rept.* **330** (2000) 263 [arXiv:hep-ph/9903387].
- [62] G. D. Kribs, T. Plehn, M. Spannowsky and T. M. P. Tait, *Phys. Rev. D* **76** (2007) 075016 [arXiv:0706.3718 [hep-ph]].
- [63] S. Heinemeyer, *Int. J. Mod. Phys. A* **21** 2659 (2006) [arXiv:hep-ph/0407244];
 A. Djouadi, *Phys. Rept.* **459** (2008) 1. arXiv:hep-ph/0503173.
- [64] T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak, G. Weiglein and K. Williams, *Pramana* **69** (2007) 861 [arXiv:hep-ph/0611373].
- [65] M. Carena, S. Heinemeyer, C. Wagner and G. Weiglein, *Eur. Phys. J. C* **26** (2003) 601 [arXiv:hep-ph/0202167].
- [66] Tevatron Electroweak Working Group and CDF Collaboration and D0 Collaboration, arXiv:0808.1089 [hep-ex].
- [67] S. Heinemeyer, W. Hollik and G. Weiglein, *JHEP* **0006** (2000) 009 [arXiv:hep-ph/9909540].
- [68] M. S. Carena, J. R. Ellis, A. Pilaftsis and C. E. M. Wagner, *Phys. Lett. B* **495** (2000) 155 [arXiv:hep-ph/0009212].
- [69] M. S. Carena, J. R. Ellis, A. Pilaftsis and C. E. M. Wagner, *Nucl. Phys. B* **586** (2000) 92 [arXiv:hep-ph/0003180].
- [70] K. E. Williams and G. Weiglein, *Phys. Lett. B* **660** (2008) 217 [arXiv:0710.5320 [hep-ph]].
- [71] P. Bock, *JHEP* **0701** (2007) 080 [arXiv:hep-ex/0405072].
- [72] T. Junk, *Nucl. Instrum. Meth. A* **434** (1999) 435 [arXiv:hep-ex/9902006].