

Editorial Manager(tm) for Geo-Marine Letters
Manuscript Draft

Manuscript Number:

Title: Along-slope oceanographic processes and sedimentary products around the Iberian margin

Article Type: SI Baiona

Corresponding Author: Francisco Javier J. Hernández Molina, SI Baiona

Corresponding Author's Institution: FACULTAD DE CIENCIAS DEL MAR

First Author: Francisco Javier J. Hernández Molina, SI Baiona

Order of Authors: Francisco Javier J. Hernández Molina, SI Baiona;Nuno Serra;Dorrik A.V. Stow;Estefanía Llave;Gemma Ercilla;David Van Rooij

Abstract: The present contribution represents the first attempt to comprehensively describe regional along-slope processes and their sedimentary impacts around the Iberian margin, combining numerically simulated bottom currents with existing knowledge of contourite depositional and erosive features. The obtained correlation links the circulation of water masses with the main contourite depositional systems (CDS) and estimates other potential areas where new CDS could be found. Their study should be of great interest not only because of the stratigraphic, sedimentological, palaeoceanographic and palaeoclimatological significance, but also because of their relation with possible specific deep marine geohabitats and/or mineral and energy resources.

Suggested Reviewers: Angelo Camerlenghi

ICREA, Istitució Catalana de Recerca i Estudis Avançats c/o GRC Geociències Mari, Departament d'Estratigrafia , Paleontologia i Geociències Marines, Universitat de Barcelona / Facultat de Geologia C/ Martí i Franquès, s/n, E-08028 Barcelona, Spain
acamerlenghi@ub.edu

Specialist on contourite processes

Miguel Bruno Mejías

Profesor Titular del Área de Física Aplicada, Dpto. Física Aplicada, University of Cadiz. CASEM, Avda República Saharaui s/n. Puerto Real, 11510. Spain
miguel.bruno@uca.es

Specialist on oceanographic processes around Iberia

Along-slope oceanographic processes and sedimentary products around the Iberian margin

Hernández-Molina, F.J.¹, Serra, N.², Stow, D.A.V.³, Llave, E.⁴, Ercilla, G.⁵, Van Rooij, D.⁶

1. Facultad de Ciencias del Mar, Universidad de Vigo, 36200 Vigo (Spain). fjhernan@uvigo.es

2. Institut für Meereskunde, University of Hamburg, Bundesstr. 53, 20146 Hamburg (Germany). nuno.serra@zmaw.de

3. IPE-ECOSSE, Heriot Watt Univ. Edinburgh EH14 4AS (UK). dorrik.Stow@pet.hw.ac.uk

4. Instituto Geológico y Minero de España, Ríos Rosas, 23, 28003 Madrid (Spain). e.llave@igme.es

5. Instituto de Ciencias del Mar CMIMA-CSIC. Paseo Marítimo de la Barceloneta. 37-49 08003. Barcelona (Spain). gemma@cmima.csic.es

6. Renard Centre of Marine Geology (RCMG). Ghent University, Krijgslaan 281 S8. B-9000 Gent (Belgium). david.vanrooij@ugent.be

Abstract

The present contribution represents the first attempt to comprehensively describe regional along-slope processes and their sedimentary impacts around the Iberian margin, combining numerically simulated bottom currents with existing knowledge of contourite depositional and erosive features. The obtained correlation links the circulation of water masses with the main contourite depositional systems (CDS) and estimates other potential areas where new CDS could be found. Their study should be of great interest not only because of the stratigraphic, sedimentological, palaeoceanographic and palaeoclimatological significance, but also because of their relation with possible specific deep marine geohabitats and/or mineral and energy resources.

Key words: *Water-mass circulation, simulated bottom current velocities, along-slope processes, contourites, Iberian margin.*

Introduction

Along-slope oceanographic processes related to near-bottom currents generate large erosive and depositional features (drifts), which together shape Contourite Depositional Systems (CDS) or mixed contourite-turbidite systems, where bottom currents have interacted with down-slope processes (Faugères *et al.*, 1999; Stow *et al.*, 2002a; Viana and Rebesco, 2007; Hernández-Molina *et al.*, 2008a,b; Rebesco and Camerlenghi, 2008). However, when the water masses flowing along the continental slope are energetic enough, they can mask the effects of down-slope processes and generate complex CDS's of huge dimensions (10^3 - 10^6 km²) and sedimentary thicknesses (> 2000 m) comparable in extent with large Turbidite Depositional Systems (Zhenzhong *et al.*, 1998; Faugères *et al.*, 1999; Stow *et al.*, 2002a). Some of the better-known cases are those located along-slope of the Gulf of Cadiz (Nelson *et al.*, 1993, 1999; Llave *et al.*, 2001, 2007; Hernández-Molina *et al.*, 2003, 2006; Mulder *et al.*, 2003, 2006; Habgood *et al.*, 2003), Porcupine Seabight (Van Rooij *et al.*, 2003; Øvrebø *et al.*, 2006); offshore southern Greenland (Hunter *et al.*, 2007); offshore Brazil (Viana *et al.*, 2002a, b; Viana and Rebesco, 2007) and along the Argentine margin (Hernández-Molina *et al.*, 2009a).

The study of along-slope processes and their deposits in general, and the CDS's in particular, has represented one of the most active lines of research in Marine Geology during the last decade, mainly because of their very great stratigraphic, sedimentological, palaeoceanographic, and palaeoclimatological significance; their close link with sediment instability on continental slopes; and because of their direct relation with possible mineral and energy resources (Pickering *et al.*, 1989; Stow *et al.*, 2002b; Rebesco, 2005; Viana and Rebesco, 2007; Rebesco and Camerlenghi, 2008; Hernández-Molina *et al.*, 2010a).

1 International projects, such as the oceanic drilling programmes **DSDP** (*Deep Sea Drilling Project*), **ODP**
2 (*Ocean Drilling Program*) and **IODP** (*Integrated Ocean Drilling Program*) have corroborated this importance,
3 by underlining their very common occurrence along many ocean margins and in deep ocean basins.
4 Cumulatively, this work has also served to demonstrate that our understanding of such systems is still in its
5 infancy.
6

7
8 The Iberian continental margins have had a very complex and varied origin and geodynamic and
9 sedimentary evolution. For several decades, many studies have been carried out on the structure and
10 evolution of these margins (e.g. Boillot *et al.*, 1972, 1975, 1987; Derégnaucourt and Boillot, 1982; Comas
11 and Maldonado, 1988; Boillot 1979; Nelson y Maldonado, 1990; Maldonado and Nelson, 1988, 1999,
12 Maldonado and Comas, 1992; Estrada *et al.*, 1997; Pérez-Belzuz *et al.*, 1997; Pérez-Belzuz, 1999;
13 Maldonado *et al.*, 1999; Vazquez, 2001; Iglesias, 2009, among many others). The circulation of water
14 masses around Iberia leads to the development of along-slope currents generating, in turn, contourite
15 erosive and depositional features that form extensive, complex and often poorly known CDS's of large
16 dimensions and sediment thickness in different geologic contexts. These have a valuable sedimentary
17 record of their geological evolution, since the Iberian margins lie under the influence of several different
18 water masses, some of which impinge upon the seafloor with relatively high velocity and interact along the
19 middle and upper continental slope.
20
21

22
23 Although, there are many scattered bottom current measurements along the margin, there is still a major
24 lack of long-term records as well as areas with very few or no measurements. Whereas mean velocities are
25 commonly relatively low (5-15 cm/s), these are greatly exceeded in some places, reaching normal values >
26 80 cm/s and rarely almost 300 cm/s, as at the exit of the Strait of Gibraltar (Ambar and Howe, 1979; Iorga
27 and Lozier, 1999; Candela, 2001). Clearly, therefore, bottom currents are a major controlling factor on
28 along-slope sedimentation and in shaping the deep seafloor (Stow *et al.*, 2009). They are semi-permanent
29 features of deep ocean circulation, which vary in location and velocity over a range of timescales. They
30 operate at a range of scales, from the construction of small bedforms (surface lineation, crag and tail
31 structures, small ripples, etc) to the maintenance of large-scale drifts (Stow *et al.*, 2008), and even the
32 generation of large-scale erosional features, such as erosional terraces, abraded surfaces, channels, moat,
33 and furrows (Hernández-Molina *et al.*, 2008b). Stow *et al.* (2009) published recently a *bedform-velocity*
34 *matrix*, which facilitates the estimation of bottom current velocity based on bedform type, but can also be
35 applied in reverse to deduce the likely contourite features under a particular bottom current velocity
36 regime.
37
38

39
40 As part of this special volume of the Baiona International Congress on "*Deep-Water Circulation:*
41 *Processes and Products*" (Baiona, Spain, 16-18 June 2010) this paper represents the first attempt to
42 comprehensively describe along-slope processes and products along the Iberian margin (Fig. 1). There are
43 three main objectives: (a) to combine previous work and new regional data on along-slope processes and
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 their sedimentary impacts around Iberia; (b) to combine for the first time numerically simulated bottom
2 currents with existing knowledge of contourite depositional and erosive features; and (c) to use these data
3 to better link the circulation of water masses with the main CDS's, both known and as yet unexplored, and
4 to consider their future implications.
5
6
7

8 **Methods and data base** 9

10 This work synthesizes a large amount of both our own and published data from the Iberian margin in the
11 following broad areas.
12
13

14 *a) Characterization of water masses and their dynamics.* 15 16

17 A compilation of data and results about the water masses around the Iberian margins and their
18 dynamics, both in the Mediterranean Sea and in the Atlantic Ocean, has been realised and simplified for
19 this contribution. Physical oceanographic compilation was gathered from bibliographic sources, based on
20 data and ideas from many authors (Fig. 2 and 3). A compilation of water masses like the one presented
21 here is extremely difficult and complex since there is a great heterogeneity in both data coverage and
22 terminology. Therefore, our compilation should be considered as a preliminary sketch for evaluating the
23 along-slope processes around Iberia and further improvements will undoubtedly be necessary to update
24 and improve this first compilation. A special emphasis have been realized related the Mediterranean water
25 masses, especially to the Mediterranean Outflow Water (MOW), since many along-slope processes are
26 related to its circulation.
27
28
29
30
31
32
33
34

35 *b) Numerical bottom current circulation* 36 37

38 The numerical ocean simulations were performed with the MIT general circulation model (Marshall et
39 al., 1997), which is a primitive-equation z-coordinate ocean model. The computational domain included the
40 Northeast Atlantic and Western Mediterranean regions and extended from 9°E to 24°W and from 30°N to
41 48°N. In the horizontal, a resolution of 1/30° (approx. 2.8 km) was employed and 140 levels were used in
42 the vertical, with resolutions varying from 5m in the upper ocean to 100m in the deep ocean. The model
43 was initialized with temperature and salinity values from the January climatology of the World Ocean Atlas
44 2005 (Boyer et al., 2005) and bottom topography from the ETOPO2 database. Surface fluxes of momentum,
45 heat and freshwater used to force the ocean model were computed internally using bulk formulae and the
46 6-hourly atmospheric state from the NCEP reanalysis 1 (Kalnay et al., 1996). Since the configuration
47 featured a limited area, volume-balanced open boundary conditions were constructed from a 1/6°
48 resolution Atlantic solution of the same model forced by the same NCEP dataset (Serra et al., 2010a) and
49 applied at the four boundaries. Unresolved vertical mixing was parameterized by the KPP formulation
50 (Large et al., 1994) and horizontal mixing by a biharmonic operator with coefficients of $1 \times 10^3 \text{ m}^4 \text{ s}^{-1}$ and
51 $5 \times 10^8 \text{ m}^4 \text{ s}^{-1}$ for diffusion and viscosity, respectively. The model was integrated for the 6-year period 1990-
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1995 and the analysed daily-averaged results correspond to the 3-year period 1993-1995. The present simulations were previously analyzed in Serra et al. (2010b) with focus on the vertical structure of the model generated Mediterranean Water eddies and their interaction with the upper ocean layers. The good agreement with observations reported in that work built confidence on the realism of the simulations, which in the present work are analysed for near-bottom properties.

c) *Along-slope (contourite) sea bottom feature characterization*

We have compiled a broad database and from separate publications, noting details around Iberia of contourite depositional and erosive features, bedforms, substrate and current velocity as available. These data have enabled us to: a) identify where known CDS's are located, b) determine the major common features and their differences, outlining their role in the recent slope evolution; c) combine along-slope (contourite) sea bottom feature characterization with numerically simulated bottom currents.

Characterization of contourite features on sea-bottom has been executed including data from the Western Mediterranean Sea (Canals, 1985; Velasco *et al.*, 1996; Frigola *et al.*, 2007; 2008. Esteras *et al.*, 2000; Ercilla *et al.*, 2002), and the Atlantic Iberian margins (Madelain, 1970; Kenyon and Belderson, 1973; Gonthier *et al.*, 1984; Stow *et al.*, 1986, 2002c; Nelson *et al.*, 1993; 1999; Buitrago *et al.*, 2001; Llave *et al.*, 2001, 2005, 2006, 2007; Alves *et al.*, 2003; Habgood *et al.*, 2003; Hernández-Molina *et al.*, 2003, 2006, 2009b, 2010; Mulder *et al.*, 2003, 2006; Van Rooij *et al.*, 2003, 2010a; Øvrebø *et al.*, 2006; Hanquiez *et al.*, 2007; Ercilla *et al.*, 2008a and b; 2009, 2010; Marchès *et al.*, 2007; Garcia *et al.*, 2009; Iglesias, 2009; Bender *et al.*, 2010; Mena *et al.*, 2010). These data include: a) *Bathymetric data*, mainly regional bathymetric data from mono- and multi beam echosounders; b) *Sidescan sonar data*, available from some areas and providing useful backscatter imagery of the sea bottom; c) *Seismic reflection data*, including low-resolution reflection Multichannel seismic (MCS) reflection profiles; medium/high-resolution seismic profiles (from Sparker & Airgun systems); very high-resolution seismic profiles using mainly 3.5-kHz system and Topographic Parametric Sounder (TOPAS); d) *Core and borehole data* from different surveys, incorporated into the study of some areas, for groundtruthing seismic facies patterns, and in order to analyze the timing of the processes responsible for the development of the depositional units and to establish the chronology of the units; e) *Submarine photographs* as locally available (mainly in the Gulf of Cadiz); f) *Physical oceanographic data* taken on several of our own cruises comprising CTD and XBT profiles, which yield temperature and salinity information. Compilation and interpretation of this oceanographic data and its correlation with the morphologic features have already been published for some areas (García, 2002; Mulder *et al.*, 2003; 2006; Hernández-Molina *et al.*, 2006; 2009c).

Terminology

There is a growing consensus within the marine science community about the terminology to use for these along-slope sedimentary systems. We follow this terminology as far as possible. The term *bottom current* is preferred as the general term for those semi-permanent deep-water currents capable of eroding,

1 transporting, and depositing sediments on the sea-floor (Rebesco and Camerlenghi, 2008). Bottom currents
2 are the result of both THC and the major wind-driven circulation pattern of the oceans. Generally, these
3 currents are semi-permanent in nature with a net flow along-slope, but can be extremely variable in
4 direction and velocity, plus exhibit giant eddies, and local downslope, upslope or oblique-to-slope flow,
5 especially near the basin's entrance or exit gateways (Rebesco and Camerlenghi, 2008).
6

7
8 The term *contourite* is now generally accepted as the term for those sediments deposited or substantially
9 reworked by bottom currents and contour currents *sensu stricto*. The major accumulations of contourite
10 deposits are referred to as *drifts* or *contourite drifts*, for which several classifications have been proposed
11 mainly based on their morphologic, sedimentological and seismic characteristics (McCave and Tucholke,
12 1986; Faugères and Stow, 1993; Faugères et al., 1993, 1999; Rebesco and Stow, 2001; Stow et al., 2002b;
13 Rebesco, 2005; Rebesco and Camerlenghi, 2008). An association of various drifts and related erosional
14 features has been termed a *contourite depositional system* (CDS), by analogy with and of equal importance
15 to *turbidite depositional systems* (Stow et al., 1986, 2002b; Hernández-Molina et al., 2003; 2006, 2008a). In
16 the same way, where different CDS's are connected laterally (and vertically) and associated with the same
17 water mass in the same or adjacent basins, we can consider this as a *contourite depositional complex* (CDC,
18 Hernández-Molina et al., 2008a).
19
20
21
22
23
24
25
26
27
28
29

30 **Circulation of water masses around Iberia: oceanographic processes**

31 *a) Water masses and dynamics around Iberia.*

32
33
34
35 Around the Iberian margins, both in the Mediterranean Sea and in the Atlantic Ocean, there are several
36 water masses flowing at different depths and in the same or opposite directions, which generate important
37 along-slope sedimentary processes at the seafloor.
38
39

40
41 In the Mediterranean Sea, there are several places where dense water masses are generated (Fig. 2),
42 such that the Western Mediterranean Sea off the Iberian Peninsula comprises three main water masses
43 (Millot, 1999, 2009; Candela, 2001). 1) The deepest parts of the basin are filled by the *Western*
44 *Mediterranean Deep Water* (WMDW, Fig. 3 A), generated in the *Gulf of Lion* where convective processes
45 are known to reach depths up to 2000 m. In general, the WMDW flows below 500 m following the contour
46 of the lower slope, continental rise and abyssal plain around the Iberian Peninsula and Balearic Islands,
47 although in the Western Basin of the Alboran Sea, this water mass mainly flows along the Moroccan margin
48 (Fig. 3). 2) Between 500 and 150 m water depth, the basin is occupied by *Levantine Intermediate Water*
49 (LIW, Fig. 3 B), formed by convection in the Eastern Mediterranean and having passed through the *Sicilian*
50 *Strait*. Around Iberia, the LIW preferably flows along the upper and middle continental slope (Fig. 3). 3) The
51 upper layer (< 150 m) is formed by the inflow of Atlantic Water through the Gibraltar Strait (*Modified*
52 *Atlantic Water*, MAW, Fig. 3 C). Although these aforementioned water masses are generally recognised,
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Millot (2009) very recently has considered a more complex structure and dynamics of five water masses
2 along the western Mediterranean, including light versus dense Mediterranean water masses.

3 On the other hand, around the Atlantic Iberian margin, five main water masses have been more clearly
4 identified (Fig. 3A, B and C) (e.g. Iorga & Lozier, 1999; González-Pola, 2006; Serra et al., 2010b, among
5 others). These are, from bottom to top: 1) the Lower Deep Water (LDW) (mainly composed of Antarctic
6 Bottom Water, AABW, Fig. 3 A), flowing regionally below 4000 m mainly across the abyssal plains; 2) the
7 North Atlantic Deep Water (NADW, Fig. 3 A), flowing in different directions between 4000 and 2200 m
8 depth; 3) the Labrador Sea Water (LSW, Fig. 3 B), circulating regionally towards the SW between 2200 and
9 1500 m depth (main core at ~1800 m); 4) the Mediterranean Outflow Water (MOW, Fig. 3 B), located
10 between 1500 and 600 m depth, with the main core at ~1200 m flowing to the N and W along the middle
11 slope around the Iberian margin into the Bay of Biscay; and 5) the North Atlantic Central Water (NACW, Fig.
12 3 C) flowing in a complex circulation pattern with different currents and directions between 600 m and the
13 surface (main core at ~350 m).

14 Amongst these, the MOW is the most important in terms of along-slope processes and CDS development.
15 After its exit through the Gibraltar Gateway, the MOW represents an intermediate water mass, warm and
16 very saline which flows to the NW along the middle slope (Fig. 3B) under the *Atlantic Inflow* (AI) and above
17 the NADW (Zenk, 1975; Thorpe, 1975; Gardner and Kidd, 1983; Ochoa and Bray, 1991; Baringer and Price,
18 1999). It represents a flux of around 1.78 Sv through the Gibraltar Gateway, composed of both LIW and
19 WMDW (Bryden and Stommel, 1984; Bryden *et al.*, 1994), and generates important along-slope
20 sedimentary processes along the Atlantic margin (Serra et al., 2010b). In the Gulf of Cadiz, it flows between
21 500 and 1400 m depth with a velocity close to 300 cm/s at the Strait (Ambar and Howe, 1979) and around
22 80-100 cm/s at Cape San Vicente latitude (Cherubin *et al.*, 2000). Its distribution is conditioned by the
23 complex morphology of the continental slope, which generates two main cores, between 500-700 m depth
24 (Upper core or *Mediterranean Upper Water*, MU), and between 800 and 1400 m depth (Lower Core or
25 *Mediterranean Lower Water*, ML). The ML is further subdivided into three branches (Fig. 3B) (Madelain,
26 1970; Zenk, 1975; Ambar and Howe, 1979; Johnson and Stevens, 2000; Borenäs *et al.*, 2002). In the western
27 sectors, the interaction of these branches with the seafloor generates big *meddies* (Richardson *et al.*, 2000).

28 After exiting the Gulf of Cadiz, the MOW shows three principal branches (Fig. 3B): the main one flows to
29 the North, the second to the W, and the third to the S reaching the Canary Islands, and then veering
30 towards the W (Iorga and Lozier, 1999a; Slater, 2003). The northern branch flows along the middle slope of
31 the Portuguese margin, being divided in two by the influence of the Galicia Bank (Fig. 3B). These two
32 branches return to converge and subsequently circulate to the E in the Gulf of Biscay following the
33 continental slope contour (Fig. 3B). The MOW reaches the *Porcupine Bank* and partly circulates to the N
34 along the *Rockall Trough* until reaching the Norwegian Sea (Iorga and Lozier, 1999a; Slater, 2003).

b) *Simulated bottom velocities around Iberia.*

Analysing the simulated average bottom velocity (cm/s) and standard bottom velocity (cm/s) from Figures 4 and 5, one of the principal observations is that the regional circulation of the aforementioned water masses has a great impact at the seafloor in specific parts of both the Western Mediterranean Sea and Atlantic Ocean margins around Iberia. In our discussion below, we adopt the following notation for mean bottom current velocities, while recognising that maximum values may periodically reach two or three times the average:

- Low velocity: 0-10 cm/s. Little effect on seafloor, inhibits deposition, smooth surfaces, lineation.
- Medium velocity: 11-20 cm/s. Transports clay and silt, fine sand rippled surfaces and lineation.
- High velocity: 21-40 cm/s. Transports fine-medium sand, larger ripples, grooves/ridges.
- Very high velocity: > 40 cm/s. Transports sand, severe winnowing, variety of bedforms and erosion.

Within the Western Mediterranean Sea the greater velocities are found mainly on shelves and over the abyssal plains (Fig. 4 and 5), whereas most of the slope domains are swept by waters masses with low velocities (<4-5 cm/s). Nevertheless, in this general context the continental slopes that bound the Alboran Sea are an exception. Here the WMDW and LIW have slightly higher velocities associated with submarine scarps, local seamounts, and both slopes and abyssal plains of the western basin (Fig. 4 and 5). Along the Moroccan slope major velocities are along the middle and lower slope remain quite low (> 6 cm/s, Fig. 4 and 5), due to the LIW and WMDW circulation respectively. Along the northern Spanish slope, the higher velocities are also of similar low magnitudes and mainly located on the upper slope associated with the inflow of Atlantic Water through the Gibraltar Strait.

On the continental shelves of the Western Mediterranean Sea bottom simulated current velocities values are generally just over 5 cm/s (but locally over 10 cm/s), not only around the continent, but also on the continental shelf around the Balearic Islands (Fig. 4 and 5).

Simulated bottom current velocities on the abyssal plains are generally lower, especially under the 2000 m water depth due to the circulation of the WDMW. Average values are 3-4 cm/s, with relatively higher velocities mainly located on the lower slope around the Balearic Promontory and also in borders around the Algero-Provençal abyssal plain. Here the bottom-current velocities reach 10 cm/s, especially along the African margin (Fig. 4 and 5). Again, in this general context, the abyssal plain of Western Alboran Basins has several areas with bottom current velocities values up to 5 cm/s. These areas look like circular and are possibly related to eddies from the WMDW circulation (Fig. 4 and 5).

The Strait of Gibraltar itself has very high bottom current velocities related to the overflow of the MOW, up to a maximum recorded velocity of 2.8 m/s near the seafloor. Throughout the Gateway region velocities

are everywhere over 10 cm/s, and range from > 90 cm/s near the exit of the Straits to > 50 cm/s several hundred kilometres downstream in the Gulf of Cadiz (Fig. 4 and 5).

Along the Atlantic Ocean margin around Iberia, the highest simulated bottom current velocities are identified over the western and northern shelves, and especially along the middle slope (Fig. 4 and 5). Along-slope processes are therefore dominant along the middle slope of Gulf of Cadiz related to circulation of both the Upper (MU) and Lower (ML) strands. Here, simulated bottom current velocities are over 10 cm/s, and commonly with values of 50-90 cm/s. Over the rest of the Atlantic Iberian slope, the highest velocities are also identified along the middle slope (Fig. 4), with common values > 10 cm/s (and locally > 50 cm/s) associated with the northern branch of the MOW along the Portuguese, Galicia and Cantabrian slope. The upper and lower slope, in general, has simulated bottom current velocities with lower velocities of about < 4-5 cm /s, although where associated with large seamounts and banks (such as the Extremadura Spur; Oporto, Vigo and Vasco de Gama seamounts; and the Galicia Bank), velocities > 9-10 cm/s can be found.

Bottom-current velocities across the Atlantic abyssal plains are quite variable and can be high to very high. In the Madeira, the Horseshoe, the Tagus, and the Iberia abyssal plains, the highest velocities are mainly located along their eastern boundaries (e.g. at the base of the Madeira-Tore Rise or base of the Galicia Bank) often reaching 10 cm/s, and locally > 50 cm/s. More rarely, locally high values are found along their western boundaries (Fig. 4 and 5) or, within the Union basin, along its northern flank. Bottom current processes at this scale in these abyssal plains are associated with the LDW north-eastern circulation. Compared with these aforementioned abyssal plains, bottom current velocities over the Agadir, Sena, Biscay and Porcupine abyssal plains (Fig. 4 and 5) are generally much lower (< 5-8 cm/s) to extremely low (2-4 cm/s).

Deep gateways, such as the Discovery gap, Tagus valley and Theta Gap, are essential in controlling circulation and water-mass exchange in the abyssal regions (Fig. 4 and 5), and relatively higher bottom current velocities are also identified within them.

In the oceanic domain, away from the abyssal plains, the simulated bottom-current velocities are usually low (< 3 cm/s), but in this part of the north Atlantic there are numerous seamounts which represent important obstacles for the water-mass circulation and high bottom current velocities (10-20 times in magnitude their average velocity) are also identified around their flanks, reaching locally values > 50 cm/s (Fig. 4 and 5). In fact, one important observation is that, depending of the seamount location the position of the highest bottom current velocities is different. In this sense, the Ampere and Coral Patch Seamounts have the highest velocities along their northern flanks; the Madeira Island, DSM, USM and SSM seamounts have the highest velocities on their SE flanks; and although around the Atalante and Charcot Seamounts there are generally low bottom current velocities, the highest values are identified on their northern flanks. Over irregularities in oceanic crust located in the NW sector of Figures 4 and 5, south from the King's

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Trough, the highest bottom current velocities are related to the western and south-western flanks of oceanic highs. North from the King's Trough the distribution of the highest simulated bottom-current velocities is less clear.

Circulation of water masses around Iberia: products

Circulation of water-masses around the Iberia margin, in both the Mediterranean Sea and Atlantic Ocean, has produced a number of individual contourite drifts, and several different contourite depositional systems (CDS), each involving two or more drifts and associated erosional elements (Fig. 6). Within the Western Mediterranean Sea, three systems are documented under the influence of the LIW and WMDW (Fig. 6 and 7).

- 1) Along the NE Catalanian slope (Fig. 6), Canals (1985) defined the possible occurrence of contourite features at Rosas between 1200 and 2300 m water depth, and between the Cap de Creus and slightly south of the La Fonera Canyons. Here one or two distinct drifts are evident, each prograding upslope (Canals, 1985).
- 2) NW of Menorca, on the lower slope at depths up to 2000 m (Fig. 6), there is a broad, low-relief but distinct contourite drift. It is a plastered drift about 150 km in length and 25 km wide with a contourite moat along its seaward margin (Maufret, 1979; Velasco *et al.*, 1996) (Fig. 7). The drift is mud dominated and has about 100 m of relief above the adjacent sea floor. This contourite drift was generated by deep bottom currents belonging to the southward branch of the WMDW flow, which borders the Valencia Trough from north to south following a cyclonic pattern at depths of 2000m (Frigola *et al.*, 2007; 2008).
- 3) The Ceuta contourite drift is located in the southwestern Alboran Sea (SW Mediterranean), close to the Strait of Gibraltar (Fig. 6), running parallel to the Moroccan middle slope at a water depth of 200–700 m (Ercilla *et al.*, 2002). This drift is probably the best known drift in the Western Mediterranean to date. It is mainly characterized as an elongated plastered drift deposited over a slope terrace, with a broad lenticular geometry. It is around 100 km long, 28 km wide, with a relief of up to 400 m above from the surrounding sea-floor (up to 700 milliseconds two-way travel time, Fig. 7). It is formed under a westward flowing, simple tabular deep-water mass, producing deposition on the middle slope, and erosion at the toe of the slope (Ercilla *et al.*, 2002). The Ceuta Drift is composed of Pliocene and Quaternary deposits over the basal surface. The Quaternary sedimentary record has been divided into five prograding seaward units, with seismically stratified facies that produce reflections with high lateral continuity and amplitude, converging both seaward and landward (Fig. 7). The drift is composed mainly of muds, intercalated with 40–50 cm thick layers of sandy muds bounded by sharp surfaces and thinner (10 cm thick) silty clay layers with gradual contacts.

1 Within the Strait of Gibraltar the major defined processes are erosive, since the bottom current is high
2 enough to prevent deposition (Kelling and Stanley, 1972; Stanley et al., 1975; Serrano et al., 2005).
3 Although this area has been surveyed in several cruises, no distinctive contourite features have been
4 published thus far, apart from several large depressions and large channels that might be associated with
5 the MOW circulation (Esteras et al., 2000).
6

7
8 Along the Atlantic Iberian margins the following drifts and CDS's are known to have been generated (Fig.
9 6 and 7):
10

- 11 1) The interaction of the MOW with the middle slope of the Gulf of Cadiz margin has developed one of the
12 most extensive and complex CDS ever described, extending around the west Iberian margin (Fig. 6).
13 Many authors have highlighted this interaction and have characterized its erosive and depositional
14 features along the middle slope (e.g. Madelain, 1970; Kenyon and Belderson, 1973; Gonthier *et al.*,
15 1984; Nelson *et al.*, 1993; 1999; Llave *et al.*, 2001, 2006, 2007, 2010; Stow et al., 2002c; Alves et al.,
16 2003; Habgood et al., 2003; Hernández-Molina *et al.*, 2003, 2006; Mulder *et al.*, 2003, 2006; Hanquiez *et*
17 *al.*, 2007; Marchès *et al.*, 2007). This CDS comprises both large depositional and erosional features (Fig.
18 7), conditioned by a strong current with speeds reaching nearly 300 cm s^{-1} close to the Strait of
19 Gibraltar, slowing to $\sim 80 \text{ cm s}^{-1}$ at Cape St. Vincent (Kenyon and Belderson, 1973; Ambar and Howe,
20 1979; Cherubin et al., 2000). The main depositional features are sedimentary wave fields, sedimentary
21 lobes, mixed drifts, plastered drifts, elongated mounded and separated drifts, and sheeted drifts. The
22 main erosional features are contourite channels, furrows, marginal valleys and moats. All of them have a
23 specific location along the margin, and their distribution defines five morphosedimentary sectors within
24 the CDS (details are to be found in Hernández-Molina et al. 2003, 2006 and Llave et al., 2007). The
25 development of each of these five sectors at any time is related to a systematic deceleration of the
26 MOW as it flows westwards from the Strait, due to its interaction with margin bathymetry, and to the
27 effects of Coriolis force. In general, the drifts are mainly composed of muddy, silty and sandy sediments,
28 with a mixed terrigenous (the dominant component) and biogenic composition (Gonthier et al., 1984;
29 Stow et al., 1986, 2002c). By contrast, within the large contourite channels, sand and gravel are found
30 (Nelson et al., 1993, 1999) as well as many erosional features (Hernández-Molina et al., 2006; Garcia et
31 al., 2009). In the proximal sector close to the Strait of Gibraltar, an exceptionally thick sandy-sheeted
32 drift ($\sim 815 \text{ m}$ thick) is located, with sand layers that average $12\text{--}15 \text{ m}$ in thickness (minimum 1.5 m ,
33 maximum 40 m) (Buitrago et al., 2001; Llave et al., 2007).
34
35 2) Along the western Portuguese margin, Alves et al. (2003) have described the occurrence of contourite
36 features (Fig. 6), both along the middle slope as well as the continental rise. Middle slope drifts are
37 associated with the interaction of the MOW with different sectors of the middle slope of the Atlantic
38 Iberian margin (Fig. 7), but, on the contrary, drifts located on the rise have been attributed to the local
39 influence of AABW (LDW) north-eastern circulation (Alves *et al.*, 2003).
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
- 3) Contourite features have been recently defined around the Galicia Bank (Fig. 6), and those around the highs and topographic irregularities of the Transitional Zone, Deep Galicia Margin, and Northwestern Flank of the Galicia Bank (Ercilla et al., 2006, 2009, 2010). The main features described include moats, erosive and constructional furrows, abraded surfaces, large drifts (mainly elongated separated, and plastered) and sediment waves (Fig. 7). They are genetically associated with several water masses: MOW, LSW, NADW, and LDW, flowing at different depths and velocities and in different directions (Ercilla et al., 2010). In addition, along the Galicia slope between 1700 and 2300 m water depth (Fig. 6), elongated separated drifts mainly composed of muddy sediments have been described recently (Bender et al., 2010; Mena et al., 2010). Their genesis is attributed to the local slope morphology interacting with the Deep Intermediate Water mass (DIW), which flows parallel to the slope, as well as their proximity to shelf export pathways (Bender et al., 2010; Mena et al., 2010).
- 4) At Ortegual Spur, a CDS has been identified very recently, related to the MOW circulation along slope (Fig. 6) between 500-600 m and about 1500-1600 m (Hernández-Molina et al., 2009c). The interaction of the impinging MOW from west with the Ortegual Spur has generated a contourite depositional system (CDS) composed of erosive and depositional features (Fig. 7). The erosive features comprise terraces of tens of meters of relief and several hundreds of meters long, and moats of tens of meters of relief and of hundreds meters wide. The main depositional features (drifts) include plastered to mounded, elongated types (tens of meters thick). Sediment samples collected from the near surface of these drifts contain fine sands to muddy sandy (Hernández-Molina et al., 2009c), as well as planktonic foraminiferal tests as part of a biogenic sandy contourite.
- 5) The northernmost large CDS identified around the Iberian margin is located on the Cantabrian margin (Fig. 7), at the Le Danois Bank or “Cachucho” (Ercilla *et al.*, 2008a and b; Iglesias, 2009; Van Rooij et al., 2010a). It is unique with respect to the known sedimentary systems along the upper slope of the Biscay margin. Whereas the steep Biscay slopes are dominated by downslope processes, the Le Danois CDS has been generated by along-slope processes due to the MOW circulation, conditioned by seafloor irregularities and two topographic highs; the large Le Danois Bank and the smaller Vizco High. This has allowed the development of the present-day depositional and erosive features, such as respectively elongated mounded and separated drifts, plastered drifts, moats and slide scars (Van Rooij et al., 2010a).

54 Discussion

55 Numerical simulation of bottom currents

56
57
58 The circulation of water masses around Iberia leads to the development of along-slope currents and to
59 the generation of contourite erosive and depositional features, in some places forming complex CDS's of
60
61
62
63
64
65

1 large dimensions and sediment thickness (Hernández-Molina et al., 2009b, 2010b). The general link
2 between bottom current and deposits are well established and, although they are not well known in detail,
3 their sedimentary record will yield important information regarding their geological evolution and
4 palaeoclimate variation. Most of the drifts and CDS's described thus far are located on the slope. In the
5 Mediterranean Sea they are located on the middle (Rosas and Alboran CDS) and lower (Menorca and
6 Alboran CDS) slope due to the influence of the WMDW and LIW. Whereas on the Iberia Atlantic margin
7 they are mostly located on the middle slope associated with the remarkable influence of the MOW. This is
8 the case of the CDS's located in the Gulf of Cadiz, Portuguese slope, Galicia Margin, Ortegá Spur, and Le
9 Danois Bank. The influence of MOW stretches far outside the present study area to the N, to the Porcupine
10 Seabight and other parts of the NW European margin. Very few examples of contourite features are
11 associated in the Iberia Atlantic margin with other water masses, only locally in the Portuguese margin
12 (Alves et al., 2003) and the surrounding areas of the Galicia Bank (Ercilla et al., 2009, 2010).

13
14
15
16
17
18
19
20 This paper represents the first attempt to combine numerically simulated bottom currents with existing
21 data on contourite features. Immediately it can be demonstrated that the highest simulated bottom
22 currents are very coincident with the distribution of contourite drifts and CDS's (Fig. 8). In addition, there
23 are other areas with high simulated bottom current velocities but no previous description of any CDS.
24 Nevertheless, some large features related to the effect of bottom-current circulation have been published,
25 such as around the southern and southern east flanks of the Madeira Island and within the Agadir slope,
26 where large sedimentary wave fields were noted (Wynn et al., 2000).

27
28
29
30
31
32
33 Furthermore, numerically simulated bottom currents allow the identification of other areas where
34 contourite processes can also be found, not only associated to the MOW but also to other water masses.
35 For example, two major areas in deeper water can be identified (Fig. 8): a) the continental slopes of the
36 Alboran Sea and Atlantic Iberian margins; and b) the abyssal plains. In the Algero-Provençal abyssal plain,
37 bottom-current velocities are up of 10 cm/s, especially in its southern boundary along the African margin.
38 Strong bottom-current velocities in the Atlantic abyssal plains can also be identified. In the Madeira, the
39 Tagus, and the Iberia abyssal plains, higher velocities are located in their eastern boundaries, reaching 10
40 cm/s, and locally > 50 cm/s. Along-slope processes at this scale are mainly associated with the WMDW and
41 LIW along the Alboran Sea, and with both the MOW and the LDW in the Atlantic. In all these areas, there is
42 good potential for the recognition of contourite features in the future.

43
44
45
46
47
48
49
50
51 We recognise several important controls exerted by margin topography on bottom current velocities:

52
53
54 (a) *Oceanic gateways* are essential in controlling water-mass exchange and also the bottom-current
55 velocities and pathways between the abyssal plains. The Discovery gap, Tagus valley and Theta Gap are
56 each important deep gateways for deep-water circulation and each are associated with higher than normal
57 bottom current velocities, favouring the occurrence of contourite features (Fig. 8). The Gibraltar Gateway
58 has a very special role on the Mediterranean-Atlantic water-mass exchange and results in the highest
59
60
61
62
63
64
65

1 bottom current velocities known anywhere (nearly 300 cm/s). It also constitutes one of the most important
2 oceanic gateways in the world allowing the overflow and circulation of MOW to the Atlantic Ocean (Serra,
3 2004; Legg et al., 2009).

4 (b) *Oceanic gateways* with high velocities are typical of erosional bottom-current systems, characterized by
5 mainly linear bedforms displaying moderate to large seafloor relief resulting from scour and incision, as
6 well as by irregular scour hollows, sub-circular to circular holes, sand-gravel pavements and bare rock
7 surfaces (Stow et al., 2009).

8 (c) *Slopes, banks and promontories* are all instrumental in forcing higher than normal bottom current
9 velocities. The highest velocities observed around Iberia are found when the impinging flow increases its
10 velocity related to increased slope angles of prominent banks (such as the Galicia, Guadalquivir, or Le
11 Danois Banks), individual promontories (such as the Balearic or the Extremadura promontories), and spurs
12 (such as Ortegá Spur).

13 (d) *Seamounts* also represent important obstacles for water-mass circulation and high bottom current
14 velocities (up to 10-20 times the average velocity) can be identified around their flanks (Fig. 8), reaching
15 values > 50 cm/s. For this reason, notable contourite features are found enhanced around small and large
16 seamounts in the Alboran Sea (Palomino et al., 2010), along the Iberia Atlantic margin in the Gulf of Cadiz
17 (Hernández-Molina et al., 2006; García et al., 2009), and along the Galicia margin (Ercilla et al., 2009, 2010).

18 The collation and study of numerically simulated bottom currents can therefore determine new areas
19 where along-processes may be important and new large contourite depositional systems could be
20 developed. There are three principal reasons why they have not so far been identified: (i) they are masked
21 by the deposits of dominant down-slope processes, especially in some areas at the base of the slope,
22 continental rise or proximal areas of abyssal plains (Fig. 8); (ii) contourite deposition is inhibited or removed
23 by the frequent and/or erosive action of downslope processes; and (iii) good descriptive knowledge of
24 contourites in marine studies has only fairly recently been established, and much earlier traditional work
25 did not consider them. Certainly, there are more contourite systems to be found along the Iberian margins
26 so that their importance in the deep-sea context will be even greater than now. The new maps of simulated
27 bottom currents presented here will help define where new drifts and CDS's may be found, even within the
28 camouflage of major downslope systems (Fig. 9).

29 Although we have shown that the numerically simulated bottom current data are relatively coincident
30 with existing bottom current and sediment data, we should be cautious in its use and applications. In some
31 areas, for example, the simulated velocities do not coincide with present data collected by current meter or
32 inferred by bedforms. For example, this is the case on the slope north and northeast of Menorca Island,
33 where high bottom-currents were previously estimated (Mauffret et al., 1982; Maldonado et al., 1985;
34 Palanques et al., 1995; Velasco et al., 1996; Acosta, 2005). Here, near-bottom current measurements at
35 1800 m water depth in the Gulf of Lion deep margin where WMDW formation takes place (Millot and
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Monaco, 1984), gave mean values of 20 cm s^{-1} (maximum of 50 cm s^{-1}), although the simulated velocities are
2 very low indeed. A second example is in the middle slope of the Gulf of Cadiz, where the numerically
3 simulated bottom currents probably underestimate considerably the real values, because MOW reaches
4 nearly 300 cm s^{-1} close to the Strait of Gibraltar, slowing regionally to $\sim 80 \text{ cm s}^{-1}$ at Cape St. Vincent (Kenyon
5 and Belderson, 1973; Ambar and Howe, 1979; Cherubin et al., 2000), but having a great variability (Serra,
6 2004; Serra et al., 2010b), and increasing locally considerably due to seafloor irregularities (Hernández-
7 Molina et al., 2006; Gracia et al., 2009; Stow et al., 2009). In spite of these inconsistencies, the numerically
8 simulated bottom currents remain a very important first stage estimate for the margin as a whole.

13 We further note that the numerically simulated bottom current data provided here gives an average
14 value for the velocity, which takes into account some of the natural variability that occurs in nature. In fact,
15 bottom current velocities vary considerably over a range of timescales (Gross and Williams, 1991), whereas
16 other processes can model the speed and the instantaneous direction (Rebesco y Camerlenghi, 2008). This
17 natural variability may occur on a scale of 24 h due to deepwater tidal, internal wave or soliton influence
18 (Rebesco y Camerlenghi, 2008; Stow et al., 2009), a few weeks due to eddies, vortices, episodic benthic
19 storm events (Hollister, 1993; Stow et al., 2009; Serra et al., 2010), seasonally by winter convection (Canals
20 et al., 2006), and from tens to thousands of years due to other natural variations in bottom current
21 generation and transport (Rebesco y Camerlenghi, 2008; Stow et al., 2009). Nevertheless, the bottom
22 current velocity presented in this work (Fig. 5) estimates the areas where highest mean bottom current
23 velocities are located considering all this possible variability, and therefore is fully informative of the
24 regional view of where highest mean bottom current velocities should be located.

37 *Processes, products and their implications.*

40 Large contourite depositional and erosional features, as described here along the Iberia margin, are
41 generated by relatively stable hydrological conditions leading to long-term bottom water flows (Rebesco
42 and Camerlenghi, 2008). The present model of water masses circulation and its dynamics around Iberia
43 started to be developed after the opening of the Gibraltar Gateway at the end of the Miocene (5.4-5.6 Ma)
44 (Berggren and Hollister, 1974; Mulder and Parry, 1977; Maldonado *et al.*, 1999; García-Castellanos et al.,
45 2009), when the complete isolation of the Mediterranean Sea and global effects of the Messinian salinity
46 crisis ended (Ryan *et al.*, 1973; Hsü *et al.*, 1978; Duggen *et al.*, 2003). However, the major effects of MOW
47 were probably not felt until the Gateway had deepened sufficiently to allow its major outflow around 4.0-
48 4.2 Ma (Hernández-Molina et al., 2009d; Llave *et al.*, this volume). According to several authors, the most
49 recent hydrodynamic model was fully established later still at the base of the Quaternary (~ 2.4 - 2.6 Ma)
50 contemporary to the global cooling that initiates the glacial-interglacial cyclicity characteristic of the
51 Pleistocene (Thunell *et al.*, 1991). So far, the studies carried out on the CDS's of Mar de Alboran, Gulf of
52 Cadiz, and Le Danois has allowed to identify a similar evolution between them, determining coeval
53
54
55
56
57
58
59
60
61
62
63
64
65

1 palaeoceanographic changes in water masses in the past (Ercilla et al., 2002; Llave *et al.*, 2001, 2006;
2 Hernández-Molina *et al.*, 2002, 2006; Stow *et al.*, 2002c; Van Rooij et al., 2010a). Based on these works,
3 there was a further change in the nature of contourite depositional and erosive features coeval with the
4 Middle Pleistocene Revolution (MPR, 0.9 Ma) due to the switch to a “full glacial” mode with 100 ka cyclicity
5 (Llave, 2003; Llave et al., 2007; Van Rooij et al., 2010a). Since then, the influence of the environmental
6 changes (climatic and eustatic) in water masses circulation and their influence in the marine sedimentation
7 is partially known (e.g. Grousset *et al.*, 1988; Vergnaud-Grazzini *et al.*, 1989; Nelson *et al.*, 1993; Zanh et al.,
8 1999; Cacho *et al.*, 2000; Shackleton et al., 2000; Schönfeld and Zahn, 2000; Schönfeld et al., 2003; Llave et
9 *al.*, 2006, 2007; Voelker *et al.*, 2006; Frigola et al., 2007; 2008; Lebreiro et al., 2009, 2010; Voelker and
10 Lebreiro, 2010; Bender et al., 2010; Mena et al., 2010, among others).

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Once the present along-slope oceanographic processes were established we point out below some important implications based on the combination of numerically simulated bottom current with existing knowledge of contourite features.

1. *Continental Margin studies.* Recently, Weaver *et al.* (2000), Mienert and Weaver (2003) and Benetti (2006) have characterized the Atlantic continental margins based on their sedimentary processes. However, their model in common with earlier work, significantly underplays the role of along-slope processes on margin evolution. It is largely for this reason that our data for the Iberian margins do not conform well to the models that these authors suggest. The along-slope processes around Iberia are generated mainly by the MOW circulation in certain areas (e.g. *Gulf of Cadiz*, *Portuguese margin*, *Galicia margin* and *Cantabrian margin*). The interaction of these water masses has generated extensive contourite drifts and CDS's, especially during the Quaternary.
2. *Depositional controls.* The development of both depositional and erosional contourite features not only depends of the bottom-current velocity (as above) but also on several other important controls including: (a) *Local morphology of the margin.* The particular inherited geological features that shape the margin morphology and the various slope irregularities (as noted above), providing the best place for local water mass interaction with the seabed. (b) *Sediment supply.* There are a great variety of sediment inputs to drift generation – downslope supply from submarine canyons, slides and debris flows; suspended sediments stirred up at the shelf break or pirated by the water masses and deposited along slope; sand spillover from the adjacent shelf and hinterland; local and more distal erosion of the impinging water mass along its path flow. (c) *Local oceanographic behaviour of the MOW.* Some slope irregularities represent specific obstacles for the water mass circulation. The presence of such features can generate increased velocity and produce local turbulence, eddies, cores, filaments, secondary flows, etc, all of which affect local erosive and depositional contourite features (Hernández-Molina et al., 2008b; García et al., 2009). (d) *Sea-level and climate change.* One area of further work we are presently

following is to better elucidate the role of sea-level and climate change in controlling along-slope processes and products.

- 1
2
3 3. *Geohazards*. A combination of simulated bottom velocities around Iberia and seafloor mapping of
4 bottom-current bedforms along the margin can be of great significance in potential hazard mapping. In
5 addition to the simulated bottom velocities, there is a direct link between bottom current velocity and
6 bedform based on the bedform-velocity matrix recently published by Stow et al. (2009). Together these
7 data can provide information on likely seafloor current velocities and hence their impact on erosion and
8 chaffing of seafloor structures, submarine cables and pipelines. Furthermore, there is a growing
9 awareness of the link between bottom current activity and slope instabilities leading to major
10 downslope slumping and sliding.
- 11
12
13
14
15
16
17 4. *Water mass structure*. There are two specific elements of water mass structure that have important
18 implications for along-slope systems. (a) The development and maintenance of large gyres (eddies of
19 Meddies), which although very variable, serve to concentrate the highest bottom current energy in time
20 and space (Serra, 2004; Serra et al., 2010b). This can lead to periodic benthic storm events interspersed
21 with normal flow conditions. (b) The fact that some of the water masses circulating around the Iberia
22 margin have remarkable density contrasts. This in turn leads to the development of internal waves
23 and/or solitons along the interface between these water masses (Cacchione et al., 2002; Apel, 2004).
24 Any disturbance penetrating the pycnocline is capable of generating internal waves, but their generation
25 usually is conditioned by tidal influence (Apel, 2000; 2004). Energy associated with these internal waves
26 is locally important around the Iberia margin and can explain the reason why the simulated velocities
27 are higher than in surrounding areas, as in the case of the Western Alboran basin. The Camarinal and
28 Spartel Sills produce within the Strait of Gibraltar solitons with amplitudes of 50 to 100 m, wavelengths
29 of 2 to 4 km (Farmer and Armi, 1988; Armi and Farmer, 1988; Brandt et al., 1996; Jackson, 2004) and can
30 reach at least 200 km into the western Mediterranean Sea and exist for more than two days before
31 decaying toward background level (Apel, 2000; Jackson, 2004). This is also the case along the
32 northwestern coast of the Iberian Peninsula and around the Galicia Bank where large internal wave
33 occurrences take place (Correia, 2003; Jackson, 2004). The interaction of internal waves or solitons with
34 the sea-bottom affects depositional and erosion processes. In particular, the density interface
35 associated with internal waves has been postulated as one of the major mechanisms in the production
36 and maintenance of intermediate and bottom nepheloid layers (McCave, 1986; Dickson and McCave,
37 1986; Cacchione et al., 2002; Puig et al., 2004).
- 38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55 5. *Biota and bottom currents*. There is a relationship between specific biotic communities in deepwater, the
56 nature of bottom-water flow, the substrate sediment type, and any localized geological processes
57 (Howe et al., 2004; OSPAR, 2006; Reveillaud et al., 2008; Van Rooij et al., 2010b). This is evidenced, for
58 example, by the finding of chemosynthetic communities associated with carbonate mounds and mud
59
60
61
62
63
64
65

1 volcanoes; and cold-water coral (CWC) ecosystems under the path of bottom currents. For example, the
2 discovery of cold-water coral mounds along the Galicia Bank (Ercilla et al., 2009; 2010); the Bay of Biscay
3 and the Cantabrian margins, and the Porcupine Seabight (Van Rooij et al., 2003; Huvenne et al., 2009;
4 Van Rooij et al., 2010b) has been clearly associated with the dynamics of MOW. Additionally, the density
5 of the MOW seems to be most beneficial for benthic ecosystems in order to keep nutrients in
6 suspension and may provide a “connectivity” link for the dispersal and distribution of CWC larvae (Dullo
7 et al., 2008). Strong bottom currents are required in order to deliver re-suspended nutrients to the cold-
8 water coral ecosystems (Roberts et al., 2006). Therefore, bottom currents are one of the environmental
9 factors that influence both the growth and maintenance of deepwater biotic communities. They are also
10 one of the risk factors leading to substratum erosion and collapse, and to ecosystem silting up (Van
11 Rooij et al., 2010b).

12
13
14
15
16
17
18
19 6. *Mineral and energy resources.* Contourite systems are recognized as having a mainly unexplored
20 potential for both mineral and energy resources (Teleki *et al.*, 1987; Earney, 1990; Seibol y Berger, 1993;
21 Viana and Rebesco, 2007; Shanmugam, 2007; Rebesco and Camerlenghi, 2008; Viana, 2008; 2010). For
22 example, there are some specific mineral resources, such as ferromanganese nodules fields, which
23 depend directly on the sea-floor being swept by strong bottom currents (Van Andel et al., 1973; Kennet,
24 1982; Cronan, 2003). It was for this reason that Faugères and Stow (1993) suggested *manganiferrous*
25 *contourites* as a typical marine contourite facies. They are mainly associated with contourite channels
26 and moats (Faugères and Stow, 1993; Faugères *et al.*, 1993); or around Banks and seamounts (Kennet,
27 1982; Cronan, 2003), where higher bottom current velocities usually exist, as has been described for the
28 Gulf of Cadiz (Melières, et al., 1070; González et al., 2009a,b; 2010). The highest velocity bottom
29 currents have been demonstrated as fully capable of winnowing, cleaning and transporting sandy
30 material and of building extensive sandy sheeted drifts, such as that in the proximal Gulf of Cadiz.
31 Where buried in the subsurface, these provide a potential reservoir target for oil and gas resources.
32
33
34
35
36
37
38
39
40
41
42
43
44

45 **Conclusions**

46
47 The circulation of water masses around Iberia leads to the development of along-slope currents
48 generating, in turn, contourite erosive and depositional features that form individual contourite drifts,
49 erosional elements and complex CDS's of large dimensions and sediment thickness in a range of different
50 geologic contexts. These have a valuable sedimentary record of their geological evolution linked to
51 palaeoceanographic, paleoclimate and sea-level changes through time.
52
53
54
55

56 The regional simulated bottom current velocities of the water mass circulation around Iberia have
57 strong impact on the seafloor. Two principal areas of influence are noted: (a) the continental slopes of the
58 Alboran Sea and of the Atlantic Iberian margins; and (b) the abyssal plains in the Western Mediterranean
59 and eastern Atlantic. In the Algero-Provençal abyssal plain, bottom-current velocities are up of 10 cm/s,
60
61
62
63
64
65

1 especially on its southern boundary along the African margin. Strong bottom-current velocities in the
2 Atlantic abyssal plains can also be identified. In the Madeira, the Tagus, and the Iberia abyssal plains, higher
3 velocities are located in their eastern boundaries, reaching 10 cm/s, and locally > 50 cm/s. Along-slope
4 processes at this scale are mainly associated with the WMDW and LIW in the Alboran Sea, and with both
5 the MOW and the LDW in the Atlantic. Deep gateways are essential in controlling water-mass exchange and
6 also the bottom-current velocities and pathways between the abyssal plains. In addition, seamounts
7 represent important obstacles for the water-mass circulation and high bottom current velocities (10-20
8 times the normal velocity) are also identified around their flanks, reaching values > 50 cm/s.
9
10
11

12 Simulated bottom current velocities around Iberia, are very useful for a better understanding of the
13 detailed nature of bottom currents and of how they erode, transport and deposit sediment. From this study
14 it is clear that the role of bottom currents in shaping continental margins is generally significantly
15 underestimated, and that other parts of most margins will have still unknown and unexplored contourite
16 systems. Controls on along-slope sedimentary systems include the bottom current structure and velocity,
17 seafloor morphology and irregularities, sediment supply, sea-level and climate change. The combination of
18 simulated bottom velocities with seafloor mapping of bedforms and the bedform-velocity matrix recently
19 published by Stow et al. (2009) yields important information for geohazard mapping with respect to slope
20 instability and for the location of seafloor installations, cables and pipelines. It is also of great significance
21 because of their direct relation with specific deep marine geohabitats and mineral and energy resources.
22
23
24
25
26
27
28
29
30
31
32
33

34 Acknowledgements

35
36 The Spanish “Comisión Interministerial de Ciencia y Tecnología” supports this research through the Project CTM
37 2008-06399-C04/MAR (CONTOURIBER) and CTM2009-14157-C02. This work has been carried out as part of one
38 research stages funded by the ‘Mobility Award’ from the Spanish Ministry of Education and Science, which enable F^o.
39 Javier Hernández-Molina to work at Heriot-Watt University (UK) (Reference PR2009-0343). This research has also been
40 performed within the framework of the ESF MICROSISTEMS, EC FP6 HERMES, and EC FP7 HERMIONE. The numerical
41 simulations were developed in the frame of the Portuguese FCT project PPCDT/MAR/58384/2004. D. Vaan Rooij is a
42 post-doctoral fellow of the FWO-Flanders. The authors would like to thank Dr. Ricardo León (Spanish Geological
43 Survey, IGME, Spain) for his help in the bathymetric base map used in most of the figures. We also thanks to
44 Euromargins project MVSEIS and in particular to Luis Pinheiro, 2009 (Aveiro Univ., Portugal) for allowing us to use a
45 fragment of the seismic line 114 from Figure 7F. We are very grateful to REPSOL and TGS-NOPEC for allowing us to use
46 an unpublished data set of seismic records from the Gulf of Cadiz.
47
48
49

50 References

- 51
52 Acosta J (2005) El Promontorio Balear: Morfología submarina y recubrimiento sedimentario. Ph.D. thesis,
53 Univ. de Barcelona e Instituto Español de Oceanografía, Barcelona. 154 pp
54
55 Alves TM, Gawthorpe RL, Hunt DW, Monteiro JH (2003) Cenozoic tectono-sedimentary evolution of the
56 western Iberian margin. *Mar Geol* 195:75-108
57
58 Ambar I, Howe MR (1979) Observations of the Mediterranean outflow-II. The deep circulation in the vicinity
59 of the Gulf of Cadiz. *Deep-Sea Res* 26 A:555-568
60
61 Ambar I, Fiúza AFG (1994) Some features of the Portugal current system: a poleward slope undercurrent,
62
63
64
65

1 an upwelling-related summer southward flow and an autumn–winter poleward coastal surface
2 current. In: Katsaros KB, Fiúza AFG, Ambar I, (eds) Proceedings of the Second International Conference
3 on Air–Sea Interaction and on Meteorology and Oceanography of the Coastal Zone, Am Meteor Soc,
4 pp 286-287

5 Apel JR (2000) Solitons near Gibraltar: Views from the European Remote Sensing Satellites. Global Ocean
6 Assoc?? pp 23

7 Apel JR (2004) Oceanic Internal Waves and Solitons. In: Jackson CR, Apel JR (eds) Synthetic Aperture Radar
8 Marine User's Manual. U.S. Department of Commerce, National Oceanic and Atmospheric
9 Administration, pp 189-206

10 Arhan M, Colin de Verdière A, Mémery L (1994) The Eastern Boundary of the Subtropical North Atlantic. Am
11 Meteor Soc, 24:1295-1316.

12 Armi L, Farmer DM (1988) The flow of Mediterranean water through the Strait of Gibraltar. Prog Oceanogr
13 21:1-105

14 Arnone RA, Wiesenburg DE, Saunder KD (1990) The origin and characteristic of the Algerian Current. Jour
15 Geophys Res 95:1587-1598

16 Baringer MO, Price JF (1999) A review of the physical oceanography of the Mediterranean Outflow. Mar
17 Geol 155: 63-82

18 Bender VB, Hanebuth TJJ, Baumann KH (2010) Detrital input, marine productivity, deep-water currents and
19 sea level alternatively influencing sedimentation on the Galician continental margin over the past 30
20 ka BP. Geo-Temas 11:19-20

21 Benetti S (2006) Late Quaternary Sedimentary Processes along the western North Atlantic margin.
22 Unpublished PhD Thesis. Univ. Southampton (UK), pp 188

23 Berggren WA, Hollister CD (1974) Paleogeography, paleobiogeography and the history of the circulation in
24 the Atlantic Ocean. In: Studies in Paleooceanography. Soc Econ Paleontol Mineral Spec Pap 20:126-186

25 Boillot G, Dupeuble PA, Hennequin-Marchand I, Lamboy M, Lepetre JP (1972) Carte géologique du plateau
26 continental nord-espagnol entre le canyon de Capbreton et canyon d'Aviles. Bull Soc Geol France 7,
27 XV, 2-3:367-391

28 Boillot G, Dupeuble PA, Musellec P (1975) Carte géologique du plateau continental nord-portugais. Bull Soc.
29 Géol France (7)17:462-480

30 Boillot G, Dupeuble PA, Malod J (1979) Subduction and tectonics on the continental margin off northern
31 Spain. Mar Geol 32:53-70

32 Boillot G, Malod J-A, Dupeuble P-A, CYBERE Group (1987) Mesozoic evolution of Ortegual Spur, North Galicia
33 margin: Comparison with adjacent margins. In: Boillot G, Winterer EL, Meyer AW *et al*, (eds). Proc init
34 Repts (Pt. A) ODP 103, Texas A&M University, pp 107-119

35 Borenäs KM, Wahlin AK, Ambar I, Serra N (2002) The Mediterranean outflow splitting- a comparison
36 between theoretical models and CANIGO data. Deep-Sea Res II, 49:4195-4205

37 Boyer T, Levitus S, Garcia H, Locarnini R, Stephens C, Antonov J (2005) Objective analyses of annual,
38 seasonal, and monthly temperature and salinity for the World Ocean on a 0.25_ grid. Int J Climatol
39 25(7):931-945

40 Brandt P, Alpers W, Backhaus JO (1996) Study of the generation and propagation of internal waves in the
41 Strait of Gibraltar using a numerical model and synthetic aperture radar images of the European ERS 1
42 satellite. J Geophys Res 101:14,237-14,252

43 Bryden HL, Stommel HM (1984) Limiting processes that determine basic features of the circulation in the
44 Mediterranean Sea. Oceanol Acta 7:289-296

45 Bryden HL, Candela J, Kinder TH (1994) Exchange through the Strait of Gibraltar. Progr Oceanog 33:201-248

46 Buitrago J, García C, Cajebread-Brow J, Jiménez A, Martínez del Olmo W (2001) Contouritas: Un Excelente
47 Almacén Casi Desconocido (Golfo de Cádiz, SO de España). 1er Congreso Técnico Exploración y
48 Producción REPSOL-YPF (Madrid, 24-27, Septiembre), pp 10

- Cacchione DA, Pratson LF, Ogston AS (2002) The shaping of continental slopes by internal tides, *Science* 269:724-727
- 1 Cacho I, Grimalt JO, Sierro FJ, Shackleton NJ, Canals M (2000) Evidence of enhanced Mediterranean
2 thermohaline circulation during rapid climatic coolings. *Earth Planet Sci Lett* 183:417-429
3
- 4 Canals M (1985) Estructura sedimentaria y evolución morfológico del talud y el glacis continentales del
5 Golfo de León: fenómenos de desestabilización de la cobertera sedimentaria plio-cuaternaria. Tesis
6 Doct., Univ. Barcelona, pp 618
7
- 8 Canals M, Puig P, de Madron XD, Heussner S, Palanques A, Fabres J (2006) Flushing submarine canyons.
9 *Nature* 444:354-357
10
- 11 Candela J (2001) Mediterranean Water and Global Circulation. In: Sedler G, Church J, Gould J (eds) *Ocean
12 Circulation and Climate observing and Modelling the Global Ocean*. Ademic Press, 419-429
13
- 14 Cano N, García JM (1991) Corrientes en el litoral Malagueño. Baja frecuencia. *Boletino Instituto Español de
15 Oceanografía* 1(2):59-77
16
- 17 Caralp MH (1988) Late Glacial to Recent Deep-sea Benthic Foraminifera from the Northeastern Atlantic
18 (Cadiz Gulf) and Western Mediterranean (Alboran Sea): Paleoceanographic Results.
19 *Mar, Micropaleontol* 13:265-289
20
- 21 Caralp MH (1992) Paléohydrologie des bassins profonds nord-marocain (Est et Quest Gibraltar) au
22 Quaternaire terminal: apport des foraminifères benthiques. *Bull Soc Geol France* 163(2):169-178
23
- 24 Cherubin L, Carton X, Paillet J, Morel Y, Serpatte A (2000) Instability of the Mediterranean Water
25 undercurrents southwest of Portugal: effects of baroclinicity and topography. *Oceanolog Acta* 23
26 (5):551-573
27
- 28 Comas MC, Maldonado A (1988) Late Cenozoic sedimentary facies and processes in the Iberian Abyssal
29 Plain, Site 637, ODP Leg 103. In: Boillot G, Winterer EL, *et al*, (eds) *Proc ODP Sc Results 103*, College
30 Station, TX pp 635-655
31
- 32 Correia SML (2003) Observation of internal waves using multisensor satellite data off the Iberian Peninsula,
33 University of Lisbon, Institute of Oceanography, (in Portuguese) pp 164
34
- 35 Cronan DS (2003) Manganese Nodules. *Encycl Ocean Sciences*, 1526-1533.
36
- 37 Danialt N, Mazem JM, Arhan M (1994) Circulation and mixing of Mediterranean Water West of the Iberian
38 Peninsula. *Deep-Sea Res* 41:1685-1714
39
- 40 Díaz del Río G (2006) Patrones hidrológicos y variabilidad temporal de la circulación oceánica en la region
41 plataforma/talud del Golfo Artabro, noroeste de Galicia. Ph Degree. Univeridad de Vigo. Unpublished,
42 pp 188
43
- 44 Díaz del Río, G., González, N., Marcote, D., 1998. The intermediate Mediterranean water inflow along the
45 northern slope of the Iberian Peninsula. *Oceanologica Acta*, 21(2), 157-163.
46
- 47 Dickson RR, McCave N (1986) Nepheloid layers on the continental slope west of Porcupine Bank. *Deep-Sea
48 Res* 33(6): 91-818
49
- 50 Dickson RR, Gould WJ, Muller TJ, Maillard C (1985) Estimates of the mean circulation in the deep (> 2000 m)
51 layer of the eastern North Atlantic. *Prog Oceanogr* 14:103-127
52
- 53 Duggen S, Hoernle K, Van den Bogaard P, Rüpke L, Morgan JP (2003) Deep roots of the Messiniam salinity crisis.
54 *Nature*, 422:602-606
55
- 56 Dullo C, Flögel S, Rüggeberg A (2008) Cold-water coral growth in relation to the hydrography of the Celtic
57 and Nordic European continental margin. *Mar Ecol Prog Ser* 371:165-176
58
- 59 Earney CF (1990) *Marine Mineral Resources*. Routledge, Ocean Manage Policy Ser pp 387
60
- 61 Ercilla G, Baraza J, Alonso B, Estrada F, Casas D, Farrán M (2002) The Ceuta Drift, The Alboran Sea
62 (Southwestern Mediterranean), Quaternary. *Atlas of Deep-Water Contourite Systems: Modern Drifts
63 and Ancient Series, Seismic and Sedimentary Chacacteristics*. In: Stow DAV, Pudsey CJ, Faugeres JC
64
65

(eds). IGCP Publications, pp 155-170

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Ercilla G, García-Gil S, Estrada F, Gràcia E, Vizcaino A, Vázquez T, Díaz S, Vilas F, Casas D, Alonso B, Dañobeitia J, Farran M (2008a) High resolution seismic stratigraphy of the Galicia Bank Region and neighbouring abyssal plains (NW Iberian continental margin). *Mar. Geol.* 249: 108-127.
- Ercilla G, Casas D, Estrada F, Vázquez T, Iglesias J, García M, Gómez M, Acosta J, Gallart J, Maestro-González A, Marconi Team (2008b) Morphosedimentary features and recent depositional architectural model of the Cantabrian continental margin. *Mar Geol* 247(1-2):61-83
- Ercilla G, Casas D, Somoza L, Vázquez JT, Iglesias J, García-Gil S, Medialdea T, León R, Estrada F, ERGAP Project and Cruise Teams (2009) Cartografiando la dinámica sedimentaria de la región del Banco de Galicia. *Nuevas Contribuciones al Margen Ibérico*: 201-204
- Ercilla G, Casas D, Iglesias J, Vázquez JT, Somoza L, León R, Medialdea T, Juan C, García M (2010) Contourites in the Galicia Bank region (NW Iberian Atlantic). *Geo-Temas* 11:33-34
- Esteras M, Izquierdo J, Sandoval NG, Mamad A (2000) Evolución morfológica y estratigráfica plio-cuaternaria del Umbral de Camarinal (Estrecho de Gibraltar) basada en sondeos marinos. *Revista de la Sociedad Geológica de España*, 13(3-4):539-550
- Estrada F, Ercilla G, Alonso B (1997) Pliocene-Quaternary tectonic-sedimentary evolution of the NE Alboran Sea (SW Mediterranean Sea). *Tectonophysics* 282:423-442
- Farmer DM, Armi L (1988) The flow of Mediterranean water through the Strait of Gibraltar, *Prog. Oceanogr* 21:1-105
- Faugères JC, Stow DAV (1993) Bottom current controlled sedimentation: a synthesis of the contourite problem. *SedGeol* 82:287-297
- Faugères JC, Stow DAV, Imbert P, Viana A (1999) Seismic features diagnostic of contourite drifts. *Mar Geol* 162:1-38
- Faugères JC, Mezerais ML, Stow DAV (1993) Contourite drift types and their distribution in the North and South Atlantic Ocean basins. *Sed Geol* 82:189-203
- Fiúza AFG (1984) Hidrologia e dinâmica das águas costeiras de Portugal. PhD Thesis. Universidade de Lisboa, pp 294
- Fiúza AFG, Hamann M, Ambar I, Díaz del Río GD, González N, Cabanas JM (1998) Water masses and their circulation off western Iberia during May 1993. *Deep-Sea Res* 45:1127-1160
- Frigola J, Moreno A, Cacho I, Canals M, Sierro FJ, Flores JA, Grimalt JO, Hodell DA, Curtis JH (2007) Holocene climate variability in the western Mediterranean region from a deepwater sediment record *Paleoceanography*, 22, PA2209, doi:10.1029/2006PA001307
- Frigola J, Moreno A, Cacho I, Canals M, Sierro FJ, Flores JA, Grimalt JO (2008) Evidence of abrupt changes in Western Mediterranean Deep Water circulation during the last 50 kyr: A high-resolution marine record from the Balearic Sea. *Quat Int* 22:88-104
- García Lafuente J, Sánchez Garrido JC, Díaz del Río G, Criado Aldeanueva F, Marcote D, Sánchez Román A (2008) Low-frequency variability of the Mediterranean undercurrent off Galicia, northwestern Iberian Peninsula. In: Gómez Gesteira MG, de Castro M, Rosón G et al. (eds). *J Mar Syst* 74(1-4):351-363
- García-Castellanos D, Estrada F, Jiménez-Munt I, Gorini C, Fernández M, Vergés J, De Vicente R (2009) Catastrophic flood of the Mediterranean after the Messinian salinity crisis. *Nature*, Vol 462, doi: 10.1038/08555, 778-781
- García M (2002) Caracterización Morfológica del Sistema de Canales y Valles Submarinos del Talud Medio del Golfo de Cádiz (SO de la Península Ibérica): Implicaciones Oceanográficas. Tesis de Licenciatura, Facultad de Ciencias del Mar, Univ. Cádiz, pp 114
- García M, Hernández-Molina FJ, Llave E, Stow DAV, León R, Fernández-Puga MC, Díaz del Río V, Somoza L (2009) Contourite erosive features caused by the Mediterranean Outflow Water in the Gulf of Cadiz: Quaternary tectonic and oceanographic implications. *Mar Geol* 257:24-40
- Gardner J, Kidd RB (1987) Sedimentary processes on the northwestern Iberian continental margin viewed

by long-range side-scan sonar and seismic data. *J Sed Petrol* 57: 397-407

- 1 Gardner J, Kidd R.B (1983) Sedimentary processes on the Iberian continental margin viewed by long-range
2 side-scan sonar. 1: Gulf of Cadiz. *Oceanolog Acta* 6(3):245-254
- 3 Garrett C, Thompson K, Blanchard W (1990) Sea-level flips. *Nature* 348:292
- 4 Gascard JC, Richez C (1985) Water masses and circulation in the Western Alboran Sea and in the Straits of
5 Gibraltar. *Prog Oceanogr* 15:157-216
- 6
- 7 Gonthier EG, Faugères JC, Stow DAV (1984) Contourite facies of the Faro Drift, Gulf of Cadiz. In: Stow DAV,
8 Piper DJW (eds). *Fine-Grained Sediments, Deep-Water Processes and Facies*. *Geol Soc Sp Publ* 15:275-
9 292
- 10
- 11 González-Pola, C (2006) Variabilidad climática en la región sureste del Golfo de Vizcaya, Universidad de
12 Oviedo, Oviedo, pp 196
- 13
- 14 González FJ, Somoza L, Lunar R, Martínez-Frías J, Martín Rubí JA, Torres T, Ortiz JE, Díaz del Río V, Pinheiro
15 L, Magalhaes VH (2009) Hydrocarbon-derived ferromanganese nodules in carbonate-mud mounds
16 from the Gulf of Cadiz: mud-breccia sediments and clasts as nucleation sites. *Mar Geol* 261(1-4): 64-81
- 17
- 18 González FJ, Somoza L, Lunar R, Martínez-Frías J, Martín Rubí JA, Torres T, Ortiz JE, Díaz del Río V (2010a)
19 Internal features, mineralogy and geochemistry of Fe-Mn nodules from the Gulf of Cadiz: The role of
20 the Mediterranean Outflow Water Undercurrent (doi:10.1016/j.jmarsys.2009.10.010). *Mar Syst* 80(3-
21 4):203-218.
- 22
- 23 González FJ, Somoza L, León R. Medialdea T (2010b) Ferromanganese deposits associated to the Cadiz
24 Contourite Channel: imprints of the Mediterranean Outflow Water. *Geo-Temas* 11:55-56
- 25
- 26 Gross TF, Williams AJ (1991) Characterisation of deep-sea storms. *Mar Geol* 99:281-301
- 27
- 28 Grousset F, Joron JL, Biscaye PE, Latouche C, Treuil M, Maillet N, Faugères JC, Gonthier E (1988)
29 Mediterranean outflow through the Strait of Gibraltar since 18.000 years B.P. *Mineralogical and*
30 *Geochemical agreement*. *Geo-Mar Lett* 8(1):25-34
- 31
- 32 Habgood EL, Kenyon NH, Masson DG, Akhmetzhanov A, Weaver PPE, Gardner J, Mulder T (2003) Deep-
33 water sediment wave fields, bottom current sand channels and gravity flow channel-lobe system: Gulf
34 of Cadiz, NE Atlantic. *Sedimentology* 50:483-510
- 35
- 36 Hanquiez V, Mulder T, Lecroart P, Gonthier E, Marchès E, Voisset M (2007) High resolution seafloor images
37 in the Gulf of Cadiz, Iberian margin. *Mar Geol* 28:42-59
- 38
- 39 Haynes R, Barton, ED (1990) A poleward flow along the Atlantic coast of the Iberian Peninsula. *J*
40 *Geophysical Research*, 95, 11425-11441.
- 41
- 42 Herburn, G.W., La Violette, P.E., 1990. Variations in the structure of the anticyclonic gyres found in the
43 Alboran Sea. *Journal of Geophys Res* 95:1599-1613
- 44
- 45 Hernández-Molina FJ, Somoza L, Vázquez JT, Lobo F, Fernández-Puga MC, Llave E, Díaz-del-Río V (2002)
46 Quaternary stratigraphic stacking patterns on the continental shelves of the southern Iberian
47 Peninsula: Their relationship with global climate and palaeoceanographic changes. *Quat Int* 92(1): 5-23
- 48
- 49 Hernández-Molina FJ, Llave E, Somoza L, Fernández-Puga MC, Maestro A, León R, Barnolas A, Medialdea T,
50 García M, Vázquez JT, Díaz del Río V, Fernández-Salas LM, Lobo F, Alveirinho Dias JM, Rodero J,
51 Gardner J (2003) Looking for clues to paleoceanographic imprints: a diagnosis of the gulf of Cadiz
52 contourite depositional systems. *Geology* 31(1): 19-22
- 53
- 54 Hernández-Molina FJ, Llave E, Stow DAV, García M, Somoza L, Vázquez JT, Lobo FJ, Maestro A, Díaz del Río
55 V, León R, Medialdea T, Gardner J. (2006) The contourite depositional system of the Gulf of Cádiz: A
56 sedimentary model related to the bottom current activity of the Mediterranean outflow water and its
57 interaction with the continental margin. *Deep Sea Res II*, 53:1420-1463
- 58
- 59 Hernández-Molina FJ, Maldonado A, Stow DAV (2008a) Abyssal Plain contourites. In: Rebesco M,
60 Camerlenghi A (eds) *Contourites*. *Developments in Sedimentology* 60. Elsevier, pp 347-377
- 61
- 62 Hernández-Molina FJ, Llave E, Stow DAV (2008b) Continental slope contourites. In: Rebesco M,
63 Camerlenghi A (eds) *Contourites*. *Developments in Sedimentology* 60. Elsevier, pp 379-407
- 64
- 65

- 1 Hernández-Molina FJ, Paterlini M, Violante R, Marshall P, de Isasi M, Somoza L, Rebesco M (2009a) A
2 contourite depositional system in the Argentine slope: An exceptional record of the influence of
3 Antarctic water masses. *Geology* 37:507-510
- 4 Hernández-Molina FJ (2009b) Los Sistemas Deposicionales Contorníticos: ejemplos alrededor del Margen
5 Ibérico. *Nuevas Contribuciones al Margen Ibérico. 6th Symposium on the Atlantic Iberian Margin (MIA
6 09)*. Volume of abstract, xxi-xxiv.
- 7 Hernández-Molina FJ, Nombela MA, Van Rooij D, Roson G, Ercilla G, Urgorri V, Llave E, Francés G, De Mol L,
8 Estrada F, León R, Mena A, Pérez Arlucea M, Alejo I. Jane G, VanReusel A (2009c) The Ortegale Spur
9 Contourite Depositional System (Bay of Biscay): the implications of the Mediterranean Outflow Waters
10 in sedimentary processes and cold-water coral ecosystems. *Nuevas Contribuciones al Margen Ibérico.
11 6th Symposium on the Atlantic Iberian Margin (MIA 09)*. Volume of abstract, 281-284.
- 12 Hernández-Molina, F.J., Matias, H., Llave, E., Stow, D.A.V. (2009d). Onset of contourite deposition in the
13 Gulf of Cadiz: preliminary results. *Nuevas Contribuciones al Margen Ibérico. 6th Symposium on the
14 Atlantic Iberian Margin (MIA 09)*. Volume of abstract, 277-280.
- 15 Hernández-Molina FJ, Serra N, Stow DAV (2010b) Along-slope oceanographic processes and sedimentary
16 products around Iberia. *Geo-Temas* 11:69-70
- 17 Hernández-Molina FJ, Stow DAV, Llave E, Rebesco M, Ercilla G, Van Rooij D, Mena A, Vázquez JT, Voelker A
18 (eds), (2010a). *Deep-Water Circulation: Processes and Products*. International Congress. Baiona 16, 17
19 & 18 June. *Geo-Temas* 11, pp 204
- 20 Hollister CD (1993) The concept of deep-sea contourites. *Sed Geol* 82:1-7
- 21 Howe JA, Shimmield TM, Diaz R (2004) Deep-water sedimentary environments of the northwestern
22 Weddell Sea and South Sandwich Islands, Antarctica. *Deep-Sea Res II* 51: 1,489-1,514
- 23 Hsü KJ, Montader L, Bernoulli D, Cita MB, Erickson A, Garrison RE, Kidd RB, Mélières F, Muller C, Wright R
24 (1978) History of the Mediterranean salinity crisis. In: Hsü KJ, Montadert L, et al. (eds) *Initial Reports of
25 the Deep Sea Drilling Project, 42 (Part 1)* U.S. Govt. Printing Office, Washington, D.C. pp 1053-1078
- 26 Hunter SE, Wilkinson D, Stanford J, Stow DAV, Bacon S, Akhmetzhanov AM, Kenyon NH (2007) The Eirik
27 Drift: a longterm barometer of north Atlantic deepwater flux south of Cape Farewell, Greenland. In:
28 Viana A, Rebesco M (eds) *Economic and Paleoceanographic Importance of Contourites*. *Geol Soc
29 London, Sp Publ*, pp 245-264
- 30 Huthnance M, van Aken HM, White M, Barton ED, Le Cann B, Coelho EF, Fanjul EA, Miller P, Vitorino J
31 (2002) Ocean margin exchange – water flux estimates. *J Mar Syst* 32(1–3):107-137
- 32 Huvenne VAI, Van Rooij D, De Mol B, Thierens M, O'Donnell R, Foubert A (2009) Sediment dynamics and
33 palaeo-environmental context at key stages in the Challenger cold-water coral mound formation: Clues
34 from sediment deposits at the mound base. *Deep Sea Res I*, 56:2263-2280
- 35 Iglesias J (2009) *Sedimentation on the Cantabrian Continental Margin from Late Oligocene to Quaternary*.
36 Ph D. Universidad de Vigo, pp 185
- 37 Iorga M, Lozier MS (1999) Signatures of the Mediterranean outflow from a North Atlantic climatology. 1.
38 Salinity and density fields. *J Geophys Res* 194:25985-26029
- 39 Jackson CR (ed) (2004) *An Atlas of Oceanic Internal Solitary Waves (February 2004)* by Global Ocean
40 Associates Prepared for Office of Naval Research – Code 322 PO, pp 560
- 41 Johnson J, Stevens I (2000) A fine resolution model of the eastern North Atlantic between the Azores, the
42 Canary Islands and the Gibraltar Strait. *Deep-Sea Res I*, 47:875-899
- 43 Kalnay E, Kanamitsu M, Kistler R, Collins W, Deave D, Gandin L, Iredell M, Saha S, White G, woollen J, Zhu Y,
44 Chelliah M, Ebisuzaki W, Higgins W, Janowiak J, Mo KC, Ropelewski Cm wang J, Leetmaa A, Reynolds R,
45 Jenne R, Joseph D (1996) The NCEP/NCAR 40-year reanalysis project. *B Am Meteor Soc* 77:437-470
- 46 Kelling G, Stanley DJ (1972) Sedimentary evidence of bottom current activity, Strait of Gibraltar region. *Mar
47 Geol* 13:M51-M60
- 48 Kennett JP (1982) *Marine Geology*. Prentice Hall, pp 813
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- Kenyon NH, Belderson RH (1973) Bed forms of the Mediterranean undercurrent observed with side-scan sonar. *Sed Geol* 9:77-99
- 1 Large W, McWilliams J, Doney S (1994) Ocean vertical mixing: a review and a model with a nonlocal
2 boundary layer parameterization. *Rev Geophys* 32:363-403
3
- 4 Lebreiro SM (2010) Circum-Iberia turbidites and contourites: enhancement of processes by AMOC
5 millennial-scale climate changes. *Geo-Temas* 11:93-94
6
- 7 Lebreiro S, Voelker A, Vizcaino A, Abrantes F, Alt-Epping U, Jung S, Thouveny N, Gracia E (2009) Sediment
8 instability on the Portuguese margin under abrupt glacial climate changes (last 60 kyr). *Quat Sci Rev*
9 *28*:3211-3223.
- 10 Legg S, Briegleb B, Chang Y, Chassignet EP, Danabasoglu G, Ezer T, Gordon AL, Griffies S, Hallberg R, Jackson
11 L, Large W, Özgökmen TM, Peters H, Price J, Riemenschneider U, Wu W, Xu X, Yang J (2009) Improving
12 Oceanic overflow representation in climate models. The Gravity Current Entrainment Climate Process
13 Team. *Am Meteorol Soc* 657-670
14
- 15 Llave E (2003) Análisis morfosedimentario y estratigráfico de los depósitos contorníticos del Golfo de Cádiz:
16 Implicaciones paleoceanográficas. Tesis Doctoral, Facultad de Ciencias del Mar, Univ. Cádiz, pp 343
17
- 18 Llave E, Hernández-Molina FJ, Somoza L, Díaz del Río V, Stow DAV, Maestro A, Alveirinho Dias JM (2001)
19 Seismic stacking pattern of the Faro-Albufeira contourite system (Gulf of Cadiz): a Quaternary record
20 of paleoceanographic and tectonic influences. *Mar Geophys Res* 22: 487-508
21
- 22 Llave E, Hernández-Molina FJ, Stow DAV, Somoza L, Díaz del Río V (2005) The Contourite Depositional
23 System in the Gulf of Cadiz: An Example of Drifts with Reservoir Potential Characteristics - 25
24 Aniversario de la Asociación de Geólogos y Geofísicos Españoles del Petróleo. AAGEP, Sp Publ pp 53-73
25
- 26 Llave E, Schönfeld J, Hernández-Molina FJ, Mulder T, Somoza L, Díaz del Río V, Sánchez-Almazo I (2006)
27 High-resolution stratigraphy of the Mediterranean outflow contourite system in the Gulf of Cadiz
28 during the late Pleistocene: the impact of Heinrich events. *Mar Geol* 227:241-262
29
- 30 Llave E, Hernández-Molina FJ, Somoza L, Stow DAV, Díaz del Río V (2007) Quaternary evolution of the
31 contourite depositional system in the Gulf of Cadiz. In: Viana A, Rebesco M (eds) *Economic and*
32 *Paleoceanographic Importance of Contourites*. *Geol Soc London Sp Publ* 276, pp 49-79
33
- 34 Llave E, Matias H, Hernández-Molina FJ, Ercilla G, Stow DAV, Medialdea T (2010) Pliocene and Quaternary
35 seismic stacking pattern and distribution of contourites in the Algarve margin (Northern Gulf of Cadiz,
36 Spain). *Geo-Temas* 11: 03-104
37
- 38 Madelain F (1970) Influence de la topographie du fond sur l'écoulement méditerranéen entre le Détroit de
39 Gibraltar et le Cap Saint-Vincent. *Cah Océanogr* 22: 3-61
40
- 41 Maldonado A, Nelson CH (1988) Dos ejemplos de márgenes continentales de la Península Ibérica: El margen
42 del Ebro y el Golfo de Cádiz. *Revista de la Sociedad Geológica de España* 1(3-4):317-325
43
- 44 Maldonado A, Comas MC (eds) (1992) *Geology and geophysics of the Alboran Sea*. *Geo-Mar Lett* 12(2-3) pp
45 188
- 46 Maldonado A, Nelson CH (eds) (1999) *Marine Geology of the Gulf of Cadiz*. *Mar Geol* 155(1/2) pp 248
47
- 48 Maldonado A, Got H, Monaco A, O'Connell S, Mirabile L (1985) Valencia Fan (northwestern
49 Mediterranean): Distal deposition fan variant. *Mar Geol* 62:295-319
50
- 51 Maldonado A, Somoza L, Pallares L (1999) The Betic orogen and the Iberian-African boundary in the Gulf of
52 Cadiz, geological evolution central North Atlantic. *Mar Geol* 155:9-43
53
- 54 Marchès E, Mulder T, Cremer M, Bonnel C, Hanquiez V, Gonthier E, Lecroart P (2007) Contourite drift
55 construction influenced by capture of Mediterranean Outflow Water deep-sea current by the Portimao
56 submarine canyon (Gulf of Cadiz, South Portugal). *Mar Geol* 242:247-260
57
- 58 Marshall J, Adcroft A, Hill C, Perelman L, Heisey C (1997) A finite-volume, incompressible Navier Stokes
59 model for studies of the ocean on parallel computers. *J Geophys Res* 102:5753-5766
60
- 61 Mauffret A (1979) Etude géodynamique de la marge des Illes Baléares. *Mem Soc Geol Fr LVI(132)*:1-96
62
- 63 Mauffret A, Labarbarie M, Montadert L (1982) Les affleurements de series sedimentaires pre-pliocenes
64
65

dans le bassin Méditerranéen nord-occidental. *Mar Geol* 45:159-175

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Mauritzen C Morel Y, Paillet J (2001) On the influence of Mediterranean water on the central waters of the North Atlantic Ocean. *Deep Sea Res* 44:1543-1574
- Mazé JP, Arhan M, Mercier H (1997) Volume budget of the eastern boundary layer off the Iberian Peninsula. *Deep Sea Res* 44:1543-1574
- McCartney MS, Talley LD (1982) The Subpolar Mode Water of the North Atlantic Ocean. *J Phys Oceanogr* 12:1169-1188
- McCartney MS (1992) Recirculating components to the deep boundary current of the northern North Atlantic. *Prog Oceanogr* 29:283-383
- McCave IN (1986) Local and global aspects of the bottom nepheloid layers in the world ocean. *Neth J Sea Res* 20:167-181
- McCave IN, Tucholke BE (1986) Deep current controlled sedimentation in the western North Atlantic. In: Vogt PR, Tucholke BE (eds), *The Geology of North America, Vol. M, The Western North Atlantic Region*. Geol Soc Am Boulder (CO), pp 451-468
- Melières F (1974) Recherches sur la dynamique sédimentaire du Golfe de Cadix (Espagne). These de Doctoral, Univ. Paris A, pp 235
- Melières F, Nesteroff WD, Lancelot Y (1970) Etude photographique des fonds du Golfe de Cádiz. *Cah Océanogr* 22:63-72
- Mena A, Francés G, Pérez-Arlucea M, Hanebuth TJJ, Nombela MA (2010) Sedimentary evolution of the Galicia Inner Basin during the last 70 kyr. *Geo-Temas* 11:125-126
- Mienert J, Weaver PPE (2003) *European Margin Sediment Dynamics*. Springer, Berlin, pp 323
- Millot C (1999) Circulation in the Western Mediterranean Sea. *J Mar Syst* 20:423-442
- Millot C (2009) Another description of the Mediterranean Sea outflow. *Prog Oceanogr* 82:101-124
- Millot C, Monaco A (1984) Deep strong currents and sediment transport in the northwestern Mediterranean Sea. *Geo Mar Lett* 4:13-17
- Mulder CJ, Parry GR (1977) Late Tertiary evolution of the Albran Sea at the eastern entrance of the Straits of Gibraltar. In: Biju-Duval B, Montadert L (eds) *International symposium on the structural history of the Mediterranean Basins*. Split (Yugoslavia), Editions Technip, Paris, pp 401-410
- Mulder T, Voisset M, Lecroart P, Le Drenzen E, Gonthier E, Hanquiez V, Faugères J-C, Habgood E, Hernandez-Molina FJ, Estrada F, Llave E, Poirier D, Gorini C, Fuchey Y, Volker A, Freitas P, Lobo Sanchez F, Fernandez LM, Morel J (2003) The Gulf of Cadix: an unstable giant contouritic levee. *Geo-Mar Lett* 23(1):7-18
- Mulder T, Lecroart P, Hanquiez V, Marches E, Gonthier E, Guedes JC, Thiébot E, Jaaidi B, Kenyon N, Voisset M, Perez C, Sayago M, Fuchey Y, Bujan S (2006) The western part of the Gulf of Cadix: contour currents and turbidity currents interactions. *Geo-Mar Lett* 26:31-41
- Nelson CH, Maldonado A (eds) (1990) *The Ebro Continental Margin, Northwestern Mediterranean*. *Mar Geol* 95(3/4) pp 286
- Nelson CH, Baraza J, Maldonado A (1993) Mediterranean undercurrent sandy contourites, Gulf of Cadix. Spain. *Sed Geol* 82:103-131
- Nelson CH, Baraza J, Rodero J, Maldonado A, Escutia C, Barber Jr JH (1999) Influence of the Atlantic inflow and Mediterranean outflow currents on Late Pleistocene and Holocene sedimentary facies of Gulf of Cadix continental margin. *Mar Geol* 155:99-129
- Ochoa J, Bray NA (1991) Water mass exchange in the Gulf of Cadix. *Deep Sea Res* 38(1): 465-S503
- OSPAR Commission (2000) *Quality Status Report 2000. Region IV. Bay of Biscay and Iberian Coast*. OSPAR commission London, pp 132
- OSPAR (2006) *Initial OSPAR List of Threatened and/or Declining Species and Habitats* (www.ospar.org).
- Øvrebø LK, Houghton PDW, Shannon PM (2006) A record of fluctuating bottom currents on the slopes west

of the Porcupine Bank, offshore Ireland — implications for Late Quaternary climate forcing. *Mar Geol* 225:279-309

- 1 Paillet J, Mercier H (1997) An inverse model of the eastern North Atlantic general circulation and
2 thermocline ventilation. *Deep Sea Res Part I-Oceanogr Res Pap* 44(8):1293-1328
3
- 4 Paillet J, Arhan M, MacCartney M S (1998) Spreading of Labrador Sea Water in the eastern North Atlantic. *J*
5 *Geophys Res* 103(C5):10223-10239
6
- 7 Paillet J, Le Cann B, Serpette A, Morel Y, Carton X (1999) Real-time tracking of a Galician Meddy. *Geophys*
8 *Res Lett* 26:1877-1880
9
- 10 Palanques A, Kenyon NH, Alonso B, Limonov A (1995) Erosional and depositional patterns in the Valencia
11 channel mouth: An example of a modern channellobe transition zone. *Mar Geophys Res* 17:503-517
- 12 Palomino D, Vázquez JT, Ercilla G, Alonso B, López-González N, Díaz del Río V, Fernández-Salas LM (2010)
13 Influence of water masses on the seabed morphology around the Seamounts of the Motril Marginal
14 Shelf (Alboran Sea, Western Mediterranean). *Geo-Temas* 11:131-132
15
- 16 Parrilla G, Kinder TH (1987) Oceanografía física del mar de Alborán. *Boletín Instituto Español de*
17 *Oceanografía* 4:133-165
18
- 19 Pérez-Belzuz F (1999) Geología del Margen y Cuenca del Mar de Alborán durante el Plio-Cuaternario:
20 Sedimentación y tectónica. PhD Thesis, University of Barcelona, pp 538
21
- 22 Pérez-Belzuz F, Alonso B, Ercilla G (1997) History of mud diapirism and trigger mechanisms in the Western
23 Alboran Sea. *Tectonophysics* 282:399-422
24
- 25 Perkins H, Kinders T, Violette P (1990) The Atlantic inflowing the Western Alboran Sea. *J Phys Oceanogr*
26 20:242-263
27
- 28 Pickering KT, Hiscott RN, Hein FJ (1989) Deep Marine Environments. *Clastic Sedimentation and Tectonics*.
29 Unwin Hyman Ltd. London, pp 416
30
- 31 Pingree RD, Le Cann B (1990) Structure, strength and seasonality of the slope currents in the Bay of Biscay
32 region. *J Mar Biol Assoc U K* 70:857-885
33
- 34 Pollard RT, Pu S (1985) Structure and circulation of the upper Atlantic Ocean northeast of the Azores. *Progr*
35 *Oceanogr* 14:443-462
36
- 37 Pollard S, Griffiths CR, Cunningham SA, Read JF, Perez FF, Ríos AF (1996) Vivaldi 1991 - A study of the
38 formation, circulation and ventilation of Eastern North Atlantic Central Water. *Progr Oceanogr* 37:167-
39 192
40
- 41 Puig P, Palanques A, Guillén J, El Khatab M (2004) Role of internal waves in the generation of nepheloid
42 layers on the northwestern Alboran slope: Implications for continental margin shaping. *J Geophys Res*
43 109, C09011, doi:10.1029/2004JC002394
44
- 45 Rebesco M (2005) Contourites. In: Richard C, Selley RC, Cocks LRM, Plimer IR (eds) *Encyclopedia of Geology*,
46 vol 4, Elsevier, London, pp 513-527
47
- 48 Rebesco M, Stow D (2001) Seismic expression of contourites and related deposits: a preface. *Mar Geophys*
49 *Res* 22:303-308
50
- 51 Reid JL (1994) On the total contribution of the Mediterranean Sea outflow to the Norwegian-greenland Sea.
52 *Deep Sea Res* 26A:1199-1223
53
- 54 Reveillaud J, Freiwald A, Van Rooij D, Le Guilloux E, Altuna A, Foubert A, Vanreusel A, Olu-Le Roy K, Henriët
55 J-P (2008) The distribution of scleractinian corals in the Bay of Biscay, NE Atlantic. *Facies* 54:317-331
56
- 57 Richardson PL, Bowe, AS, Zenk W (2000) A census of Meddies tracked by floats. *Progr Oceanogr* 45:209-250
58
- 59 Roberts JM, Wheeler AJ, Freiwald A (2006) Reefs of the Deep: The Biology and Geology of Cold-Water Coral
60 Ecosystems. *Science* 312:543-547
61
- 62 Ryan WBF, Hsü KJ, Cita MB, Dumitricia P, Lort J, Maync W, Nesteroff WD, Pautot G, Stradner H, Wezel FC
63 (1973) Western Alboran Basin-Site 121. *Initial Rep DSDP* 13: 43-89
64
65

- 1 Sánchez R, Ruiz Villarreal M, González Pola C, Diaz del Río G, Cabanas JM, González N, Rodríguez M, Alonso
2 J, Plaza F, Lavín A (2006) Labrador Sea Water in the Bay of Biscay: analysis of historical data (1990-
3 2005). X International Symposium on Oceanography of the Bay of Biscay, April 19-21 2006. Vigo,
4 Galicia. Spain. Abstract volume, 1-4.
- 5 Sarnthein M, Statterger K, Dreger D, Erlenkeuser H, Grootes P, Haupt BJ, Jung S, Kiefer T, Kuhnt W,
6 Pflaumann U, Schäfer-Neth C, Schulz H, Schulz M, Seidov D, Simstich J, van Kreveld S, Vogelsang E,
7 Völker A, Weinelt M (2000) Fundamental Modes and Abrupt Changes in North Atlantic Circulation and
8 Climate over the last 60 ky – Concepts, Reconstruction and Numerical Modeling. In: Schäfer P, Ritzrau
9 W, Schlüter M, Thiede J (eds) The Northern North Atlantic: A Changing Environment, Springer, Berlin,
10 365-410
- 11 Schönfeld J, Zahn R (2000) Late Glacial to Holocene history of the Mediterranean Outflow. Evidence from
12 benthic foraminiferal assemblages and stable isotopes at the Portuguese margin. *Palaeogeogr*
13 *Palaeoclimatol Palaeoecol* 159:85-111
- 14 Schönfeld J, Zahn R, Abreu L (2003) Surface and deep water response to rapid climate changes at the
15 Western Iberian Margin. *Global Planet Change* 36:237-264
- 16 Seibold E, Berger WH (1993) *The Sea Floor. An Introduction to Marine Geology*. 2d Edition. Springer Verlag,
17 pp 336
- 18 Serra N (2004) Observations and numerical modelling of the Mediterranean Outflow. Ph.D. Thesis,
19 University of Lisbon, pp 234
- 20 Serra N, Käse RH, Köhl A, Stammer D, Quadfasel D (2010a) On the low-frequency phase relation between
21 the Denmark Strait and the Faroe-Bank Channel overflows. *Tellus* 62A:530-550
- 22 Serra N, Ambar I, Boutov D (2010b) Surface expression of Mediterranean Water dipoles and their
23 contribution to the shelf/slope – open ocean exchange- *Ocean Sci* 6:191-209
- 24 Serrano A, Maestro A, Nozal F (Coord) (2005) Memoria del mapa Geomorfológico de España a escala
25 1:1000.000: Geomorfología de la zona submarina española. Spanish Geological Survey (Instituto
26 Tecnológico Geominero de España, ITGE), pp 232
- 27 Shackleton NJ, Hall MA, Vincent E (2000) Phase relationships between millennial scale events 64,000 to
28 24,000 years ago. *Paleoceanography* 15:565-569
- 29 Shanmugam G (2007) Deep-water processes and facies models: implications for sandstone petroleum
30 reservoirs. *Handbook of Petroleum Exploration and Production* 5, Elsevier, Amsterdam, pp 476
- 31 Slater DR (2003) The transport of Mediterranean water in the north Atlantic Ocean. PhD Thesis, Univ. of
32 Southampton., pp 155
- 33 Stanley DJ, Kelling G, Vera JA, Slieng H (1975) Sands in the Alborán Sea: A model of input in a deep marine
34 basin. *Earth Sci* 15:51
- 35 Stow DAV, Faugères JC, Gonthier E (1986) Facies distribution and textural variations in Faro drift
36 contourites: velocity fluctuation and drift growth. *Mar Geol* 72:71-100
- 37 Stow DAV, Pudsey CJ, Howe JA, Faugères JC, Viana A (eds) (2002a) Deep-water contourite systems: Modern
38 drifts and ancient series, seismic and sedimentary characteristics. *Geol Soc London Memoir* 22, pp 464
- 39 Stow DAV, Faugères JC, Howe JA, Pudsey CJ, Viana A (2002b) Contourites, bottom currents and deep-sea
40 sediment drifts: current state-of-the-art. *Geol. Soc. London Memoir* 22: -20
- 41 Stow DAV, Faugères JC, Gonthier E, Llave E, Hernández-Molina FJ, Somoza L, Díaz del Río V (2002c) Faro-
42 Albufeira drift complex, northern Gulf of Cadiz. *Geol Soc London Memoir* 22: 37-154
- 43 Stow DAV, Hunter S, Wilkinson D, Hernández-Molina FJ (2008) The nature of contourite deposition. In:
44 Rebesco M, Camerlenghi A (eds) *Contourites. Developments in Sedimentology*, 60. Elsevier, pp 143-156
- 45 Stow DAV, Hernández-Molina F.J, Llave E, Sayago M, Díaz del Río V, Branson A (2009) Bedform-velocity
46 matrix: the estimation of bottom current velocity from bedform observations. *Geology* 37(4):327-330
- 47 Teleki PG, Dabson MR, Moore JR, Von Stackelberg V (1987) *Marine Minerals. Advances in Research and*
48 *Resource Assessment*. D. Reidel Publishing Company. Nato ASI Series 194, pp 588
- 49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Thorpe SA (1975) Variability of the Mediterranean in the Gulf of Cadiz. *Deep Sea Res* 23:711-727
- Thunell R, Rio D, Sprovieri R, Vergnaud-Grazzini C (1991) An overview of the post-Messinian paleoenvironmental history of the Mediterranean. *Paleoceanography* 6(1):143-164
- Valencia V, Franco J, Borja A, Fontan A (2004) Hydrography of the southeastern Bay of Biscay. In: Borja A, Collins, MB (eds) *Oceanography and Marine Environment of the Basque Country*. Elsevier Oceanography Series, Amsterdam, pp 159-193
- Van Aken HM (2000) The hydrography of the mid-latitude Northeast Atlantic Ocean II: The intermediate water masses. *Deep Sea Res I* 47:789-824
- Van Aken HM (2002) Surface currents in the Bay of Biscay as observed with drifters between 1995 and 1999. *Deep Sea Res I* 49:1071-1086
- Van Andel Tj H, Thiede J, Sclater JG, Hay WW (1977) Depositional History of the South Atlantic during the last 125 Million Years. *J Geol* 85:651-98
- Van Rooij D, De Mol B, Huvenne V, Ivanov M, Henriët JP (2003) Seismic evidence of current-controlled sedimentation in the Belgica mound province, upper Porcupine slope, southwest of Ireland. *Marine Geology* 195: 31-53
- Van Rooij D, Iglesias J, Hernandez-Molina FJ, Ercilla G, Gomez-Ballesteros M, De Hauwere A, Casas D, Llave E, Garcia Gil S, Henriët JP, Acosta J (2010a) The Le Danois Contourite Depositional System off the North Iberian margin, Spain. *Mar Geol* 274(1-4):1-20
- Van Rooij D, De Mol L, Le Guilloux E, Réveillaud J, Hernández-Molina FJ, Llave E, Léon R, Estrada F, Mienis F, Moeremans R, Blamart D, Vanreusel A, Henriët JP (2010b) Influence of the Mediterranean Outflow Water on benthic ecosystems: answers and questions after a decade of observations. *Geo-Temas* 11:179-180
- Vázquez JT (2001) Estructura del margen septentrional del Mar de Alboran. Tesis Doctoral. Universidad Complutense. Madrid, pp 422
- Velasco JPB, Baraza J, Canals M, Balón J (1996) La depresión periférica y el lomo contourítico de Menorca: evidencias de la actividad de corrientes de fondo al N del talud Balear. *Geogaceta* 20(2):359-362
- Vernaud-Grazzini C, Caralp M-H, Faugères JC, Gonthier E, Grousset F, Pujol C, Saliège JF (1989) Mediterranean outflow through the Strait of Gibraltar since 18.000 years B.P. *Oceanological Acta* 12:305-324
- Viana A (2008) Economic Relevance of Contourites. In: Rebesco M, Camerlenghi A (eds) *Contourites. Developments in Sedimentology* 60. Elsevier, pp 493-510
- Viana A (2010) The economic importance of contourites. *Geo-Temas* 11:181-182
- Viana A, Rebesco M (eds) 2007 Economic and Paleoceanographic Significance of Contourites. *Geol Soc London Sp Publ* 276, pp 350
- Viana AR, Almeida Jr. W, Almeida CFW (2002a) Upper slope sands – the late Quaternary shallow-water sandy contourites of Campos Basin SW Atlantic margin. In: Stow DAV, Pudsey CJ, Howe JA, Faugères JC, Viana AR (eds) *Deep-Water Contourite Systems: Modern Drifts and Ancient Series, Seismic and Sedimentary Characteristics*. *Geol Soc London Mem* 22, pp 261-270
- Viana AR, Hercos CM, Almeida Jr. W, Magalhães JLC, Andrade SB (2002b) Evidence of bottom current Influence on the Neogene to Quaternary sedimentation along the Northern Campos Slope, SW Atlantic Margin. In: Stow DAV, Pudsey CJ, Howe JA, Faugères JC, Viana AR (eds) *Deep-Water Contourite Systems: Modern Drifts and Ancient Series, Seismic and Sedimentary Characteristics*. *Geol Soc London Mem* 22, pp 249-259
- Voelker A, Lebreiro S, Schönfeld J, Cacho I, Exlenkenser H, Abrantes F (2006) Mediterranean outflow strengthenings during Northern Hemisphere Coolings: a salt sources for the glacial Atlantic?. *Earth Planet Sci Lett* 245:39-55
- Voelker AHL, Lebreiro S (2010) Millennial-scale changes in deep water properties at the mid-depth western Iberian margin linked to Mediterranean Outflow Water activity *Geo-Temas* 11: 85-186.

- Weaver PPE, Wynn RB, Kenyon NH, Evans J (2000) Continental margin sedimentation, with special reference to north-east Atlantic margin. *Sedimentology* 47:239-256
- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Wüst G. (1936) Schichtung und Zirkulation des Atlantischen Ozeans: Das Bodenwasser und die Gliederung der Atlantischen Tiefsee. *Wissenschaftliche Ergebnisse der Deutschen Atlantischen Expedition "Meteor" 1925-1927 vol. VI part 1: 3-107*
- Wüst G (1961) Das Bodenwasser und die Vertikalzirkulation des Mittelländischen Meeres. *Deutsche Hydrographische Zeitschrift* 14:81-92
- Wynn, R.B., Masson, D.G., Stow, D.A.V., Weaver, P.P.E., 2000. The Northwest African slope apron: a modern analogue for deep-water systems with complex seafloor topography. *Mar Pet Geol* 17:253-265
- Zahn R, Comas MC, Klaus A (eds) (1999) *Proceedings of the Ocean Drilling Program, Scientific Results. 161*, College Station, TX (Ocean Drilling Program), [doi:10.2973/odp.proc.sr.161.1999](https://doi.org/10.2973/odp.proc.sr.161.1999).
- Zenk W (1975) On the Mediterranean outflow west of Gibraltar. *"Meteor" Forsch.-Ergebnisse*, 16:23-34
- Zenk W, Armi L (1990) The complex spreading patterns of Mediterranean water off the Portuguese continental slope. *Deep Sea Res* 37:1805-1823
- Zhenzhong G, Eriksson KA, He YB, Luo SS, Guo JH (1998) *Deep-Water Traction Current Deposits- A Study of Internal Tides, Internal Waves, Contour Currents and Their Deposits*. Science Press, Beijing, New York, pp 128

1 **Figures**

2
3 **Figure captions**

4
5
6 Figure 1. Area Location and bathymetric map with continental margins and abyssal plains around Iberia
7 [Base bathymetric map from R. Leon (Pers. Comm.), realised from satellite data from Smith and
8 Sandwell, 1997]. Legend of the main physiographic reference points, in alphabetical order: ASM =
9 Ampere Seamount; ArSM= Armorican Seamount; BH = Barbate high; BSM= Biscay Seamount; CK=
10 Cantabrian Knoll; CPSM= Coral Patch Seamount; DSM = Dragon Seamount; FSM= Finisterre
11 Seamount; GaB= Galicia Bank; GB = Guadalquivir Bank; GK= Gascone Knoll; GoB= Goringe Bank;
12 JoSM= Jovellanos Seamount; JSM = Josephine Seamount; HSM = Hirondele Seamount; LCSM= La
13 Coruña Seamount; LSM = Lion Seamount; NASM= North Atlante Seamount; NCSM= North Charcot
14 Seamount; OSM= Oporto Seamount; SASM= South Atlante Seamount; SCSM= South Charcot
15 Seamount; SSM = Sea Seamount; TSM= Tore Seamount; USM = Unicorn Seamount; VGSM= Vasco
16 de Gama Seamount; VSM= Vigo Seamount.

17
18
19
20
21 Figure 2. Mediterranean Sea sketch with the general superficial water mass circulation, the winter
22 convection areas, and the main water mass formation zones (*Adapted from Arnone et al., 1990 and*
23 *Candela, 2001*).

24
25
26
27 Figure 3. Figure 3. Water masses compilation both the Western Mediterranean and North Atlantic Sea
28 gathered from bibliographic sources: A) Deep water circulation; B) Intermediate water Circulation
29 and C) Superficial water circulation (see the text for further explication). This compilation is based
30 on data and ideas from all the following authors: Wüst, 1936, 1961; Madelain, 1970; Kelling and
31 Stanley, 1972; Kenyon and Belderson, 1973; Melières, 1974; Stanley et al., 1975; Thorpe, 1975;
32 Zenk, 1975; Ambar and Howe, 1979; McCartney and Talley, 1982; Gardner and Kidd, 1983, 1987;
33 Bryden and Stommel, 1984; Gascard and Richez, 1984; Fiúza, 1984; Dickson et al., 1985; Pollard and
34 Pu, 1985; Parrilla and Kinder, 1987; Caralp, 1988, 1992; Garrett et al., 1990; Haynes and Barton,
35 1990; Herburn and La Violette, 1990; Perkins et al., 1990; Pingree and Le Cann, 1990; Zenk and
36 Armi, 1990; Arham et al., 1994; Cano and García, 1991; Ochoa and Bray, 1991; McCartney, 1992;
37 Nelson et al., 1993, 1999; Ambar and Fiúza, 1994; Bryden et al., 1994; Danialt et al., 1994; Reid,
38 1994; Pollard et al., 1996; Mazé et al., 1997; Paillet and Mercier, 1997; Díaz del Río et al., 1998;
39 Fiuza et al., 1998; Paillet et al., 1998, 1999; 2001; Baringer and Price, 1999; Ioga and Lozier, 1999;
40 Millot, 1999; Cherubin et al., 2000; Johnson and Stevens, 2000; OSPAR 2000; Richardson et al.,
41 2000; Sarnthein et al., 2000; Weaver et al., 2000; Candela, 2001; Van Aken, 2000, 2002; Mauritzen
42 et al., 2001; Borenäs et al., 2002; Hernández-Molina et al., 2002, 2006; Huthnance et al., 2002;
43 Slater, 2003; Serra, 2004; Valencia et al., 2004; Díaz del Río, 2006; González Pola, 2006; Sánchez et
44 al., 2006; García Lafuente et al., 2008; García et al., 2009; Legg et al., 2009; Serra et al., 2010b; Van
45 Rooij et al., 2010a; Stow et al., in preparation). Legend of the main water masses, in alphabetical
46 order: AABW= Antarctic Bottom Water; AC= Atlantic Current; AI= Atlantic Inflow; LDW= Lower Deep
47 Water; LADW= Labrador Deep Water; LIW= Levantine Intermediate Water; LSW= Labrador Sea
48 Water; IPC= Iberian Polar Current; MAW= Modified Atlantic Water; MOW= Mediterranean Outflow
49 Water; NAC= North Atlantic Current; NACW/ENACW= North Atlantic Central Water; NADW= North
50 Atlantic Deep Water; SAIW = Subarctic Intermediate Water; and WMDW= Western Mediterranean
51 Deep Water.

52
53
54
55
56
57
58 Figure 4. Simulated bottom velocities (cm/s) around Iberia.

59
60 Figure 5. Time average of simulated bottom velocities (cm/s) around Iberia.

61
62
63
64
65

Figure 6. Main area locations with dominate *along-slope* (contourite) processes and identified Contourite Depositional Systems (CDS) within the continental margins and abyssal plains around Iberia.


Figure 7. Examples of contourite features (depositional & erosive) generated within the identified Contourite Depositional Systems (CDS): A) Airgun seismic profile across the Menorca CDS (Velasco et al., 1996); B) Airgun seismic profile oblique to the Ceuta plastered drift across its central sector (Ercilla et al., 2002); C) Uninterpreted multichannel seismic-reflection (MCS) profile across the sandy drifts closer to the Strait of Gibraltar and along the middle slope of the Gulf of Cadiz CDS (line S-81A, provided by REPSOL-YPF Oil Company); D) Sparker seismic profiles showing examples of a plastered drift along the upper and middle slope of the Gulf of Cadiz CDS (Hernández-Molina et al., 2006; Llave et al., 2007, 2010); E) Uninterpreted MCS profile across the Faro-Albufeira mounded elongated and separated drift and the Alvarez Cabral moat on the middle slope of the Gulf of Cadiz CDS (data courtesy of TGS-NOPEC Geophysical Company ASA). It represents a classic example of middle-slope contourite deposition (Hernández-Molina et al., 2006; Llave et al., 2007, 2010); F) Uninterpreted MCS profile across the possible plastered drifts deposits defined by Alves et al. (2003) on the Portuguesse middle slope (data courtesy of Luis Pinheiro, Pers. Comm.); G) & H) Uninterpreted Airgun seismic profiles at Galicia Bank CDS, showing examples of elongated mounded & separated drift at the base of the slope of the Galicia Bank Plateau (G) and a plastered drift deposits on top of that Bank (H); I) High-resolution sparker profile across the Ortegaleira Spur CDS; J) High-resolution sparker profile across the Le Danois elongated mounded & separated drift at the base of the Le Danois Bank (Van Rooij et al., 2010); K) & L) Sparker seismic profiles on the Porcupine Bank indicating the seismic features and unit geometry on the eastern slope of the Porcupine Seabight (Van Rooij et al., 2003). For the approximate location of the profiles, see Fig. 6.

Figure 8. Combining numerically simulated bottom currents with existing knowledge of contourite features around Iberia and major gravitational areas.

Combining numerically simulated bottom currents with existing regional knowledge of the main areas locations with dominant *along-slope* (contourite) and *across-slope* (mass wasting, turbiditic, & gravitational) processes identified within the continental margins and abyssal plains around Iberia. Across-slope processes and associated products compilation is mainly based on data and ideas from Weaver et al. (2000) and Serrano et al. (2005).

Figure 9. Potential areas where in theory new contourites features (depositional & erosive) could be presented, within the continental margins and abyssal plains around Iberia. Also the best deep and intermediate water areas with major interest for future explorations are selected.

Figure
[Click here to download high resolution image](#)



Figure

[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)




Figure
[Click here to download high resolution image](#)





Figure
[Click here to download high resolution image](#)



- Previously described CDS
- Potential intermediate water areas where contourite features could be presented
- Potential deep water areas where contourite features could be presented
- Intermediate water areas with major interest for future explorations
- Deep water areas with major interest for future explorations