

Observation of the Baryonic Flavor-Changing Neutral Current Decay $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$

T. Aaltonen,²² B. Álvarez González,^{10,w} S. Amerio,^{42a} D. Amidei,³³ A. Anastassov,³⁷ A. Annovi,¹⁸ J. Antos,¹³ G. Apollinari,¹⁶ J. A. Appel,¹⁶ A. Apresyan,⁴⁷ T. Arisawa,⁵⁶ A. Artikov,¹⁴ J. Asaadi,⁵² W. Ashmanskas,¹⁶ B. Auerbach,⁵⁹ A. Aurisano,⁵² F. Azfar,⁴¹ W. Badgett,¹⁶ A. Barbaro-Galtieri,²⁷ V. E. Barnes,⁴⁷ B. A. Barnett,²⁴ P. Barria,^{45c,45a} P. Bartos,¹³ M. Baucé,^{42b,42a} G. Bauer,³¹ F. Bedeschi,^{45a} D. Beecher,²⁹ S. Behari,²⁴ G. Bellettini,^{45b,45a} J. Bellinger,⁵⁸ D. Benjamin,¹⁵ A. Beretvas,¹⁶ A. Bhatti,⁴⁹ M. Binkley,^{16,a} D. Bisello,^{42b,42a} I. Bizjak,^{29,cc} K. R. Bland,⁵ C. Blocker,⁷ B. Blumenfeld,²⁴ A. Bocci,¹⁵ A. Bodek,⁴⁸ D. Bortoletto,⁴⁷ J. Boudreau,⁴⁶ A. Boveia,¹² B. Brau,^{16,b} L. Brigliadori,^{6b,6a} A. Brisuda,¹³ C. Bromberg,³⁴ E. Brucken,²² M. Bucciantonio,^{45b,45a} J. Budagov,¹⁴ H. S. Budd,⁴⁸ S. Budd,²³ K. Burkett,¹⁶ G. Busetto,^{42b,42a} P. Bussey,²⁰ A. Buzatu,³² S. Cabrera,^{15,y} C. Calancha,³⁰ S. Camarda,⁴ M. Campanelli,³⁴ M. Campbell,³³ F. Canelli,^{12,16} A. Canepa,⁴⁴ B. Carls,²³ D. Carlsmith,⁵⁸ R. Carosi,^{45a} S. Carrillo,^{17,l} S. Carron,¹⁶ B. Casal,¹⁰ M. Casarsa,¹⁶ A. Castro,^{6b,6a} P. Catastini,¹⁶ D. Cauz,^{53a} V. Cavaliere,^{45c,45a} M. Cavalli-Sforza,⁴ A. Cerri,^{27,g} L. Cerrito,^{29,r} Y. C. Chen,¹ M. Chertok,⁸ G. Chiarelli,^{45a} G. Chlachidze,¹⁶ F. Chlebana,¹⁶ K. Cho,²⁶ D. Chokheli,¹⁴ J. P. Chou,²¹ W. H. Chung,⁵⁸ Y. S. Chung,⁴⁸ C. I. Ciobanu,⁴³ M. A. Ciocci,^{45c,45a} A. Clark,¹⁹ D. Clark,⁷ G. Compostella,^{42b,42a} M. E. Convery,¹⁶ J. Conway,⁸ M. Corbo,⁴³ M. Cordelli,¹⁸ C. A. Cox,⁸ D. J. Cox,⁸ F. Crescioli,^{45b,45a} C. Cuenca Almenar,⁵⁹ J. Cuevas,^{10,w} R. Culbertson,¹⁶ D. Dagenhart,¹⁶ N. d'Ascenzo,^{43,u} M. Datta,¹⁶ P. de Barbaro,⁴⁸ S. De Cecco,^{50a} G. De Lorenzo,⁴ M. Dell'Orso,^{45b,45a} C. Deluca,⁴ L. Demortier,⁴⁹ J. Deng,^{15,d} M. Deninno,^{6a} F. Devoto,²² M. d'Errico,^{42b,42a} A. Di Canto,^{45b,45a} B. Di Ruzza,^{45a} J. R. Dittmann,⁵ M. D'Onofrio,²⁸ S. Donati,^{45b,45a} P. Dong,¹⁶ T. Dorigo,^{42a} K. Ebina,⁵⁶ A. Elagin,⁵² A. Eppig,³³ R. Erbacher,⁸ D. Errede,²³ S. Errede,²³ N. Ershaidat,^{43,bb} R. Eusebi,⁵² H. C. Fang,²⁷ S. Farrington,⁴¹ M. Feindt,²⁵ J. P. Fernandez,³⁰ C. Ferrazza,^{45d,45a} R. Field,¹⁷ G. Flanagan,^{47,s} R. Forrest,⁸ M. J. Frank,⁵ M. Franklin,²¹ J. C. Freeman,¹⁶ I. Furic,¹⁷ M. Gallinaro,⁴⁹ J. Galyardt,¹¹ J. E. Garcia,¹⁹ A. F. Garfinkel,⁴⁷ P. Garosi,^{45c,45a} H. Gerberich,²³ E. Gerchtein,¹⁶ S. Giagu,^{50b,50a} V. Giakoumopoulou,³ P. Giannetti,^{45a} K. Gibson,⁴⁶ C. M. Ginsburg,¹⁶ N. Giokaris,³ P. Giromini,¹⁸ M. Giunta,^{45a} G. Giurgiu,²⁴ V. Glagolev,¹⁴ D. Glenzinski,¹⁶ M. Gold,³⁶ D. Goldin,⁵² N. Goldschmidt,¹⁷ A. Golossanov,¹⁶ G. Gomez,¹⁰ G. Gomez-Ceballos,³¹ M. Goncharov,³¹ O. González,³⁰ I. Gorelov,³⁶ A. T. Goshaw,¹⁵ K. Goulianos,⁴⁹ A. Gresele,^{42a} S. Grinstein,⁴ C. Grosso-Pilcher,¹² R. C. Group,¹⁶ J. Guimaraes da Costa,²¹ Z. Gunay-Unalan,³⁴ C. Haber,²⁷ S. R. Hahn,¹⁶ E. Halkiadakis,⁵¹ A. Hamaguchi,⁴⁰ J. Y. Han,⁴⁸ F. Happacher,¹⁸ K. Hara,⁵⁴ D. Hare,⁵¹ M. Hare,⁵⁵ R. F. Harr,⁵⁷ K. Hatakeyama,⁵ C. Hays,⁴¹ M. Heck,²⁵ J. Heinrich,⁴⁴ M. Herndon,⁵⁸ S. Hewamanage,⁵ D. Hidas,⁵¹ A. Hocker,¹⁶ W. Hopkins,^{16,h} D. Horn,²⁵ S. Hou,¹ R. E. Hughes,³⁸ M. Hurwitz,¹² U. Husemann,⁵⁹ N. Hussain,³² M. Hussein,³⁴ J. Huston,³⁴ G. Introzzi,^{45a} M. Iori,^{50b,50a} A. Ivanov,^{8,p} E. James,¹⁶ D. Jang,¹¹ B. Jayatilaka,¹⁵ E. J. Jeon,²⁶ M. K. Jha,^{6a} S. Jindariani,¹⁶ W. Johnson,⁸ M. Jones,⁴⁷ K. K. Joo,²⁶ S. Y. Jun,¹¹ T. R. Junk,¹⁶ T. Kamon,⁵² P. E. Karchin,⁵⁷ Y. Kato,^{40,o} W. Ketchum,¹² J. Keung,⁴⁴ V. Khotilovich,⁵² B. Kilminster,¹⁶ D. H. Kim,²⁶ H. S. Kim,²⁶ H. W. Kim,²⁶ J. E. Kim,²⁶ M. J. Kim,¹⁸ S. B. Kim,²⁶ S. H. Kim,⁵⁴ Y. K. Kim,¹² N. Kimura,⁵⁶ S. Klimentenko,¹⁷ K. Kondo,⁵⁶ D. J. Kong,²⁶ J. Konigsberg,¹⁷ A. Korytov,¹⁷ A. V. Kotwal,¹⁵ M. Kreps,²⁵ J. Kroll,⁴⁴ D. Krop,¹² N. Krumnack,^{5,m} M. Kruse,¹⁵ V. Krutelyov,^{52,e} T. Kuhr,²⁵ M. Kurata,⁵⁴ S. Kwang,¹² A. T. Laasanen,⁴⁷ S. Lami,^{45a} S. Lammel,¹⁶ M. Lancaster,²⁹ R. L. Lander,⁸ K. Lannon,^{38,v} A. Lath,⁵¹ G. Latino,^{45c,45a} I. Lazzizzera,^{42a} T. LeCompte,² E. Lee,⁵² H. S. Lee,¹² J. S. Lee,²⁶ S. W. Lee,^{52,x} S. Leo,^{45b,45a} S. Leone,^{45a} J. D. Lewis,¹⁶ C.-J. Lin,²⁷ J. Linacre,⁴¹ M. Lindgren,¹⁶ E. Lipeles,⁴⁴ A. Lister,¹⁹ D. O. Litvintsev,¹⁶ C. Liu,⁴⁶ Q. Liu,⁴⁷ T. Liu,¹⁶ S. Lockwitz,⁵⁹ N. S. Lockyer,⁴⁴ A. Loginov,⁵⁹ D. Lucchesi,^{42b,42a} J. Lueck,²⁵ P. Lujan,²⁷ P. Lukens,¹⁶ G. Lungu,⁴⁹ J. Lys,²⁷ R. Lysak,¹³ R. Madrak,¹⁶ K. Maeshima,¹⁶ K. Makhoul,³¹ P. Maksimovic,²⁴ S. Malik,⁴⁹ G. Manca,^{28,c} A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁷ C. Marino,²⁵ M. Martínez,⁴ R. Martínez-Ballarín,³⁰ P. Mastrandrea,^{50a} M. Mathis,²⁴ M. E. Mattson,⁵⁷ P. Mazzanti,^{6a} K. S. McFarland,⁴⁸ P. McIntyre,⁵² R. McNulty,^{28,j} A. Mehta,²⁸ P. Mehtala,²² A. Menzione,^{45a} C. Mesropian,⁴⁹ T. Miao,¹⁶ D. Mietlicki,³³ A. Mitra,¹ H. Miyake,⁵⁴ S. Moed,²¹ N. Moggi,^{6a} M. N. Mondragon,^{16,l} C. S. Moon,²⁶ R. Moore,¹⁶ M. J. Morello,¹⁶ J. Morlock,²⁵ P. Movilla Fernandez,¹⁶ A. Mukherjee,¹⁶ Th. Muller,²⁵ P. Murat,¹⁶ M. Mussini,^{6b,6a} J. Nachtman,^{16,n} Y. Nagai,⁵⁴ J. Naganoma,⁵⁶ I. Nakano,³⁹ A. Napier,⁵⁵ J. Nett,⁵⁸ C. Neu,^{44,aa} M. S. Neubauer,²³ J. Nielsen,^{27,f} L. Nodulman,² O. Norriella,²³ E. Nurse,²⁹ L. Oakes,⁴¹ S. H. Oh,¹⁵ Y. D. Oh,²⁶ I. Oksuzian,¹⁷ T. Okusawa,⁴⁰ R. Orava,²² L. Ortolan,⁴ S. Pagan Griso,^{42b,42a} C. Pagliarone,^{53a} E. Palencia,^{10,g} V. Papadimitriou,¹⁶ A. A. Paramonov,² J. Patrick,¹⁶ G. Pauletta,^{53b,53a} M. Paulini,¹¹ C. Paus,³¹ D. E. Pellett,⁸ A. Penzo,^{53a} T. J. Phillips,¹⁵ G. Piacentino,^{45a} E. Pianori,⁴⁴ J. Pilot,³⁸ K. Pitts,²³ C. Plager,⁹ L. Pondrom,⁵⁸ K. Potamianos,⁴⁷ O. Poukhov,^{14,a} F. Prokoshin,^{14,z} A. Pronko,¹⁶ F. Ptohos,^{18,i} E. Pueschel,¹¹ G. Punzi,^{45b,45a} J. Pursley,⁵⁸ A. Rahaman,⁴⁶ V. Ramakrishnan,⁵⁸ N. Ranjan,⁴⁷ I. Redondo,³⁰ P. Renton,⁴¹ M. Rescigno,^{50a} F. Rimondi,^{6b,6a} L. Ristori,^{45a,16} A. Robson,²⁰ T. Rodrigo,¹⁰ T. Rodriguez,⁴⁴ E. Rogers,²³

S. Rolli,⁵⁵ R. Roser,¹⁶ M. Rossi,^{53a} F. Ruffini,^{45c,45a} A. Ruiz,¹⁰ J. Russ,¹¹ V. Rusu,¹⁶ A. Safonov,⁵² W. K. Sakumoto,⁴⁸ L. Santi,^{53b,53a} L. Sartori,^{45a} K. Sato,⁵⁴ V. Saveliev,^{43,u} A. Savoy-Navarro,⁴³ P. Schlabach,¹⁶ A. Schmidt,²⁵ E. E. Schmidt,¹⁶ M. P. Schmidt,^{59,a} M. Schmitt,³⁷ T. Schwarz,⁸ L. Scodellaro,¹⁰ A. Scribano,^{45c,45a} F. Scuri,^{45a} A. Sedov,⁴⁷ S. Seidel,³⁶ Y. Seiya,⁴⁰ A. Semenov,¹⁴ F. Sforza,^{45b,45a} A. Sfyrila,²³ S. Z. Shalhout,⁸ T. Shears,²⁸ P. F. Shepard,⁴⁶ M. Shimojima,^{54,t} S. Shiraishi,¹² M. Shochet,¹² I. Shreyber,³⁵ A. Simonenko,¹⁴ P. Sinervo,³² A. Sissakian,^{14,a} K. Sliwa,⁵⁵ J. R. Smith,⁸ F. D. Snider,¹⁶ A. Soha,¹⁶ S. Somalwar,⁵¹ V. Sorin,⁴ P. Squillacioti,¹⁶ M. Stanitzki,⁵⁹ R. St. Denis,²⁰ B. Stelzer,³² O. Stelzer-Chilton,³² D. Stentz,³⁷ J. Strologas,³⁶ G. L. Strycker,³³ Y. Sudo,⁵⁴ A. Sukhanov,¹⁷ I. Suslov,¹⁴ K. Takemasa,⁵⁴ Y. Takeuchi,⁵⁴ J. Tang,¹² M. Tecchio,³³ P. K. Teng,¹ J. Thom,^{16,h} J. Thome,¹¹ G. A. Thompson,²³ E. Thomson,⁴⁴ P. Ttito-Guzmán,³⁰ S. Tkaczyk,¹⁶ D. Toback,⁵² S. Tokar,¹³ K. Tollefson,³⁴ T. Tomura,⁵⁴ D. Tonelli,¹⁶ S. Torre,¹⁸ D. Torretta,¹⁶ P. Totaro,^{53b,53a} M. Trovato,^{45d,45a} Y. Tu,⁴⁴ N. Turini,^{45c,45a} F. Ukegawa,⁵⁴ S. Uozumi,²⁶ A. Varganov,³³ E. Vataga,^{45d,45a} F. Vázquez,^{17,1} G. Velev,¹⁶ C. Vellidis,³ M. Vidal,³⁰ I. Vila,¹⁰ R. Vilar,¹⁰ M. Vogel,³⁶ G. Volpi,^{45b,45a} P. Wagner,⁴⁴ R. L. Wagner,¹⁶ T. Wakisaka,⁴⁰ R. Wallny,⁹ S. M. Wang,¹ A. Warburton,³² D. Waters,²⁹ M. Weinberger,⁵² H. Wenzel,¹⁶ W. C. Wester III,¹⁶ B. Whitehouse,⁵⁵ D. Whiteson,^{44,d} A. B. Wicklund,² E. Wicklund,¹⁶ S. Wilbur,¹² F. Wick,²⁵ H. H. Williams,⁴⁴ J. S. Wilson,³⁸ P. Wilson,¹⁶ B. L. Winer,³⁸ P. Wittich,^{16,h} S. Wolbers,¹⁶ H. Wolfe,³⁸ T. Wright,³³ X. Wu,¹⁹ Z. Wu,⁵ K. Yamamoto,⁴⁰ J. Yamaoka,¹⁵ T. Yang,¹⁶ U. K. Yang,^{12,q} Y. C. Yang,²⁶ W.-M. Yao,²⁷ G. P. Yeh,¹⁶ K. Yi,^{16,n} J. Yoh,¹⁶ K. Yorita,⁵⁶ T. Yoshida,^{40,k} G. B. Yu,¹⁵ I. Yu,²⁶ S. S. Yu,¹⁶ J. C. Yun,¹⁶ A. Zanetti,^{53a} Y. Zeng,¹⁵ and S. Zucchelli^{6b,6a}

(CDF Collaboration)

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*²*Argonne National Laboratory, Argonne, Illinois 60439, USA*³*University of Athens, 157 71 Athens, Greece*⁴*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*⁵*Baylor University, Waco, Texas 76798, USA*^{6a}*Istituto Nazionale di Fisica Nucleare Bologna, I-40127 Bologna, Italy*^{6b}*University of Bologna, I-40127 Bologna, Italy*⁷*Brandeis University, Waltham, Massachusetts 02254, USA*⁸*University of California, Davis, Davis, California 95616, USA*⁹*University of California, Los Angeles, Los Angeles, California 90024, USA*¹⁰*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*¹¹*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*¹²*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*¹³*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*¹⁴*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*¹⁵*Duke University, Durham, North Carolina 27708, USA*¹⁶*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*¹⁷*University of Florida, Gainesville, Florida 32611, USA*¹⁸*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*¹⁹*University of Geneva, CH-1211 Geneva 4, Switzerland*²⁰*Glasgow University, Glasgow G12 8QQ, United Kingdom*²¹*Harvard University, Cambridge, Massachusetts 02138, USA*²²*Division of High Energy Physics, Department of Physics,**University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*²³*University of Illinois, Urbana, Illinois 61801, USA*²⁴*The Johns Hopkins University, Baltimore, Maryland 21218, USA*²⁵*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*²⁶*Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757, Korea; Chonbuk National University, Jeonju 561-756, Korea*²⁷*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*²⁸*University of Liverpool, Liverpool L69 7ZE, United Kingdom*²⁹*University College London, London WC1E 6BT, United Kingdom*³⁰*Centro de Investigaciones Energéticas Medioambientales y Tecnológicas, E-28040 Madrid, Spain*³¹*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

- ³²*Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3*
- ³³*University of Michigan, Ann Arbor, Michigan 48109, USA*
- ³⁴*Michigan State University, East Lansing, Michigan 48824, USA*
- ³⁵*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
- ³⁶*University of New Mexico, Albuquerque, New Mexico 87131, USA*
- ³⁷*Northwestern University, Evanston, Illinois 60208, USA*
- ³⁸*The Ohio State University, Columbus, Ohio 43210, USA*
- ³⁹*Okayama University, Okayama 700-8530, Japan*
- ⁴⁰*Osaka City University, Osaka 588, Japan*
- ⁴¹*University of Oxford, Oxford OX1 3RH, United Kingdom*
- ^{42a}*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*
- ^{42b}*University of Padova, I-35131 Padova, Italy*
- ⁴³*LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France*
- ⁴⁴*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ^{45a}*Istituto Nazionale di Fisica Nucleare Pisa, I-56127 Pisa, Italy*
- ^{45b}*University of Pisa, I-56127 Pisa, Italy*
- ^{45c}*University of Siena and I-56127 Pisa, Italy*
- ^{45d}*Scuola Normale Superiore, I-56127 Pisa, Italy*
- ⁴⁶*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- ⁴⁷*Purdue University, West Lafayette, Indiana 47907, USA*
- ⁴⁸*University of Rochester, Rochester, New York 14627, USA*
- ⁴⁹*The Rockefeller University, New York, New York 10065, USA*
- ^{50a}*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, I-00185 Roma, Italy*
- ^{50b}*Sapienza Università di Roma, I-00185 Roma, Italy*
- ⁵¹*Rutgers University, Piscataway, New Jersey 08855, USA*
- ⁵²*Texas A&M University, College Station, Texas 77843, USA*
- ^{53a}*Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, I-33100 Udine, Italy*
- ^{53b}*University of Trieste/Udine, I-33100 Udine, Italy*
- ⁵⁴*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- ⁵⁵*Tufts University, Medford, Massachusetts 02155, USA*
- ⁵⁶*Waseda University, Tokyo 169, Japan*
- ⁵⁷*Wayne State University, Detroit, Michigan 48201, USA*
- ⁵⁸*University of Wisconsin, Madison, Wisconsin 53706, USA*
- ⁵⁹*Yale University, New Haven, Connecticut 06520, USA*
- (Received 18 July 2011; published 8 November 2011)

We report the first observation of the baryonic flavor-changing neutral current decay $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ with 24 signal events and a statistical significance of 5.8 Gaussian standard deviations. This measurement uses a $p\bar{p}$ collisions data sample corresponding to 6.8 fb^{-1} at $\sqrt{s} = 1.96 \text{ TeV}$ collected by the CDF II detector at the Tevatron collider. The total and differential branching ratios for $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ are measured. We find $\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-) = [1.73 \pm 0.42(\text{stat}) \pm (\text{syst})] \times 10^{-6}$. We also report the first measurement of the differential branching ratio of $B_s^0 \rightarrow \phi \mu^+ \mu^-$, using 49 signal events. In addition, we report branching ratios for $B^+ \rightarrow K^+ \mu^+ \mu^-$, $B^0 \rightarrow K^0 \mu^+ \mu^-$, and $B \rightarrow K^*(892) \mu^+ \mu^-$ decays.

DOI: 10.1103/PhysRevLett.107.201802

PACS numbers: 13.30.Ce, 12.15.Mm, 14.20.Mr

Rare decays of hadrons containing bottom quarks through the process $b \rightarrow s \mu^+ \mu^-$, where b is a bottom quark and s is a strange quark, occur in the standard model (SM) with $\mathcal{O}(10^{-6})$ branching ratios [1,2]. The b and s quarks carry the same charge but different flavor, so this process is a flavor-changing neutral current (FCNC) decay. FCNC decays are suppressed at tree level in the SM, and must occur through higher order, and more suppressed, loop amplitudes. Their suppressed nature and clean experimental signature, along with reliable theoretical predictions for their rates [1,3,4], make them excellent search

channels for new physics. With multibody final states, these decays offer sensitivity to new physics in a number of kinematic distributions in addition to the total branching ratio. In this Letter, we report measurements of the total branching ratios of FCNC decays, as well as their differential branching ratios as a function of $q^2 \equiv M_{\mu\mu}^2 c^2$, where $M_{\mu\mu}$ is the dimuon invariant mass. Exclusive decays of $B \rightarrow K^{(*)} \mu^+ \mu^-$ have been observed by BABAR [5], Belle [6], and CDF [7]. The CDF experiment also recently reported the observation of $B_s^0 \rightarrow \phi(1020) \mu^+ \mu^-$ [7].

No significant departure from the SM has been found thus far.

In addition, the study of the baryonic $b \rightarrow s\mu^+\mu^-$ decays is very important, since the baryonic FCNC decays are sensitive to the helicity structure of effective Hamiltonian, which is lost in the hadronization of the mesonic decays [8]. Although the theoretical calculations of the exclusive baryonic $b \rightarrow s\mu^+\mu^-$ decays have large uncertainties compared to the mesonic decays due to additional degrees of freedom in the baryon bound states, the measurements of the total and the differential branching ratios can help the improvement of the theoretical treatments. One can also compare the measurements of the mesonic $b \rightarrow s\mu^+\mu^-$ decays with the baryonic decays, which follow the common quark transition. Measurements of both mesonic and baryonic FCNC decays therefore provide additional tests of the SM and its extensions. However, no b baryon FCNC decay has been observed and there are few experimental constraints on their decay rates. The $\Lambda_b^0 \rightarrow \Lambda\mu^+\mu^-$ decay is considered promising in this respect [8–11] and experimentally accessible since the branching ratio is predicted as $(4.0 \pm 1.2) \times 10^{-6}$ [10].

The data sample used in the measurements reported in this Letter corresponds to an integrated luminosity of 6.8 fb^{-1} from $p\bar{p}$ collisions at a center-of-mass energy of $\sqrt{s} = 1.96 \text{ TeV}$ collected with the CDF II detector between March 2002 and June 2010. The $\Lambda_b^0 \rightarrow \Lambda\mu^+\mu^-$ decay is reconstructed and measurements are made of the total branching ratio and the differential branching ratio as a function of q^2 . Besides the updated branching ratios of $B_s^0 \rightarrow \phi\mu^+\mu^-$, $B^+ \rightarrow K^+\mu^+\mu^-$, and $B^0 \rightarrow K^*(892)^0\mu^+\mu^-$, we report the branching ratios of $B^0 \rightarrow K^0\mu^+\mu^-$ and $B^+ \rightarrow K^*(892)^+\mu^+\mu^-$, which are measured for the first time in hadron collisions. We also report the first measurement of the differential branching ratio as a function of q^2 of $B_s^0 \rightarrow \phi\mu^+\mu^-$. To cancel the dominant systematic uncertainties, decay rates for each rare channel $H_b \rightarrow h\mu^+\mu^-$ are measured relative to the corresponding resonant channel $H_b \rightarrow J/\psi h$ with $J/\psi \rightarrow \mu^+\mu^-$, used as a normalization, where H_b represents the b hadron and h stands for Λ , ϕ , K^+ , K_S^0 , K^{*0} , and K^{*+} . Charge conjugation is implied throughout the Letter.

The reconstruction of the exclusive b hadron events starts with a dimuon sample selected by the online trigger system [12] of the CDF II detector [13]. The trigger system utilizes information from muon detectors and the central outer tracker [14]. Muon chambers CMU and CMX [15] cover $|\eta| < 0.6$ and $0.6 < |\eta| < 1.0$, respectively, [16]. The CMP muon chamber covers $|\eta| < 0.6$ and is located behind the CMU and an additional steel absorber. The dimuon trigger requires a pair of oppositely charged particles with a momentum transverse to the beam line $p_T \geq 1.5 \text{ GeV}/c$, which are matched to track segments in the CMU or CMX chambers. At least one of the muon tracks is required to have a CMU track segment. The trigger also requires that

the dimuon pair satisfies either $L_{xy} > 100 \mu\text{m}$, where the transverse decay length L_{xy} is the flight distance between the dimuon vertex and the event primary vertex [17], or $p_T > 3.0 \text{ GeV}/c$ and matched segments in both CMU and CMP chambers for one of the muon candidates.

Offline event selection starts with the triggered dimuon pairs. Each offline track is required to satisfy more stringent requirements on the number of hits used to reconstruct the track. The dimuon selection requirements used in the trigger are repeated with the higher quality offline tracks. The decay length and invariant mass of each dimuon pair are calculated after a vertex fit using the muon tracks. Dimuon pairs are classified according to their invariant mass $M_{\mu\mu}$. Dimuons from FCNC b hadron decays are required to be inconsistent with decaying from J/ψ ($\psi(2S)$) mesons by requiring q^2 values outside the window of $8.68(12.86) < q^2 < 10.09(14.18) \text{ GeV}^2/c^2$ [7]. The J/ψ candidates are required to have $M_{\mu\mu}$ within $50 \text{ MeV}/c^2$ of the known J/ψ mass [18].

The $\Lambda_b^0 \rightarrow \Lambda\mu^+\mu^-$ candidates are selected by combining the dimuon pairs with Λ baryons reconstructed from decays $\Lambda \rightarrow p\pi^-$. The $p\pi^-$ pairs are required to have invariant mass consistent with the known Λ mass [18], $p_T \geq 1.0 \text{ GeV}/c$, and a vertex displaced from the dimuon vertex. The transverse momentum of the Λ_b^0 candidate is required to be greater than $4.0 \text{ GeV}/c$. Candidates with an invariant mass calculated from two or three daughter particles compatible with J/ψ , $\psi(2S)$, D^0 , D^+ , D_s^+ , or Λ_c masses are rejected to remove backgrounds from these charm-hadron decays [7]. The $B_s^0 \rightarrow \phi\mu^+\mu^-$ candidates are reconstructed from dimuons together with a pair of oppositely-charge kaons consistent with a ϕ decay with a selection similar to that of $\Lambda_b^0 \rightarrow \Lambda\mu^+\mu^-$. The $B^{0,+} \rightarrow \mathcal{K}^{0,+}\mu^+\mu^-$ candidates, where $\mathcal{K}^{0,+}$ is one of $\{K^+, K_S^0, K^{*0}, K^{*+}\}$, are formed from a dimuon combined with up to three charged tracks. The K_S^0 meson is reconstructed in its $\pi^+\pi^-$ final state by requiring the dipion mass to be consistent with the known K_S^0 mass [18]. Details about the reconstruction of the decays of $K^{*0} \rightarrow K^+\pi^-$ and $\phi \rightarrow K^+K^-$ can be found in Ref. [7]. Cross feed between $\Lambda_b^0 \rightarrow \Lambda\mu^+\mu^-$ and $B^0 \rightarrow K_S^0\mu^+\mu^-$ is suppressed by evaluating the momentum imbalance of Λ and K_S^0 daughters [19]. We utilize the correlation between invariant mass and the asymmetry $\alpha \equiv (q_L^+ - q_L^-)/(q_L^+ + q_L^-)$, where $q_L^{\pm(-)}$ is the longitudinal momentum of the positive (negative) decay product relative to the direction of the Λ or K_S^0 . We reject candidates that satisfy $-0.26 < -1.9M(K_S^0) + |\alpha| < -0.15$ for $K_S^0\mu^+\mu^-$ and $4.73 < 3.6M(\Lambda) + |\alpha| < 4.78$ for $\Lambda\mu^+\mu^-$. We remove 76 (90)% of the cross feed while the signal loss is 11 (7)% for $\Lambda\mu^+\mu^-$ ($K_S^0\mu^+\mu^-$). A residual cross-feed contamination of 0.1% (0.6%) to the $\Lambda\mu^+\mu^-$ ($K_S^0\mu^+\mu^-$) signal is considered as a systematic uncertainty. To further optimize the event selection, an artificial neural network (NN) classifier is trained using simulated signal events and




FIG. 1 (color online). Invariant mass of (a) $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$, (b) $B_s^0 \rightarrow \phi \mu^+ \mu^-$, (c) $B^+ \rightarrow K^+ \mu^+ \mu^-$, (d) $B^0 \rightarrow K^{*0} \mu^+ \mu^-$, (e) $B^0 \rightarrow K_S^0 \mu^+ \mu^-$, (f) $B^+ \rightarrow K^{*+} \mu^+ \mu^-$, with fit results overlaid. The histograms are the data. Solid, dashed-dotted, and dotted curves show the total fit, the signal PDF, and the background PDF, respectively.

background events taken from H_b mass sidebands ($0.1\text{--}0.36 \text{ GeV}/c^2$ far from the known H_b mass) in data. Some kinematical distributions of the simulated signal, e.g., the transverse momentum of b hadron, and the energy depositions of muon candidates in the electromagnetic and hadron calorimeters, are corrected using scale factors extracted by comparing simulation to data in the normalization channels. We use 70% of the sideband events for the training, and use the remaining events to check that the NN does not bias or over suppress the mass distribution. The optimized NN threshold is determined to maximize the average expected significance of the branching ratio, using many kinematic observables including transverse momentum, invariant mass, vertex fit qualities, and muon identification qualities [7].

TABLE I. Summary of observed yields, the statistical significance s , and the relative efficiency ε_{rel} .

Mode	$N_{h\mu^+\mu^-}$	$s(\sigma)$	$N_{J/\psi h}$	ε_{rel}
$\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$	24 ± 5	5.8	1740 ± 50	0.33 ± 0.01
$B_s^0 \rightarrow \phi \mu^+ \mu^-$	49 ± 7	9.0	4560 ± 80	0.56 ± 0.01
$B^+ \rightarrow K^+ \mu^+ \mu^-$	234 ± 19	13.7	$72\,200 \pm 300$	0.41 ± 0.01
$B^0 \rightarrow K^{*0} \mu^+ \mu^-$	164 ± 15	13.7	$28\,300 \pm 200$	0.45 ± 0.02
$B^0 \rightarrow K_S^0 \mu^+ \mu^-$	28 ± 9	3.5	9470 ± 90	0.47 ± 0.01
$B^+ \rightarrow K^{*+} \mu^+ \mu^-$	20 ± 6	3.5	4560 ± 80	0.38 ± 0.02

The signal yield of the $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ candidates is obtained by an unbinned maximum likelihood fit to the Λ_b^0 invariant mass distribution with the signal probability density function (PDF) parametrized by Gaussian distributions using simulated signals and the background PDF modeled by a linear function. We fix the Λ_b^0 mass width for the rare decay while it is floated for the normalization channel. Different mass width between data and the simulated signal is corrected by measured mass width ratio of the normalization channel between data and the simulated signal. The signal region is defined within $\pm 40 \text{ MeV}/c^2$ from the world average Λ_b^0 mass [7]. The statistical significance is obtained through a likelihood-ratio test between the signal plus background and background-only hypotheses interpreted assuming it distributed as a χ^2 variable. The invariant mass distribution of the $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ candidates is shown in Fig. 1. In the signal region, we observe 24 ± 5 events from $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ decays while the total number of the signal candidates is 34. The statistical significance of the signal s corresponds to 5.8 Gaussian standard deviations. The signal yields of $B_s^0 \rightarrow \phi \mu^+ \mu^-$ and other FCNC B meson decays are obtained by a similar procedure as that of $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$. Each channel uses independent NN weight and PDF. The fit range for B^+ and B^0 decays is from 5.18 to 5.70 GeV/c^2 to avoid the region of 5.0–5.18 GeV/c^2 , which is dominated by the feed-down background from multibody decays of b hadrons. While the contribution from charmless H_b decays is negligible due to the muon identification, we estimate a 1% cross talk between $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ and $B_s^0 \rightarrow \phi \mu^+ \mu^-$ using simulation, and correct for it. Invariant mass distributions of $B_s^0 \rightarrow \phi \mu^+ \mu^-$ and other FCNC B meson decays are shown in Fig. 1 and signal yields are listed in Table I.

The branching ratios of $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$, $B_s^0 \rightarrow \phi \mu^+ \mu^-$, and $B \rightarrow K^{(*)} \mu^+ \mu^-$ are calculated by comparing their signal event yield to that of the normalization decay modes $\Lambda_b^0 \rightarrow J/\psi \Lambda$, $B_s^0 \rightarrow J/\psi \phi$, and $B \rightarrow J/\psi K^{(*)}$, where $J/\psi \rightarrow \mu^+ \mu^-$, after the reconstruction efficiency correction:

TABLE II. Measured branching ratios of rare modes. First (second) uncertainty is statistical (systematic). The last two values are for the isospin average.

Mode	Relative $\mathcal{B}(10^{-3})$	Absolute $\mathcal{B}(10^{-6})$
$\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$	$2.45 \pm 0.59 \pm 0.29$	$1.73 \pm 0.42 \pm 0.55$
$B_s^0 \rightarrow \phi \mu^+ \mu^-$	$1.13 \pm 0.19 \pm 0.07$	$1.47 \pm 0.24 \pm 0.46$
$B^+ \rightarrow K^+ \mu^+ \mu^-$	$0.46 \pm 0.04 \pm 0.02$	$0.46 \pm 0.04 \pm 0.02$
$B^0 \rightarrow K^{*0} \mu^+ \mu^-$	$0.77 \pm 0.08 \pm 0.03$	$1.02 \pm 0.10 \pm 0.06$
$B^0 \rightarrow K^0 \mu^+ \mu^-$	$0.37 \pm 0.12 \pm 0.02$	$0.32 \pm 0.10 \pm 0.02$
$B^+ \rightarrow K^{*+} \mu^+ \mu^-$	$0.67 \pm 0.22 \pm 0.04$	$0.95 \pm 0.32 \pm 0.08$
$B \rightarrow K \mu^+ \mu^-$...	$0.42 \pm 0.04 \pm 0.02$
$B \rightarrow K^* \mu^+ \mu^-$...	$1.01 \pm 0.10 \pm 0.05$




FIG. 2 (color online). Differential branching ratios of (a) $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$, (b) $B_s^0 \rightarrow \phi \mu^+ \mu^-$, (c) $B^+ \rightarrow K^+ \mu^+ \mu^-$, and (d) $B^0 \rightarrow K^0 \mu^+ \mu^-$. The points are the fit result. The solid curves are the SM expectation [1,10,21]. The dashed line in the $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ plot is the SM prediction normalized to our total branching ratio measurement. The hatched regions are the charmonium veto regions.

$$\frac{\mathcal{B}(H_b \rightarrow h \mu^+ \mu^-)}{\mathcal{B}(H_b \rightarrow J/\psi h)} = \frac{N_{h\mu^+\mu^-}}{N_{J/\psi h}} \frac{\mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)}{\varepsilon_{\text{rel}}}, \quad (1)$$

where $N_{h\mu^+\mu^-}$ is the $h\mu^+\mu^-$ yield, $N_{J/\psi h}$ is the $J/\psi h$ yield for the normalization channel, and $\varepsilon_{\text{rel}} \equiv \varepsilon_{h\mu^+\mu^-}/\varepsilon_{J/\psi h}$ is the relative reconstruction efficiency determined from the simulation. The calculated relative and absolute branching ratios are listed in Table II. The absolute branching ratios are obtained using world averages of the $J/\psi h$ decay rates [18]. The branching ratios of $B^0 \rightarrow K^0 \mu^+ \mu^-$ and $B^+ \rightarrow K^+ \mu^+ \mu^-$ are measured for the first time in hadron collisions.

The dominant sources of systematic uncertainty are the scale-factor reweighting of the simulated signal (the trigger efficiency near the threshold) which ranges from 0.5% to 4.0% (0.8% to 7.2%), depending on the channel. We estimate the former uncertainty from the comparison of the

relative efficiencies with and without reweighting and the latter uncertainty from the different p_T requirements for each trigger. In the $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ case we consider an additional uncertainty of 6.6% due to the unknown $\Lambda_b^0 \rightarrow J/\psi \Lambda$ polarization.

For the absolute branching ratio measurements we assign the uncertainties on the world average $\mathcal{B}(H_b \rightarrow J/\psi h)$ [18] or the most recent measurement [20]. Contributions from other sources (e.g., background PDF shape or the decay model of the simulated event) are minor (0.3%–3.4%).

The combined branching ratio is calculated by assuming isospin symmetry and using the B^+ and B^0 total widths [18]. These numbers are consistent with our previous results [7], B -factory measurements [5,6], and theoretical expectations [9,10].

We also measure differential branching ratios with respect to q^2 . We divide the signal region into six bins in q^2 . We fit the signal yield in each q^2 bin. In each fit, we fix the mean of the H_b mass and the background slope to the value from the global fit, so that only the signal fraction is allowed to vary in the fit. Figure 2 shows the differential branching ratios for $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$, $B_s^0 \rightarrow \phi \mu^+ \mu^-$, and $B \rightarrow K^{(*)} \mu^+ \mu^-$. For illustration, we superimpose the SM expectations, which are based on the formula in Ref. [1], with the form factors in Ref. [21], except for the case of $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ decays which follows Ref. [10]. The cusp at $q^2 \sim 7 \text{ GeV}^2/c^2$ is due to a change in parameter approximations. Tables III and IV summarize the differential branching ratio measurements. The two bottom rows in each table show the results for the semi-inclusive bins, which are included with ranges covering theoretically well-controlled regions.

In summary, we have updated our previous analysis of the flavor-changing neutral current decays $b \rightarrow s \mu^+ \mu^-$ using data corresponding to an integrated luminosity of 6.8 fb^{-1} and adding new decay channels. We report the first observation of $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ and measure the total and differential branching ratios of this decay with respect to q^2 . We also measure the total and differential branching ratios of $B \rightarrow K^{(*)} \mu^+ \mu^-$ and $B_s^0 \rightarrow \phi \mu^+ \mu^-$, with respect to q^2 . All measurements are consistent and competitive

TABLE III. Differential branching ratios of $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$, $B^+ \rightarrow K^+ \mu^+ \mu^-$, $B^0 \rightarrow K^0 \mu^+ \mu^-$, combined $B \rightarrow K \mu^+ \mu^-$, in units of 10^{-7} . The q_{max}^2 is 20.30 (23.00) GeV^2/c^2 for $\Lambda \mu^+ \mu^-$ ($K \mu^+ \mu^-$). The first (second) uncertainty is statistical (systematic).

q^2 (GeV^2/c^2)	$\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$	$B^+ \rightarrow K^+ \mu^+ \mu^-$	$B^0 \rightarrow K^0 \mu^+ \mu^-$	$B \rightarrow K \mu^+ \mu^-$
[0.00, 2.00)	$0.15 \pm 2.01 \pm 0.05$	$0.36 \pm 0.11 \pm 0.03$	$0.312 \pm 0.372 \pm 0.024$	$0.33 \pm 0.10 \pm 0.02$
[2.00, 4.30)	$1.84 \pm 1.66 \pm 0.59$	$0.80 \pm 0.15 \pm 0.05$	$0.929 \pm 0.485 \pm 0.070$	$0.77 \pm 0.14 \pm 0.05$
[4.30, 8.68)	$-0.20 \pm 1.64 \pm 0.08$	$1.18 \pm 0.19 \pm 0.09$	$0.663 \pm 0.510 \pm 0.052$	$1.05 \pm 0.17 \pm 0.07$
[10.09, 12.86)	$2.97 \pm 1.47 \pm 0.95$	$0.68 \pm 0.12 \pm 0.05$	$-0.030 \pm 0.223 \pm 0.005$	$0.48 \pm 0.10 \pm 0.03$
[14.18, 16.00)	$0.96 \pm 0.73 \pm 0.31$	$0.53 \pm 0.10 \pm 0.03$	$0.726 \pm 0.257 \pm 0.055$	$0.52 \pm 0.09 \pm 0.03$
[16.00, q_{max}^2)	$6.97 \pm 1.88 \pm 2.23$	$0.48 \pm 0.11 \pm 0.03$	$0.214 \pm 0.182 \pm 0.016$	$0.38 \pm 0.09 \pm 0.02$
[0.00, 4.30)	$2.65 \pm 2.52 \pm 0.85$	$1.13 \pm 0.19 \pm 0.08$	$1.268 \pm 0.622 \pm 0.096$	$1.07 \pm 0.17 \pm 0.07$
[1.00, 6.00)	$1.27 \pm 2.08 \pm 0.41$	$1.41 \pm 0.20 \pm 0.10$	$0.980 \pm 0.614 \pm 0.076$	$1.29 \pm 0.18 \pm 0.08$

TABLE IV. Differential branching ratios of $B_s^0 \rightarrow \phi \mu^+ \mu^-$, $B^0 \rightarrow K^{*0} \mu^+ \mu^-$, $B^+ \rightarrow K^{*+} \mu^+ \mu^-$, and combined $B \rightarrow K^* \mu^+ \mu^-$, in units of 10^{-7} . The q_{\max}^2 is 18.90 (19.30) GeV^2/c^2 for $\phi \mu^+ \mu^-$ ($K^* \mu^+ \mu^-$). The first (second) uncertainty is statistical (systematic).

q^2 (GeV^2/c^2)	$B_s^0 \rightarrow \phi \mu^+ \mu^-$	$B^0 \rightarrow K^{*0} \mu^+ \mu^-$	$B^+ \rightarrow K^{*+} \mu^+ \mu^-$	$B \rightarrow K^* \mu^+ \mu^-$
[0.00, 2.00)	$2.78 \pm 0.95 \pm 0.89$	$1.80 \pm 0.36 \pm 0.11$	$1.30 \pm 0.98 \pm 0.14$	$1.73 \pm 0.33 \pm 0.10$
[2.00, 4.30)	$0.58 \pm 0.55 \pm 0.19$	$0.84 \pm 0.28 \pm 0.06$	$0.71 \pm 1.00 \pm 0.15$	$0.82 \pm 0.26 \pm 0.06$
[4.30, 8.68)	$1.34 \pm 0.83 \pm 0.43$	$1.73 \pm 0.43 \pm 0.15$	$1.71 \pm 1.58 \pm 0.49$	$1.72 \pm 0.41 \pm 0.14$
[10.09, 12.86)	$2.98 \pm 0.95 \pm 0.95$	$1.77 \pm 0.36 \pm 0.12$	$1.97 \pm 0.99 \pm 0.22$	$1.77 \pm 0.34 \pm 0.11$
[14.18, 16.00)	$1.86 \pm 0.66 \pm 0.59$	$1.34 \pm 0.26 \pm 0.08$	$0.52 \pm 0.61 \pm 0.09$	$1.21 \pm 0.24 \pm 0.07$
[16.00, q_{\max}^2)	$2.32 \pm 0.76 \pm 0.74$	$0.97 \pm 0.26 \pm 0.07$	$1.57 \pm 0.96 \pm 0.17$	$0.88 \pm 0.22 \pm 0.05$
[0.00, 4.30)	$3.30 \pm 1.09 \pm 1.05$	$2.60 \pm 0.45 \pm 0.17$	$2.01 \pm 1.39 \pm 0.27$	$2.53 \pm 0.43 \pm 0.15$
[1.00, 6.00)	$1.14 \pm 0.79 \pm 0.36$	$1.42 \pm 0.41 \pm 0.12$	$2.57 \pm 1.61 \pm 0.40$	$1.48 \pm 0.39 \pm 0.12$

with other results, and the differential measurements of $B_s^0 \rightarrow \phi \mu^+ \mu^-$ and $\Lambda_b^0 \rightarrow \Lambda \mu^+ \mu^-$ are the first such measurements. At present there is no evidence of discrepancy from the SM prediction.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science, and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A. P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

^aDeceased.

^bVisitor from University of Massachusetts Amherst, Amherst, MA 01003, USA.

^cVisitor from Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy.

^dVisitor from University of California Irvine, Irvine, CA 92697, USA.

^eVisitor from University of California Santa Barbara, Santa Barbara, CA 93106, USA.

^fVisitor from University of CA Santa Cruz, Santa Cruz, CA 95064, USA.

^gVisitor from CERN, CH-1211 Geneva, Switzerland.

^hVisitor from Cornell University, Ithaca, NY 14853, USA.

ⁱVisitor from University of Cyprus, Nicosia CY-1678, Cyprus.

^jVisitor from University College Dublin, Dublin 4, Ireland.

^kVisitor from University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017.

^lVisitor from Universidad Iberoamericana, Mexico D.F., Mexico.

^mVisitor from Iowa State University, Ames, IA 50011, USA.

ⁿVisitor from University of Iowa, Iowa City, IA 52242, USA.

^oVisitor from Kinki University, Higashi-Osaka City, Japan 577-8502.

^pVisitor from, Kansas State University, Manhattan, KS 66506, USA.

^qVisitor from University of Manchester, Manchester M13 9PL, United Kingdom.

^rVisitor from Queen Mary, University of London, London, E1 4NS, United Kingdom.

^sVisitor from Muons, Inc., Batavia, IL 60510, USA.

^tVisitor from Nagasaki Institute of Applied Science, Nagasaki, Japan.

^uVisitor from National Research Nuclear University, Moscow, Russia.

^vVisitor from University of Notre Dame, Notre Dame, IN 46556, USA.

^wVisitor from Universidad de Oviedo, E-33007 Oviedo, Spain.

^xVisitor from Texas Tech University, Lubbock, TX 79609, USA.

^yVisitor from IFIC(CSIC-Universitat de Valencia), 56071 Valencia, Spain.

^zVisitor from Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile.

^{aa}Visitor from University of Virginia, Charlottesville, VA 22906, USA.

^{bb}Visitor from Yarmouk University, Irbid 211-63, Jordan.

^{cc}On leave from J. Stefan Institute, Ljubljana, Sloveni.

- [1] A. Ali *et al.*, *Phys. Rev. D* **61**, 074024 (2000).
- [2] D. Melikhov, N. Nikitin, and S. Simula, *Phys. Rev. D* **57**, 6814 (1998).
- [3] Q. Chan and Y.-H. Gao, *Nucl. Phys.* **B845**, 179 (2011).
- [4] Y.-M. Wang, M.J. Aslam, and C.-D. Lu, *Eur. Phys. J. C* **59**, 847 (2009).
- [5] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **102**, 091803 (2009).
- [6] J. T. Wei *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **103**, 171801 (2009).

- [7] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **79**, 011104 (2009); T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **106**, 161801 (2011).
- [8] Y.-M. Wang, Y. Li, and C.-D. Lu, *Eur. Phys. J. C* **59**, 861 (2009).
- [9] C.-H. Chen and C.Q. Geng, *Phys. Rev. D* **64**, 074001 (2001).
- [10] T.M. Aliev, K. Azizi, and M. Savci, *Phys. Rev. D* **81**, 056006 (2010).
- [11] M. J. Aslam, Y.-M. Wang, and C.-D. Lu, *Phys. Rev. D* **78**, 114032 (2008).
- [12] E.J. Thomson *et al.*, *IEEE Trans. Nucl. Sci.* **49**, 1063 (2002).
- [13] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 032001 (2005).
- [14] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **102**, 242002 (2009), and references therein.
- [15] G. Ascoli *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **268**, 33 (1988); T. Dorigo (CDF Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **461**, 560 (2001).
- [16] We use a cylindrical coordinate system in which θ is the polar angle with respect to the proton beam line and pseudorapidity $\eta \equiv -\ln(\tan\theta/2)$.
- [17] W. Ashmanskas *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **518**, 532 (2004).
- [18] K. Nakamura *et al.* (Particle Data Group), *J. Phys. G* **37**, 075021 (2010).
- [19] J. Podolanski and R. Armenteros, *Philos. Mag.* **45**, 13 (1954).
- [20] V.M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **84**, 031102 (2011).
- [21] P. Ball and R. Zwicky, *Phys. Rev. D* **71**, 014015 (2005); P. Ball and R. Zwicky, *Phys. Rev. D* **71**, 014029 (2005).