

Effects of commercial enrichment diets on the nutritional value of the rotifer (*Brachionus plicatilis*)

M.J. Fernandez-Reiriz, U. Labarta and M.J. Ferreiro

Instituto de Investigaciones Marinas, C.S.I.C., Eduardo Cabello, 6, 36208 Vigo, Spain

Correspondence to: M.J. Fernandez-Reiriz, Instituto de Investigaciones Marinas, C.S.I.C., Eduardo Cabello, 6, 36208 Vigo, Spain.

ABSTRACT

Fernandez-Reiriz, M.J., Labarta, U. and Ferreiro, M.J., 1993. Effects of commercial enrichment diets on the nutritional value of the rotifer (*Brachionus plicatilis*). *Aquaculture*. 112: 195-206.

The rotifer, *Brachionus plicatilis*, pre-fed on baker's yeast, was enriched for 6 h with three commercial products from Artemia Systems S.A., namely Protein Selco (microcapsules), Dry Selco (microparticles) and Super Selco (an emulsion containing high levels of n-3 HUFA). The biochemical composition (protein, carbohydrate, total lipid, lipid classes and fatty acids) and dry weight of the rotifers before and after the enrichment experiments were studied. Two of the enrichers, Dry Selco and Super Selco, are rich in lipid but poor in protein and carbohydrate. Protein Selco contains lipid as well as protein and carbohydrate. The biochemical composition and the dry weight of the rotifers were changed after 6 h of enrichment. The percentage of protein and lipid increased with all of the three enrichers and the percentage of carbohydrate decreased slightly. Rotifer dry weight increased in rotifers enriched on Protein Selco.

Phospholipids, sterol esters+ waxes, triacylglycerols and free fatty acids increased after the enrichment with the three enrichers while sterols slightly decreased when the rotifers were fed on Protein Selco. The fatty acid content of the rotifers increased after the enrichment with all of the enrichers and this increase was particularly apparent in the case of n-3 HUFAs (20:5n-3 and 22:6n-3).

INTRODUCTION

The rotifer, *Brachionus plicatilis*, has been widely used as a food during the first development stages of fishes and crustaceans since the first experiments on its culture were published by Ito (1960). Since then, the nutritional quality of the rotifer has been studied by many authors (Watanabe et al., 1978, 1983; Ben-Amotz et al., 1987; Dendrinis and Thorpe, 1987; Fernandez-Reiriz et al., 1987; Lubzens et al., 1987). However, fish larvae fed on rotifers often exhibit slow growth rates and high mortalities which has been attributed to PUFA deficiencies (Owen et al., 1975; Cowey, 1976; Howell, 1979; Scott and Middleton, 1979; Watanabe et al., 1983; Ben-Amotz et al., 1987; Dendrinis and Thorpe, 1987; James et al., 1987; Kissil and Koven, 1990). The chemical composition of the rotifer has been shown to be altered by diet (Watanabe et al., 1983; Frolov et al., 1991). Furthermore, a relationship between the fatty acid content in the diet and the rotifer has also been reported (Watanabe et al., 1978, 1983; Scott and Middleton, 1979; Lubzens et al., 1985, 1987; Estevez and Planas, 1988; Stottrup, 1989; Frolov et al., 1991).

Mass culture of rotifers is usually carried out on baker's yeast, *Saccharomyces cerevisiae*. However, rotifers fed exclusively on baker's yeast are often nutritionally deficient when used as food for fish larvae. Many attempts have been made to improve the nutritional quality of the rotifers in the last few years, and two enrichment methods have been described: the "indirect method" which attempts to improve the biochemical composition of the yeast by changing its culture medium (Imada et al., 1979) and the "direct method" which supplies the rotifers with baker's yeast plus some specific nutritional compounds, mainly emulsions of marine oils (Watanabe et al., 1983). During the last few years, various diets such as microalgae, lipid emulsions, microparticulate or microcapsules containing lipids or lipids plus proteins and carbohydrates have been used to enrich rotifers previously fed on baker's yeast (Gatesoupe and Luquet, 1981; Walford and Lam, 1987; Leger et al., 1989; Nichols et al., 1989; Rodriguez-Rainuzzo et al., 1989; Kissil and Koven, 1990; Linares et al., 1990).

This paper reports the chemical composition (proteins, carbohydrates, total lipids, lipid classes and fatty acids) of the rotifer, *Brachionus plicatilis*, after 3 and 6 h of enrichment with three commercial products: Protein Selco, Dry Selco and Super Selco from Artemia Systems. The composition of the enriched and non-enriched rotifers is

compared in order to establish the suitability of these live organisms as potential food for fish larvae, particularly turbot larvae.

MATERIAL AND METHODS

Culture of rotifers

The rotifers, *Brachionus plicatilis*, previously fed on baker's yeast (1 g/10⁶ rotifers day⁻¹) were enriched for 3 and 6 h with three commercial products from Artemia Systems: Protein Selco (microcapsules), Dry Selco (microparticles) and Super Selco (an emulsion containing high levels of n-3 HUFA) .

The enrichment experiments were carried out in 171 tanks with a density of 600 rotifers per ml. The enrichment was made by adding 2 doses of 0.189 g/l to each culture. The first dose was added to the cultures at the beginning of the experiment and the second after 3 h. The enrichments were carried out at a temperature of 20°C. Two replicates were made for each experiment.

The cultures were sampled at the beginning of the experiments and after 3 and 6 h of enrichment for various biochemical analyses. Volumes of water with a quantity of 1 x 10⁶ rotifers were filtered out. The samples were then gently washed with distilled water to eliminate both salt and possible residues of enricher. They were frozen in liquid nitrogen and later freeze-dried and frozen until corresponding biochemical analyses were performed.

Analytical methods

Protein was assayed as described by Lowry et al. (1951) after hydrolysis with 0.5 N NaOH for 24 h at 30°C. Total carbohydrates were quantified as glucose by the phenol-sulfuric acid method (Strickland and Parsons, 1968). Lipids were extracted by the method of Bligh and Dyer (1959) as modified by Fernandez-Reiriz et al. (1989). Total lipids were determined gravimetrically. Lipid classes were studied by thin layer chromatography (TLC)-densitometry. Silicagel 60W plates (Merck 16486) with a size of 20 x 20 cm and a layer thickness of 0.25 mm were used. The chromatographic stain

was made by following the technique described by Freeman and West (1966). Samples were applied on the plate with the help of an automatic TLC sampler (Camag 27220). The plates were stained with a 10% CuSO₄ solution in 0.85% H₃PO₄ by heating to 180°C (Bitman and Wood, 1982). For the quantitative analysis, cholesterol palmitate, cholesterol, palmitic acid and tripalmitin (Sigma) were used as standards for sterol esters + waxes, sterols, free fatty acids and triacylglycerides, respectively. For phospholipids a standard obtained from rotifers was used. The plates were scanned with a Shimadzu CS9000-densitometer, using a beam monochromatic 370 nm of 0.4 x 0.4 mm working in the zig-zag mode, reading the whole spot, and with automatic autozero for base line correction.

Fatty acids from total lipid were transesterified to methyl esters with methanolic hydrogen chloride as described by Christie (1982) and subsequently analyzed by gas chromatography as previously described by Fernandez-Reiriz et al. (1989), with the only exception that a PTV cold injector (Perkin-Elmer), operating in the solvent elimination mode as described Herraiz et al. (1987) was used. Nonadecanoic acid was used as an internal standard and a response factor was calculated for each fatty acid in order to perform quantitative analysis.

The caloric content of rotifers was calculated using the conversion factors given by Brett and Groves (1979).

A Fisher PLSD test was applied for statistical analyses at a confidence level of 95% (P<0.05).

RESULTS

Dry weight and proximate composition (proteins, carbohydrates and total lipids)

The dry weight (as ng/ind.) and proximate composition (as % dry weight) of *B. plicatilis* and the enricher are shown in Table 1. Only one of the three enrichers, Protein Selco, had appreciable quantities of protein and carbohydrate (around 30% each) with about 30% of lipid. Dry Selco and particularly Super Selco contained mainly lipid (56.1 and 94.2%, respectively). With respect to the rotifers, the percentages of protein and

lipid increased significantly ($P < 0.05$) after 6 h of enrichment with all of the three products. The percentages of carbohydrate decreased significantly ($P < 0.05$) after the enrichment.

The dry weight of the rotifers fed on Protein Selco increased significantly ($P < 0.05$) after 6 h (453.5 rig/inn.). This increase represents 22.7% of the value found in non-enriched rotifers. The dry weight decreased significantly ($P < 0.05$) 14.3% (316.5 ng/ind.) if the enrichment was made on Super Selco. No significant difference ($P > 0.05$) was found in the dry weight of the rotifers fed on Dry Selco. These differences in dry weight will affect the absolute values of the biochemical compounds. In this sense, the content of protein and carbohydrate of Protein Selco-enriched rotifers increased significantly ($P < 0.05$) 29.7% and 14.3% with regard to the values found in non-enriched rotifers but decreased significantly ($P < 0.05$) if the rotifers were enriched with Super Selco (6.1 and 33.0%, respectively). In Dry Selco-enriched rotifers, the protein content increased significantly ($P < 0.05$) 7.7% and carbohydrate decreased significantly ($P < 0.05$) 10.9%. The lipid content of the rotifers increased significantly ($P < 0.05$) in the three experiments (116.5, 74.9 and 62.396, for Protein, Dry and Super Selco, respectively).

Lipid classes

Table 2 shows the composition (mg/g dry weight) of lipid classes of the three enrichers and the rotifers before and after the enrichment experiments. Protein Selco contained alkyl glyceryl ethers, triacylglycerols, free fatty acids and phospholipids. Dry Selco contained alkyl glyceryl ethers, phospholipids and free fatty acids and Super Selco alkyl glyceryl ethers and phospholipids. Alkyl glyceryl ethers were the major lipid class in the three enrichers although these lipids were not found in the rotifers even after 6 h of enrichment. Phospholipids were the major lipid class in the non-enriched rotifers, followed by sterol esters + waxes, triacylglycerols (TAG), free fatty acids (FFA) and sterols. Phospholipids, sterol esters + waxes, TAG and FFA increased significantly ($P < 0.05$) after the enrichment in the three experiments while sterols decreased significantly ($P < 0.05$) when the rotifers were fed on Protein Selco.

Fatty acid composition

Table 3 shows the fatty acid composition (mg fatty acid/g dry weight) of the three enrichers and *B. plicatilis* without enrichment and after enrichment for 3 and 6 h. Super Selco had the highest quantity of unsaturated fatty acids. Dry Selco had the highest content of saturated fatty acids. Protein Selco had the lowest fatty acid content. In the non-enriched rotifers, the major fatty Acids were 18:1(n-9), 16:1(n-7), 18:2(n-6), 16:0, 18:0, 20:5(n-3), 22:6(n-3), 20:1(n-9), 18:1(n-7), 18:1(n-11), 20:3(n-6) 22:5(n-3), 20:2(n-6) and 20:4 (n-6). After the enrichment, the same fatty acids were found but the relative percentages were different depending on the enricher. In Protein Selco-enriched rotifers, the values of each fatty acid increased significantly ($P < 0.05$) after 3 h and did not change after 6 h with the exception of 20:5(n-3) which showed a significant increase ($P < 0.05$). In Dry Selco and Super Selco-enriched rotifers, the fatty acids increased significantly ($P < 0.05$) during the enrichment period with the highest values detected after 6 h.

Caloric content

Table 4 shows the caloric content of the rotifers before and after the enrichment. The highest total caloric content was observed after 6 h, with values of 2.00, 1.62 and 1.42 Cal/ 10^3 rotifers, for Protein, Dry and Super Selco, respectively.

DISCUSSION

At the end of the experiments the dry weight of rotifers enriched with Protein Selco increased significantly ($P < 0.05$). In the case of Super Selco-enriched rotifers, there was a significant decrease ($P < 0.05$) in dry weight after the enrichment. No significant difference ($P > 0.05$) was found in the dry weight of Dry Selco-enriched rotifers. These results suggest that Dry and Super Selco could be given to the rotifers together with baker's yeast to avoid a weight loss. Protein Selco-enriched rotifers showed a significant increase ($P < 0.05$) in the absolute values of protein and carbohydrate. In Dry Selco-enriched rotifers, proteins increased significantly ($P < 0.05$) and carbohydrates decreased significantly ($P < 0.05$). Super Selco-enriched rotifers depleted their own reserves probably due to the lack of protein and carbohydrate in the enricher.

The role of lipid as an energy reserve during the first larval stages of fishes has been reported by many authors (Ehrlich, 1974; Love, 1980; Kanazawa, 1985).

Triacylglycerols are considered as the main energy reserve (Hakanson, 1989; Tandler et al., 1989) while phospholipids and even proteins seem to be used when there is an energy deficiency (Kimata, 1983; Fyhn et al., 1987; Tandler et al., 1989).

Planas et al. (1991) reported a decrease in protein (41%), carbohydrate (32%) and lipid (5.1%) in larvae starved from 2 to 5 days. When newly hatched larvae were fed on Selco-enriched rotifers (an emulsion with high levels of lipid), there was a decrease in lipid (triglycerides, sterol esters + waxes, free fatty acids and phospholipids) and protein (Planas et al., 1992a) in the larvae. When the larvae were fed on Protein Selco-enriched rotifers, an increase in protein (31%) and carbohydrate (100%) and a decrease in lipid (2%) were observed between days 2 and 5 posthatching, and between days 5 and 7 both proteins and lipids increased (5.1 and 18%, respectively) (Planas et al., 1991).

Therefore, the protein and lipid content of the larval food seems to be an important factor in determining its nutritional value.

Once the larvae have consumed their yolk reserves, they catabolize mainly triglycerides, waxes and phospholipids (Planas et al., 1991). After enrichment with any of the diets tested in this study the rotifers accumulate mainly energetic lipids (triacylglycerols and sterol esters + waxes) and structural lipids.

The suitability of a larval diet is also related to its fatty acid content. Gatesoupe et al. (1977), Léger et al. (1979) and Le Milinaire et al. (1983) have reported dietary requirements of n-3 HUFA of between 0.55 and 1.8% of the dry weight of the food for larval and juvenile turbot. Nevertheless, Gatesoupe and Le Milinaire (1984) reported very low survival rates in turbot larvae after feeding rotifers containing 1.6% n-3 HUFA. Our results show that the n-3 HUFA content in non-enriched rotifers was very low (0.4%) but increased to the required levels after 3 h of enrichment with Super or Protein Selco and after 6 h with Dry Selco. Planas et al. (1991, 1992b) have shown that the fatty acids depleted between days 2 and 5 were mainly the n-3 (n-3 HUFA), n-9 (18:1 n-9) and 16:0. In our experiments, the fatty acid content of the rotifers was enhanced by the enrichment and the increase was particularly clear in the case of n-3 HUFA (20:5n-3 and 22:6n-3).

The requirements for other fatty acids such as 18:3(n-3) and 20:4(n-6) have been studied in juvenile turbot but are unknown for the larvae. Léger et al. (1979) reported a requirement of 1% 18:3(n-3) for turbot juveniles and Linares and Henderson (1991) indicated that turbot juveniles require dietary 20:4(n-6) for incorporation into phosphatidylinositol. The three enrichers tested in our experiments improved the levels of 18:3(n-3) and 20:4(n-6) in the rotifers and the values were highest in the case of Super Selco-enriched rotifers.

Further studies will be carried out in order to verify the effectiveness of these enrichers in larval culture.

ACKNOWLEDGEMENTS

We would like to thank Dr. M. Planas who kindly read the manuscript and gave valuable suggestions, and J.L. Garrido, Ana Ayala, Auxi Jorge and Lourdes Nieto for help with analytical procedures. This work was supported by the contract-project "Larval Nutrition of Turbot" with the fish farm Cultivo de Peces S.A. (CULTIPECSA).

REFERENCES

- Ben-Amotz, A., Fishler, R. and Schneller, A., 1987, Chemical composition of dietary species of marine unicellular algae and rotifers with emphasis on fatty acids. *Mar. Biol.*, 95: 31-36.
- Bitman, J. and Wood, D.L., 1982. An improved copper reagent for quantitative densitometric thin-layer chromatography of lipids. *J. Chromatogr.*, 5: 1155-1162.
- Bligh, E.G. and Dyer, W.L., 1959. A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.*, 37: 911-917.
- Brett, J.R. and Groves, T.D.D., 1979. Physiological energetics. In: W.S. Hoar, L. Buckley and J.R. Brett (Editors), *Fish Physiology*. Vol. VIII. Academic Press, New York, pp. 279-352.
- Christie, W.W., 1982. *Lipid Analysis*, 2nd edn. Pergamon Press, Oxford, 207 pp.
- Cowey, C.B., 1976. Use of synthetic diets and biochemical criteria in the assessment of nutrient requirements of fish. *J. Fish. Res. Board. Can.*, 33: 1040-1045.

- Dendrinou, P. and Thorpe, J.P., 1987. Experiments on the artificial regulation of the amino acid and fatty acid contents of food organisms to meet the assessed nutritional requirements of larval, post-larval and juvenile Dover sole (*Solea solea* L.). *Aquaculture*, 61: 121-154.
- Ehrlich, K.F., 1974. Chemical changes during growth and starvation of larval *Pleuronectes platessa*. *Mar. Biol.*, 24: 39-48.
- Estevez, A. and Planas, M., 1988. Efecto de diferentes fuentes alimenticias en la composition en acidos grasos del rotifero *Brachionus plicatilis* O.F. Muller (1786). *Inv. Pesq.*, 52 (1): 67-76.
- Fernandez-Reiriz, M.J., Ferreiro, M.J., Labarta, U. and Munilla, R., 1987. Influencia de dietas enriquecedoras en la calidad nutritiva de *Brachionus plicatilis* y *Artemia* sp. *Cuad. Marisq. Publ. Tee.*, 12: 579-584.
- Fernandez-Reiriz, M.J., Perez-Camacho, A., Ferreiro, M.J., Blanco, J., Planas, M., Campos, M.J. and Labarta, U., 1989. Biomass production and variation in the biochemical profile (total protein, carbohydrates, RNA, lipids and fatty acids) of seven species of marine microalgae. *Aquaculture*, 83: 17-37.
- Freeman, C.P. and West, D., 1966. Complete separation of lipid classes on a single thin-layer plate. *J. Lipid Res.*, 7: 324-327.
- Frolov, A.V., Pankov, S.L., Geradze, K.N. and Pankova, S.A., 1991. Influence of salinity on the biochemical composition of rotifer *Brachionus plicatilis* (Muller). Aspects of adaptation. *Comp. Biochem. Physiol.*, 99A: 541-550.
- Fyhn, H.J., Serigstad, B. and Mangor-Jensen, A., 1987. Free amino acids in developing eggs and yolk-sac larvae of the cod, *Gadus morhua* L. *Sarsia*, 72: 363-365.
- Gatesoupe, F.J. and Le Milinaire, C., 1984. Dietary value adaptation of live food organism for covering the nutritional requirements of marine fish larvae. In: *Acts of Norwegian-French Workshop on Aquaculture*. Brest, 5-8 December. pp. 231-243.
- Gatesoupe, F.J. and Luquet, P., 1981. Practical diet for mass culture of the rotifer *Brachionus plicatilis*. Application to larval rearing of sea bass, *Dicentrarchus labrax*. *Aquaculture*, 22: 149-163.
- Gatesoupe, F.J., Leger, C., Metailler, R. and Luquet, P., 1977. Lipid diet of the turbot (*Scophthalmus maximus* L.). 1. Influence of the chain length of ω 3 fatty acids. *Ann. Hydrobiol.*, 8: 89-98.
- Hakanson, J.L., 1989. Analysis of lipid components for determining the condition of anchovy larvae, *Engraulis mordax*. *Mar. Biol.*, 102: 143-151.

- Herraiz, M., Reglero, G., Loyola, E. and Herraiz, T., 1987. Sampling of volatile components using a PTV in the solvent split mode. *J. High Resolut. Chromatogr. Chromatogr. Comm.*, 10: 598-602.
- Howell, B.R., 1979. Experiments on the rearing of larval turbot, *Scophthalmus maximus* L. *Aquaculture*, 18: 215-227.
- Imada, O., Kageyama, Y., Watanabe, T., Kitajima, C., Fujita, S. and Yone, Y., 1979. Development of a new yeast as a culture medium for living feeds used in the production of fish feed. *Bull. Jpn. Sot. Sci. Fish.*, 45: 955-959.
- Ito, T., 1960. On the culture of the mixohaline rotifer *Brachionus plicatilis* O.F. Muller, in sea water. *Rep. Fat. Fish., Prefect. Univ. Mie*, 3: 708-740.
- James, Ch., Dias, P. and Salman, A.E., 1987. The use of marine yeast (*Candida* sp.) and baker's yeast (*Sacharomyces cerevisiae*) in combination with *Chlorella* sp. for mass culture of the rotifer *Brachionusplicatilis*. *Hydrobiologia*, 147: 263-268.
- Kanazawa, A., 1985. Essential fatty acid and lipid requirement of fish. In: C.B. Cowey, A.M. Mackie and J.G. Bell (Editors), *Nutrition and Feeding in Fish*. Academic Press, London, pp. 281-298.
- Kimata, M., 1983. Changes of chemical composition during early development in the red sea bream *Chrysophrys major* egg and larvae. *J. Fat. Mar. Sci. Technol. Tokai Univ.*, 16: 213-223
- Kissil, G. and Koven, W., 1990. Preparation of oils, enhanced in highly unsaturated fatty acid (HUFA) content, by low temperature crystallization separation, for rotifer (*Brachionusplicatilis*) enrichment. *Aquaculture*, 88: 69-74.
- Leger, C., Gatesoupe, F.J., Metailler, R., Luquet, P. and Fremont, L., 1979. Effect of dietary fatty acids differing by chain lengths and w series on the growth and lipid composition of turbot (*Scophthalmus maximus* L.). *Comp. Biochem. Physiol.*, 64B: 345-350.
- Leger, Ph., Grymonpre, D., Van Ballaer, E. and Sorgeloos, P., 1989. Advances in the enrichment of rotifers and *Artemia* as food sources in marine larviculture. In: *EAS Spec. Publ. No. 10, Eur. Aquacult. Sot., Bredene, Belgium*, pp. 141-142.
- Le Milinaire, C., Gatesoupe, F.J. and Stephan, G., 1983. Approche du besoin quantitatif en acidez gras long polyinsatures de la serie n-3 chez la larve de turbot (*Scophthalmus Maximus* L.). *C.R. Acad. Sci. Paris*, 296: 917-920.

- Linares, F. and Henderson, R.J., 1991. Incorporation of ^{14}C -labelled polyunsaturated fatty acids by juvenile turbot, *Scophthalmus maximus* (L.) in vivo. *J. Fish Biol.*, 38: 335-347.
- Linares, F., Bellver, P., Teijeiro, M.C., Perez Benavente, G. and Amoedo, F., 1990. Influencia de la utilizacion de enriquecimiento en acidos grasos sobre la composicion de rotifero (*Brachionus plicatilis*) y *Artemia* sp. y su evaluacion en condiciones de ayuno. In: *Actas III Congreso Nat. Acuicultura*, Santiago de Compostela (España), pp. 675-680.
- Love, R.M., 1980. Feeding and starving. In: R.M. Love (Editor), *The Chemical Biology of Fishes*, Vol. 2. Academic Press, London, pp. 133-229.
- Lowry, O.H., Rosebrough, N.J. and Fair, A.L., 1951. Protein measurement with the folin phenol reagent. *J. Biol. Chem.*, 193: 265-275.
- Lubzens, E., Marko, A. and Tietz, A., 1985. De novo synthesis of fatty acids in rotifer, *Brachionus plicatilis*. *Aquaculture*, 47: 27-37.
- Lubzens, E., Rothbard, S., Blumenthal, A., Kolodny, G., Perry, B., Olund, B., Wax, Y. and Farbstein, H., 1987. Possible use of *Brachionus plicatilis* (O.F. Muller) as food for freshwater cyprinid larvae. *Aquaculture*, 60: 143-155.
- Nichols, P.D., Holdsworth, D.G., Volkman, J.K., Daintith, M. and Allanson, S., 1989. High incorporation of essential fatty acids by the rotifer *Brachionus plicatilis* fed on the prymnesiophyte alga *Paulova lutheri*. *Aust. J. Mar. Freshwater Res.*, 40: 645-655.
- Owen, J.M., Adron, J.W., Middleton, C. and Cowey, C.B., 1975. Elongation and desaturation of dietary fatty acids in turbot, *Scophthalmus maximus* L., and rainbow trout, *Salmo gairdnerii* Rich. *Lipids*, 10: 528-531.
- Planas, M., Estevez, A. and Garrido, J.L., 1991. Energy metabolism during early ontogenesis of turbot (*Scophthalmus maximus*) and the effect of starvation. In: P. Lavens, P. Sorgeloos, E. Jaspers and F. Ollevier (Editors), *Larvi'91, Fish & Crustacean Larviculture Symposium*. European Aquaculture Society, Spec. Publ. No. 15, Gent, Belgium, pp. 210-212.
- Planas, M., Labarta, U., Fernandez-Reiriz, M.J., Ferreira, M.J., Munilla, R. and Garrido, J.L., 1992a. Chemical changes during early development in turbot (*Scophthalmus maximus*) eggs and larvae. In: B.T. Walther and H.J. Fyhn (Editors), *Physiology and Biochemistry of Marine Fish Larvae*. University of Bergen, Bergen, Norway (in press).

- Planas, M., Garrido, J.L., Labarta, U., Ferreiro, M.J., Fernandez-Reiriz, M.J. and Munilla, R. 1992b. Changes of fatty acid composition during early development in turbot (*Scophthalmus maximus*) eggs and larvae. In: B.T. Walther and H.J. Fyhn (Editors), *Physiology and Biochemistry of Marine Fish Larvae*. University of Bergen, Bergen, Norway (in press).
- Rodriguez-Rainuzzo, J., Olsen, Y. and Roselund, G., 1989. The effect of enrichment diets on the fatty acids composition of the rotifer *Brachionus plicatilis*. *Aquaculture*, 79: 157-161.
- Scott, A.P. and Middleton, L., 1979. Unicellular algae as a food for turbot (*Scophthalmus maximus* L.) larvae. The importance of dietary long-chain polyunsaturated fatty acids. *Aquaculture*, 18: 227-241.
- Stottrup, J., 1989. PUFA-requirements in marine fish larvae: results on a trial performed on first-feeding turbot larvae. ICES Working Group on Mass Rearing of Juvenile Marine Fish. Montpellier, 15-19 June.
- Strickland, J.D. and Parsons, T.R., 1968. A practical handbook of sea water analysis. *Bull. Fish. Res. Board Can.*, 167: 173-174.
- Tandler, A., Watanabe, T., Satoh, S. and Fukusho, K., 1989. The effect of food deprivation on the fatty acid profile of red seabream (*Pagrus major*) larvae. *Br. J. Nutr.*, 62: 349-361.
- Walford, J. and Lam, T.J., 1987. Effect of feeding with microcapsules on the content of essential fatty acids in live foods for the larvae of marine fishes. *Aquaculture*, 61: 219-229.
- Watanabe, T., Kitajama, C., Arakawa, T., Fukusho, K. and Fujita, S., 1978. Nutritional quality of rotifer, *Brachionus plicatilis* as a living feed from the viewpoint of essential fatty acids for fish. *Bull. Jpn. Soc. Sci. Fish.*, 44: 1109-1114 (in Japanese).
- Watanabe, T., Kitajama, C. and Fujita, S., 1983. Nutritional values of live organisms used in Japan for mass propagation of fish: a review. *Aquaculture*, 34: 115-143.

TABLE 1

Proximate composition (% dry weight) of the three enrichers and rotifers before and after enrichment and dry weight (ng/rotifer) of the rotifers ^{1,2}

Parameter	Initial rotifers	Protein Selco enriched			Dry Selco enriched			Super Selco enriched		
		Enricher	Rotifers	Rotifers	Enricher	Rotifers	Rotifers	Enricher	Rotifers	Rotifers
			3h	6h		3h	6h		3h	6h
Protein	36.06±0.52 ^a	32.70±1.12	34.54±0.16	38.11±0.97 ^b	4.44±0.23	34.14±1.69	37.16±2.06 ^b	0.00±0.00	36.16±1.88	39.54±0.67 ^c
Carbohydrate	16.65±0.83 ^a	33.59±0.90	16.02±1.46	15.50±2.07 ^b	1.76±0.02	15.45±1.90	14.20±0.94 ^c	0.00±0.00	15.80±1.33	13.03±0.76 ^d
Lipid	10.4±10.97 ^a	30.29±0.75	13.89±1.50	18.48±1.70 ^b	56.14±2.13	14.25±1.80	17.54±2.75 ^b	94.20±3.20	15.66±1.80	19.85±1.90 ^c
Dry weight	369.5±9.0 ^a		419.5±4.5 ^c	453.5±4.8 ^b		400.0±7.0	386.0±6.2 ^a		330.5±3.0	316.5±3.7

¹Mean ± s.d. of two replicate groups.²Means within the same rows with different superscript letters are significantly different (P< 0.05)

TABLE 2

Lipid class content (mg/g dry weight) of the three enrichers and rotifers before and after enrichment ^{1,2}

Lipid class	Initial rotifers	Protein Selco enriched			Dry Selco enriched			Super Selco enriched		
		Enricher	Rotifers	Rotifers	Enricher	Rotifers	Rotifers	Enricher	Rotifers	Rotifers
			3h	6h		3h	6h		3h	6h
Phospholipids	48.3.3±0.9 ^a	17.9±1.2	65.0±1.3	54.7±1.4 ^b	59.4±2.1	63.2±0.5	65.9±0.7 ^c	49.4±2.1	57.1±0.9	91.1±1.2 ^d

Sterol esters waxes	21.9±1.1 ^a	ND	28.6±1.0	25.4±1.0 ^b	ND	28.4±0.5	36.1±0.5 ^c	ND	39.2±0.6	45.6±1.0 ^d
Triacylglycerols	18.7±1.2 ^a	32.9±1.1	17.5±1.1	45.6±0.9 ^b	ND	18.1±0.6	38.7±0.4 ^c	ND	28.8±0.4	51.1±0.9 ^d
Free fatty acids	9.7±0.2 ^a	30.6±0.9	14.8±0.2	21.2±0.9 ^b	36.7±1.2	20.6±0.7	20.6±0.2 ^c	ND	28.7±0.3	42.2±0.1 ^d
Sterols	3.7±0.3 ^a	ND	2.7±0.1	2.9±0.1 ^b	ND	4.1±0.1	4.0±0.1 ^c	ND	3.9±0.1	3.7±0.1 ^d
Akyl glyceryl ethers	ND	204.0±3.1	ND	ND	411.0±4.6	ND	ND	687.1±5.6	ND	ND

¹Mean ± s.d. of two replicate groups. ND indicates not detected.

²Means within the same row with different superscript letters are significantly different (P<0.05).

TABLE 3

Fatty acid composition (mg/g dry weight) of three enrichers and rotifers before and after enrichment ^{1,2}

Fatty acid	Initial rotifers	Protein Selco enriched			Dry Selco enriched			Super Selco enriched		
		Enricher	Rotifers 3h	Rotifers 6h	Enricher	Rotifers 3h	Rotifers 6h	Enricher	Rotifers 3h	Rotifers 6h
14:0	0.57±0.01 ^a	5.30±0.02	1.70±0.42 ^b	1.70±0.04	8.40±0.03	1.20±0.43	1.79±0.23 ^b	4.20±0.03	1.05±0.45	1.05±0.96 ^c
16:0	4.20±0.18 ^a	15.50±0.10	6.10±0.50 ^b	6.22±0.10	26.70±0.60	4.62±1.80	6.51±0.88 ^b	8.80±0.90	4.17±1.99	4.38±0.90 ^a

16:1(n-9)	0.24±0.03 a	0.00±0.00	0.40±0.01 ^b	0.00±0.00	0.00±0.00	0.23±0.33	0.17±0.02 ^a	0.00±0.00	0.26±0.37	0.00±0.00 ^c
16:1(n-7)	6.65±0.36 a	7.50±0.20	8.49±0.70 ^b	8.26±1.24	9.70±0.90	6.35±2.33	7.74±1.35 ^c	0.00±0.00	6.07±1.44	6.08±1.23 ^a
17:0	0.20±0.03 a	1.40±0.01	0.55±0.01 ^b	0.47±0.07	1.70±0.01	0.41±0.07	0.53±0.08 ^b	2.30±1.01	0.36±0.06	0.40±0.30 ^c
17:1(n-7)	0.40±0.01 a	1.00±0.01	0.62±0.02 ^b	0.53±0.14	2.50±0.20	0.41±0.06	0.29±0.04 ^a	2.80±0.91	0.26±0.37	0.00±0.00 ^c
18:0	2.84±0.13 a	6.00±0.50	3.60±0.10 ^b	3.44±0.58	8.30±0.90	2.87±1.02	3.56±0.39 ^b	16.00±0.99	3.29±0.96	4.00±0.99 ^b
18:1(n-11)	1.08±0.05 a	0.00±0.00	1.18±0.05 ^a	0.99±0.26	0.00±0.00	1.06±0.41	1.18±0.06 ^a	0.00±0.00	1.03±0.55	1.05±0.63 ^a
18:1(n-9)	9.53±0.4 ^a	14.20±0.70	12.77±1.12 ^b	12.49±2.22	22.20±1.10	9.41±2.77	12.10±1.35 ^b	27.60±2.34	10.21±2.64	15.90±2.51 ^c
18:1(n-7)	1.24±0.09 a	2.80±0.08	1.55±0.04 ^a	1.72±0.18	6.90±0.30	1.32±0.86	1.63±0.13 ^a	8.50±1.10	1.65±0.91	2.30±0.56 ^b
18:2(n-6)	5.30±0.2 ^a	10.60±0.09	6.78±0.17 ^b	6.55±1.36	10.90±1.1	05.66±1.79	7.80±1.11 ^c	6.10±0.90	5.23±1.92	6.41±1.23 ^b
18:3(n-3)	0.36±0.01 a	1.60±0.05	0.55±0.50 ^b	0.52±0.00	2.30±0.07	0.21±0.29	0.61±0.02 ^{bc}	2.50±0.99	0.26±0.36	0.90±0.02 ^c
20:1(n-9)	1.29±0.04 a	5.20±0.01	1.75±0.21 ^{ab}	2.16±0.49	4.70±0.99	1.34±0.37	1.90±0.10 ^{ab}	7.30±1.10	1.64±0.49	2.20±0.99 ^b
20:2(n-6)	0.60±0.01 a	0.00±0.05	0.76±0.37 ^a	0.66±0.24	0.00±0.00	0.31±0.44	0.74±0.06 ^a	0.00±0.00	0.59±0.21	0.60±0.32 ^a

20:3(n-6)	1.00±0.03 a	0.00±0.00	1.08±0.10 ^a	0.93±0.34	0.00±0.00	1.02±0.29	1.04±0.19 ^a	0.00±0.00	1.15±0.16	0.80±0.23 ^a
20:4(n-6)	0.58±0.02 a	2.70±0.04	0.81±1.04 ^a	0.93±0.04	0.00±0.00	0.63±0.25	0.88±0.55 ^a	7.30±1.26	0.95±0.54	1.40±0.05 ^b
20:S(n-3)	2.25±0.13 a	35.10±0.70	6.28±1.04 ^b	8.20±0.99 ^c	57.40±2.00	4.27±1.89	8.38±0.06 ^c	96.70±3.20	6.90±0.99	13.80±1.23 ^d
22:1(n-9)	0.33±0.01 a	0.00±0.00	0.73±0.41 ^b	0.49±0.30	0.00±0.00	0.26±0.06	0.59±0.06 ^b	3.90±1.30	0.31±0.43	0.40±0.03 ^a
22:2(n-6)	0.33±0.05 a	0.00±0.00	0.00±0.00 ^b	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00 ^b	0.00±0.00	0.21±0.03	0.00±0.00 ^b
22:5(n-3)	0.85±0.04 a	5.10±0.20	1.71±0.13 ^b	1.96±0.16	7.40±0.05	1.22±0.00	1.77±0.20 ^b	16.40±1.56	1.53±0.97	2.90±0.36 ^c
22:6(n-3)	1.31±0.03 a	34.70±0.90	5.22±0.11 ^{ab}	6.52±0.26	54.60±2.10	2.90±0.22	5.59±0.71 ^b	114.60±2.30	6.67±2.36	12.80±2.96 ^c
24:1(n-9)	0.55±0.01 a	1.20±0.09	0.56±0.16 ^a	0.81±0.50	2.90±0.30	0.27±0.06	0.71±0.10 ^a	4.30±1.20	0.32±0.45	0.70±0.01 ^a
Σ Saturated	7.81±0.32 a	28.30±0.51	11.95±0.51 ^b	11.83±0.59	45.10±1.08	9.10±2.11	12.39±0.99 ^c	31.30±1.68	8.87±2.25	9.83±1.67 ^d
Σ Monoenes	21.31±1.0 6 ^a	31.90±0.74	28.05±1.40 ^b	27.45±2.65	48.90±1.79	20.65±3.78	26.3±1.92 ^c	54.40±3.44	21.75±3.32	28.63±3.08 ^b
Σ	12.58±0.4 9 ^a	89.80±1.16	23.19±1.62 ^b	26.27±1.75	132.60±3.10	16.22±2.69	26.81±1.45 ^c	243.60±4.61	23.49±3.41	39.61±3.47 ^d

Polyunsat.										
$\Sigma n-6$	7.81±0.30 ^a	13.30±0.11	9.43±1.12 ^b	9.07±1.42	10.90±1.10	7.62±1.88	10.46±1.25 ^c	13.40±1.54	8.13±2.01	9.21±1.29 ^b
n-3 HUFA	4.41±0.21 ^a	74.90±1.14	13.21±1.05 ^b	16.68±1.03	119.40±2.00	8.39±1.90	15.74±0.21 ^c	227.70±4.23	15.10±2.74	29.50±3.22 ^d
n-3 ± n-6	0.61±0.01 ^a	5.76±0.13	1.46±0.12 ^b	1.90±0.29	11.17±0.10	1.13±0.22	1.56±0.12 ^b	17.18±0.10	1.89±0.45	3.30±0.33 ^c

¹Mean ± s.d. of two replicate groups.

²Means within the same rows with different superscript letters are significantly different (P<0.05).

TABLE 4

Caloric content of the rotifers (Cal/10³ rotifers) before and after enrichment1

Treatment	Protein	Carbohydrate	Lipid	Total
0h	0.75	0.25	0.34	1.34
3h				
Protein Selco	0.82	0.28	0.50	1.60
Dry Selco	0.77	0.25	0.49	1.51
Super Selco	0.68	0.21	0.45	1.34
6h				

Protein Selco	0.98	0.29	0.73	2.00
Dry Selco	0.81	0.22	0.59	1.62
Super Selco	0.71	0.17	0.54	1.42

¹The conversion factors used to calculate these values were 5.65, 4.10 and 8.66 cal/mg for protein, carbohydrate, and lipid, respectively (Brett and Groves, 1979).