

## Graphical Abstract


### Gold-catalyzed direct cycloketalization of acetonide-tethered alkynes in the presence of water

Leave this area blank for abstract info.

Benito Alcaide,<sup>a,\*</sup> Pedro Almendros,<sup>b,\*</sup> and Rocío Carrascosa<sup>a</sup>

<sup>a</sup> Grupo de Lactamas y Heterociclos Bioactivos, Departamento de Química Orgánica I, Unidad Asociada al CSIC, Facultad de Química, Universidad Complutense de Madrid, 28040 Madrid

<sup>b</sup> Instituto de Química Orgánica General, IQOG, Consejo Superior de Investigaciones Científicas, CSIC, Juan de la Cierva 3, 28006 Madrid, Spain





# Gold-catalyzed direct cycloketalization of acetonide-tethered alkynes in the presence of water

Benito Alcaide,<sup>a,\*</sup> Pedro Almendros,<sup>b,\*</sup> and Rocío Carrascosa<sup>a</sup>

<sup>a</sup> Grupo de Lactamas y Heterociclos Bioactivos, Departamento de Química Orgánica I, Unidad Asociada al CSIC, Facultad de Química, Universidad Complutense de Madrid, 28040 Madrid

<sup>b</sup> Instituto de Química Orgánica General, IQOG, Consejo Superior de Investigaciones Científicas, CSIC, Juan de la Cierva 3, 28006 Madrid, Spain

## ARTICLE INFO

### Article history:

Received  
Received in revised form  
Accepted  
Available online

### Keywords:

Acetals  
Alkynes  
Cyclization  
Gold  
Regioselectivity

## ABSTRACT

A methodology for the direct preparation of bridged acetals from acetonide-tethered alkynes under gold catalysis in the presence of water has been developed. The bicyclic ring structures bearing a bridged five-membered ring arise from the regioselective bis-oxycyclization by initial attack of the oxygen atom to the internal alkyne carbon.


2011 Elsevier Ltd. All rights reserved.

## 1. Introduction

Over the course of the last decade, efforts aimed at maximizing the total efficiency of a given organic transformation has been increased.<sup>1</sup> The use of gold salts has gained a lot of attention in the recent times because of their powerful soft Lewis acidic nature.<sup>2</sup> In particular, the recently developed noble metal activation of alkynes toward simultaneous attack by two contiguous oxygen nucleophiles is an important C–O bond-forming reaction.<sup>3,4</sup> However, these reports involves the use of free hydroxy groups; being missed the direct bis(oxycyclization) sequence of acetonide-tethered alkynes until we merged into this field.<sup>5</sup> In connection with our current research interest in the preparation of biologically relevant heterocycles,<sup>6</sup> we wish to report now details of the direct cycloketalization of acetonide-tethered alkynes to bridged bicyclic acetals which is carried out using gold catalysis in the presence of water.

## 2. Results and discussion

Starting substrates, alkynyldioxolanes **1a–h**, **2a–e**, and **3a–d** (Figure 1) required for our study were prepared from (*R*)-2,3-*O*-isopropylidene-glyceraldehyde, from (*S*)-(2,2-dimethyl-1,3-dioxolan-4-yl)methanol, from (*S*)-2-(2,2-dimethyl-1,3-dioxolan-4-yl)ethanol, or from 2-*O*-benzyl-D-threitol.




**Figure 1.** Structures of cyclization precursors, alkynyldioxolanes **1a–h**, **2a–e**, and **3a–d**. PMP = 4-MeOC<sub>6</sub>H<sub>4</sub>.

Alkynyldioxolanes **1a–h** were prepared as shown in Scheme 1. Alkynyldioxolanes **1a** and **1b** were prepared from (*S*)-(+)-1,2-isopropylidene-glycerol through *O*-propargylation and Sonogashira coupling. Precursors **1c–h** were prepared from (*R*)-2,3-*O*-isopropylidene-glyceraldehyde via metal-mediated carbonyl-allylation followed by *O*-propargylation, deuteration, Sonogashira and Cadiot–Chodkiewicz coupling, as appropriate.


\* Corresponding author. Tel.: (+34) 91-3944314; fax: (+34) 91-3944103; e-mail: alcaideb@quim.ucm.es (B. Alcaide)

\* Corresponding author. Tel.: (+34) 91-5618806; fax: (+34) 91-5644853; e-mail: Palmendros@iqog.csic.es (P. Almendros)




**Scheme 1.** Reagents and conditions: i) Propargyl bromide,  $\text{CH}_2\text{Cl}_2/\text{NaOH}$  (aq. 50%), TBAI, RT, 24 h. ii) 2 mol% CuI, 1 mol%  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ , ArI,  $\text{Et}_3\text{N}$ , acetonitrile, RT, 15 h. iii) In, THF/ $\text{NH}_4\text{Cl}$  (aq. sat.), RT, 15 h. iv) BuLi,  $\text{D}_2\text{O}$ , THF,  $-78^\circ\text{C}$  to RT. v) NBS, AcOAg, acetone. vi) Phenylacetylene, CuCl,  $\text{NH}_2\text{OH}$ ,  $\text{EtNH}_2$ ,  $\text{CH}_2\text{Cl}_2/\text{MeOH}$ ,  $0^\circ\text{C}$  to RT. vii) 1-Bromopent-2-yne,  $\text{CH}_2\text{Cl}_2/\text{NaOH}$  (aq. 50%), TBAI, RT, 24 h.

Alkyndioxolanes **2a–e** were prepared as shown in Scheme 2. Alkyndioxolanes **2a–b** were prepared from (4S)-(+)-4-(2-hydroxyethyl)-2,2-dimethyl-1,3-dioxolane through *O*-propargylation. Precursors **2c–e** were prepared from 2-*O*-benzyl-D-threitol<sup>7</sup> via *O*-isopropylidensation<sup>8</sup> followed by *O*-propargylation and Sonogashira coupling.




**Scheme 2.** Reagents and conditions: i) Propargyl bromide or (3-bromoprop-1-ynyl)benzene,  $\text{CH}_2\text{Cl}_2/\text{NaOH}$  (aq. 50%), TBAI, RT, 24 h. In, THF/ $\text{NH}_4\text{Cl}$  (aq. sat.), RT, 15 h. ii) 10 mol% pyridinium *p*-toluenesulfonate, 2,2-dimethoxypropane, RT, 2 h. iii) Propargyl bromide or 1-bromobut-2-yne, NaH, TBAI, THF, RT. iv) 2 mol% CuI, 1 mol%  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ , ArI,  $\text{Et}_3\text{N}$ , acetonitrile, RT, 15 h. TBAI = Tetrabutylammonium iodide.

As shown in Scheme 3, alkyndioxolanes **3a–d** were prepared from (R)-2,3-*O*-isopropylidenedeglyceraldehyde through reductive amination followed by protection and Sonogashira coupling, as appropriate.



**Table 1.** Selective direct bis(oxycyclization) reaction of acetonide-tethered alkyne **1a** under modified metal-catalyzed conditions.<sup>a</sup>




**Scheme 3.** Reagents and conditions: i) Propargyl amine,  $\text{MgSO}_4$ , RT, 15 h. ii)  $\text{NaBH}_4$ , methanol,  $0^\circ\text{C}$ , 30 min. iii) Benzylbromide,  $\text{K}_2\text{CO}_3$ , acetonitrile, RT. iv) Methyl chloroformate,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ , RT, 2h. v) 2 mol% CuI, 1 mol%  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ , ArI,  $\text{Et}_3\text{N}$ , acetonitrile, RT, 15 h.

Using directly a dioxolane ring instead of the deprotected 1,2-diol moiety as the starting material, would be a significant breakthrough for metal-catalyzed alkyne-cycloketalization in terms of cost-effectiveness. Initial experiments were carried out using alkyndioxolane **1a** as a model substrate. To optimise the reaction method, different parameters involved in the metal-catalyzed reactions that could affect the formation of the desired product were tested. Attempts of the cyclization reaction of **1a** using AgOTf,  $\text{FeCl}_3$ , and  $[\text{PtCl}_2(\text{CH}_2=\text{CH}_2)]_2$  catalysts failed. Nicely, it was found that  $[\text{AuClPPH}_3]/\text{AgOTf}$  along with a Brønsted acid (PTSA) in  $\text{CH}_2\text{Cl}_2$  was a competent catalytic system for this purpose (see Table).<sup>9,10</sup> Performing the reaction at  $80^\circ\text{C}$  on a sealed tube accelerated the reaction. However, the conversion to the corresponding tricyclic acetal **4a** could not be satisfied in high yield. Solvent screening demonstrated that dichloromethane was the best choice in the reaction. The use of a more Lewis acidic gold salt ( $\text{AuCl}_3$ ) as catalyst alone without PTSA was also tested. Thus, we treated alkyne acetonide **1a** with 5 mol% of  $\text{AuCl}_3$ . However, acetal **4a** was obtained in a poor 25% yield. After identifying an appropriate catalyst, we optimized other critical reaction parameters, such as the addition of additives. In particular, the transformation was strongly influenced by the presence of water. Gratifyingly, it was found that alkyne acetonide **1a** on exposure to the system  $[\text{AuClPPH}_3]$  (2.5 mol%)/AgOTf (2.5 mol%)/PTSA (10 mol%)/ $\text{H}_2\text{O}$  (100 mol%) in dichloromethane at  $80^\circ\text{C}$  on a sealed tube, directly afforded bicyclic ketal **4a** through a regio- and diastereoselective bis(oxycyclization) (see Table 1). Probably, the presence of water favors the hydrolysis of intermediate **10** in the mechanistic proposal of Scheme 4. Taking into account that there are only a few gold-catalyzed transformations which tolerate the presence of water,<sup>11</sup> there would be certain merit using water as an alternative solvent for alkyne-cycloketalization.<sup>12</sup> To test for conditions for the reaction as a suspension in water, we try our optimized catalytic system. Notably, substrate **1a** is completely insoluble in aqueous media. After some experimentation, we observed that the cyclization reaction did not proceed “on water”. Thus, the use of biphasic water/dichloromethane mixtures was necessary in order to improve the solubility of the substrates; being (4 water: 1 dichloromethane) the best solvent system. The presence of a phase transfer agent<sup>13</sup> gave rise to a positive influence on the yield of bridged acetal **4a**.<sup>14</sup> The reaction time has a crucial role in the amount of the product obtained: the longer the reaction time, the better the yield, with a best result of 90% at 48 hours, because a short reaction time of 2 hours leads to a 20% yield.

Catalyst (mol %)	Additive (mol %)	t (h)	Acid	T (°C)	Solvent	Yield (%) <sup>f</sup>
[PtCl <sub>2</sub> (CH <sub>2</sub> =CH <sub>2</sub> ) <sub>2</sub> (2.5)	TDMPP (5)	24	–	20	DCM	–
AgOTf (5)	–	24	–	20	DCM	– <sup>b</sup>
FeCl <sub>3</sub> (10)	–	24	–	20	DCM	–
AuCl <sub>3</sub> (5)	–	24	PTSA	80	DCM	25
[AuClPPH <sub>3</sub> ]/AgOTf (2.5)	–	24	PTSA	20	DCM	45
[AuClPPH <sub>3</sub> ]/AgOTf (2.5)	H <sub>2</sub> O (100)	3	–	80	DCM	75
[AuClPPH <sub>3</sub> ]/AgOTf (2.5)	H <sub>2</sub> O (100)	48	FeCl <sub>3</sub>	80	DCM	50
[AuClPPH <sub>3</sub> ]/AgOTf (2.5)	H <sub>2</sub> O (100)	4	PTSA	80	THF	64
[AuClPPH <sub>3</sub> ]/AgOTf (2.5)	H <sub>2</sub> O (100)	3	PTSA	80	DCE	90
[AuClPPH <sub>3</sub> ]/AgOTf (2.5)	H <sub>2</sub> O (100)	5	PTSA	80	toluene	760
[AuClPPH <sub>3</sub> ]/AgOTf (2.5)	TBAI (10)	48	PTSA	80	4H <sub>2</sub> O : 1DCM	90
[AuClPPH <sub>3</sub> ]/AgOTf (2.5)	H <sub>2</sub> O (100)	3	PTSA	80	DCM	100

<sup>a</sup> Yield of pure, isolated product with correct analytical and spectral data. PTSA = *p*-Toluenesulfonic acid. TDMPP = tris(2,6 dimethoxyphenyl)phosphine. DCM = Dichloromethane. THF = Tetrahydrofuran. DCE = 1,2-Dichloroethane. TBAI = Tetrabutylammonium iodide. <sup>b</sup> Taking into account this result, AgOTf cannot be considered as a co-catalyst because its action is restricted to form the cationic gold species by anion exchange.<sup>7</sup>


Having established the optimal reaction conditions, we explored the scope of the methodology by subjecting a range of (prop-2-ynyloxy)methyl-tethered dioxolanes **1b–h** to direct alkynyl-cycloketalization and the results are shown in Scheme 4. In both cases (addition of one equivalent of water as well as the presence of large amount of water), the crude reaction mixtures are extremely clean and the acetals are the only products detected. Tolerance toward a variety of substituents (aliphatic, aromatic, and alkynyl groups) on the acetylenic end was demonstrated by obtaining the corresponding enantiopure bridged acetals **4** in good yields. Except for ethyl-substituted alkynyl dioxolane **1h**, bicyclic ketals **4** were obtained as single regio- and diastereomers through a 6-*exo*/5-*exo* bis-oxycyclization by initial attack of the oxygen atom to the internal alkyne carbon. The exposure of ethyl-substituted alkyne **1h** to these reactions conditions afforded the 7-*endo*/5-*exo* bis-oxycyclization bridged acetal **5h** as major adduct, being the corresponding 6-*exo*/5-*exo* adduct **4h** the minor component. From these experiments, it can be concluded that this catalytic system in the presence of water enables regioselective control of the intramolecular ketalization reaction between the dioxolane unit and a range of alkynes separated by three atoms. Besides, the method using one equivalent of water is superior to the method using water as main solvent, in terms of yields and reaction time.



**Scheme 4.** Preparation of 3,6,8-trioxabicyclo[3.2.1]octanes **4**. PMP = 4-MeOC<sub>6</sub>H<sub>4</sub>. PTSA = *p*-Toluenesulfonic acid. TBAI = Tetrabutylammonium iodide.


We further examined the scope of this methodology by investigating (prop-2-ynyloxy)ethyl-tethered dioxolanes **2a–e**, alkynyl dioxolane substrates with a four-atom chain at the junction between the acetonide group and the alkyne. We were pleased to observe that these modifications had little impact on the regiochemical control of the reaction (total regioselectivity was observed in this series). Thus, the increase of the distance

between the reactive moieties in one atom favored the 7-*exo*/5-*exo* bis-oxycyclization of the acetonide group towards the internal alkyne carbon, affording proximal adducts **6** (Scheme 5).



**Scheme 5.** Preparation of 3,8,9-trioxabicyclo[4.2.1]nonanes **6**. PMP = 4-MeOC<sub>6</sub>H<sub>4</sub>. PTSA = *p*-Toluenesulfonic acid. TBAI = Tetrabutylammonium iodide.


Aza-analogues of bridged acetals **4–6** have been recently identified as inhibitors of drug-resistant *Candida albicans* strains.<sup>15</sup> Thus, the replacement of the cycloether oxygen for a nitrogen atom was accomplished next. Similar trends to the oxo series were observed under the same gold-catalyzed conditions in the presence of water. Cyclization of aminoalkynes **3a–d** possessing a chiral dioxolane ring results in the regio- and stereoselective formation of 6,8-dioxa-3-azabicyclo[3.2.1]octanes **7** (Scheme 6). Bicyclic acetals **7** were obtained as exclusive reaction products, except for **3c**. Although substitution at the final position of the aminoalkyne had little effect upon the regioselectivity of the reaction, alkynyl dioxolane **7c** underwent cycloketalization reaction to afford a mixture of proximal and distal adducts **7c** and **8c**.




**Scheme 6.** Preparation of 6,8-dioxa-3-azabicyclo[3.2.1]octanes **7**. Thf = 2-Thiophenyl. PTSA = *p*-Toluenesulfonic acid. TBAI = Tetrabutylammonium iodide.

Our proposed mechanism is shown in Scheme 7. Initially, one of the oxygen atoms of the alkynyldioxolanes **1–3** attacks through (6 + *n*)-*exo* alkoxyauration the gold-activated alkyne **9** to

form a vinyl-gold complex **10**, which reacts with water through acetonide hydrolysis and a deauration process, to give intermediate **11** (right circle) and releasing the gold catalyst. The reaction does not stop at this stage; instead methylenic oxacycles **11** can be further cyclized to give the functionalized bridged acetals **4–6** (left circle). The second catalytic cycle starts with the coordination of the alkene group by the gold salt, on forming the  $\pi$ -complex **12**, which after 5-*exo* cyclization would form the ate complex **13**. Deauration linked to proton transfer liberates adducts **4–6** with concomitant regeneration of the gold catalyst, closing the second catalytic cycle (Scheme 7). Because we had never isolated **11**, it is also possible to think in a Brønsted acid activation of the transient methylenic intermediate **11**. However, the successful gold-catalyzed cyclo-ketalization reaction of alkynylacetone in the absence of Brønsted acid additive, may discard this assumption. With the aim of trapping the organometal intermediate to confirm the mechanism of this reaction, we performed deuterium labeling studies with deuterium oxide. When the gold-catalyzed alkyne cyclo-ketalization reaction of substrate **1e** was carried out in presence of D<sub>2</sub>O (4 heavy water: 1 dichloromethane), adduct **4e** with additional deuterium incorporation (50% D) at the two protons of the benzylic carbon was achieved (Scheme 8). The fact that the gold-catalyzed conversion of alkyne **1e** into bridged acetal **4e** in the presence of D<sub>2</sub>O partially afforded double deuterated [D]-**4e**, suggests that deuterolysis of the carbon-gold bonds in species **10** and **13** has occurred.



**Scheme 7.** A conceivable mechanistic explanation for the gold-catalyzed bis(oxycyclization) of alkynyldioxolanes **1–3**.



**Scheme 8.** Gold-catalyzed bis(oxycyclization) of alkynyldioxolane **1e** in a heavy water medium.

### 3. Conclusions

In conclusion, we have developed a convenient methodology for the gold-catalyzed direct synthesis of bridged bicyclic acetals from alkynyldioxolanes. The presence of water was beneficial for the transformation. The method using one equivalent of water is superior to the method using water as main solvent, in terms of yields and reaction time. A conceivable mechanism for the achievement of bridged acetals may imply a regioselective bis-oxycyclization by initial attack of the oxygen atom of the dioxolane moiety to the internal alkyne carbon.

## 4. Experimental section

### 4.1. General information

<sup>1</sup>H NMR and <sup>13</sup>C NMR spectra were recorded on a Bruker Avance AVIII-700 with cryoprobe, Bruker AMX-500, Bruker Avance-300, Varian VRX-300S or Bruker AC-200. NMR spectra were recorded in CDCl<sub>3</sub> solutions, except otherwise stated. Chemical shifts are given in ppm relative to TMS (<sup>1</sup>H, 0.0 ppm), or CDCl<sub>3</sub> (<sup>13</sup>C, 77.0 ppm). Low and high resolution mass spectra were taken on an AGILENT 6520 Accurate-Mass QTOF LC/MS spectrometer using the electronic impact (EI) or electrospray modes (ES) unless otherwise stated. IR spectra were recorded on a Bruker Tensor 27 spectrometer. Specific rotation [ $\alpha$ ]<sub>D</sub> is given in 10<sup>-1</sup> deg cm<sup>2</sup> g<sup>-1</sup> at 20 °C, and the concentration (*c*) is expressed in g per 100 mL. All commercially available compounds were used without further purification.

### 4.2. Method A. General procedure for the gold-catalyzed bis(heterocyclization) reaction of alkynyldioxolanes **1, 2 and 3** using one equivalent of water. Preparation of bridged acetals **4, 5, 6 and 7**.

[AuClPPh<sub>3</sub>] (0.0093 mmol), AgOTf (0.0093 mmol), and *p*-toluenesulfonic acid (0.037 mmol) were sequentially added to a stirred solution of the corresponding alkynyldioxolane **1** (0.37 mmol) in dichloromethane (0.37 mL). The resulting mixture was heated in a sealed tube at 80 °C for 3 h. The reaction was allowed to cool to room temperature and filtered through a pack of celite. The filtrate was extracted with dichloromethane (3 x 5 mL), and the combined extracts were washed twice with brine. The organic layer was dried (MgSO<sub>4</sub>) and concentrated under reduced pressure. Chromatography of the residue gave analytically pure adducts. Spectroscopic and analytical data for pure forms of **4, 5, 6 and 7** follow.<sup>16</sup>

### 4.3. Method B. General procedure for the gold-catalyzed bis(heterocyclization) reaction of alkynyldioxolanes **1, 2 and 3** using water as main solvent. Preparation of bridged acetals **4, 5, 6 and 7**.

[AuClPPh<sub>3</sub>] (0.0093 mmol), AgOTf (0.0093 mmol), *p*-toluenesulfonic acid (0.037 mmol), and tetrabutylammonium iodide (0.037 mmol) were sequentially added to a stirred solution of the corresponding alkynyldioxolane **1** (0.37 mmol) in a biphasic water (4 mL)/dichloromethane (1 mL) mixture. The resulting mixture was heated in a sealed tube at 80 °C for 48 h. The reaction was allowed to cool to room temperature and filtered through a pack of celite. The filtrate was extracted with dichloromethane (3 x 5 mL), and the combined extracts were washed twice with brine. The organic layer was dried (MgSO<sub>4</sub>) and concentrated under reduced pressure. Chromatography of the residue gave analytically pure adducts. Spectroscopic and analytical data for pure forms of **4, 5, 6 and 7** follow.

**4.3.1. Bicyclic acetal (–)-4a.** From 129 mg (0.76 mmol) of alkynyldioxolane (+)-**1a**, and after chromatography of the residue using petroleum ether/diethyl ether (3:1) as eluent gave the ketal (–)-**4a** [method A : 99 mg, 100%; method B : 89 mg, 90%] as a colorless oil; [ $\alpha$ ]<sub>D</sub> = –0.5 (*c* = 10.0 in CHCl<sub>3</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>, 25 °C):  $\delta$  = 4.42 (m, 1H), 4.23 (d, *J* = 6.6 Hz, 1H), 3.87 (m, 2H), 3.60 (dd, *J* = 11.4, 1.3 Hz, 1H), 3.54 and 3.49 (d, *J* = 11.1 Hz, each 1H), 1.36 (s, 3H); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>, 25 °C):  $\delta$  = 104.9, 74.6, 72.6, 68.5, 67.9, 19.3; IR (CHCl<sub>3</sub>):  $\nu$  = 2924, 1101, 693 cm<sup>-1</sup>; HRMS (ES): calcd for C<sub>6</sub>H<sub>11</sub>O<sub>3</sub> [*M* + H]<sup>+</sup>: 131.0708; found: 131.0705.

**4.3.2. Bicyclic acetal (–)-4d.** From 60 mg (0.25 mmol) of alkynyldioxolane (+)-**1c**, and after chromatography of the residue using hexanes/ethyl acetate (7:1) as eluent gave the ketal (–)-**4d** [method A : 38 mg, 76%; method B : 33 mg, 66%] as a yellow oil;

$[\alpha]_D = -0.6$  ( $c = 2.5$  in  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (300 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 5.96$  (dd,  $J = 17.7$ , 11.0 Hz, 1H), 5.00 (m, 2H), 4.38 (m, 2H), 3.73 (dd,  $J = 7.0$ , 5.7 Hz, 1H), 3.62 and 3.56 (d,  $J = 11.0$  Hz, each 1H), 3.53 (s, 1H), 1.33 (t, 2H), 1.07 and 1.06 (s, each 3H);  $^{13}\text{C NMR}$  (75 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 144.3$ , 112.3, 104.3, 84.3, 75.6, 74.2, 66.4, 38.8, 24.7, 23.8, 18.6 (t,  $J = 19.6$  Hz,  $\text{CH}_2\text{D}$ ); IR ( $\text{CHCl}_3$ ):  $\nu = 1094$ , 653  $\text{cm}^{-1}$ ; HRMS (ES): calcd for  $\text{C}_{11}\text{H}_{18}\text{DO}_3$  [ $M + \text{H}$ ] $^+$ : 200.1396; found: 200.1391.

**4.3.3. Bicyclic acetal (+)-4f.** From 55 mg (0.17 mmol) of alkynyldioxolane (–)-**1f**, and after chromatography of the residue using hexanes/ethyl acetate (2:1) as eluent gave the ketal (+)-**4f** [method A : 30 mg, 65%; method B : 27 mg, 58%] as a colorless oil;  $[\alpha]_D = +0.3$  ( $c = 1.8$  in  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (300 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 8.54$  (d,  $J = 3.8$  Hz, 1H), 7.62 (td,  $J = 7.6$ , 0.4 Hz, 1H), 7.34 (d,  $J = 7.6$  Hz, 1H), 7.16 (m, 1H), 5.94 (dd,  $J = 17.7$ , 11.0 Hz, 1H), 4.97 (m, 2H), 4.40 (m, 2H), 3.68 (m, 1H), 3.71 and 3.63 (d,  $J = 11.1$  Hz, each 1H), 3.52 (s, 1H), 3.22 and 3.17 (d,  $J = 14.0$  Hz, each 1H), 1.04 and 1.03 (s, each 3H);  $^{13}\text{C NMR}$  (75 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 155.6$ , 148.9, 144.2, 136.3, 124.9, 121.8, 112.3, 105.0, 84.3, 75.7, 73.2, 66.5, 42.2, 38.9, 24.7, 23.8; IR ( $\text{CHCl}_3$ ):  $\nu = 1179$ , 1045  $\text{cm}^{-1}$ ; HRMS (ES): calcd for  $\text{C}_{16}\text{H}_{22}\text{NO}_3$  [ $M + \text{H}$ ] $^+$ : 276.1600; found: 276.1603.

**4.3.4. Bicyclic acetal (+)-6a.** From 151 mg (0.82 mmol) of alkynyldioxolane (+)-**2a**, and after chromatography of the residue using petroleum ether/diethyl ether (2:1) as eluent gave the ketal (+)-**6a** [method A : 66 mg, 56%; method B : 53 mg, 45%] as a colorless oil;  $[\alpha]_D = -0.01$  ( $c = 2.4$  in  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (300 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 4.60$  (td,  $J = 6.9$ , 2.9 Hz, 1H), 4.11 (t,  $J = 7.0$  Hz, 1H), 4.04 and 3.69 (ddd,  $J = 12.4$ , 6.1, 3.1 Hz, each 1H), 3.98 (dd,  $J = 6.9$ , 2.9 Hz, 1H), 3.78 and 3.39 (d,  $J = 12.4$  Hz, each 1H), 2.13 (m, 1H), 1.87 (dddd,  $J = 10.2$ , 6.1, 0.9 Hz, 1H), 1.35 (s, 3H);  $^{13}\text{C NMR}$  (75 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 110.6$ , 79.0, 74.2, 73.1, 68.1, 36.7, 21.5; IR ( $\text{CHCl}_3$ ):  $\nu = 1190$   $\text{cm}^{-1}$ ; HRMS (ES): calcd for  $\text{C}_7\text{H}_{13}\text{O}_3$  [ $M + \text{H}$ ] $^+$ : 145.0865; found: 145.0868.

**4.3.5. Bicyclic acetal (–)-7b.** From 56 mg (0.25 mmol) of alkynyldioxolane (–)-**3b**, and after chromatography of the residue using hexanes/ethyl acetate (1:1) as eluent gave the ketal (–)-**7b** [method A : 37 mg, 80%; method B : 34 mg, 73%] as a colorless oil;  $[\alpha]_D = -0.1$  ( $c = 1.4$  in  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (300 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 4.46$  (m, 1H), 3.85 (m, 3H), 3.77 and 2.92 (d,  $J = 5.6$  Hz, 1H), 3.69 (s, 3H), 2.92 (d,  $J = 12.8$  Hz, 1H), 1.44 (s, 3H);  $^{13}\text{C NMR}$  (75 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 159.5$ , 107.0, 75.9, 71.2, 55.6, 54.9, 49.8, 24.1; IR ( $\text{CHCl}_3$ ):  $\nu = 1744$   $\text{cm}^{-1}$ ; HRMS (ES): calcd for  $\text{C}_8\text{H}_{14}\text{NO}_4$  [ $M + \text{H}$ ] $^+$ : 188.0923; found: 188.0924.

**4.4. Procedure for the gold-catalyzed cyclization of alkynyldioxolane (–)-1e in a heavy water medium. Preparation of acetal (–)-[D]-4e.**  $[\text{AuClPPH}_3]$  (0.0065 mmol),  $\text{AgOTf}$  (0.0065 mmol), *p*-toluenesulfonic acid (0.025 mmol), and tetrabutylammonium iodide (0.025 mmol) were sequentially added to a stirred solution of the alkynyldioxolane (–)-**1e** (82 mg, 0.26 mmol) in a biphasic water (0.4 mL)/dichloromethane (0.1 mL) mixture. The resulting mixture was heated in a sealed tube at 80 °C for 48 h. The reaction was allowed to cool to room temperature and filtered through a pack of celite. The filtrate was extracted with dichloromethane (3 x 2 mL), and the combined extracts were washed twice with brine. The organic layer was dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure. Chromatography of the residue eluting with hexanes/ethyl acetate (10:1) gave 41 mg (57%) of analytically pure adduct (–)-[D]-**4e** as a colorless oil;  $[\alpha]_D = -0.4$  ( $c = 1.0$  in  $\text{CHCl}_3$ );  $^1\text{H NMR}$  (300 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 7.27$  (m, 5H), 5.94 (dd,  $J = 17.6$ , 10.8 Hz, 1H), 4.98 (m, 2H), 4.40 (m, 2H), 3.69 (dd,  $J = 7.0$ , 5.6 Hz, 1H), 3.61 and 3.54 (d,  $J = 11.0$  Hz, each 1H), 3.51 (s, 1H), 3.02 and 2.92 (d,  $J = 14.2$  Hz, each 1H), 1.05 and 1.04 (s, each 3H);  $^{13}\text{C NMR}$  (75 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta = 144.5$ ,

130.4, 128.4, 127.0, 112.6, 84.6, 77.5, 75.9, 73.4, 66.7, 40.1, 39.1, 25.0, 24.0; IR ( $\text{CHCl}_3$ ):  $\nu = 1184$   $\text{cm}^{-1}$ ; HRMS (ES): calcd for  $\text{C}_{17}\text{H}_{22}\text{DO}_3$  [ $M + \text{H}$ ] $^+$ : 276.1709; found: 276.1705.

## Acknowledgments

Support for this work by the DGI-MICINN (Project CTQ2009-09318) and Comunidad Autónoma de Madrid (Project S2009/PPQ-1752) are gratefully acknowledged. R. C. thanks MEC for a predoctoral grant.

## References and notes

- For recent reviews and monographs, see: (a) Dunn, P. J. *Chem. Soc. Rev.* **2012**, *41*, 1452; (b) Sheldon, R. A. *Chem. Soc. Rev.* **2012**, *41*, 1437; (c) Ranu, B. C.; Saha, A.; Dey, R. *Curr. Opin. Drug. Disc.* **2010**, *13*, 658; (d) Uozumi, Y. *Synlett* **2010**, 1988; (e) Beach, E. S.; Cui, Z.; Anastas, P. T. *Energy Environ. Sci.* **2009**, *2*, 1038; (f) *Green Catalysis*, Anastas, P. T., Ed., Wiley-VCH: Weinheim, 2009; (g) Horváth, I. T.; Anastas, P. T. *Chem. Rev.* **2007**, *107*, 2169.
- For selected reviews on gold catalysis, see: (a) Rudolph, M.; Hashmi, A. S. K. *Chem. Soc. Rev.* **2012**, *41*, 2448; (b) Corma, A.; Leyva-Pérez, A.; Sabater, M. J. *Chem. Rev.* **2011**, *111*, 1657; (c) Krause, N.; Winter, C. *Chem. Rev.* **2011**, *111*, 1994; (d) Alcaide, B.; Almendros, P.; Alonso, J. M. *Org. Biomol. Chem.* **2011**, *9*, 4405; (e) Hashmi, A. S. K. *Angew. Chem. Int. Ed.* **2010**, *49*, 5232; (f) *Chem. Rev.* **2008**, *108*, issue 8, Lipshutz, B.; Yamamoto, Y. Eds.; (g) *Chem. Soc. Rev.* **2008**, *37*, issue 9, Hutchings, G. J.; Brust, M.; Schmidbaur, H. Eds.; (h) Bongers, N.; Krause, N. *Angew. Chem. Int. Ed.* **2008**, *47*, 2178; (i) Hashmi, A. S. K.; Hutchings, G. J. *Angew. Chem. Int. Ed.* **2006**, *45*, 7896; (j) Muzart, J. *Tetrahedron* **2008**, *6*, 5815; (k) Hashmi, A. S. K. *Chem. Rev.* **2007**, *107*, 3180; (l) Zhang, L.; Sun, J.; Kozmin, S. A. *Adv. Synth. Catal.* **2006**, *348*, 2271.
- For the gold-catalyzed access to bicyclic ketals from terminal alkyne diols, see: (a) Antonietti, S.; Genin, E.; Michelet, V.; Genêt, J.-P. *J. Am. Chem. Soc.* **2005**, *127*, 9976. For the platinum-catalyzed synthesis of bicyclic ketals, see: (b) Diéguez-Vázquez, A.; Tzschucke, C. C.; Lam, W. Y.; Ley, S. V. *Angew. Chem. Int. Ed.* **2008**, *47*, 209. For the iridium-catalyzed access to bicyclic ketals from alkyne diols, see: (c) Benedetti, E.; Simonneau, A.; Hours, A.; Amouri, H.; Penoni, A.; Palmisano, G.; Malacria, M.; Goddard, J.-P.; Fensterbank, L. *Adv. Synth. Catal.* **2011**, *353*, 1908.
- For the gold-catalyzed access to tricyclic cage-like ketals from diyne-diols and external nucleophiles, see: (a) Hashmi, A. S. K.; Bührle, M.; Wölfle, M.; Rudolph, M.; Wietek, M.; Rominger, F.; Frey, W. *Chem. Eur. J.* **2010**, *16*, 9846. For the gold-catalyzed access to monocyclic ketals from alkynes and diols, see: (b) Corma, A.; Ruiz, V. R.; Leyva-Pérez, A.; Sabater, M. J. *Adv. Synth. Catal.* **2010**, *352*, 1701.
- For a preliminary communication of part of this work, see: Alcaide, B.; Almendros, P.; Carrascosa, R.; Torres, M. R. *Adv. Synth. Catal.* **2010**, *352*, 1277.
- (a) Alcaide, B.; Almendros, P.; Alonso, J. M.; Fernández, I. *Chem. Commun.* **2012**, *48*, 6604; (b) Alcaide, B.; Almendros, P.; Aragoncillo, C.; Gómez-Campillos, G.; Arnó, M.; Domingo, L. R. *ChemPlusChem* **2012**, *77*, 563; (c) Alcaide, B.; Almendros, P.; Luna, A.; Gómez-Campillos, G.; Torres, M. R. *J. Org. Chem.* **2012**, *77*, 3549; (d) Alcaide, B.; Almendros, P.; Carrascosa, R. *Chem. Eur. J.* **2011**, *17*, 4968; (e) Alcaide, B.; Almendros, P.; Quirós, M. T. *Adv. Synth. Catal.* **2011**, *353*, 585; (f) Alcaide, B.; Almendros, P.; Carrascosa, R.; Martínez del Campo, T. *Chem. Eur. J.* **2010**, *15*, 13243.
- Steuer, B.; Lieberknecht, W. A.; Jager, V.; Bullock, J. P.; Hegedus, L. S. *Org. Synth.* **1996**, *74*, 1.
- Hale, K. J.; Manaviar, S.; George, J. H.; Walters, M. A.; Dalby, S. M. *Org. Lett.* **2009**, *11*, 733.
- After control experiments it was demonstrated that the silver salt alone does not catalyze the reaction of interest. Thus, it is believed that the role of the silver salt just consist in the activation of the precatalyst: (a) Gaillard, S.; Bosson, J.; Ramón, R. S.; Nun, P.; Slawin, A. M. Z.; S. Nolan, P. *Chem. Eur. J.* **2010**, *16*, 13729. However, despite proving that the silver activators themselves are unreactive, there are examples of gold(I)-catalyzed reactions

- where Ag(I) salts effected either activity or selectivity: (b) Tarselli, M. A.; Chianese, A. R.; Lee, S. J.; Gagné, M. R. *Angew. Chem. Int. Ed.* **2007**, *46*, 6670; (c) Wang, D.; Cai, R.; Sharma, S.; Jirak, J.; Thummanapelli, S. K.; Akhmedov, N. G.; Zhang, H.; Liu, X.; Petersen, J. L.; Shi, X. *J. Am. Chem. Soc.* **2012**, *134*, 9012.
10. The reaction of alkynyldioxolane **1a** in aqueous medium under gold catalysis in the absence of acid additive (PTSA) proceeded to afford the corresponding acetal **2a** in just a slightly lower yield. For a seminal review on gold and protons, see: (a) Hashmi, A. S. K. *Catal. Today* **2007**, *122*, 211. For a selected report on gold and protons, see: (b) Hashmi, A. S. K.; Schwarz, L.; Rubenbauer, P.; Blanco, M. C. *Adv. Synth. Catal.* **2006**, *348*, 705.
  11. (a) Minkler, S. R. K.; Lipshutz, B. H.; Krause, N. *Angew. Chem. Int. Ed.* **2011**, *50*, 7820; (b) Wang, Z. J.; Brown, C. J.; Bergman, R. G.; Raymond, K. N.; Toste, F. D. *J. Am. Chem. Soc.* **2011**, *133*, 7358; (c) Poonoth, M.; Krause, N. *J. Org. Chem.* **2011**, *76*, 1934; (d) Yang, C.-Y.; Lin, M.-S.; Liao, H.-H.; Liu, R.-S. *Chem. Eur. J.* **2010**, *16*, 2696; (e) Winter, C.; Krause, N. *Green Chem.* **2009**, *11*, 1309; (f) Ye, D.; Zhang, X.; Zhou, Y.; Zhang, D.; Zhang, L.; Wang, H.; Jiang, H.; Liu, H. *Adv. Synth. Catal.* **2009**, *351*, 2770; (g) Marion, N.; Ramon, R. S.; Nolan, S. P. *J. Am. Chem. Soc.* **2009**, *131*, 448; (h) Oh, C. H.; Karmakar, S. *J. Org. Chem.* **2009**, *74*, 370; (i) Cuenca, A. B.; Mancha, G.; Asensio, G.; Medio-Simón, M. *Chem.-Eur. J.* **2008**, *14*, 1518; (j) Yang, C.-Y.; Lin, G.-Y.; Liao, H.-Y.; Datta, S.; Liu, R.-S. *J. Org. Chem.* **2008**, *73*, 4907; (k) Zhang, C.; Cui, D.-M.; Yao, L.-Y.; Wang, B.-S.; Hu, Y.-Z.; Hayashi, T. *J. Org. Chem.* **2008**, *73*, 7811; (l) Tang, J.-M.; Liu, T.-A.; Liu, R.-S. *J. Org. Chem.* **2008**, *73*, 8479; (m) Genin, E.; Leseurre, L.; Toullec, Y. P.; Genêt, J.-P.; Michelet, V. *Synlett* **2007**, 1780; (n) Jung, H. H.; Floreancig, P. E. *J. Org. Chem.* **2007**, *72*, 7359; (o) Yan, B.; Liu, Y. *Org. Lett.* **2007**, *9*, 4323; (p) Zhou, C.-Y.; Chan, P. W. H.; Che, C.-M. *Org. Lett.* **2006**, *8*, 325; (q) Casado, R.; Contel, M.; Laguna, M.; Romero, P.; Sanz, S. *J. Am. Chem. Soc.* **2003**, *125*, 11925; (r) Wei, C.; Li, C.-J. *J. Am. Chem. Soc.* **2003**, *125*, 9584; (s) Mizushima, E.; Sato, K.; Hayashi, T.; Tanaka, M. *Angew. Chem., Int. Ed.* **2002**, *41*, 4563.
  12. The appealing properties of organic reactions in aqueous media include their synthetic advantages as well as unique reactivity and selectivity that are not often attained under dry conditions, making then profitable in many cases. Thus, water-based organic reactions have been gaining in popularity in synthetic chemistry, particularly for its abundant availability, and for its inexpensiveness. For selected reviews on organic reactions in aqueous media, see: (a) Simon, M.-O.; Li, C.-J. *Chem. Soc. Rev.* **2012**, *41*, 1415; (b) Butler, R. N.; Coyne, A. G. *Chem. Rev.* **2010**, *110*, 6302; (c) Lombardo, M.; Trombini, C. *Curr. Opin. Drug. Disc.* **2010**, *13*, 717; (d) Chanda, A.; Fokin, V. V. *Chem. Rev.* **2009**, *109*, 725; (e) *Organic Reactions in Water: Principles, Strategies and Applications*, Linström, U. M., Ed., Blackwell: Oxford, **2007**; (f) Li, C. J.; Chen, L. *Chem. Soc. Rev.* **2006**, *35*, 68; (g) Pirrung, M. C. *Chem. Eur. J.* **2006**, *12*, 1312; (h) Li, C. J. *Chem. Rev.* **2005**, *105*, 3095; (i) Lindström, U. M. *Chem. Rev.* **2002**, *102*, 2751; (j) Manabe, K.; Kobayashi, S. *Chem. Eur. J.* **2002**, *8*, 4095; (k) Ribe, S.; Wipf, P. *Chem. Commun.* **2001**, 299; (l) Li, C. J.; Chan, T. H. *Tetrahedron* **1999**, *55*, 11149; (m) Paquette, L. A. In *Green Chemistry: Frontiers in Benign Chemical Synthesis and Processing*, Anastas, P. T.; Williamson, T. C., Eds., Oxford University Press: New York, **1998**.
  13. The improvement of the reaction upon the addition of the phase transfer catalyst is due to an easiness of the migration of a reactant from one phase into another phase where reaction occurs.
  14. Stereogenic acetals are ubiquitous in natural products. For example, the bridged dioxabicyclo[n.2.1]cycloalkane scaffold is an important structural organization present in a number of pheromones and biologically active products. For recent examples, see: (a) Luo, X.; Pu, J.-X.; Xiao, W.-L.; Zhao, Y.; Gao, X.-M.; Li, X.-N.; Zhang, H.-B.; Wang, Y.-Y.; Li, Y.; Sun, H.-D. *J. Nat. Prod.* **2010**, *73*, 1112; (b) Mascitti, V.; Préville, C. *Org. Lett.* **2010**, *12*, 2940; (c) Luo, J.; Wang, J.-S.; Wang, X.-B.; Luo, J.-G.; Kong, L.-Y. *J. Nat. Prod.* **2010**, *73*, 835.
  15. Trabocchi, A.; Mannino, C.; Machetti, F.; De Bernardis, F.; Arancia, S.; Cauda, R.; Cassone, A.; Guarna, A. *J. Med. Chem.* **2010**, *53*, 2502.
  16. Full spectroscopic and analytical data for compounds not included in this Experimental Section are described in the Supplementary data.

#### Supplementary data

Supplementary data associated with this article can be found in online version at doi:10.1016/j.tet.2011.

Click here to remove instruction text...

Autores: Alcaide, B.; Almendros, P.; Carrascosa, R.

Título: Gold-Catalyzed Direct Cycloketalization of Acetonide-Tethered Alkynes in the Presence of Water

Revista: *Tetrahedron* **2012**, *68*, 9391-9396