

## Correspondence

Michael R. DeBaun, Vanderbilt-Meharry Sickle Cell Disease Center of Excellence, 2525 West End Ave, Ste 750, Nashville, TN 37203.

Email: m.debaun@vmc.org

DOI 10.1002/ajh.25616

## ORCID

Michael R. DeBaun  <https://orcid.org/0000-0002-0574-1604>

## REFERENCES

1. Abdullahi SU, DeBaun MR, Jordan LC, Rodeghier M, Galadanci NA. Stroke recurrence in Nigerian children with sickle cell disease: evidence for a secondary stroke prevention trial. *Pediatr Neurol.* 2019;95:73-78.
2. Piel FB, Hay SI, Gupta S, Weatherall DJ, Williams TN. Global burden of sickle cell anaemia in children under five, 2010-2050: modelling based on demographics, excess mortality, and interventions. *PLoS Med.* 2013;10:e1001484.
3. Anie KA, Egunjobi FE, Akinyanju OO. Psychosocial impact of sickle cell disorder: perspectives from a Nigerian setting. *Globalization Health.* 2010;6:2.
4. Enniful-Eghan H, Moore RH, Ichord R, Smith-Whitley K, Kwiatkowski JL. Transcranial Doppler ultrasonography and prophylactic transfusion program is effective in preventing overt stroke in children with sickle cell disease. *The Journal of Pediatrics.* 2010;157(3):479-484.
5. Galadanci NA, Umar Abdullahi S, Vance LD, Musa Tabari A, Ali S, Belonwu R, Salihu A, Amal Galadanci A, Wudil Jibir B, Bello-Manga H, Neville K. Feasibility trial for primary stroke prevention in children with sickle cell anemia in Nigeria (SPIN trial). *American journal of hematology.* 2017;92(8):780-788.
6. Amendah DD, Mvundura M, Kavanagh PL, Sprinz PG, Grosse SD. Sickle cell disease -related pediatric medical expenditures in the US. *American Journal of Preventive medicine.* 2010;38(4):S550-6.
7. Reeves SL, Madden B, Freed GL, Dombkowski KJ. Transcranial doppler screening among children and adolescents with sickle cell anemia. *JAMA Pediatr.* 2016;170:550-556.
8. Therrell BL Jr, Lloyd-Puryear MA, Eckman JR, Mann MY. Newborn screening for sickle cell diseases in the United States: a review of data spanning 2 decades. *Semin Perinatol.* 2015;39:238-251.

Received: 9 July 2019 | Revised: 13 August 2019 | Accepted: 15 August 2019

DOI: 10.1002/ajh.25619

## Long-term treatment results of Polish pediatric and adolescent patients enrolled in the ALL IC-BFM 2002 trial

To the Editor:

In 2002, the Polish Pediatric Leukemia/Lymphoma Study Group joined the ALL IC-BFM 2002 trial, which was intended for countries with inadequate skills and resources for PCR-based MRD monitoring.<sup>1</sup> On the basis of the pioneering findings of the BFM group regarding measurement of early response to prednisone in peripheral blood on

day eight and percentage of bone marrow blasts on day 15, all the patients could be stratified in risk groups by accessible and inexpensive methods. Considering inequalities in resources and facilities among the participating countries, it seems justified to us to report on the treatment results of this trial in patients with ALL treated in 14 Polish pediatric oncology centers.

Between November 2002 and November 2011, 1872 children aged 1-17 with newly diagnosed ALL were consecutively enrolled in the ALL IC-BFM 2002 protocol according to a registered trial at ClinicalTrials.gov (NCT00764907). Infants younger than 12 months were excluded. The patients were diagnosed according to cytomorphologic criteria (more than 25% lymphoblasts present in bone marrow), immunophenotyping (EGIL criteria) and cytogenetic and FISH/molecular genetics examination of bone marrow specimens. A central review on the national level of morphologic, flow cytometric and genetic results was required. The prognostic factors of ALL IC-BFM 2002 included: age, white blood cell count (WBC) at diagnosis, response to prednisone at day 8, results of bone marrow evaluation on day 15 or 33 of the therapy and, finally, the presence of *BCR/ABL1*, and *KMT2A/AFF1* specific genetic aberrations. At diagnosis, the patients were stratified into three groups: standard, intermediate, and high-risk groups. Risk group definitions and treatment plan are presented in the paper by Stary et al.<sup>1</sup> OS and EFS curves were estimated according to Kaplan-Meier with Greenwood Standard Error (SE), and compared using a two-tailed log-rank test. Cumulative incidence (CI) curves for the events were estimated according to risk groups, and were compared using the Gray test. All statistical calculations were performed using STATISTICA13. The study was approved by the local institutional review board and a written informed consent from the guardians was a prerequisite.

A total of 1872 patients (56.6% males and 43.4% females) were treated under the ALL IC-BFM 2002 Protocol. A median follow up time for the entire group was 6.4 years (Q1 = 4.32, Q3 = 8.53). The median age at diagnosis was 5.3 years, ranging from more than 1 to 17 years of age. Most children (55%) were in the age range 1 to <6 years, while the group aged 6 to <10 years accounts for 19% of the cohort, 10 to <15 years was 17%, and those older than 15 years constitute 9%. For WBC counts at diagnosis below  $20 \times 10^9/l$  was revealed in 64.3% of the children while in 10.1% WBC was more than  $100 \times 10^9/l$ . CNS status 1 in 88%, 2-7% and 5% showed signs of CNS involvement. Mediastinal involvement from the leukemic process was found in 7.8%, testicular involvement was shown in 20 boys (1.9%). The immunophenotype in most children was revealed as preB cell lineage (89%) and T cell lineage was shown in 11%. The conventional cytogenetic analysis was performed in all subjects with newly diagnosed ALL. Chromosome preparations of BM cells were informative for analysis in 1056 (56.4%) patients. Chromosome aberrations were revealed in 523 (27.9%) cases. Hypodiploidy was presented in 22 (2.1%) patients. Investigations in fusion genes *BCR/ABL1*, *KMT2A/AFF1* were required under the ALL IC-BFM 2002 protocol as primary and obligatory tests. The t (9;22)(q34;q11.2) translocation that produced the fusion gene *BCR/ABL1* was detected in 65 (3.5%) pediatric patients. The rearrangements of *KMT2A* were presented in

42 (2.2%) children and in five children the *KMT2A/AFF1* fusion gene was confirmed.

According to risk stratification, 32.6% of the children were qualified for a standard-risk group (SR), 48.1% were qualified as intermediate-risk (IR), and 19.3% as high-risk (HR). After 7 days of steroid therapy, 10.9% of patients were assessed as prednisone poor responders. Bone marrow on day 15 M1 (<5% blasts) was revealed in 68.6%, on M2 (5% to <25% blasts) in 19.7%, and on M3 (≥25% blasts) in 9.9% patients. Complete remission (CR1) of the disease on day 33 was achieved in 96% of the patients, while 65 patients (4%) achieved CR1 between days 52-96, and were classified as late responders and they were stratified to the HR group. Five children did not achieve CR1 by day 96 (before first HR2 block) and they continued therapy on individual decisions.


A total of 268 (14.3%) deaths were noted in the entire group, including 30 (11.2%) deaths in the SR group, 116 (43.2%) in IR, and 122 (45.5%) deaths in HR. Given the size of the risk groups, the death rate observed in the SR group was 4.77%, 13.47% in IR, and 32.87% in the HR group. When patients who underwent BMT are excluded, the death rate in the HR group is as high as 39.3%. Cumulative death risk in the SR group at 5 years was estimated at 4.4%, 14.2% in the IR

group, and 37.3% in HR (Figure S1). In 36 cases, deaths happened during the induction therapy, before CR1 was achieved, and the estimated death rate at induction was 1.92% (0.82% for the SR group, 2.23% for IR, and 3.04% for the HR group). A total of 81 deaths were noted at the CR1 phase (death rate of 4.55%, with 1.19% in SR, 3.26% in IR, and 13.77% in the HR group). Deaths related to relapse were noted in 106 children.

The incidence of death at CR was significantly higher in children aged ≥10 years vs younger children (9.3% vs 2.6%;  $P < .001$ ) and in T-ALL vs BCP-ALL (9% v 3.5%;  $P < .001$ ). But, differences were not found between girls and boys (4.0% v 4.5%;  $P = .7$ ). The main causes of deaths are presented in Table S1.

Overall survival (OS) for 1872 children with ALL treated under the ALL IC-BFM 2002 protocol in Poland was 86% after 5 years, whereas the EFS rate was 79%. The EFS for the SR group ( $n = 608$ ) was 90%, 80% for IR ( $n = 899$ ), and 62% for HR ( $n = 361$ ) at 5 years, and at 10 years EFS was 88%, 79%, and 59%, respectively (Figure 1).

Relapses occurred in 275 children (14.7%), of which 51 cases occurred in the SR group (8.4% of the patients who were stratified for the SR group), 147 cases in IR (16.4%), and 77 in HR (21.2%). Cumulative relapse risk for the SR group was 8% (SE 1.1), whereas the IR and



**FIGURE 1** Overall survival for entire cohort of children with ALL A, and by risk groups B, and event free survival C, D, respectively

HR groups showed 16.5% (SE 1.2) and 25.9% (SE 2.0) respectively (Figure S2).

Treatments using standardized regimens or protocols have led to unprecedented improvements in survival of children with cancer, but most published regimens are based on therapies developed and delivered in high-income countries.<sup>2</sup> The ALL IC-BFM 2002 trial was designed to be conducted in countries with inadequate skills and resources for PCR-based MRD monitoring enabling, however, the stratification and treatment of children with ALL according to a BFM-based backbone protocol.<sup>1</sup> The EFS in our cohort at 5 years is estimated to be as high as 79%, and OS of 86% at 5 years. These results are clearly comparable to treatment outcomes presented by several major study groups or institutions, since over the last two decades the world's leading leukemia groups have achieved 5-year OS rates of approximately 90% with 2% to 3% deaths resulting from toxicity in childhood ALL.<sup>3</sup>

A major concern regarding BFM-type chemotherapy for less experienced groups with limited resources was the potential risk of excessive therapy related mortality (TRM). Indeed, we observed a 4.55% rate of deaths upon CR, ranging from 1.19% in the SR to 13.77% in the HR patients. Several study groups from countries with high health expenditure have reported that 2-5% of children with ALL die from other causes than relapse, and 1-2% of patients die during the induction phase.<sup>4</sup>

Our study demonstrates a death rate for both induction before CR1 (1.92%) and during the CR1 phase (4.55%). This may result from the long duration of the analysis and a change in the standards of supportive therapy. The major leading causes of death, similarly to other reports, were sepsis, progressive ALL and SCT-related complications. Infections, both bacterial and fungal, resulted in 24.3% of deaths in our cohort. In comparison, Möricke et al.<sup>5</sup> have reported 69% cases of death as infection-related. The risk of life-threatening infection increased during the induction phase.

The cumulative incidence of relapse in our patients after 5 years for the whole cohort was 15% (SE 0.8), while the individual risk groups showed 8% (SE 1.1) for SR, 16.5% (SE 1.2) for IR, and 25.9% (SE 2.0) for HR. These results are nearly identical with those obtained by the ALL BFM95 study: 16.2% overall (SR-7.8%, SE 1.0; MR-16.8%, SE 1.2, and HR-38.6%, SE 3.6).<sup>6</sup>

In conclusion, ALL IC-BFM 2002 proved as a highly effective protocol for treatment of children with ALL in Polish pediatric oncohematology centers. The design of the study allowed us to achieve outcomes comparable to those reported by study groups in several countries with high health-related expenditures. The study based on inexpensive and accessible stratification criteria was highly successful.

## ACKNOWLEDGMENTS

We thank all staff members of the collaborating institutes. Special thanks are due to data manager Radosław Nowakowski.

## FUNDING

This work was supported by the National Centre for Research and Development, Poland [grant number: STRATEGMED3/304586/5/

NCBR/2017]. The funders had no role in study design, data collection, data analysis, interpretation, or writing of the report. The corresponding author had full access to all the data reported in the manuscript and had final responsibility for the decision to submit for publication.

## CONFLICT OF INTEREST

The authors declare no potential conflict of interest.

## AUTHOR CONTRIBUTIONS

J.R.K. planned the study and coordinated the study in Poland, J.Z., K.D., M.S. contributed to study coordination and data revision in Poland, M.L. was responsible for the genetic analyses. J.Z., M.M., M.R., W.B., M.Ć., B.K., J.O.L., J.W., K.D., E.D., M.N., J.T., W.M., M.W., A.K., T.S., M.K.R., A.K., M.W., T.U., T.O., G.S.M., A.M.M., G.K., J.S. collected the clinical data. J.R.K., J.Z., M.L., K.D., M.S. analyzed and interpreted the data, wrote, and supervised the paper. All authors read and approved the final manuscript.

## ETHICS STATEMENT

The ethics committee of Medical University of Lublin, Poland. The committee's reference number: KE-0254/178/2002.

Jerzy R. Kowalczyk<sup>1</sup> , Joanna Zawitkowska<sup>1</sup>, Monika Lejman<sup>2</sup>, Katarzyna Drabko<sup>1</sup>, Marzena Samardakiewicz<sup>1,3</sup>, Michał Matysiak<sup>4</sup>, Michał Romiszewski<sup>4</sup>, Walentyna Balwierz<sup>5</sup>, Magdalena Ćwiklińska<sup>5</sup>, Bernarda Kazanowska<sup>6</sup>, Joanna Owoc-Lempach<sup>6</sup>, Jacek Wachowiak<sup>7</sup>, Katarzyna Derwich<sup>7</sup>, Elżbieta Adamkiewicz-Drożyńska<sup>8</sup>, Maciej Niedźwiecki<sup>8</sup>, Joanna Trelińska<sup>9</sup>, Wojciech Młynarski<sup>9</sup>, Mariusz Wysocki<sup>10</sup>, Andrzej Kołtan<sup>10</sup>, Tomasz Szczepański<sup>11</sup>, Maryna Krawczuk-Rybak<sup>12</sup>, Anna Kitszel<sup>12</sup>, Maria Wiecezorek<sup>13</sup>, Tomasz Urański<sup>13</sup>, Tomasz Ociepa<sup>13</sup>, Grażyna Sobol-Milejska<sup>14</sup>, Agnieszka Mizia-Malarz<sup>14</sup>, Grażyna Karolczyk<sup>15</sup>, Jan Stary<sup>16</sup>

<sup>1</sup>Department of Pediatric Hematology, Oncology and Transplantology, Medical University of Lublin, Lublin, Poland

<sup>2</sup>Department of Pediatric Hematology, Oncology and Transplantology, Laboratory of Genetic Diagnostics, Medical University of Lublin, Lublin, Poland

<sup>3</sup>Department of Applied Psychology, Medical University of Lublin, Lublin, Poland

<sup>4</sup>Department of Hematology and Pediatrics, Children's Hospital, Warsaw, Poland

<sup>5</sup>Department of Pediatric Oncology and Hematology, Children's University Hospital, Jagiellonian University, Kraków, Poland

<sup>6</sup>Department of Pediatric Transplantology, Oncology, Hematology, Medical University of Wrocław, Wrocław, Poland

<sup>7</sup>Department of Pediatric Oncology, Hematology and Transplantology, Medical University of Poznań, Poznań, Poland

<sup>8</sup>Department of Pediatrics, Hematology, Oncology and Endocrinology,  
Medical University of Gdańsk, Gdańsk, Poland

<sup>9</sup>Department of Pediatrics, Oncology, Hematology and Diabetology,  
Medical University of Łódź, Łódź, Poland

<sup>10</sup>Department of Pediatrics, Hematology and Oncology, Collegium  
Medicum of Bydgoszcz, Bydgoszcz, Poland

<sup>11</sup>Department of Pediatrics, Hematology and Oncology, Medical  
University of Zabrze, Zabrze, Poland

<sup>12</sup>Department of Pediatric Oncology, Hematology, Medical University of  
Białystok, Białystok, Poland

<sup>13</sup>Department of Pediatrics, Hematology and Oncology, Medical  
University of Szczecin, Szczecin, Poland

<sup>14</sup>Department of Pediatric Oncology, Hematology and Chemotherapy,  
Medical University of Katowice, Katowice, Poland

<sup>15</sup>Department of Pediatric Oncology and Hematology, Children's  
Hospital, Kielce, Poland

<sup>16</sup>Department of Pediatric Hematology and Oncology, Charles University  
and University Hospital Motol, Prague, Czech Republic

#### Correspondence

Jerzy R. Kowalczyk, Department of Paediatric Haematology,  
Oncology and Transplantology, Medical University of Lublin, Antoni  
Gębala Street 6, 20-093 Lublin, Poland.

Email: jkowalcz@dsk.lublin.pl

DOI 10.1002/ajh.25619

#### ORCID

Jerzy R. Kowalczyk  <https://orcid.org/0000-0003-3080-0845>

#### REFERENCES

1. Stary J, Zimmermann M, Campbell M, et al. Intensive chemotherapy for childhood acute lymphoblastic leukemia: results of the randomized intercontinental trial ALL IC-BFM 2002. *J Clin Oncol.* 2014;32(3):174-184.
2. Cui L, Li ZG, Chai YH, et al. Outcome of children with newly diagnosed acute lymphoblastic leukemia treated with CCLG-ALL 2008: The first nation-wide prospective multicenter study in China. *Am J Hematol.* 2018;93(7):913-920.
3. Hunger SP, Lu X, Devidas M, et al. Improved survival for children and adolescents with acute lymphoblastic leukemia between 1990 and 2005: a report from the children's oncology group. *J Clin Oncol.* 2012;30(14):1663-1669.
4. Christensen MS, Heyman M, Mottonen M, et al. Treatment-related death in childhood acute lymphoblastic leukaemia in the Nordic countries: 1992-2001. *Br J Haematol.* 2005;131(1):50-58.
5. Moricke A, Zimmermann M, Valsecchi MG, et al. Dexamethasone vs prednisone in induction treatment of pediatric ALL: results of the randomized trial AIEOP-BFM ALL 2000. *Blood.* 2016;127(17):2101-2112.
6. Moricke A, Reiter A, Zimmermann M, et al. Risk-adjusted therapy of acute lymphoblastic leukemia can decrease treatment burden and improve survival: treatment results of 2169 unselected pediatric and adolescent patients enrolled in the trial ALL-BFM 95. *Blood.* 2008;111(9):4477-4489.

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.