

The complete mitochondrial genome of the subterranean crustacean *Metacrangonyx longipes* (Amphipoda): a unique gene order and extremely short control region

MARIA DEL MAR BAUZÀ-RIBOT¹, DAMIÀ JAUME², CARLOS JUAN¹, & JOAN PONS^{2§}

¹*Departament de Biologia, Universitat de les Illes Balears, Palma de Mallorca, Balearic Islands, Spain.*

²*IMEDEA (CSIC-UIB), Instituto Mediterráneo de Estudios Avanzados, Esporles, Balearic Islands, Spain*

Correspondence: Joan Pons, IMEDEA (CSIC-UIB), Instituto Mediterráneo de Estudios Avanzados, Miquel Marquès 21, Esporles 07190, Balearic Islands, Spain. Fax: +34 971 611761, Email: joan.ponspons@uib.es

Abstract

Metazoan mitochondrial genomes usually consist of the same gene set, but some taxonomic groups show a considerable variety in gene order and nucleotide composition. The mitochondrial genomes of 37 crustaceans are currently known. Within the malacostracan superorder Peracarida, only three partial mitogenome sequences and the complete sequence of *Ligia oceanica* (Isopoda) are available. Frequent translocation events have changed the mitochondrial gene order in crustaceans, providing an opportunity to study the patterns and mechanisms of mitogenome rearrangement and to determine their impact on phylogenetic reconstructions. Here we report the first complete nucleotide sequence of an amphipod species, *Metacrangonyx longipes*, belonging to a phylogenetically enigmatic family occurring in continental subterranean waters. The genome has 14,113 base pairs (bp) and contains the usual 13 protein coding genes and two rRNA subunits, but only 21 out of the typical 22 tRNA genes of Metazoa. This is the shortest mitogenome described thus far for a crustacean and also one of the richest in AT (76.03%). The genome compactness results from a very small control region of 76 bp, the occurrence of frequent gene overlap, and the absence of large non-coding fragments. Six of the protein-coding genes have unusual start codons. Comparison of individual protein coding genes with the sequences known for other crustaceans suggests that *nad2*, *nad6*, *nad4L* and *atp8* show the highest divergence rates. *Metacrangonyx longipes* shows a unique crustacean mitogenome gene order, differing even from the condition found in *Parhyale hawaiiensis* (Amphipoda), whose coding sequence has also been completed in the present study.

Keywords: *Metacrangonyx longipes*, Amphipoda, mitochondrial genome, control region, gene order

Introduction

The mitochondrial genome (mitogenome) of metazoans generally comprises a circular double-stranded DNA molecule of 12-20 kb with a highly conserved gene content. It includes 13 protein-coding, two ribosomal and up to 22 transfer RNA genes (Wolstenholme 1992). The Crustacea have more than 52,000 described species, with a range in body plan not matched in any other group of metazoans (Martin and Davis 2001). They include the six recognized classes: Branchiopoda, Cephalocarida, Malacostraca, Maxillopoda, Ostracoda and Remipedia (Martin and Davis 2001). The mitogenome sequences of 37 species of Crustacea have been completed thus far (<http://www.ncbi.nlm.nih.gov/genomes>), of which 15 correspond to malacostracan decapods (Carapelli et al. 2007; Yang and Yang 2008). Within the malacostracan peracarid order Amphipoda, only a partial mitogenome sequence is currently available in sequence databases: that of *Parhyale hawaiiensis* (Dana 1853), although it lacks of about 3 Kb including the *rrnS* gene and parts of *rrnL* and *nad2*, and also the control region (Cook et al. 2005). In addition, in the peracarid order Isopoda one entire (*Ligia oceanica*) and two partial mitogenomes (*Armadillidium vulgare* and *Idotea baltica*) are known (Kilpert and Podsiadlowski 2006; Podsiadlowski and Bartolomaeus 2006; Marcadé et al. 2007). The taxon sampling set for crustacean mitogenomes is quite poor because only 30 out of about 800 known crustacean families are represented (Martin and Davis 2001). Despite this, two important insights into pancrustacean phylogenetics are based on mitogenome data. First, phylogenetic analyses of protein-coding genes (PCGs) including the more intensively sampled mitochondrial genomes of Hexapoda suggest a mutual paraphyly of Crustacea and Hexapoda (Carapelli et al. 2007; Cook et al. 2005). Second, frequent translocation events have apparently changed the mitochondrial gene order in crustaceans compared with the putative ancestral pancrustacean pattern (Kilpert and Podsiadlowski 2006; Yang and Yang 2008). This gene order has been proposed as ancestral based on a common inversion of a *trnL2* gene present in a large number of crustaceans and insects, that subsequently translocated from a location inferred to be the primitive state as it is found in chelicerates, myriapods, onychophorans, tardigrades, as well as in Pogonophora, Annelida, Echiura, and Mollusca (Boore et al. 1995; 1998). Gene order is not conserved within the superorder Peracarida (for which only information on Isopoda and Amphipoda is currently available), nor is it even conserved within the Isopoda. Despite those differences the mitogenome of the isopods *Ligia oceanica* (Linnaeus, 1767), *Idotea baltica* (Pallas, 1772) and *Armadillidium vulgare* (Latreille, 1804) share some gene rearrangements (i.e. putative isopod synapomorphies), compared with the arthropod pattern and that of the amphipod *Parhyale hawaiiensis* (Kilpert and Podsiadlowski 2006).

The Metacrangonyctidae (Boutin and Messouli 1988a) represent a small family of amphipod

crustaceans with two genera: *Metacrangonyx* Chevreux, 1909 (17 species) and *Longipodacrangonyx* Boutin and Messouli, 1988 (monotypic). All members of the family occur only in continental subterranean waters and represent a phylogenetically enigmatic lineage of marine origin showing an extremely disjunct geographic distribution. Two species are found in the Dominican Republic (Hispaniola, Jaume and Christenson 2001), one from Fuerteventura in the Canary Islands (Stock and Rondé-Broekhuizen 1986), 11 from Morocco (Balazuc and Ruffo 1953; Ruffo 1954; Karaman and Pesce 1979; Boutin and Messouli 1988a; 1988b; Oulbaz et al. 1988; Messouli et al. 1991), one from Elba Island, Italy (Stoch 1997), one from the Balearic Islands (Chevreux 1909; Margalef 1952) and two from the Middle East (Ruffo 1982; Karaman 1989). Whereas most species live in interstitial freshwater associated with springs, wells or alluvial sediments, some taxa occur in brackish or athalassohaline waters. Only *Metacangronyx longipes* Chevreux, 1909 from the Balearics and the two Hispaniolan species are ordinary cave dwellers, living in fresh to marine subterranean waters (Jaume and Christenson 2001).

It has been proposed that the Metacrangonyctidae derive from marine littoral ancestors that colonized the continental ground waters during episodes of marine regression (Boutin and Coineau 1990). Although first supposed to be no older than the opening of the northern Atlantic ocean (Boutin 1994), the discovery of *Metacrangonyx* in the Greater Antilles (Jaume and Christenson 2001) suggests a much older origin for the genus: at least before the opening of the northern Atlantic (110 million years before present). Thus its current distribution would be the result of vicariance by plate tectonics and of peripatric speciation associated with episodes of regression in the paleocoastline of Tethys.

We present here the first complete sequence of a mitochondrial genome of an amphipod. We have used the mitogenome of *Metacrangonyx longipes* to compare its gene order with those of other crustaceans, as well as its nucleotide composition and tRNA structure. We especially focus comparisons on other peracarids such as the amphipod *Parhyale hawaiiensis*, for which we have almost completed the whole mitogenome (except approximately 500 bp of the control region that has not been sequenced because of technical problems), and the isopods *Ligia oceanica*, *Idotea baltica* and *Armadillidium vulgare*.

Materials and Methods

Sampling and DNA extraction

A 3 mm long specimen of *Metacrangonyx longipes* preserved in absolute ethanol was used for DNA extraction by means of the DNeasy Tissue Kit (Qiagen, Hilden, Germany) following the

manufacturer's protocol for total genomic DNA purification. The specimen was collected in Cala Varques cave (Mallorca Island, Spain) during fall 2007.

PCR primers and conditions

Gene fragments at opposing ends of the mitochondrial genome were amplified using standard protocols outlined elsewhere (Balke et al. 2005) and universal primers (Table I). Based on the sequence obtained, we designed species-specific long primers (Table I) of about 25-29 bp targeted at the *cox1* / *rrnL* genes to amplify two long fragments of about 4.5 and 10 Kb covering the whole circular mitochondrial genome. Long-range polymerase chain reaction (PCR) amplifications were performed using TaKaRa LA *Taq*TM polymerase (Takara Bio Inc., Tokyo, Japan) according to the manufacturer's specifications. The general reaction mixture for each 50 μ L was: 5 μ L of 10 \times LA PCR buffer, 5 μ L of 25 mM MgCl₂, 8 μ L of dNTP mixture (2.5 mM each), 2.5 μ L forward primer (10 μ M), 2.5 μ L reverse primer (10 μ M), 0.5 μ L Takara LA *Taq*TM (5 U/ μ L), 24.5 μ L distilled H₂O and 2 μ L of genomic DNA. PCR cycles were as follows: after an initial denaturation step of 94 $^{\circ}$ C for 90 s, 14 cycles were performed at 94 $^{\circ}$ C for 30 s, 57-62 $^{\circ}$ C (depending on primers) for 30 s and 68 $^{\circ}$ C for 5-15 min depending on the expected fragment size. This was followed by 16 cycles at 94 $^{\circ}$ C for 30 s, 57-62 $^{\circ}$ C for 30 s and 68 $^{\circ}$ C for 15 min (increasing by 15 s each cycle) and a final extension for 10 min at 72 $^{\circ}$ C.

Cloning and sequencing

Long mitochondrial fragments were digested independently with *Dra*I, *Rsa*I and *Taq*I restriction enzymes according to the manufacturer's specifications. DNA digestions showed fragments ranging from 150 pb to 1.5 Kb when checked on 2% agarose gels. DNA fragments from the three digestions were pooled and purified using the MiniElute PCR Purification Kit (Qiagen) and then cloned into a pJET blunt cloning vector (Fermentas, Glen Burnie, MD, USA) according to the specifications of the manufacturer. One-shot competent *E. coli* cells from Invitrogen (Madison, WI) were used for transformation. Ninety-six recombinant colonies were screened by PCR amplification for inserts of a minimum of 300 bp, and 63 were sequenced in both directions using the pJET vector sequencing primers. Sequences obtained from clones were then used to design specific primers to sequence the long PCR fragments directly by primer walking (list of primers available upon request) to obtain a full contig of the mitogenome. Additional primers were designed to close particular gaps in the sequence. The forward and reverse strands of small PCR amplicons or the long PCR fragments were cycle-sequenced using the ABI Prism BigDye Terminator Cycle Sequencing Kit on an ABI

3100 automated sequencer (Applied Biosystems, Foster City, CA).

Gene annotation and sequence analysis

Analyses of the quality of chromatograms and contig construction to obtain the whole mitochondrial sequence were performed with the software CodonCode Aligner v2.0 (CodonCode Corp., Denham, MA). Ambiguous nucleotide positions were validated by direct checking of the chromatograms. Preliminary gene identification was determined by BLAST searching on GenBank databases (<http://www.ncbi.nlm.nih.gov>) and making multiple alignments to other crustacean nucleotide and amino acid sequences (see Additional File 1 for a list of species and accession numbers). Definitive annotations were performed using the DOGMA webserver (Dual Organellar GenoMe Annotator; <http://bugmaster.jgi-psf.org/dogma>). The 5' and 3' ends of protein and ribosomal genes were refined manually by comparison with the complete genes of other crustaceans. Transfer RNA genes were determined with tRNAscan-SE Search Server v1.21 (<http://lowelab.ucsc.edu/tRNAscan-SE>) using a tRNA covariance model (Lowe and Eddy 1997) and by inspection of anti-codon sequences and the predicted secondary structures. Nucleotide frequencies of protein coding and RNA genes were calculated with the DAMBE software package (Xia and Xie 2001), while the effective number of codons was determined according to INCA v1.20 (Supek and Vlahovicek 2004).

Divergence in protein coding genes

Mean nucleotide divergences of individual PCGs were estimated from pairwise comparisons among the complete mitogenomes of crustaceans and subsequently compared with the values obtained for 35 species representing all major Hexapoda orders for which there are data available. MEGA v4.0.2 (Tamura et al. 2007) was used to calculate corrected distances using the Maximum Composite Likelihood model (Tamura et al. 2004) and among-sites rate variation following a gamma distribution with a shape parameter of 0.4 as estimated in RAxML v7.0.4 (Stamatakis 2006). Gapped positions were not considered in the analysis of each pairwise comparison. Mean divergence values were normalized by dividing the value obtained for each gene by the value of the gene with the highest rate.

Gene rearrangement analyses

We used the program CREx (Bernt et al. 2007) to deduce gene rearrangement scenarios in crustacean mitogenomes based on the detection of strong interval trees (STIs) on the CREx

webserver (<http://pacosy.informatik.uni-leipzig.de/crex>). The STIs reflect genes that appear consecutively in several of the input gene orders, i.e. given two gene orders, a set of genes is a common interval if the genes in that set appear consecutively in both gene orders. A certain subset of all common intervals, the “strong common intervals” can be represented as the nodes of a special type of tree. The descendants of a node (strong common interval) are simply the strong common intervals that it includes entirely. If the descendants of a node appear in the same order in both input gene orders, the node is called “linear increasing” (+); if the children of a node appear in exactly the opposite order, it is “linear decreasing” (-); otherwise the node is called prime (Bernt et al. 2007).

Results and Discussion

Genome organization

The mitochondrial sequence of *M. longipes* has an overall length of 14,113 bp [EMBL accession number: AM944817] and shows the usual circular organization found in most metazoans (Fig. 1). To our knowledge, this is the smallest mitogenome described so far for a crustacean: close to that of *Tigriopus californicus* (Copepoda, Harpacticoida, Maxillopoda; 14,546 bp) (Machida et al. 2002). Gene annotation reveals the presence of the typical 13 PCGs and the two rRNA subunits of metazoan mitochondrial genomes (Table II), but only 21 tRNA genes instead of the typical 22; this is similar to the condition found in the isopod *Ligia oceanica* (Kilpert and Podsiadlowski 2006). The compactness of the genome is due to the occurrence of frequent gene overlap, since more than 20 genes share borders. These overlapping regions range in size from just one bp (several cases) to a maximum of 63 bp (in the gene coding for tRNA-Val, that overlaps with 44 bp of the 5' end of *rrnL* and with 19 bp of the 3' end of *rrnS*). Small non-coding sequences or intergenic spacers (range 1-17 bp; see Table II) are also evident in the mtDNA. A further region of non-coding DNA comprising 76 bp, placed between the *rrnS* and *cob* genes and with an AT content of 84.22%, presumably corresponds to the control region and contains the origin of mtDNA replication. The region has a putative secondary structure folding into a hairpin, with a stem of 15 paired nucleotides plus a short loop of four nucleotides (Fig. 2). This is similar to other stem-loop structures known to occur in insect mitochondrial control regions (Zhang et al. 1995) and that are presumed to be the origin of replication of mtDNA. The 3'-flanking sequence around the stem region shows the conserved motif GACT present also in the isopod *Ligia oceanica* and the hoplocarid malacostracan *Squilla mantis* Linnaeus, 1758 (Kilpert and Podsiadlowski 2006), but the TATA element found in many hexapods at the 5'-flanking region is here replaced by an AATT motif. The low level of non-coding sequences found in the mitogenome of *M. longipes* (< 1%) and the occurrence of frequent gene overlap are indicative of an extremely compact mitogenome.

Protein coding genes: nucleotide composition and codon usage

The AT content of the protein genes of *Metacrangonyx longipes* is 75.33% (A=31.25%, C=11.34%, G=13.33%, T=44.08%), while that of the complete mitogenome (+ strand) is 76.03%; this is one of the highest percentages reported in crustaceans and similar to those frequently found in Hexapoda mitochondrial genomes. *Argulus americanus* C. B. Wilson, 1902 (Branchiura, Maxillopoda) has the highest AT content found so far in any crustacean at 77.80% (Machida et al. 2002), while the nearly complete amphipod mitochondrial sequence of *Parhyale hawaiiensis* reaches 73.66% (Cook et al. 2005 and our own data).

Six of the 13 protein-coding genes of *M. longipes* display unusual start codons for an arthropod mtDNA. The codon ATT is present in five genes (Table II), including *cox1*, which starts with ACG in other malacostracans (Kilpert and Podsiadlowski 2006 and references therein). In addition, the gene *atp8* starts with the non-canonical codon ATC (Table II). In turn, three of the PCGs show truncated stop codons (Table II). The genes for *nad2*, *nad4* and *cox2* end in a single T. As shown elsewhere, these truncated stop codons are likely to be completed by post-transcriptional polyadenylation, with final transcripts having functional UAA terminal codons (Ojala et al. 1981).

The *M. longipes* mitogenome shows a clear bias in nucleotide frequencies, with similar values in both strands (Table III). Strand bias reflected by GC skew (Perna and Kocher 1995) is slightly negative but close to zero in the genes encoded by the + strand, in contrast to the peracarid isopods studied so far, which show positive values (Kilpert and Podsiadlowski 2006). This has been attributed to the occurrence of inversions of the control region containing the replication origin in *Ligia oceanica* and *Idotea baltica*, since most crustaceans show moderate to high negative GC skews in the + strand (Hassanin et al. 2005; Hassanin 2006; Kilpert and Podsiadlowski 2006).

The effective number of codons (ENC), was calculated for the PCGs. This is a simple measure of codon usage, ranging from 20 when only one codon is used for each amino acid, to 61 (or even 62 in the invertebrate mitochondrial genetic codes since UGA codes there for tryptophan) when all synonymous codons are equally in use (Wright 1990). In the *M. longipes* mitogenome the PCGs show low ENC values (35.38 ± 2.84), so they use about half of the possible codons. There is a positive correlation between ENC values and GC content in third codon positions ($r^2 = 0.464$; $P < 0.01$), as described elsewhere (Kilpert and Podsiadlowski 2006). However, the genes *nad2*, *nad3* and *nad6* use a lower number of effective codons than could be expected from their relatively high GC content at third codon positions (18% for *nad2* and 23% for both *nad3* and *nad6*). Compared with isopods, *M. longipes* displays a considerably lower mean number of effective codons (and hence lower GC content at synonymous sites) and shows values similar to those found in the

amphipod *Parhyale hawaiiensis* and the more AT-rich mitogenome of *Argulus americanus* (Branchiura, Maxillopoda) (Machida et al. 2002).

Divergence in protein coding genes

We used the complete dataset of mtDNA sequences of crustacean taxa plus a representation of all major Hexapodan orders for which data are available (35 taxa, see Additional File 1) to assess the relative divergence of individual PCGs. The genes showing lower corrected divergences across Crustacea and Hexapoda were *cox1*, *cox2*, *cox3* and *cob*, while *nad2*, *nad6*, *nad4L* and *atp8* displayed about twice the mean divergence values (Fig. 3). There seems to be an association between gene variation and length and, perhaps, strand location, because shorter genes, often present on the – strand (such as *atp8* and *nad4L*), are the most divergent. Nevertheless, *nad2* is placed on the + strand in Hexapoda and in most crustaceans also shows a high substitution rate. As noted elsewhere (Cameron and Whiting 2007; Salvato et al. 2008), both variability and codon usage analyses of individual PCGs of Isoptera and Lepidoptera reveal that some of the genes most used in molecular systematics, such as *cox1* and *cox2*, have the lowest variability, while the neglected genes *nad2*, *nad3*, *nad4* and *nad6* may prove to be very useful for systematics given their variability and informative nature. Our results show that this could be extended to crustaceans, which show an underlying substitution pattern similar to hexapods at protein coding genes.

Transfer RNA genes

We identified 17 tRNA genes in a general search on the *M. longipes* mitogenome using tRNAscan-SE, and other four (*trnS1*, *trnN*, *trnF* and *trnV*) were inferred from less stringent specific searches in non-coding regions (COVE score cut-off -20). Despite this, the *trnS2* gene (tRNA-Ser_{AGN}) was not found, although it could almost completely overlap with either the *trnG* or *trnW* genes (COVE scores of +0.30 and -3.54, respectively). The *trnS2* gene shows unusual characteristics in many arthropods, such as the lack of the DHU arm (Kilpert and Podsiadlowski 2006 and references therein). In addition, in *M. longipes* the tRNA-Thr shows an unusual secondary structure, lacking completely the TΨC arm, whereas the tRNA-Gln lacks the loop normally present at this arm (Fig. 4). Nucleotide mismatches were evident in the acceptor stem for tRNA-Gln, tRNA-Arg and tRNA-Ile, and in the anticodon stem for tRNA-Lys (Fig. 4). Many cases of mismatches in stems have been described in mitochondrial tRNAs, and are supposed to be modified by RNA editing (Ojala et al. 1981; Yokobori and Paabo 1995; Masta and Boore 2004; Kilpert and Podsiadlowski 2006). The tRNA genes are present in both strands although most of them (13) are located in the + strand

(Table II, Fig. 1).

Ribosomal RNA genes

The *rrnS* and *rrnL* genes are approximately 695 and 1137 bp in length, respectively (Table II), and around 78% AT rich, thus being considerably shorter than in other crustaceans. This further explains the extreme compactness of the *M. longipes* mitogenome. The *rrnL* gene of *M. longipes* is closest to those of the amphipods *Parhyale hawaiiensis* and *Niphargus rhenorhodanensis* Schellenberg, 1937 (75% sequence identity), while the *rrnS* gene does not show any significant similarity to the sequences of other crustaceans. Not enough information on crustacean 12S and 16S rRNAs secondary structure is available to attempt reconstructing their structure based on comparative analyses.

Gene order

Metacrangonyx longipes shows a mitochondrial gene order not found in any other crustacean analysed so far (Fig. 5). Although the pancrustacean position of *trnL2* between *cox1* and *cox2* is conserved (Boore et al. 1995), many rearrangements in the *Metacrangonyx* genome compared with the ancestral pattern can be deduced (Boore et al. 1995; 1998). At least three transpositions involving genes *trnR*, *trnG* and *trnC* separately, two shifts of strand (reversals) -one involving the gene *cob* and another the segment including *trnP* and *trnT*-, and three complex tandem duplications with subsequent random losses (TDRL) are needed to account for the pattern observed in *M. longipes* compared to the pancrustacean ancestral pattern using heuristic analyses of strong common intervals with CREx (Additional File 2). Alternatively, one single reversal of the ancestral pancrustacean segment including *cob nad6 trnP trnT*, followed by a new reversal of the gene *nad6* from the - to the + strand, plus three TDRLs could have produced the *M. longipes* mitogenome gene order. The gene order also differs from the pattern found in the only other amphipod analysed thus far, *Parhyale hawaiiensis* (Cook et al. 2005 and our own data). We have almost completed the sequence for the mitogenome of this species except for a short part of the gene *rrnS* and the control region [EMBL: FM957525 and FM957526], annotating the genes for *trnV*, partial *rrnS*, *trnM*, *trnY*, *trnC*, and locating the gene *trnH* between *nad5* and *nad4*, which was absent in the previous annotation (Cook et al. 2005). In addition, based on tRNAscan results, we reannotated the tRNA genes *trnW* and *trnG* previously annotated as *trnC* and *trnW*, respectively (Cook et al. 2005; accession number AY639937). In *Parhyale hawaiiensis*, at least 10 of the tRNA genes show positional changes with respect to the pancrustacean pattern. The occurrence of identical transpositions of *trnR* and *trnG* in both *P. hawaiiensis* and *M. longipes* mitogenomes with respect to

the ancestral arrangement suggests they could have arisen in the common ancestor of amphipods. The other peracarid mitogenomes known, those of the isopods *L. oceanica* and the incomplete ones of *I. baltica* and *A. vulgare*, show quite different translocations from the assumed ancestral pancrustacean gene order (Fig. 5), with apparently no common shifts derived from the peracarid ancestor being able to explain the observed patterns (Kilpert and Podsiadlowski 2006).

Conclusions

The sequence of *Metacrangonyx longipes* introduced herein is the first complete mitogenome of a crustacean amphipod and the second for a peracarid obtained thus far, a superorder that is underrepresented in the crustacean mitochondrial genome datasets currently available. The mitogenome is very compact, with a short control region, and it appears to be the shortest mitogenome described for a crustacean. Its AT content is high (76.03%), and gene order is not conserved compared to the other four peracarids whose complete or nearly complete mitogenomes are known: the isopods *Ligia oceanica*, *Idotea baltica*, and *A. vulgare* and the amphipod *Parhyale hawaiiensis*. Common transpositions of *trnR* and *trnG* in both *P. hawaiiensis* and *M. longipes* mitogenomes with respect to the ancestral pancrustacean arrangement suggest that they were present in the common ancestor of these two amphipods. Many differences in gene order are remarkable compared to the condition displayed in isopods. Thus, no inverted strand bias of nucleotide frequencies is found in *M. longipes*, contrary to what is reported for the mitogenomes of *L. oceanica* and *I. baltica* (Kilpert and Podsiadlowski 2006). The data presented herein not only expands the sampling within the crustacean mitochondrial genomes but also will help, when congeneric species from different geographic areas are sequenced, to solve the phylogenetic position and historical biogeography of this enigmatic family found exclusively in subterranean waters.

Acknowledgements

This work has been financed by the research project CGL2006-01365 of the Spanish Ministerio de Ciencia y Innovación and European Union FEDER funds. MBR and JP are supported by FPI and “Ramon y Cajal” grants from the Ministerio de Ciencia e Innovación of Spain, respectively.

References

- Balazuc J, Ruffo S. 1953. Due nuove species del genere *Metacrangonyx* Chevreux delle acque interne del Nord Africa Francese. Memorie del Museo Civico di Storia naturale di Verona 4:25-33.
- Balke M, Ribera I, Beutel RG. 2005. The systematic position of Aspidytidae, the diversification of Dytiscoidea (Coleoptera, Adephaga) and the phylogenetic signal of third codon positions. Journal of Zoological Systematics and Evolutionary Research 43:223-242.
- Barraclough TG, Hogan JE, Vogler AP. 1999. Testing whether ecological factors promote cladogenesis in a group of tiger beetles (Coleoptera: Cicindelidae). Proceedings of the Royal Society of London B: Biological Sciences 266:1061-1067.
- Bernt M, Merkle D, Ramsch K, Fritsch G, Perseke M, Bernhard D, et al. 2007. CREx: inferring genomic rearrangements based on common intervals. Bioinformatics 23:2957-2958.
- Boore JL, Collins TM, Stanton D, Daehler LL, Brown WM. 1995. Deducing the pattern of arthropod phylogeny from mitochondrial DNA rearrangements. Nature 376:163-165.
- Boore JL, Lavrov D, Brown WM. 1998. Gene translocation links insects and crustaceans. Nature 392:667-668.
- Boutin C. 1994. Phylogeny and biogeography of metacrangonyctid amphipods in North Africa. Hydrobiologia 287:49-64.
- Boutin C, Coineau N. 1990. Regression Model, Modèle biphasé d'évolution et origine des micro-organismes stygobies interstitiels continentaux. Revue de Micropaléontologie 33:303-322.
- Boutin C, Messouli M. 1988a. *Metacrangonyx gineti* n. sp. d'une source du Haut-Atlas marocain et la famille des Metacrangonyctidae n. fam. (Crustacés Amphipodes stygobies). Vie et Milieu 38:67-84.
- Boutin C, Messouli M. 1988b. *Longipodacrangonyx maroccanus* n. gen. n. sp., nouveau représentant du groupe *Metacrangonyx* dans les eaux souterraines du Maroc. Crustaceana 13:256-271.
- Cameron SL, Whiting MF. 2007. Mitochondrial genomic comparisons of the subterranean termites from the genus *Reticulitermes* (Insecta: Isoptera: Rhinotermitidae). Genome 50:188-202.
- Carapelli A, Liò P, Nardi F, van der Wath E, Frati F. 2007. Phylogenetic analysis of mitochondrial protein coding genes confirms the reciprocal paraphyly of Hexapoda and Crustacea. BMC Evolutionary Biology 7:S8.
- Chevreux E. 1909. Amphipodes (1ère série). Archives de Zoologie Expérimentale et Générale 5:27-

- Cook CE, Yue Q, Akam M. 2005. Mitochondrial genomes suggest that hexapods and crustaceans are mutually paraphyletic. *Proceedings of the Royal Society of London B: Biological Sciences* 272:1295-1304.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek RC. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294-299.
- Hassanin A. 2006. Phylogeny of Arthropoda inferred from mitochondrial sequences: strategies for limiting the misleading effects of multiple changes in pattern and rates of substitution. *Molecular Phylogenetics and Evolution* 38:100-116.
- Hassanin A, Leger N, Deutsch J. 2005. Evidence for multiple reversals of asymmetric mutational constraints during the evolution of the mitochondrial genome of Metazoa, and consequences for phylogenetic inferences. *Systematic Biology* 54:277-298.
- Jaume D, Christenson K. 2001. Amphi-Atlantic distribution of the subterranean amphipod family Metacrangonyctidae (Gammaridea). *Contributions to Zoology* 70:99-125.
- Karaman GS. 1989. *Metacrangonyx ortali* n sp, a new subterranean member of the family Crangonyctidae, from the Dead Sea region. *Studia Marina* 20:33-49.
- Karaman GS, Pesce GL. 1979. Researches in Africa by the Zoological Institute of l'Aquila, Italy V: On three subterranean amphipods from North Africa (Amphipoda: Gammaridea). *Bulletin Zoologisch Museum, Universiteit van Amsterdam* 7:197-207.
- Kilpert F, Podsiadlowski L. 2006. The complete mitochondrial genome of the common sea slater, *Ligia oceanica* (Crustacea, Isopoda) bears a novel gene order and unusual control region features. *BMC Genomics* 7:241.
- Lowe TM, Eddy SR. 1997. tRNAscan-SE: A program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Research* 25: 955-964.
- Machida RJ, Miya MU, Nishida M, Nishida S. 2002. Complete mitochondrial DNA sequence of *Tigriopus japonicus* (Crustacea: Copepoda). *Marine Biotechnology* 4:406-417.
- Marcadé I, Cordaux R, Doublet V, Debenest C, Bouchon D, Raimond R. 2007. Structure and evolution of the atypical mitochondrial genome of *Armadillidium vulgare* (Isopoda, Crustacea). *Journal of Molecular Evolution* 65:651-659.
- Margalef R. 1952. Materiales para la hidrobiología de la isla de Menorca. *Publicaciones Instituto de Biología Aplicada* 11:5-112.

- Martin JW, Davis GE. 2001. An updated classification of the recent Crustacea. Los Angeles, CA: Natural History Museum of Los Angeles County. 123 p.
- Masta SE, Boore JL. 2004. The complete mitochondrial genome sequence of the spider *Habronattus oregonensis* reveals rearranged and extremely truncated tRNAs. *Molecular Biology and Evolution* 21:893-902.
- Messouli M, Boutin C, Coineau N. 1991. Phylogénie et biogéographie évolutive d'un groupe de Metacrangonyctidae, Crustacés Amphipodes stygobies du Maroc. I. Les espèces du groupe *panousei*. *Mémoires de Biospéologie* 25: 247-262.
- Ojala D, Montoya J, Attardi G 1981. tRNA punctuation model of RNA processing in human mitochondria. *Nature* 290:470-474.
- Oulbaz Z, Coineau N, Boutin C 1998. *Metacrangonyx knidirii* n sp et *Metacrangonyx remyi* Balazuc et Ruffo, 1953 (Crustacés Amphipodes Metacrangonyctidae) des eaux souterraines du Maroc. *Mémoires de Biospéologie* 25:35-43.
- Perna NT, Kocher TD. 1995. Patterns of nucleotide composition at fourfold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution* 41:353-358.
- Podsiadlowski L, Bartolomaeus T. 2006. Major rearrangements characterize the mitochondrial genome of the isopod *Idotea baltica* (Crustacea: Peracarida). *Molecular Phylogenetics and Evolution* 40:893-899.
- Ruffo S. 1954. *Metacrangonyx longicaudus* n sp (Amphipoda Gammaridae) delle acque sotterranee del Sahara Marocchino. *Memorie del Museo Civico di Storia naturale di Verona* 4:127-130.
- Ruffo S. 1982. Una nouvelle espèce de *Metacrangonyx* Chevreux (Amphipoda: Gammaridae) du désert du Sinaï. *Israel Journal of Zoology* 31:151-156.
- Salvato P, Simonato M, Battisti A, Negrisolò E. 2008. The complete mitochondrial genome of the bag-shelter moth *Ochrogaster lunifer* (Lepidoptera, Notodontidae). *BMC Genomics* 9: 331.
- Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22:2688-2690.
- Stoch F. 1997. *Metacrangonyx ilvanus* n sp, the first Italian representative of the family Metacrangonyctidae (Crustacea: Amphipoda). *Annals of Limnology* 33:255-262.
- Stock JH, Rondé-Broekhuizen B. 1986. A new species of *Pygocrangonyx*, an amphipod genus with African affinities, from Fuerteventura. *Bijdragen tot de dierkunde* 56:247-266.
- Supek F, Vlahovicek K. 2004. INCA: synonymous codon usage analysis and clustering by means of self-organizing map. *Bioinformatics* 14:2329-2330.

- Tamura K, Nei M, Kumar S. 2004. Prospects for inferring very large phylogenies by using the Neighbor-Joining method. *Proceedings of the National Academy of Science of the United States of America* 101:11030-11035.
- Tamura K, Dudley J, Nei M, Kumar S. 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) Software Version 4.0. *Molecular Biology and Evolution* 24: 1596-1599.
- Wolstenholme DR. 1992. Animal mitochondrial DNA: structure and evolution. *International Review of Cytology* 141:173-216.
- Wright F. 1990. The 'effective number of codons' used in a gene. *Gene* 87:23-29.
- Xia X , Xie Z. 2001. DAMBE: software package for data analysis in molecular biology and evolution. *Journal of Heredity* 92: 371-373.
- Xiong B, Kocher TD. 1991. Comparison of mitochondrial DNA sequences of seven morphospecies of black flies (Diptera: Simuliidae). *Genome* 34: 306-311.
- Yang JS, Yang WJ. 2008. The complete mitochondrial genome sequence of the hydrothermal vent galatheid crab *Shinkaia crosnieri* (Crustacea: Decapoda: Anomura): A novel arrangement and incomplete tRNA suite. *BMC Genomics* 9:257.
- Yokobori S, Paabo S. 1995. tRNA editing in metazoans. *Nature* 377: 490.
- Zhang DX, Szymura JM, Hewitt GM. 1995. Evolution and structural conservation of the control region of insect mitochondrial DNA. *Journal of Molecular Evolution* 40:382-391.

Figures

Figure 1. Map of the mitochondrial genome of *Metacrangonyx longipes*. Gray and white segments indicate genes coded on the + strand and – strand, respectively.

Figure 2. Putative secondary structure of the mitochondrial control region of *Metacrangonyx longipes*. The box indicates the conserved GACT motif present also in the isopod *Ligia oceanica* and the hoplocarid malacostracan *Squilla mantis*.

Figure 3. Mean relative corrected divergences of protein coding genes of Crustacea and Hexapoda. DNA divergences of individual genes were estimated from pairwise comparisons among the complete mitogenomes of crustaceans and 35 species representing all major Hexapoda orders.


Figure 4. Putative secondary structures of mitochondrial tRNAs in *Metacrangonyx longipes*.


Figure 5. Mitochondrial gene order in Peracarida (Isopoda + Amphipoda) mitogenomes compared with the pancrustacean ancestral pattern. Different colours are used to identify particular conserved and rearranged segments or genes. Genes underlined are present at the – strand.


Additional files


Additional File 1. Taxon names and EMBL accession numbers of the crustacean and hexapod mitogenomes used for gene annotation and gene divergence analyses.

Additional File 2. Rearrangement steps deduced using detection of strong interval trees to account for the gene order of *Metacrangonyx longipes* mitogenome compared with the ancestral pancrustacean order.









Pancrustacea putative ancestral gene order




Metacangronyx longipes (Amphipoda)




Parhyale hawaiiensis (Amphipoda)




Ligia oceanica (Isopoda)



Ideotea baltica (Isopoda)



Armadillidium vulgare (Isopoda)



CRUSTACEA

Species	Accession n°	Taxonomy
<i>Argulus americanus</i>	NC_005935	Maxillopoda Branchiura
<i>Armadillidium vulgare</i>	EF643519	Malacostraca Peracarida Isopoda
<i>Armillifer armillatus</i>	NC_005934	Pentastomida
<i>Artemia franciscana</i>	NC_001620	Branchiopoda Anostraca
<i>Callinectes sapidus</i>	NC_006281	Malacostraca Eucarida Decapoda Pleocyemata Brachyura
<i>Cherax destructor</i>	NC_011243	Malacostraca Eucarida Decapoda Pleocyemata
<i>Daphnia pulex</i>	NC_000844	Branchiopoda Anomopoda
<i>Eriocheir sinensis</i>	NC_006992	Malacostraca Eucarida Decapoda Pleocyemata Brachyura
<i>Euphausia superba</i>	AB084378	Malacostraca Eucarida Euphausiacea
<i>Fenneropenaeus chinensis</i>	NC_009679	Malacostraca Eucarida Decapoda Dendrobranchiata
<i>Geothelphusa dehaani</i>	NC_007379	Malacostraca Eucarida Decapoda Pleocyemata Brachyura
<i>Gonodactylus chiragra</i>	NC_007442	Malacostraca Hoplocarida Stomatopoda
<i>Halocaridina rubra</i>	NC_008413	Malacostraca Eucarida Decapoda Pleocyemata Caridea
<i>Harpisquilla harpax</i>	NC_006916	Malacostraca Hoplocarida Stomatopoda
<i>Hutchinsoniella macracantha</i>	NC_005937	Cephalocarida
<i>Idotea baltica</i>	DQ442915	Malacostraca Peracarida Isopoda
<i>Lepeophtheirus salmonis</i>	NC_007215	Maxillopoda Copepoda
<i>Ligia oceanica</i>	NC_008412	Malacostraca Peracarida Isopoda Oniscidea
<i>Litopenaeus vannamei</i>	NC_009626	Malacostraca Eucarida Decapoda Dendrobranchiata
<i>Lysiosquillina maculata</i>	NC_007443	Malacostraca Hoplocarida Stomatopoda
<i>Macrobrachium rosenbergii</i>	NC_006880	Malacostraca Eucarida Decapoda Pleocyemata Caridea
<i>Marsupenaeus japonicus</i>	NC_007010	Malacostraca Eucarida Decapoda Dendrobranchiata
<i>Megabalanus volcano</i>	NC_006293	Maxillopoda Thecostraca Cirripedia Thoracica
<i>Pagurus longicarpus</i>	NC_003058	Malacostraca Eucarida Decapoda Anomura
<i>Panulirus japonicus</i>	NC_004251	Malacostraca Eucarida Decapoda Pleocyemata
<i>Parhyale hawaiiensis</i>	AY639937	Malacostraca Peracarida Amphipoda
<i>Penaeus monodon</i>	NC_002184	Malacostraca Eucarida Decapoda Dendrobranchiata
<i>Pollicipes mitella</i>	NC_008742	Maxillopoda Thecostraca Cirripedia Thoracica
<i>Pollicipes polymerus</i>	NC_005936	Maxillopoda Thecostraca Cirripedia Thoracica
<i>Portunus trituberculatus</i>	NC_005037	Malacostraca Eucarida Decapoda Brachyura
<i>Pseudocarcinus gigas</i>	NC_006891	Malacostraca Eucarida Decapoda Pleocyemata Brachyura
<i>Pseudosquilla ciliata</i>	AY947836	Malacostraca Hoplocarida Stomatopoda
<i>Speleonectes tulumensis</i>	NC_005938	Remipedia
<i>Squilla empusa</i>	NC_007444	Malacostraca Hoplocarida Stomatopoda
<i>Squilla mantis</i>	NC_006081	Malacostraca Hoplocarida Stomatopoda
<i>Tetraclita japonica</i>	NC_008974	Maxillopoda Thecostraca Cirripedia Thoracica

<i>Tigriopus californicus</i>	NC_008831	Maxillopoda Copepoda
<i>Tigriopus japonicus</i>	NC_003979	Maxillopoda Copepoda Harpacticoida
<i>Triops cancriformis</i>	NC_004465	Branchiopoda Notostraca
<i>Triops longicaudatus</i>	NC_006079	Branchiopoda Notostraca
<i>Vargula hilgendorffii</i>	NC_005306	Ostracoda Myodocopa
<i>Metacrangonyx longipes</i>	AM944817	Malacostraca Peracarida Amphipoda

HEXAPODA

Species	Accession n°	Taxonomy
<i>Aleurodicus dugesii</i>	NC_005939	Insecta Hemiptera Hemimetabola
<i>Anopheles gambiae</i>	NC_002084	Insecta Diptera Holometabola
<i>Antheraea pernyi</i>	NC_004622	Insecta Lepidoptera Holometabola
<i>Apis mellifera ligustica</i>	NC_001566	Insecta Hymenoptera Holometabola
<i>Bombyx mori</i>	NC_002355	Insecta Lepidoptera Holometabola
<i>Ceratitis capitata</i>	NC_000857	Insecta Diptera Holometabola
<i>Chrysomya putoria</i>	NC_002697	Insecta Diptera Holometabola
<i>Crioceris duodecimpunctata</i>	NC_003372	Insecta Coleoptera Holometabola
<i>Drosophila melanogaster</i>	NC_001709	Insecta Diptera Holometabola
<i>Gomphiocephalus hodgsoni</i>	NC_005438	Collembola
<i>Grylotalpa orientalis</i>	NC_006678	Insecta Orthoptera Hemimetabola
<i>Haematobia irritans irritans</i>	NC_007102	Insecta Diptera Holometabola
<i>Heterodoxus macropus</i>	NC_002651	Insecta Phthiraptera Hemimetabola
<i>Homalodisca coagulata</i>	NC_006899	Insecta Hemiptera Hemimetabola
<i>Japyx solifugus</i>	NC_007214	Diplura
<i>lepidopsocid RS-2001</i>	NC_004816	Insecta Psocoptera Hemimetabola
<i>Locusta migratoria</i>	NC_001712	Insecta Orthoptera Hemimetabola
<i>Melipona bicolor</i>	NC_004529	Insecta Hymenoptera Holometabola
<i>Nesomachilis australica</i>	NC_006895	Insecta Archaeognatha Ametabola
<i>Onychiurus orientalis</i>	NC_006074	Collembola
<i>Orthetrum triangulare melania</i>	AB126005	Insecta Odonata Hemimetabola
<i>Ostrinia nubilalis</i>	NC_003367	Insecta Lepidoptera Holometabola
<i>Pachypsylla venusta</i>	NC_006157	Insecta Hemiptera Hemimetabola
<i>Periplaneta fuliginosa</i>	NC_006076	Insecta Dictyoptera Hemimetabola
<i>Philaenus spumarius</i>	NC_005944	Insecta Hemiptera Hemimetabola
<i>Podura aquatica</i>	NC_006075	Collembola

<i>Pteronarcys princeps</i>	NC_006133	Insecta Plecoptera Hemimetabola
<i>Pyrocoelia rufa</i>	NC_003970	Insecta Coleoptera Holometabola
<i>Schizaphis graminum</i>	NC_006158	Insecta Hemiptera Hemimetabola
<i>Thermobia domestica</i>	NC_006080	Insecta Thysanura Ametabola
<i>Thrips imaginis</i>	NC_004371	Insecta Thysanoptera Hemimetabola
<i>Triatoma dimidiata</i>	NC_002609	Insecta Hemiptera Hemimetabola
<i>Tribolium castaneum</i>	NC_003081	Insecta Coleoptera Holometabola
<i>Tricholepidion gertschi</i>	NC_005437	Insecta Thysanura Ametabola
<i>Xenos vesparum</i>	DQ364229	Strepsiptera Holometabola

