

# Insecticidal effects of Moroccan plant extracts on development, energy reserves and enzymatic activities of *Plodia interpunctella*

N. Bouayad<sup>1,2</sup>, K. Rharrabe<sup>3</sup>, N. N. Ghailani<sup>1</sup>, R. Jbilou<sup>1</sup>, P. Castañera<sup>2</sup> and F. Ortego<sup>2\*</sup>

<sup>1</sup> PER – Centre des Études Environnementales Méditerranéennes. Laboratoire de Biologie Appliquée & Sciences de l'Environnement. Université Abdelmalek Essaadi. Faculté des Sciences et Techniques.

BP 416. Tanger. Morocco

<sup>2</sup> Departamento de Biología Medioambiental. Centro de Investigaciones Biológicas. CSIC.

Ramiro de Maeztu, 9. 28040 Madrid. Spain

<sup>3</sup> Département des Ressources Naturelles. Faculté Polydisciplinaire-Larache. Université Abdelmalek Essaadi.

BP 745. Larache. Morocco

## Abstract

The aim of this work was to study the effects of methanol extracts of ten plant species used in traditional medicine in Morocco (*Peganum harmala*, *Ajuga iva*, *Rosmarinus officinalis*, *Lavandula stoechas*, *Lavandula dentata*, *Cistus ladanifer*, *Cistus salviaefolius*, *Cistus monspeliensis*, *Centaurium erythraea* and *Launaea arborescens*) on *Plodia interpunctella* Hübner (Lepidoptera: Pyralidae) larvae. Firstly, we studied the effects of the ingestion of these extracts at 500 ppm on post-embryonic development parameters. Most plant extracts provoked a notable decrease of larval weight 8 days after treatment (up to 33% weight loss with *C. erythraea*) and caused significant alterations on pupation (ranging from 5% to 85%) and adult emergence (below 2.5% with *R. officinalis*, *C. erythraea* and *A. iva*). The plant extracts that showed strongest effects on post-embryonic development were selected to test their effects on the following physiological parameters: larval reserve substances (at 500 ppm); and midgut activities of hydrolytic and detoxification enzymes (at 500, 750 and 1,000 ppm). All treatments provoked a significant reduction of protein and carbon hydrate larval contents, the inhibition of proteases and  $\alpha$ -amylase activities in a dose depended manner, and the induction of glutathione S-transferase and esterase (using MtB as substrate) activities, whereas the activity of cytochrome P450 monooxygenases and esterases (using 1-NA as substrate) increase or decrease depending on the extract concentration and the plant analyzed.

**Additional key words:** bioinsecticides; detoxification enzymes; digestive enzymes; reserve substances.

## Introduction

The Indianmeal moth, *Plodia interpunctella* (Hübner) (Lepidoptera: Pyralidae) is an important cosmopolitan pest of stored grain and value-added food products with a worldwide distribution. In addition to damaging a commodity by feeding, the larvae spin silken webs causing economic losses by affecting food quality. In Morocco, *P. interpunctella* is considered a major pest during the processing and storage of dried fruits such as dates (Azemat *et al.*, 2005). Currently, synthetic insecticides and methyl bromide fumigation are used for the treatment of these stored products (Azemat *et al.*, 2005). However, serious concerns have been raised about health and environmental problems associated with the widespread use of insecticides, including

development of insecticide resistant strains (White, 1995; Phillips & Throne, 2010). In addition, methyl bromide has been identified as an ozone-depleting substance and its use restricted in accordance with the Montreal Protocol (Patil *et al.*, 2004). Therefore, there is an urgent need to develop safe, convenient, environmentally friendly and low-cost alternatives.

Alkaloids, terpenoids, phenolics, flavonoids and other secondary metabolites of plants play an important role in plant-insect interaction and are commonly responsible of plant resistance to insects (Isman, 2002). These plant allelochemicals can act as attractive, repellent or toxic agents, as well as growth regulators, affecting physiological processes of insects (Ortego *et al.*, 1995; Khambay *et al.*, 2002; Liang *et al.*, 2003). The insecticidal activity of some plant derivatives has been well established, and the efficacy of a few of these active principles, such as azadirachtin, has resulted in their exploitation as crop protectants (Mordue *et al.*, 2005).

\* Corresponding author: [ortego@cib.csic.es](mailto:ortego@cib.csic.es)

Received: 30-12-11. Accepted: 22-10-12.

Plants frequently used in traditional medicine in Morocco (Bellakhdar, 1997) have shown their potential as bioinsecticides. Methanol extracts from seven plant species: *Centaurium erythraea*, *Peganum harmala*, *Ajuga iva*, *Aristolochia baetica*, *Pteridium aquilinum* and *Raphanus raphanistrum* were shown to inhibit the growth of *Tribolium castaneum* larvae (Jbilou et al., 2008). Leaf and seed extracts of *P. harmala* reduced the food intake and delayed larval development of *Schistocerca gregaria* (Abbassi et al., 2003a,b) and *T. castaneum* (Jbilou & Sayah, 2008). Likewise, Papa-christos & Stamopoulos (2002) reported that larvae of *Acanthoscelides obtectus* were susceptible to the essential oils from *Lavandula hybrida*, *Rosmarinus officinalis* and *Eucalyptus globulus*. None of these plant extracts have been tested against *P. interpunctella*. However, previous studies have shown that this species is susceptible to different allelochemicals, such as 20-hydroxyecdysone (Rharrabe et al., 2009), azadirachtin (Rharrabe et al., 2008), harmaline (Rharrabe et al., 2007) and harmine (Bouayad et al., 2012).

The mode of action of plant allelochemicals at the enzymatic level is mostly unknown, but some of them appear to disturb the digestive process by targeting insect gut digestive enzymes, such as  $\alpha$ -amylases and proteases (Duffey & Stout, 1996). Senthil Nathan et al. (2005a,b) showed that ingestion of azadirachtin by *Spodoptera litura* and *Cnaphalocrocis medinalis* caused a decrease on gut enzyme activities. The limonoids 1,7-di-O-acetylthymosin and 3,7-di-O-acetylthymosin reduced digestive protease activity in *Leptinotarsa decemlineata* (Ortego et al., 1999) and *S. exigua* (Caballero et al., 2008). Likewise, ingestion of Jatropherol-I induced a decline in midgut protease activities in *Bombyx mori* (Jing et al., 2005). Besides, the walls of the digestive tract in insects have a high content of detoxification enzymes, as a barrier to toxic allelochemicals that they may consume with the diet (Ahmad et al., 1986). A common strategy in insects is to produce these enzymes in response to the ingestion of allelochemicals, suggesting that induction plays an important role in the adaptation of insects to plant diets (Terriere, 1984). However, some plant allelochemicals are known to inhibit the activity of detoxification enzymes (Smirle et al., 1996; Jing et al., 2005), increasing the toxicity of co-occurring plant compounds (Berenbaum & Neal, 1985).

The aim of this study was to evaluate the potential of ten medicinal plants commonly found in Morocco: *P. harmala*, *A. iva*, *R. officinalis*, *C. erythraea*, *Lavandula stoechas*, *Lavandula dentata*, *Cistus ladanifer*,

*Cistus salviaefolius*, *Cistus monspeliensis*, and *Launaea arborescens* for the control of *P. interpunctella*. We assessed the effect of methanol extracts from these plants on post-embryonic development, reserve substances levels and gut enzymatic activities, in order to have some insights about the physiological responses of the insect to the ingestion of the plant extracts.

## Material and methods

### Insect rearing

*P. interpunctella* used in this study originated from a laboratory culture reared for many generations on wheat (*Triticum aestivum*) flour as a medium. All stages of the insect were kept under standard conditions at 26-30 °C, 55-65% relative humidity and a photoperiod of 16:8 (L:D). Adults were collected for oviposition and eggs were hatched in 0.25-L glass containers half full of wheat flour. In the present study, 5<sup>th</sup> instar larvae weighing  $3.4 \pm 0.2$  mg were used in all assays.

### Plant materials and extracts preparation

Methanol extracts were prepared from ten plants commonly used in traditional medicine in Morocco. The plants tested were chosen based on previous records of insecticidal activity or because they belong to plant families that are known to be rich in secondary metabolites, being all of them readily available in Tangier Region (NW of Morocco). Seeds of *P. harmala* (Zygophyllaceae) and aerial parts of *A. iva* and *R. officinalis* (Lamiaceae), *L. stoechas* and *L. dentata* (Lamiaceae), *C. ladanifer*, *C. salviaefolius* and *C. monspeliensis* (Cistaceae), *C. erythraea* (Gentianaceae) and *L. arborescens* (Compositae) were collected during February-March period. The plants were identified by Professor Ennabili Abdessalam (Institut National des Plantes Aromatiques et Médicinales de Taounate, Morocco). Plants were rinsed with distilled water, dried in an oven at 38 °C for 72 h, ground to a powder with an electrical blender and stored at room temperature in hermetically sealed plastic bags prior to extraction. Each plant sample (20 g) was extracted twice with 160 mL methanol to gain polar and medium polar metabolites by using ultrasound apparatus for 30 min. All extracts were stored at 4 °C. Before using, extracts were evaporated to dryness, and the residue was weighted

and redissolved in the same solvent at a concentration of 100 mg of crude extract mL<sup>-1</sup> of methanol.

## Bioassays

Methanol extracts were mixed with 5 g of the diet (wheat flour) at a concentration of 500, 750 or 1,000 ppm, depending on the experimental conditions (see below). The solvent was allowed to evaporate at 37 °C during 48 h.

To study the effect of the methanol extracts on post-embryonic development, 25-5<sup>th</sup> instar larvae were added to each glass vial (diameter 4 cm, height 7 cm) containing diet treated at 500 ppm. A control was prepared in the same way with methanol alone. To verify that the effect was due to plant extracts ingestion and not to a deterrence effect, another experimental group (starved) was generated by transferring larvae into clean glass vials with no food. Ten replicates were set up for the treated, starved and control larvae. Observations including larval weight, cannibalism, pupation and adult emergence were recorded. Mortality was determined by brown colour with no observable movements. Cannibalism was noted by disappearance of body larvae from Petri-dishes. Larvae and pupae were maintained at the conditions described above during the whole experiment.

For biochemical analysis, 5<sup>th</sup> instar larvae were reared individually for 8 days in diets treated with the different extracts tested. Control and starved larvae were also reared individually for 8 days as described above. For analyses of reserve substances, plant extracts were tested at 500 ppm, and each sample was composed of three cold-anesthetized larvae. For digestive and detoxification enzyme activities, plant extracts were tested at 500, 750 and 1,000 ppm, and larvae were dissected and the midguts individually stored frozen at -20 °C until needed. Ten and 20 replicates per concentration were done for reserve substances and enzymatic activities, respectively.

## Biochemical analyses

Protein content was determined according to the method of Bradford (1976). Three larvae were homogenized in 1 mL of Tris-HCl buffer (50 mM, pH 7). The homogenate was centrifuged at 9000 g for 20 min. A volume of 200 µL of extract was mixed with 2 mL of Bradford reactive. The absorbance was measured at

595 nm and protein levels calculated using bovine serum albumin as the standard.

Carbon hydrate content was quantified by the method of Roe (1955) using anthrone as reagent. Three larvae were homogenized in 100 µL of NaCl 125 mM. A volume of 25 µL was mixed with 1 mL of ethanol saturated with sodium sulphate. It was then centrifuged at 1,000 rpm for 10 min and the supernatant was discarded. The pellet was resuspended in 0.5 mL of 70% ethanol, then centrifuged as before and the supernatant was discarded. The final pellet was heated to drive off residual ethanol and then dissolved in 0.5 mL of 30% potassium hydroxide and heated at 100 °C for 15 min. The digest was allowed to cool and 1 mL of absolute ethanol was added. This was spun down by centrifugation at 1,000 rpm for 10 min. The supernatant was carefully removed and the pellet was dissolved in 0.5 mL of distilled water. Two mL of anthrone reagent (0.05% in sulfuric acid) was then added. After mixing, the samples were heated at 90 °C for 15 min. They were rapidly cooled. The absorbance was read at 620 nm and carbon hydrate levels were calculated by reference to a standard curve using glycogen.

Lipids were extracted according to the method of Van Handel (1965). Three larvae were homogenized in 1 mL of chloroform/methanol (1:1 v/v). The homogenate was centrifuged at 1,000 rpm for 10 min at 4 °C. The supernatant was mixed with 1 mL of chloroform and 0.5 mL of distilled water and centrifuged at 1,500 rpm for 1 h at 4 °C. The aqueous phase was discarded (this operation was repeated twice). The lipid extract was evaporated to dryness. Lipid content was quantified by the method of Zöllner & Krich (1962). The sample was digested with 1 mL of sulphuric acid at 100 °C for 10 min. The tube was cooled and 5 mL of sulphovanillin reagent (orthophosphoric acid/0.6% aqueous vanillin solution 4:1) was added to the mixture. After 40 min, the absorbance was measured at 530 nm and lipid level was calculated by reference to a standard curve prepared using cholesterol palmitate.

Proteolytic activities were determined using specific substrates as described by Ortego *et al.* (1996). Trypsin-like activity was assayed using 0.25 mM BApNa (N $\alpha$ -benzoyl-DL-arginine p-nitroanilide) at pH 10.5 and incubating for 1 h, chymotrypsin-like activity with 0.25 mM SA<sub>2</sub>PPpNa (N succinyl-alanine-alanine-proline-phenylalanine p-nitroanilide) at pH 10.5 and incubating for 1 h, elastase-like activity with 0.25 mM SA<sub>3</sub>pNa (N-succinyl-alanine-alanine-alanine p-nitroanilide) at pH 10.5 and incubating for 24 h, leucine

aminopeptidase-like activity with 1 mM LpNa (L-leucine p-nitroanilide) at pH 7 and incubating for 1 h, carboxypeptidase A-like activity with 1 mM HPA (hippuryl-phenylalanine) at pH 7 and incubating for 4 h, and carboxypeptidase B-like activity with 1 mM HA (hippuryl-L-arginine) at pH 7 and incubating for 4 h.

Alpha-amylase activity was measured by the method of Valencia *et al.* (2000). A volume of 50  $\mu$ L of midgut extract was added to 50  $\mu$ L of 50 mM sodium citrate buffer (NaCl 10 mM, CaCl<sub>2</sub> 20 mM), pH 8.5. To the mixture was added 100  $\mu$ L of 0.5% starch solution. The sample was incubated at 37°C during 15 min. Then 2.5 mL of iodide (0.5% I<sub>2</sub> and 5% KI) was added. It was centrifuged at 4,000 rpm for 20 min. The absorbance of the supernatant was read at 580 nm. The  $\alpha$ -amylase activity was calculated by reference to a standard curve prepared using starch solution.

Esterase activity was determined using 1-NA (1-naphthyl acetate) and MtB (S-methyl thiobutanoate) as substrates. For activity towards 1-NA, 5  $\mu$ L of midgut extract were added to 160  $\mu$ L of reaction mixture (pH 7) to give a final concentration of 0.25 mM 1-NA, and the mixture incubated 1 h at 30°C. The reaction was stopped by adding 80  $\mu$ L of 0.12% (w/v) o-dianisidine bis (diazotized) zinc double salt (Fast Blue B Salt, FBB) and 2% (w/v) sodium dodecyl sulphate (SDS). The absorbance was read at 600 nm. Specific activity was calculated by reference to the procedure described by Gomori (1953) using a standard curve of 1-naphthol accordingly. Esterase activity towards MTB was measured using a reaction mixture that contained 5  $\mu$ L of midgut extract, 0.1 M phosphate buffer pH 7.2 mM MtB and 2.5 mM DTNB (5'5 dithio-bis (2-nitrobenzoic acid)). The increment in absorbance at 412 nm was recorded during 15 min at 30°C to determine the nmol substrate hydrolyzed min<sup>-1</sup> mg<sup>-1</sup> protein, using a molar extinction coefficient of 13,600 M<sup>-1</sup> cm<sup>-1</sup> (Ellman, 1959).

Glutathione S-transferase activity was measured using 40  $\mu$ L of midgut extract, 1 mM CDNB (1-chloro-2,4-dinitrobenzene), 5 mM reduced glutathione, and 0.1 M Tris buffer pH 8. The increment in absorbance at 340 nm was recorded during 5 min at 30°C to determine the nmol substrate conjugated per min per mg protein, using a molar extinction coefficient of 9.6 mM<sup>-1</sup> cm<sup>-1</sup> (Habig *et al.*, 1974).

P450 monooxygenases activity was assayed using 50  $\mu$ L of midgut extract, 130  $\mu$ L NADPH generating system at pH 7, and 20  $\mu$ L of cytochrome C solution, and the mixture incubated 1 h at 30°C. The reaction

was stopped by adding 100  $\mu$ L of methanol, and then centrifuged at 12,000 rpm for 5 min. The absorbance was read at 550 nm to determine the nmol cytochrome c reduced min<sup>-1</sup> mg<sup>-1</sup> protein, using a molar extinction coefficient of 27.6 mM<sup>-1</sup> cm<sup>-1</sup> (Masters *et al.*, 1965).

## Statistical analysis

Statistical analyses were performed using Statistica (1997) software. Cannibalism, pupation and adult emergence data (percentages) were analyzed by one-way ANOVA, after logit transformation, followed by Dunnett's test for multiple comparisons with respect to the control. Larval weight and biochemical parameters data were analyzed using one-way ANOVA followed by Dunnett's test. The significance was tested at the  $p = 0.05$  level.

## Results

### Effects of plant extracts on post-embryonic development of *P. intepunctella*

At the beginning of the experiment, the average weight of 5<sup>th</sup> instar larvae was 3.4  $\pm$  0.2 mg. After 10 days, the weight of larvae feeding on control diet reached 7.3 mg (115% weight gain), whereas the weight of starved larvae decreased to 1.1 mg (68% weight loss) (Table 1). During the same period, the weight of larvae feeding on diet treated with the different plant extracts tested was significantly reduced when compared with the control group. In fact, most of the treatments led to a net weight loss: 32%, 18%, 21%, 6%, 33%, 32%, and 12% for larvae treated by *P. harmala*, *C. ladanifer*, *L. dentata*, *R. officinalis*, *C. erythraea*, *A. iva* and *L. stoechas*, respectively. Extracts of *R. officinalis* and *C. erythraea* induced a marked cannibalism (20-25%) between larvae, similar to that observed for starved larvae (30%) (Table 1). There was no cannibalism between larvae fed the control diet, and for the rest of the treatments was below 5%.

Pupation of control larvae was 90%, whereas it was completely abolished in starved larvae (Table 1). The percentage of pupation varied among treatments (ranging from 5% to 85%), being significantly lower than in the control for all treatments, except for *C. monspeliensis*. The higher reductions in pupation corresponding to larvae treated with *P. harmala* and *C. ladanifer*.

**Table 1.** Effect of ingestion of diet treated with plant extracts (500 ppm) on post-embryonic development of *P. interpunctella*

Treatment	Larval weight <sup>1</sup> (mg)	Cannibalism <sup>2</sup> (%)	Pupation <sup>3</sup> (%)	Adult emergence <sup>4</sup> (%)
Control	7.3 ± 0.2	0 ± 0.0	90 ± 1.2	90 ± 0.7
<i>P. harmala</i>	2.3 ± 0.1*	5 ± 0.3*	5 ± 0.3*	0 ± 0.0*
<i>C. ladanifer</i>	2.8 ± 0.1*	5 ± 0.4*	5 ± 0.3*	0 ± 0.0*
<i>L. dentata</i>	2.7 ± 0.1*	2.5 ± 0.2*	67.5 ± 1.3*	27 ± 1.0*
<i>R. officinalis</i>	3.2 ± 0.1*	25 ± 0.7*	20 ± 0.9*	2 ± 0.1*
<i>C. salviaefolius</i>	4.3 ± 0.1*	7.5 ± 0.2*	75 ± 0.9*	45 ± 1.4*
<i>C. monspeliensis</i>	4.1 ± 0.1*	5 ± 0.3*	85 ± 1.0	68 ± 1.6*
<i>C. erythraea</i>	2.3 ± 0.1*	20 ± 0.7*	14 ± 0.5*	2 ± 0.1*
<i>A. iva</i>	2.3 ± 0.1*	5 ± 0.1*	20 ± 0.9*	1.5 ± 0.2*
<i>L. stoechas</i>	3.0 ± 0.1*	5 ± 0.3*	65 ± 1.2*	26 ± 1.7*
<i>L. arborescens</i>	3.5 ± 0.1*	5 ± 0.2*	75 ± 1.3*	67.5 ± 2.0*
Starved	1.1 ± 0.1*	30 ± 1.1*	0 ± 0.0*	0 ± 0.0*

Values are the mean ± standard error of <sup>1</sup> larval weight 8 days after treatment, <sup>2</sup> cannibalism 16 days after treatment, <sup>3</sup> pupation 20 days after treatment, <sup>4</sup> adult emergence 26 days after treatment (10 replicates of 25 larvae each). \* Significantly different from control (Dunnet test,  $p \leq 0.05$ ).

In the same way, adult emergence was significantly affected by the presence of plant extracts in the larval diet (Table 1). Larval starvation and treatment with *P. harmala* and *C. ladanifer* completely inhibited adult emergence. Strong effects on adult emergence (below 2.5%) were also obtained with larvae treated with *R. officinalis*, *C. erythraea* and *A. iva*.

### Effects of plant extracts on energy reserves and enzymatic activities

The plant extracts that showed strongest effects on post-embryonic development were selected to test their effects on reserve substances levels and enzymatic activities of *P. interpunctella* larvae.

Protein content was severely reduced in larvae fed on treated diets and on starved larvae as compared with the control group; being the strongest effect for larvae treated with *C. erythraea*, *P. harmala* and *C. ladanifer* (Table 2). Plant extracts induced also a decrease in carbon hydrate level, but the reduction was smaller than that observed for starved larvae (Table 2). The lipid content was not affected by the different plants extracts tested, whereas it was significantly reduced in starved larvae (Table 2).

Our results showed that the specific activity for the hydrolysis of BApNa, SA<sub>2</sub>PPpNa, SA<sub>3</sub>pNa, LpNa, HPA, HA and starch was significantly reduced in starved larvae compared to control larvae, and all extracts tested produced a decrease in a dose depended manner (Table 3). The most remarkable reductions occurred

for trypsin-, carboxypeptidase A- and B-, and  $\alpha$ -amylase-like activities, with inhibitions over 50% for some plant extracts. The only exception was a significant increase in the hydrolysis of HPA in larvae that have ingested *R. officinalis* at 500 ppm.

On the contrary, different patterns of activity were obtained for the detoxification enzymes after ingestion of plant extracts (Table 3). Esterase activity using 1-NA as substrate increased in starved and in treated larvae at the low (500 ppm) concentration tested, but decreased when the concentration of the plant extracts was 1,000 ppm. However, esterase activity using MtB as substrate increase in starved and in treated larvae at all concentrations tested. Glutathione S-transferase increased in treated larvae, but not in starved. P-450 monooxygenase activity increase only at 500 ppm for

**Table 2.** Effect of ingestion of diet treated with plant extracts (500 ppm) on protein, carbon hydrate and lipid content of *P. interpunctella* larvae

Treatment	Protein ( $\mu\text{g}/\text{larvae}$ )	Carbon hydrate ( $\mu\text{g}/\text{larvae}$ )	Lipid ( $\mu\text{g}/\text{larvae}$ )
Control	545 ± 21	35.6 ± 4.1	846 ± 25
<i>C. erythraea</i>	117 ± 10*	11.8 ± 0.7*	852 ± 28
<i>P. harmala</i>	121 ± 11*	7.8 ± 0.7*	880 ± 25
<i>C. ladanifer</i>	133 ± 12*	11.3 ± 0.7*	870 ± 25
<i>A. iva</i>	210 ± 15*	12.4 ± 1.0*	843 ± 25
<i>R. officinalis</i>	273 ± 17*	15.8 ± 0.7*	833 ± 18
Starved	187 ± 16*	2.9 ± 0.2*	570 ± 23*

Values are the mean ± standard error of protein, carbon hydrate and lipid content (n = 10). \* Significantly different from control (Dunnet test,  $p \leq 0.05$ ).

**Table 3.** Effect of ingestion of diet treated with plant extracts on midgut hydrolytic and detoxification enzyme activities of *P. interpunctella* larvae

Specific activity <sup>1</sup>	Control	<i>C. erythraea</i>			<i>P. harmala</i>			<i>C. ladanifer</i>			<i>A. iva</i>			<i>R. officinalis</i>			Starved
		500 ppm	750 ppm	1000 ppm	500 ppm	750 ppm	1000 ppm	500 ppm	750 ppm	1000 ppm	500 ppm	750 ppm	1000 ppm	500 ppm	750 ppm	1000 ppm	
TRY (BApNa)	200±5	156±4*	126±3*	113±4*	174±3*	131±4*	98±4*	160±4*	130±5*	119±4*	188±4	158±3*	65±4*	187±6	157±4*	132±4*	83±3*
CHY (SA <sub>2</sub> PPpNa)	955±8	895±8*	810±7*	708±10*	740±7*	680±9*	643±8*	850±7*	781±7*	620±6*	860±6*	782±10*	742±8*	875±7*	741±7*	711±7*	680±8*
ELA (SA <sub>2</sub> pNa)	11.6±1.1	7.6±0.5*	6.8±0.4*	6.4±0.5*	7.7±0.5*	7.0±0.6*	6.4±0.3*	8.8±0.4*	7.5±0.6*	7.4±0.3*	8.2±0.4*	8.4±0.4*	7.4±0.3*	8.5±0.5*	7.3±0.3*	7.4±0.3*	4.7±0.4*
LAP (LpNa)	248±8	200±6*	179±5*	153±4*	203±4*	183±5*	165±6*	210±6*	177±4*	161±5*	210±6*	181±7*	170±5*	203±6*	187±7*	172±4*	92±5*
CPA (HPA)	192±4	171±5*	153±6*	146±4*	150±6*	112±5*	79±3*	176±5	150±5*	134±5*	179±4	153±5*	121±6*	224±6*	196±5	178±6	114±4*
CPB (HA)	109±5	85±4*	70±4*	63±3*	80±4*	75±5*	63±3*	62±3*	51±3*	40±3*	105±4	89±3*	73±4*	98±4	73±3*	61±3*	55±3*
α-AMY (starch)	10.0±0.4	6.0±0.2*	6.0±0.6*	5.0±0.5*	6.0±0.2*	3.2±0.3*	2.0±0.2*	6.5±0.4*	5.5±0.5*	3.0±0.2*	6.0±0.2*	4.6±0.3*	4.0±0.2*	4.9±0.1*	5.2±0.4*	4.4±0.5*	7.0±0.5*
EST (1-NA)	647±8	733±9*	682±14*	600±9*	753±8*	710±8*	629±6	724±6*	662±8	600±6*	716±9*	692±6*	612±8*	703±6*	689±11*	648±8	724±6*
EST (Mtb)	23±1	46±2*	52±3*	36±2*	41±2*	55±3*	46±2*	37±2*	42±2*	40±2*	39±2*	43±2*	39±2*	41±3*	48±3*	42±3*	33±2*
GST (CDNB)	352±5	433±8*	549±8*	645±11*	443±8*	531±8*	600±8*	466±7*	583±9*	624±9*	448±7*	526±9*	594±10*	437±8*	523±10*	584±8*	339±8*
P450 (cytochrome C)	53±2	66±2*	55±2	42±2*	68±3*	62±2	46±2	63±3*	54±3	50±2	67±3*	58±3	44±2*	62±3	56±3	51±3	47±3

<sup>1</sup> Specific activities as nmoles of substrate hydrolyzed or modified/min/mg protein for: TRY (Trypsin), CHY (Chymotrypsin), ELA (Elastase) LAP (Aminopeptidase), CPA (Carboxypeptidase A), CPB (Carboxypeptidase B), α-AMY (α-amylase), EST (Esterase), GST (Glutathion-S-Transferase), and P450 (P450 Monooxygenase). Values are the mean ± SE (n = 20). \* Significantly different from control (Dunnet test,  $p \leq 0.05$ ).

some of the plant extracts tested, whereas a significant inhibition was obtained when the larvae were exposed to *C. erythraea* or *A. iva* at 1,000 ppm.

## Discussion

The present work revealed the insecticidal effect of plants commonly found in Morocco, and corresponding to different botanical families, on *P. interpunctella* larvae. Indeed, the insecticidal activity of methanol extracts from these plants was manifested in larval weight loss, cannibalism between larvae, reduction of pupation and adult emergence. The most active plant species were *P. harmala* and *C. ladanifer*, with which total inhibition of adult emergence was obtained. A strong effect (adult emergence below 2%) was also

obtained with *A. iva*, *C. erythraea* and *R. officinalis*. Similar disturbances on *P. interpunctella* larvae have been reported after ingestion of 20-hydroxyecdysone (Rharrabe *et al.*, 2009), azadirachtin (Rharrabe *et al.*, 2008), harmaline (Rharrabe *et al.*, 2007) and harmine (Bouayad *et al.*, 2012). Interestingly, cannibalistic behaviour was observed in starved larvae and in larvae exposed to plant extracts, especially *C. erythraea* and *R. officinalis*. Cannibalism is a common phenomenon in many species of insects (Polis, 1998) that increases in stressful conditions, such as when density of populations is high or food is scarce (Via, 1999). Accordingly, the insecticidal effects of plant extracts on *P. interpunctella* larvae appear to provoke increased cannibalism in comparison with larvae reared on control diet. Likewise, Rharrabe *et al.* (2007) reported that the presence of toxic plant allelochemicals in diet

causes the appearance of this behavior among larvae of *P. interpunctella*.

After plant extract ingestion, a significant reduction of reserve substances in *P. interpunctella* larvae was observed, especially in protein and carbon hydrate contents. Overall, the depletion of energy reserves could be due to a reduction in plant consumption, or to high mobilization of these primary metabolites to compensate the metabolic stress produced by the toxic effects of the plant extracts. A depression of energy reserves has also been reported after ingestion of Jatropherol-I by *B. mori* (Jing *et al.*, 2005), neem oil by *Choristoneura rosaceana* (Smirle *et al.*, 1996), *P. harmala* leaves by *T. castaneum* (Jbilou *et al.*, 2008), and azadirachtin by some insects such as *S. litura* (Huang *et al.*, 2004). Moreover, reductions in energy reserves were reported in *P. interpunctella* after ingestion of harmine (Bouayad *et al.*, 2012), harmaline (Rharrabe *et al.*, 2007), azadirachtin (Rharrabe *et al.*, 2008) and 20-hydroxyecdysone (Rharrabe *et al.*, 2009). Remarkably, we have found that lipid reserves were not affected by ingestion of diets treated with plant extracts, whereas a reduction was observed in starved larvae. In general, the lipid content of the fat body increases continuously during the larval period of holometabolous insects to be used during metamorphosis as well as to provide reserves for the new adult to support flight and embryogenesis (Arrese *et al.*, 2001; Canavoso *et al.*, 2001). Moreover, lipid metabolism provides the energy needed during extended starvation periods (Arrese & Soulages, 2010). However, our results suggest that protein and carbohydrates provided primary sources for energy in *P. interpunctella* larvae under extreme nutritive stress conditions, whereas lipid metabolism was not mobilized. Previous studies showed a reduction of lipid reserves in *P. interpunctella* after ingestion of azadirachtin and 20-hydroxyecdysone (Rharrabe *et al.*, 2008, 2009), but not after ingestion of harmine (Bouayad *et al.*, 2012). Nevertheless, we cannot discard effects on specific classes of lipids (as triacylglycerols, diacylglycerols and phospholipids), since we have only analyzed total lipid content.

In lepidopteran insects, digestion of proteases is mainly based on serine proteases (trypsin-, chymotrypsin- and elastase-like) and exopeptidases (aminopeptidase and carboxypeptidases A- and B-like) (Terra & Ferreira, 1994), whereas carbohydrate hydrolysis is due to  $\alpha$ -amylases (Franco *et al.*, 2002). All these enzymatic activities have already been characterized in *P. interpunctella* (Baker, 1986; Oppert *et al.*, 1996).

Production of digestive enzymes has been related to food intake in different insect species (Lehane *et al.*, 1996), including larvae of *P. interpunctella* (Bouayad *et al.*, 2008). However, our results showed that  $\alpha$ -amylase and all proteolytic activities were reduced in *P. interpunctella* larvae after ingestion of diet treated with plant extracts, compared with control larvae. These results are in agreement with those reported by Senthil Nathan *et al.* (2005a,b), who demonstrated that ingestion of azadirachtin causes a decrease in digestive enzymatic activities in the midgut of *S. litura* and *C. medinalis*. In the same context, Senthil Nathan (2006) revealed that the toxic effects of *Melia azedarach* in *C. medinalis* were mediated through inhibition of digestive enzyme activities. This reduction of activity may be due to several reasons. The first hypothesis is the inhibition of the activity of the hydrolytic enzymes themselves. Indeed, several studies have already shown that plant extracts inhibit *in vitro* the activity of  $\alpha$ -amylase and proteases (Duffey & Stout, 1996). However, there are also examples of plant secondary metabolites that inhibit the activity of insect digestive enzymes *in vivo*, but not *in vitro*, suggesting alternative target sites for these compounds (Liu *et al.*, 2008, 2009). The second hypothesis is that plant secondary metabolites cause cytotoxicity in epithelial cells synthesizing these digestive enzymes. In fact, severe disorganization of proteosynthetic organelles of the midgut epithelial cells of *P. interpunctella* larvae was found after harmaline or 20-hydroxyecdysone ingestion (Rharrabe *et al.*, 2007, 2009). Marked cytotoxicity in intestinal epithelium cells has also been demonstrated in other insects caused by azadirachtin (Nasiruddin & Mordue (Luntz), 1993) and seed extract of *P. harmala* (Jbilou *et al.*, 2008). A third hypothesis may be a reduction of enzymatic activity due to a decrease in food ingestion, since we have found that the activity of digestive enzymes in starved larvae was also reduced significantly. A reduction of enzyme activities has been linked with the lack of food intake in several insect species (Baker *et al.*, 1984; Hill & Orchard, 2005). A significant decrease of  $\alpha$ -amylase activity was also described after seven days of starvation in *P. interpunctella* larvae (Rharrabe *et al.*, 2007).

Induction of detoxification metabolic system plays an important role in insects host plants adaptation (Terriere, 1984). For example, Weimin *et al.* (2003) demonstrated that P450 monooxygenases in swallowtail caterpillars that feed on furanocoumarin-containing host plants display higher activities against fura-

nocoumarins than those from species that feed on plants lacking these compounds. However, detoxification enzyme systems may also be inhibited by secondary plant metabolites, such as flavonols, isothiocyanates, polyphenols, hydroxamic acids and terpenoids (Yu, 1983; Brattsten, 1987). Our results show that the activity of detoxification enzymes in *P. interpunctella* larvae was altered after ingestion of plant extracts. Thus, GST activity increased significantly in a dose-dependent manner with all treatments. These enzymes have been reported to be induced by plant allelochemicals in a number of insects (Yu, 1983; Ortego *et al.*, 1999; Caballero *et al.*, 2008) and are considered major elements conferring resistance against insecticides (Oppenoorth *et al.*, 1977; Kao & Sun, 1991; Sun *et al.*, 2001). However, the results obtained with esterase and P450 monooxygenase activities in *P. interpunctella* larvae varied depending on the concentration of the plant extract tested and the substrate used. The increase in esterase activity at low concentrations may be due to a general induction of esterase genes in response to secondary metabolites ingestion, while the inhibition with higher concentrations, when using 1-NA as substrate, could be due to a direct inhibitory effect of these compounds on specific esterase enzymes. The inhibition of esterase activity with increasing plant extract concentrations seem to be in agreement with the results reported by Jing *et al.* (2005) who observed that esterases showed lower activity after ingestion of Jatropherol-I in *B. mori*. Similar results were obtained in *C. rosaceana* after neem oil treatment (Smirle *et al.*, 1996) and in *L. decemlineata* larvae fed with potato leaves treated with limonoids (Ortego *et al.*, 1999). A wide variety of plant chemicals have been shown to induce P-450 monooxygenase activity in insects (Scott *et al.*, 1998). An induction of these enzymes was also observed in *P. interpunctella* larvae treated with some of the plant extracts tested. However, a significant inhibition of P-450 monooxygenase activity was obtained when the larvae were exposed to *C. erythraeae* or *A. iva* at 1,000 ppm. Taking together, these results indicate that *P. interpunctella* larvae respond to the ingestion of plant extracts by the induction of detoxification enzymes, but some of the plant allelochemicals may be able to inhibit some of these enzymes. An equilibrium between induction and inhibition of detoxification enzymes may then occur that results in an increase or a reduction of the enzymatic activity depending on the concentration of the plant extract ingested. In any case, the detoxification process is not capable to counteract the

insecticidal effect of these plant extracts, resulting in the death of the larvae after a few days.

In summary, this work shows the potential of plants commonly found in Morocco for the control of *P. interpunctella*. A remarkable bioinsecticidal effect was obtained with methanol extracts from *P. harmala* and *C. ladanifer* that completely abolished adult emergence. The alteration of several physiological parameters (digestive and detoxification enzymes and energy reserves) provides some insights about the mode of action of the allelochemicals in the plant extracts.

## Acknowledgements

This work was supported by the cooperation program: CNRST (580/08)/CSIC (2007MA0011) and MAEC-AECI fellowship (N° 0000536790 I.I.E Program).

## References

- Abbassi K, Kadiri AZ, Ghaout S, 2003a. Biological effects of alkaloids extracted from three plants of Moroccan arid areas on the desert locust. *Physiol Entomol* 28: 232-236.
- Abbassi K, Mergaoui L, Kadiri AZ, Stambouli A, Ghaout S, 2003b. Effets des extraits de *Peganum harmala* (Zygophyllaceae) sur le criquet pèlerin (*Schistocerca gregaria* Forskål, 1775). *Zool Baet* 13/14: 203-217.
- Ahmad S, Brattsten LB, Mullin CA, Yu SJ, 1986. Enzymes involved in the metabolism of plant allelochemicals. In: *Molecular aspects of insect-plant associations* (Brattsten LB, Ahmad S, eds). Plenum Press, NY. pp: 73-151.
- Arrese El, Soulages JL, 2010. Insect fat body: energy, metabolism, and regulation. *Annu Rev Entomol* 55: 207-25.
- Arrese EL, Canavoso LE, Jouni ZE, Pennington JE, Tsuchida K, Wells MA, 2001. Lipid storage and mobilization in insects: current status and future directions. *Insect Biochem Mol Biol* 31: 7-17.
- Azelmat K, Sayah F, Mouhib M, Ghailani N, El Garrouj D, 2005. Effects of gamma irradiation on fourth-instar *Plodia interpunctella* (Hübner) (Lepidoptera: Pyralidae). *J Stored Prod Res* 41: 423-431.
- Baker JE, 1986. Amylase/proteinase ratios in larval midguts of ten stored-product insects. *Ent Exp Appl* 40: 41-46.
- Baker JE, Woo SM, Byrd RV, 1984. Ultrastructural features of the gut of *Sitophilus granarius* (L.) (Coleoptera: Curculionidae) with notes on distribution of proteinases and amylases in crop and midgut. *Can J Zoolog* 62: 1251-1259.
- Bellakhdar J, 1997. *La pharmacopée marocaine traditionnelle-Médecine arabe et savoirs populaires*. Ibis Press, Paris.
- Berenbaum MR, Neal JJ, 1985. Synergism between myristicin and xanthotoxin, a naturally occurring plant toxicant. *J Chem Ecol* 11: 1349-1358.



- Bouayad N, Rharrabe K, Ghailani N, Sayah F, 2008. Effects of different food commodities on development of *Plodia interpunctella* (Hübner) (Lepidoptera: Pyralidae). *J Stored Prod Res* 44: 373-378.
- Bouayad N, Rharrabe K, Lamhamdi M, Ghailani NN, Sayah F, 2012. Dietary effects of harmine, a  $\beta$ -carboline alkaloid, on development, energy reserves and  $\alpha$ -amylase activity of *Plodia interpunctella* Hübner (Lepidoptera, Pyralidae). *Saudi J Biol Sci* 19: 73-80.
- Bradford MM, 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle protein-dye binding. *Anal Biochem* 72: 248-254.
- Brattsten LB, 1987. Inductibility of metabolic insecticide defences in boll budworm caterpillars. *Pestic Bioch Physiol* 27: 13-23.
- Caballero C, López-Olguín JL, Ruiz M, Ortego F, Castañera P, 2008. Antifeedant activity and effects of terpenoids on detoxication enzymes of the beet armyworm, *Spodoptera exigua* (Hübner). *Span J Agric Res* 6: 177-184.
- Canavoso LE, Jouni ZE, Karnas KJ, Pennington JE, Wells MA, 2001. Fat metabolism in insects. *Annu Rev Nutrition* 21: 23-46.
- Duffey SS, Stout MJ, 1996. Antinutritive and toxic components of plant defense against insects. *Arch Insect Biochem Physiol* 32: 3-37.
- Ellman GL, 1959. Tissue sulfhydryl groups. *Arch Biochem Biophys* 82: 70-77.
- Franco LO, Rigden DJ, Melo RF, Grossi-De-Sá FM, 2002. Plant  $\alpha$ -amylase inhibitors and their interaction with insect  $\alpha$ -amylases. Structure, function and potential for crop protection. *Eur J Biochem* 269: 397-412.
- Gomori G, 1953. Human esterases. *J Lab Clin Med* 42: 445-453.
- Habig WH, Pabst MJ, Jakoby WB, 1974. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J Biol Chem* 249: 7130-7139.
- Hill SR, Orchard I, 2005. *In vitro* analysis of the digestive enzymes amylase and  $\alpha$ -glucosidase in the midguts of *Locusta migratoria* L. in response to the myosuppressin, SchistoFLRFamide. *J Insect Physiol* 51: 1-9.
- Huang Z, Shi P, Dai J, Du J, 2004. Protein metabolism in *Spodoptera litura* (F.) is influenced by the botanical insecticide azadirachtin. *Pest Biochem Physiol* 80: 85-93.
- Isman MB, 2002. Insect antifeedants. *Pestic Outlook* 13: 152-157.
- Jbilou R, Sayah F, 2008. Effects of *Peganum harmala* (Zygophyllaceae) seed extracts on the development of *Tribolium castaneum* (Coleoptera: Tenebrionidae). *Int J Trop Insect Sci* 27: 199-209.
- Jbilou R, Amri H, Bouayad N, Ghailani N, Ennabili A, Sayah F, 2008. Insecticidal effects of extracts of seven plant species on larval development,  $\alpha$ -amylase activity and offspring production of *Tribolium castaneum* (Herbst) (Insecta: Coleoptera: Tenebrionidae). *Bioresour Technol* 99: 959-964.
- Jing L, Fang Y, Ying X, Wenxing H, Meng X, Syed M, Syed N, Fang C, 2005. Toxic impact of ingested Jatropherol-I on selected enzymatic activities and the ultrastructure of midgut cells in silkworm, *Bombyx mori* L. *J App Entomol* 129: 98-104.
- Kao CH, Sun CN, 1991. *In vitro* degradation of some organophosphorus insecticides by susceptible and resistant diamondback moth. *Pestic Bioch Physiol* 41: 132-141.
- Khambay BPS, Beddie DG, Simmonds MSJ, 2002. An insecticidal mixture of tetramethylcyclohexenedione isomers from *Kunzea ambigua* and *Kunzea baxterii*. *Phytochemistry* 59: 69-71.
- Lehane MJ, Müller HM, Crisanti A, 1996. Mechanisms controlling the synthesis and secretion of digestive enzymes in insects. In: *Biology of the insect midgut* (Lehane MJ, Billingsley PF, eds). Chapman & Hall, London, pp: 195-205.
- Liang GM, Chen W, Liu TX, 2003. Effects of three neem-based insecticides on diamondback moth (Lepidoptera: Plutellidae). *Crop Prot* 22: 333-340.
- Liu ZL, Ho SH, Goh SH, 2008. Effect of fraxinellone on growth and digestive physiology of Asian corn borer, *Ostrinia furnacalis* Guenee. *Pestic Bioch Physiol* 91: 122-127.
- Liu ZL, Ho SH, Goh SH, 2009. Modes of action of fraxinellone against the tobacco budworm, *Heliothis virescens*. *Insect Sci* 16: 147-155.
- Masters BSS, Kamin H, Gibson QH, Williams CH, 1965. Studies on the mechanism of microsomal triphosphopyridine nucleotide-cytochrome c reductase. *J Biol Chem* 240: 921-931.
- Mordue AJL, Morgan ED, Nisbet AJ, 2005. Azadirachtin, a natural product in insect control. In: *Comprehensive molecular insect science* (Gilbert LI, Iatrou K, Gill S, eds), pp: 116-135.
- Nasiruddin M, Mordue (Luntz) AJ, 1993. The effects of azadirachtin on the midgut histology of the locusts *Schistocerca gregaria* and *Locusta migratoria*. *Tissue Cell* 25: 875-884.
- Oppenoorth FJ, Smislaert HR, Welling W, Van Der Pas LJT, Hitman KT, 1977. Insensitive acetylcholinesterase, high glutathione-S-transferase, and hydrolytic activity as resistance factors in a tetrachlorvinphos-resistant strain of house fly. *Pestic Bioch Physiol* 7: 34-47.
- Oppert B, Kramer KJ, Johnson D, Upton SJ, Mcgaughey WH, 1996. Luminal proteinases from *Plodia interpunctella* and the hydrolysis of *Bacillus thuringiensis* Cry1A(c) protoxin. *Insect Biochem Mol Biol* 26: 571-583.
- Ortego F, Rodríguez B, Castañera P, 1995. Effects of neoclerodane diterpenes from *Teucrium* on feeding behavior of Colorado potato beetle larvae. *J Chem Ecol* 21: 1375-1386.
- Ortego F, Novillo C, Castañera P, 1996. Characterization and distribution of digestive proteases of the stalk corn borer, *Sesamia nonagrioides* Lef. (Lepidoptera: Noctuidae). *Arch Insect Biochem Physiol* 33: 163-180.
- Ortego F, López-Olguín J, Ruiz M, Castañera P, 1999. Effects of toxic and deterrent terpenoids on digestive protease and detoxication enzyme activities of Colorado potato beetle larvae. *Pest Biochem Physiol* 63: 76-84.
- Papachristos DP, Stamopoulos DC, 2002. Repellent, toxic and reproduction inhibitory effects of essential oil vapours on *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). *J Stored Prod Res* 38: 117-128.

- Patil BS, Vanamal J, Hallman G, 2004. Irradiation and storage influence on bioactive components and quality of early and late season 'Rio Red' grapefruit (*Citrus paradisi* Macf.). *Posthavest Biol Technol* 34: 53-64.
- Phillips WT, Throne TE, 2010. Biorational approaches to managing stored-product insects. *Annu Rev Entomol* 55: 375-397.
- Polis GA, 1998. Exploitation competition and the evolution of interference, cannibalism, and intraguild predation in age/size-structured populations. In: *Size-structured populations* (Ebenman B, Persson L, eds). Springer, Berlin, Heidelberg, pp: 185-202.
- Rharrabe K, Bakrim A, Ghailani N, Sayah F, 2007. Bioinsecticidal effect of harmaline on *Plodia interpunctella* development (Lepidoptera: Pyralidae). *Pest Biochem Physiol* 89: 137-145.
- Rharrabe K, Amri H, Bouayad N, Sayah F, 2008. Effects of azadirachtin on post-embryonic development, energy reserves and  $\alpha$ -amylase activity of *Plodia interpunctella* Hübner (Lepidoptera: Pyralidae). *J Stored Prod Res* 44: 290-294.
- Rharrabe K, Bouayad N, Sayah F, 2009. Effects of ingested 20-hydroxyecdysone on development and midgut epithelial cells of *Plodia interpunctella* (Lepidoptera, Pyralidae). *Pest Biochem Physiol* 93: 112-119.
- Roe JH, 1955. Determination of sugar in blood and spinal fluid with anthrone reagent. *J Biol Chem* 212: 335-338.
- Scott JG, Liu N, Wen ZM, 1998. Insect cytochromes P450: diversity, insecticide resistance and tolerance to plant toxins. *Comp Biochem Physiol* 121C: 147-155.
- Senthil Nathan S, 2006. Effect of *Melia azedarach* on nutritional physiology and enzyme activities of the rice leaffolder *Cnaphalocrocis melinalis* (Guenée) (Lepidoptera: Pyralidae). *Pestic Biochem Physiol* 84: 98-108.
- Senthil Nathan S, Kalaivani K, Chung PG, 2005a. The effects of azadirachtin and nucleopolyhedrovirus on midgut enzymatic profile of *Spodoptera litura* Fab. (Lepidoptera: Noctuidae). *Pest Biochem Physiol* 83: 46-57.
- Senthil Nathan S, Kalaivani K, Murugan K, Chung PG, 2005b. The toxicity and physiological effect of neem limonoids on *Cnaphalocrocis medinalis* (Guenée) the rice leaffolder. *Pest Biochem Physiol* 81: 113-122.
- Smirle MJ, Lowery DT, Zurowski CL, 1996. Influence of neem oil on detoxication enzyme activity in the oblique-banded leafroller, *Choristoneura rosaceana*. *Pest Biochem Physiol* 56: 220-230.
- Statistica, 1997. *Statistica statsoft Inc. release 5.1*. Tulsa, OK, USA.
- Sun CN, Huang SY, Hu NT, Chung WY, 2001. Glutathione S-transferase and insect resistance to insecticides. In: *Biochemical sites of insecticide action and resistance* (Ishaaya I, ed). Springer, Berlin, pp: 239-252.
- Terra WR, Ferreira C, 1994. Insect digestive enzymes-properties, compartmentalization and function. *Comp Biochem Phys B* 109: 1-62.
- Terriere LC, 1984. Induction of detoxication enzymes in insects. *Ann Rev Entomol* 29: 71-88.
- Valencia A, Bustillo AE, Ossa GE, Chrispeels MJ, 2000. Alpha-amylases of the coffee berry borer (*Hypothenemus hampei*) and their inhibition by two plant amylase inhibitors. *Insect Biochem Mol Biol* 30: 207-213.
- Van Handel E, 1965. Micro-separation of glycogen, sugars and lipids. *Anal Biochem* 11: 266-271.
- Via S, 1999. Cannibalism facilitates the use of a novel environment in the flour beetle, *Tribolium castaneum*. *Hereditas* 82: 267-275.
- Weimin Li, Mary AS, May RB, 2003. Diversification of furanocoumarin-metabolizing cytochrome P450 monooxygenases in two papilionids: specificity and substrate encounter rate. *Proc Nat Acad Sci USA* 100(Suppl 2): 14593-14598.
- White NDG, 1995. Insects, mites and insecticides in stored grain ecosystems. In: *Stored grain ecosystem* (Jayas DS, White ND, Muir WE, eds). Marcel Dekker, NY, pp: 123-168.
- Yu SJ, 1983. Induction of detoxifying enzymes by allelochemicals and host plants in the fall armyworm. *Pestic Biochem Physiol* 19: 330-336.
- Zöllner N, Krich K, 1962. The quantitative determination of lipids (micromethod) by means of sulphosphovanillin reaction common to many natural lipids (all known plasma lipids). *Z Ges Exp Med* 135: 54.