

Crystal structure of (*E*)-3-(iodomethylene)-2,3-dihydro-[1,4]oxazino-[2,3,4-*ij*]quinolin-4-ium triiodide — iodine (2:1), [C₁₂H₉INO]I₃·0.5I₂, C₁₂H₉I₅NO

Vladimir I. Batalov^I, Dimitri G. Kim^I, Alla Dikhtiarenko^{II}, Zakariae Amghouz^{III}, Ekaterina V. Bartashevich^I and Santiago García-Granda^{*,IV}


^I Department of Organic Chemistry, South Ural State University, 76, Lenin avenue, Chelyabinsk, Russia, 454080

^{II} Departamento Química Orgánica e Inorgánica, Universidad de Oviedo, 33006 Oviedo, Spain

^{III} Servicios-Científico Técnicos, Universidad de Oviedo-CINN, 33006 Oviedo, Spain

^{IV} Departamento de Química Física y Analítica, Facultad de Química, Universidad de Oviedo - CINN, C/ Julián Clavería, 8, 33006 Oviedo, (Asturias) Spain

Received January 19, 2014, accepted May 13, 2014, available online May 28, 2014, CCDC no. 1005549



Abstract

C₁₂H₉I₅NO, triclinic, *P* $\bar{1}$ (no. 2), *a* = 8.6099(5) Å, *b* = 9.7325(3) Å, *c* = 12.2098(6) Å, α = 108.972(4)°, β = 101.112(4)°, γ = 104.315(4)°, *V* = 894.9 Å³, *Z* = 2, *R*_{gt}(*F*) = 0.0373, *wR*_{ref}(*F*²) = 0.0581, *T* = 119 K.

Table 1. Data collection and handling.

Crystal:	dark purple rhombohedrals, size 0.0197×0.0669×0.0713 mm
Wavelength:	Mo <i>K</i> _α radiation (0.71073 Å)
μ :	86.82 cm ⁻¹
Diffractionmeter, scan mode:	Xcalibur, Ruby, Gemini, ω
$2\theta_{\max}$:	62.94°
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} :	15223, 5377
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ (<i>I</i> _{obs}), 4365
<i>N</i> (<i>param</i>) _{refined} :	172
Programs:	CrysAlis PRO [11], SIR92 [12], SHELX [13], DIAMOND [14], WinGX [15], enCIFer [16]

Source of material

8-(propargyloxy)quinoline was obtained by the reaction of 8-quinolinol with propargyl bromide in acetone/K₂CO₃ with reflux. A solution of 8-(propargyloxy)quinoline (0.145 g, 1 mmol) in 5 mL of dichloromethane was mixed with 0.762 g of iodine (3 mmol) in 15 mL of dichloromethane. Dark purple single crystals

for the X-ray diffraction study were obtained after keeping the mixture at room temperature in closed flask for 48 hours, yield: 0.447 g (55 %).

Experimental details

Position of the H atoms were calculated based on geometric criteria (C–H = 0.97 Å, 0.93 Å and 0.85 Å for methylene, aromatic and methine atoms, respectively) than have been placed in their calculated position and refined isotropically using a rider model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Discussion

Tricyclic quinoline based systems with fused N-1 and C-8 centers are known as important broad-spectrum antibiotics [1]. The title compound was synthesized in the context of obtaining new heteroaromatic systems by heterocyclization with an excess of iodine [2]. The presented crystal structure can be an interesting object with I...I non-covalent interactions involving terminal triiodide atoms as donors of electrons and iodine in I₂ molecule as electrophilic species. This type of interactions refers to halogen bonding [3–10]. The asymmetric unit of the title compound consists of quinolinium derivative cation, triiodide anion and half of molecular iodine. The atoms N1 and C1–C9, defining the quinolinium part of rings, are essentially planar. The C–C bonds lengths range from 1.366(6) to 1.416(6) Å; the C–N bond distances range from 1.353(5) to 1.390(5) Å. For the exocyclic adduct, formed as a result of cyclization reaction, C–O bond distances range from 1.362(5) to 1.435(5) Å, C11–N1 bond length is 1.451(5) Å and C10–C11 bond length is 1.488(6) Å, that have single bond character. The resulting six-membered ring is partially planar, where the C10 atom of the methylene group is slightly out of plane. The crystal structure confirms the closure of the ring between O1 and N1 atoms, where formed C11–N1 bond is covalent (1.451(5) Å). The C11–C12 bond, that is associated with six-member ring, has double bond character with a bond distance of 1.324(6) Å. The C12–I1 bond (2.082(4) Å) significantly deviated from mean plane of quinolinium moiety, the C11–C12–I1 angle is 123.6(3)°. Crystal packing is formed by multiple interactions of [C₁₂H₉INO]⁺ cations, triiodide and iodine molecules. The organic cations pack could be represented as the dimeric association established by a π – π interaction. The fused rings are situated in opposite directions and lie nearly parallel to one another; the closest centroid-centroid distance is 3.51(6) Å. The dimers are surrounded by triiodide anions and iodine molecules. The geometry of triiodide is almost linear, I2–I3 and I3–I4 distances are

* Correspondence author (e-mail: sgg@uniovi.es)

2.8752(5) and 2.9572(5) Å, respectively, and I2-I3-I4 angle is 179.028(12)°. Inversion center in I₂ molecule is situated in the middle of the I5-I5 bond (2.7467(6) Å). Each of I5 atoms interacts via halogen bonding with the terminal I4 atom of the triiodide anion (I5...I4 distance is 3.444(1) Å), to give formally a [I₈]²⁻ unit [9].

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(1)	2i	-0.0079	0.2562	0.3738	0.025

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
C(1)	2i	0.0459(6)	0.2161(5)	0.3184(4)	0.020(3)	0.027(2)	0.019(2)	0.009(2)	0.007(2)	0.013(2)
C(2)	2i	0.1381(6)	0.1236(5)	0.3394(5)	0.030(3)	0.032(3)	0.028(3)	0.013(2)	0.011(2)	0.022(2)
C(3)	2i	0.2190(6)	0.0656(5)	0.2588(5)	0.022(3)	0.021(2)	0.036(3)	0.009(2)	0.003(2)	0.015(2)
C(4)	2i	0.2135(6)	0.1014(5)	0.1557(4)	0.018(2)	0.015(2)	0.027(3)	0.007(2)	0.009(2)	0.011(2)
C(5)	2i	0.2920(6)	0.0438(5)	0.0692(4)	0.018(3)	0.021(2)	0.033(3)	0.009(2)	0.011(2)	0.013(2)
C(6)	2i	0.2783(6)	0.0810(5)	-0.0310(4)	0.022(3)	0.014(2)	0.029(3)	0.005(2)	0.014(2)	0.005(2)
C(7)	2i	0.1893(6)	0.1793(5)	-0.0470(4)	0.023(3)	0.019(2)	0.017(2)	0.003(2)	0.008(2)	0.007(2)
C(8)	2i	0.1105(6)	0.2370(5)	0.0351(4)	0.014(2)	0.013(2)	0.018(2)	0.002(2)	0.004(2)	0.005(2)
C(9)	2i	0.1201(5)	0.1969(5)	0.1367(4)	0.013(2)	0.012(2)	0.017(2)	0.002(2)	0.006(2)	0.005(2)
C(10)	2i	-0.0006(6)	0.4276(5)	0.1239(4)	0.026(3)	0.018(2)	0.019(2)	0.011(2)	0.011(2)	0.008(2)
C(11)	2i	-0.0696(6)	0.3379(5)	0.1921(4)	0.021(2)	0.011(2)	0.013(2)	0.007(2)	0.005(2)	0.005(2)
C(12)	2i	-0.2053(6)	0.3329(5)	0.2286(4)	0.024(3)	0.013(2)	0.019(2)	0.010(2)	0.007(2)	0.007(2)
I(1)	2i	-0.35420(4)	0.46650(3)	0.20579(3)	0.0209(2)	0.0198(2)	0.0182(2)	0.0101(1)	0.0074(1)	0.0059(1)
I(2)	2i	0.41583(4)	0.70685(3)	0.15375(3)	0.0186(2)	0.0183(1)	0.0184(2)	0.0054(1)	0.0041(1)	0.0067(1)
I(3)	2i	0.64511(4)	0.96417(3)	0.36655(3)	0.0210(2)	0.0183(1)	0.0162(2)	0.0081(1)	0.0069(1)	0.0075(1)
I(4)	2i	0.88032(4)	1.23235(3)	0.58326(3)	0.0320(2)	0.0204(2)	0.0149(2)	0.0054(1)	0.0056(1)	0.0032(1)
I(5)	2i	0.39552(4)	0.57862(3)	0.47375(3)	0.0346(2)	0.0206(2)	0.0191(2)	0.0094(1)	0.0072(1)	0.0079(1)
N(1)	2i	0.0338(5)	0.2485(4)	0.2179(3)	0.016(2)	0.012(2)	0.016(2)	0.003(2)	0.004(2)	0.004(1)
O(1)	2i	0.0211(4)	0.3289(3)	0.0158(3)	0.026(2)	0.023(2)	0.016(2)	0.013(2)	0.010(1)	0.011(1)

Acknowledgments. We thank for financial support from the Spanish Ministerio de Economía y Competitividad under projects MAT2010-15094, grant Técnicos de Infraestructuras Científico-Tecnológicas PTA2011-4903-I to Z. A. and FEDER. A. D. Also we thank to Spanish Ministerio de Educación, Cultura y Deporte for FPU grant AP2008-03942. This work was supported by Russian Foundation for Basic Research, grant 14-03-00961a. We also like to thank Joaquín García Álvarez for the help in the realization of the synthesis.

References

- Appelbaum, P. C.; Hunter, P. A.: The fluoroquinolone antibacterials: past, present and future perspectives. *Int. J. Antimicrob. Ag.* **16** (2000) 5-15.
- Onysko, M. Y.; Lendel, V. G.: Haloheterocyclization of 2-allyl-(propargyl)oxyquinoline-3-carbaldehydes. *Chem. Heterocycl. Compd.* **43** (2007) 1020-1023.
- Desiraju, G. R.; Ho, P. S.; Kloo, L.; Legon, A. C.; Marquardt, R.; Metrangolo, P.; Politzer, P. A.; Resnati, G.; Rissanen, K.: Definition of the halogen bond. *Pure Appl. Chem.* **85** (2013) 1711-1713.
- Kim, D. G.: Halocyclization of 8-allylthioquinoline. *Chem. Heterocycl. Compd.* **33** (1997) 1133-1135.
- Paira, R.; Paira, P.; Maity, A.; Mondal, S.; Hazra, A.; Sahu, K. B.; Naskar, S.; Saha, P.; Banerjee, M.; Mondal, N. B.: Amberlite IRA 402(OH): an efficient mediator for the exclusive synthesis of fused tricyclic oxaza quinolinium salts. *Tetrahedron Lett.* **51** (2010) 3200-3204.
- Kalita, D.; Baruah, J. B.: Rearrangement of 2-Bromo-*N*-quinoline-8-yl-acetamide Leading to New Heterocycle. *J. Heterocyclic Chem.* **47** (2010) 459-462.
- van Megen, M.; Reiss, G. J.: I₆²⁻ Anion Composed of Two Asymmetric Triiodide Moieties: A Competition between Halogen and Hydrogen Bond. *Inorganics* **1** (2013) 3-13 and references cited therein.
- Slepukhin, P. A.; Batalov, V. I.; Kim, D. G.; Charushin, V. N.: A single crystal X-ray study of the products of halogen mercury cyclization of 8-allylthioquinoline. *J. Struct. Chem.* **52** (2012) 145-150.
- Svensson, P. H.; Kloo, L.: Synthesis, Structure, and Bonding in Polyiodide and Metal Iodide-Iodine Systems. *Chem. Rev.* **103** (2003) 1649-1684.
- Metrangolo, P.; Resnati, G.; Pilati, T.; Liantonio, R.; Meyer, F.: Engineering functional materials by halogen bonding. *J. Polym. Sci. Part A: Polym. Chem.* **45** (2007), 1-15.
- Agilent Technologies: CrysAlis PRO Software system, version 1.171.35.15, Agilent Technologies UK Ltd, Oxford, UK 2011.
- Altomare, A.; Cascarano, G.; Giacovazzo, C.; Guagliardi, A.; Burla, M. C.; Polidori, G.; Camalli, M.: SIR92 - a program for automatic solution of crystal structures by direct methods. *J. Appl. Cryst.* **27** (1994) 435-436.
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112-122.
- Brandenburg, K.: DIAMOND. Visual Crystal Structure Information System. Version 3.2i. Crystal Impact, Bonn, Germany 2012.
- Farrugia, L. J.: WinGX suite for small-molecule single-crystal crystallography. *J. Appl. Cryst.* **32** (1999) 837-838.
- Allen, F. H.; Johnson, O.; Shields, G. P.; Smith, B. R.; Towler, M.: CIF applications. XV. enCIFer: a program for viewing, editing and visualizing CIFs. *J. Appl. Cryst.* **37** (2004) 335-338.