

Insights into the Origin of Metazoan Filopodia and Microvilli

Arnau Sebé-Pedrós,^{†,1,2} Pawel Burkhardt,^{†,3} Núria Sánchez-Pons,¹ Stephen R. Fairclough,³ B. Franz Lang,⁴ Nicole King,^{*,3} and Iñaki Ruiz-Trillo^{*,1,2,5}

¹Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Barcelona, Catalonia, Spain

²Departament de Genètica, Universitat de Barcelona, Barcelona, Spain

³Department of Molecular and Cell Biology, University of California, Berkeley

⁴Département de Biochimie, Centre Robert-Cedergren, Université de Montréal, Montréal, Québec, Canada

⁵Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

[†]These authors contributed equally to this work.

*Corresponding authors: E-mail: nking@berkeley.edu; inaki.ruiz@multicellgenome.org.

Associate editor: David Irwin

Abstract

Filopodia are fine actin-based cellular projections used for both environmental sensing and cell motility, and they are essential organelles for metazoan cells. In this study, we reconstruct the origin of metazoan filopodia and microvilli. We first report on the evolutionary assembly of the filopodial molecular toolkit and show that homologs of many metazoan filopodial components, including fascin and myosin X, were already present in the unicellular or colonial progenitors of metazoans. Furthermore, we find that the actin crosslinking protein fascin localizes to filopodia-like structures and microvilli in the choanoflagellate *Salpingoeca rosetta*. In addition, homologs of filopodial genes in the holozoan *Capsaspora owczarzaki* are upregulated in filopodia-bearing cells relative to those that lack them. Therefore, our findings suggest that proteins essential for metazoan filopodia and microvilli are functionally conserved in unicellular and colonial holozoans and that the last common ancestor of metazoans bore a complex and specific filopodial machinery.

Key words: fascin, choanoflagellate, *Capsaspora*, Cdc42, formin evolution, gelsolin evolution, filopodia, pseudopodia.

Introduction

A dynamic cytoskeletal and membrane system is a hallmark of the eukaryotic cell. It allows cells to change cell shape to carry out motility, phagocytosis, and other key functions (Fletcher and Mullins 2010). Cell motility, in particular, is a common feature among eukaryotes that often requires specialized organelles. There are two main classes of cellular structures responsible for cell motility in eukaryotes: tubulin-based cilia and flagella, conspicuous in eukaryotes as diverse as choanoflagellates, ciliates and dinoflagellates, and actin-based filopodia and lamellipodia, which allow cells to crawl along surfaces through amoeboid movement (Soldati and Meissner 2004).

Filopodia are finger-like structures based upon 10–30 parallel bundled actin filaments whose growing/barbed ends orient toward the filopodial tip. Here, many proteins accumulate and form the so-called tip complex, which controls actin monomer addition to filament ends (Small et al. 2002; Bohil et al. 2006; Faix and Rottner 2006; Gupton and Gertler 2007; Mattila and Lappalainen 2008; Lundquist 2009; Mellor 2010; Nambiar et al. 2010). In metazoans, filopodia function as sensory and exploratory organelles and typically display active protrusive and retractile motility (Yang and Svitkina 2011). Filopodia also contribute to cell adhesion (Schäfer et al. 2010) and mediate many essential, metazoan-specific phenomena,

including growth cone guidance, wound-healing, embryonic development, and angiogenesis and they serve as precursors for dendritic spines in neurons (Magie et al. 2007; Mattila and Lappalainen 2008; Mellor 2010). In contrast, the other major type of actin-based cell protrusion (Small et al. 2002; Faix and Rottner 2006), the lamellipodium, is a flat sheet-like structure based upon a branched network of actin filaments (Mattila and Lappalainen 2008; Small et al. 2008; Vallotton and Small 2009). Filopodia often emerge from lamellipodial sheets and both structures share some molecular components, although a characteristic filopodial molecular architecture has been described (Mattila and Lappalainen 2008).

Filopodia-like structures (often also called pseudopodia) are known in cells from diverse other eukaryotic lineages. These cellular protrusions have historically been defined as “filopodia” based primarily on morphological characteristics (i.e., being long slender cellular protrusions) and the presence of actin filaments (Yang and Svitkina 2011). Although a hallmark of metazoan filopodia is their dynamic nature (i.e., protrusion and retraction), little is known about the dynamic properties of filopodia-like structures in non-metazoans (Adl et al. 2012; Cavalier-Smith 2013). Among bikonts, filopodia-like structures are found in excavates (e.g., *Naegleria gruberi*; Preston and King 2005), stramenopiles (Pawlowski 2008), and rhizarians (Cavalier-Smith 2003; Pawlowski 2008; Ota

© The Author 2013. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Open Access




FIG. 1. Phylogenetic distribution of diverse proteins associated with filopodia and related actin-based cellular protrusions. A black dot indicates the presence of clear homologs, whereas absence of a dot indicates that a homolog was not detected in that taxon. ¹Based on Boureux et al. (2007). ²Only present in the excavates *Naegleria gruberi* and *Trichomonas vaginalis*. ³Plants have a villin-like protein that, despite having the same domain architecture as metazoans, is not phylogenetically related with holozoan villin (supplementary fig. S5, Supplementary Material online). ⁴Rac1 RhoGTPases are key filopodia-inducers in amoebozoans, but not in metazoans (Dumontier et al. 2000).

Actin-Remodeling Proteins

Most actin-remodeling proteins are widespread among eukaryotes (fig. 1). This is the case with the Arp2/3 complex, a major actin-remodeling factor (Pollard 2007), which convergently elongates actin filaments from the cortical actin meshwork and is regulated, in metazoans, by WASP and cortactin

(Weaver et al. 2003). The seven subunits of this complex are present in almost all eukaryotes, suggesting this is an ancient protein network.

Formins are also involved in actin filament formation, but instead of bundling an existing actin meshwork, formins nucleate actin filaments de novo. Our phylogenetic analyses

show that Diaphanous-related formins (DRFs) (Chalkia et al. 2008), those with the domain structure GBD-FH3-FH2-DAD, are present in all Amorphea investigated as well as in Excavata (supplementary figs. S3 and S4, Supplementary Material online). Interestingly, Ena/VASP, a multifaceted actin-regulatory protein with essential roles in filopodia formation and elongation that is essential for DRF-based de novo actin nucleation (Schirenbeck et al. 2006), is exclusive to Amorphea and appears to have been secondarily lost in Fungi (fig. 1). Therefore, although DRF-like formins are present in some Excavata, it is unclear whether actin nucleation based on formins is truly an ancestral mechanism in eukaryotes, as they lack Ena/VASP.

Other proteins involved in actin remodeling include cofilin, which has depolymerizing activity, profilin, which sequesters actin monomers (Revenu et al. 2004), and several proteins, such as Eps8, Twinfilin, Villin, and other Gelsolin family proteins (supplementary fig. S5, Supplementary Material online), which are involved in capping (Mattila and Lappalainen 2008) (i.e., they stabilize filament ends by binding them and inhibiting actin monomer association or dissociation). Most of these other actin regulators are widespread among eukaryotes or Amorphea (fig. 1), except for the capping protein Eps8, which is specific to metazoans and choanoflagellates. Eps8 is a direct binding partner of ERM proteins (discussed earlier) and, together, they stimulate formation of microvilli (Zwaenepoel et al. 2012). Villin seems to be restricted to holozoans as well, although plants have a villin-like protein with similar domain architecture but with uncertain affinity to other known gelsolin-domain proteins (supplementary fig. S5, Supplementary Material online). Villin is a multi-faceted actin-remodeling protein that, in metazoans, is usually associated with microvillar formation, particularly in intestinal cells (Silacci et al. 2004; Khurana and George 2008; Nambiar et al. 2010).

In sum, we find that some actin-remodeling proteins have an ancient eukaryotic origin whereas others evolved later in evolution, either at the stem of Holozoa (villin and Eps8) or at the stem of Amorphea (Ena/VASP).

Nucleation Promoting Factors

NPFs activate the Arp2/3 complex (Goley and Welch 2006). Some important NPFs in metazoan filopodia are WASP (and the associated WASP-interacting protein [WIP]) (Veltman and Insall 2010; Kollmar et al. 2012) and Cortactin (Weaver et al. 2001; Goley and Welch 2006). In metazoans, WASP is a direct target of Cdc42 RhoGTPase, mediating the activation of the Arp2/3 complex (Takenawa and Miki 2001; Antón et al. 2007; Faix et al. 2009; Mellor 2010), which is also activated by Cortactin (Kinley et al. 2003; Ren et al. 2009), while WIP is responsible for inactivating WASP (Antón et al. 2007).

Our data show that WASP is widespread among eukaryotes (fig. 1; supplementary fig. S6, Supplementary Material online), whereas WIP is restricted to Amorphea and Cortactin to holozoans (fig. 1). It is worth mentioning that a major regulator of Cortactin, c-Src tyrosine kinase (Weaver et al. 2001, 2003), is a component of the metazoan integrin

adhesome that also originated in the holozoan lineage (Sebé-Pedrós et al. 2010; Suga et al. 2012).

Rho GTPases and Other Signaling Proteins

RhoGTPases are key regulators of actin dynamics and play important roles as major switches in filopodia formation (Ridley 2006; Ladwein and Rottner 2008; Faix et al. 2009). They act through two types of actin nucleators: WASP and DRFs (Ridley 2006). Cdc42 RhoGTPase, thought to be exclusive to opisthokonts (Boueux et al. 2007), appears to be the primary filopodia-inducing RhoGTPase in metazoans (Ridley 2006). In Dictyostelium, which has filopodia but lacks Cdc42, Rac1 GTPases induce filopodia (Vlahou and Rivero 2006), whereas in plants, the Rac1-related Rop GTPases can also stimulate actin polymerization (Boueux et al. 2007). We find that Rac1-type RhoGTPases are ancestral within Amorphea, and, interestingly, that Cdc42 is not specific to opisthokonts (Boueux et al. 2007), as it is also present in the apusozoan *T. trahens*, sister group of opisthokonts (fig. 1; supplementary fig. S7, Supplementary Material online). RIF and WRCH1, which induce filopodia in some metazoan cell types (Faix et al. 2009), are specific to Metazoa (fig. 1).

Beside RhoGTPases, the signaling proteins abLIM, Abi, Vav-1, and IRSp53-like also regulate filopodia formation. These proteins evolved during different episodes in eukaryotic evolutionary history. IRSp53-like proteins are ancestral Amorphea proteins (supplementary fig. S9, Supplementary Material online), Vav-1 is specific to holozoans (fig. 1), Abi specific to metazoans and choanoflagellates (fig. 1) and abLIM specific to metazoans (supplementary fig. S8, Supplementary Material online).

Motor Proteins: Myosins

Myosins are a large protein family present in all eukaryotes and are essential for cell trafficking along actin filaments (Richards and Cavalier-Smith 2005; Odronitz and Kollmar 2007). The MyTH4-FERM domain myosins (named according to their protein domain composition) have special relevance for filopodia function and formation. For example, the metazoan myosin X (Tuxworth et al. 2001; Berg and Cheney 2002; Nagy et al. 2008; Nambiar et al. 2010) is essential for filopodia formation (Bohil et al. 2006) and for the transport of proteins such as integrins (Breshears et al. 2010) or Ena/VASP protein to the filopodial tip (Zhang et al. 2004; Sousa and Cheney 2005). Two other MyTH4-FERM myosins, myosin VII and myosin XV, are also important for filopodia function (Breshears et al. 2010). Our analysis shows that these three myosins (X, VII, and XV) emerged at the origin of Holozoa (fig. 1).

Filopodia-like Structures in Unicellular and Colonial Relatives of Metazoans

The richness of the filopodial toolkit in unicellular and colonial holozoans prompted us to investigate the presence, abundance, and distribution of filopodia-like structures in these close relatives of metazoans. We stained for polymerized actin (using phalloidin) and tubulin (using anti-tubulin




Fig. 2. Filopodia-like structures in close relatives of metazoans. (A) *Capsaspora owczarzaki* filopodiated cells bear multiple long bundles of actin microfilaments, as revealed by staining with phalloidin (green). The cell periphery is revealed by staining with antibodies against beta-tubulin (red). SEM shows the presence of multiple long filopodia-like structures in *C. owczarzaki* filopodiated cells (B), but not in nonfilopodiated, cystic cells (C). (D) In *Salpingoeca rosetta*, attached cells bear actin microfilaments in the apical collar of microvilli and in basally positioned long cellular protrusions that resemble filopodia. *Salpingoeca rosetta* cells were stained with phalloidin (green) and antibodies against beta-tubulin (red). (E) TEM of thin sections through a choanoflagellate shows the presence of basally positioned cellular processes (indicated with black rectangle), shown in higher magnification in (F). Scale bars (A–E: 1 μ m, F: 200 nm).

antibodies) in *C. owczarzaki* and *S. rosetta*. In *C. owczarzaki*, multiple 1–20 μ m long bundles of actin microfilaments can be found in filopodiated stage cells (fig. 2A). Scanning electron microscopy (SEM) confirms the presence of multiple long filopodia-like structures in this cell stage of *C. owczarzaki* (fig. 2B), in contrast with the naked cystic cell stage (fig. 2C). In the choanoflagellate *S. rosetta* actin microfilaments were detected in two distinct sites: in the apical collar of actin-filled microvilli and in basally positioned 1–10 μ m long cellular protrusions that resemble filopodia (fig. 2D) (Leadbeater 1979; Dayel et al. 2011). Supporting the inference that the basal actin microfilaments are associated with filopodia, transmission electron microscopy (TEM) of thin sections through *S. rosetta* cells shows the presence of multiple basally positioned cellular processes (fig. 2E and F, black rectangle). Thus, our data show the presence of multiple long, actin-filled cellular projections that resemble filopodia in two close relatives of metazoans.

Fascin Localizes to Filopodia-Like Structures and Actin-Filled Microvilli in *S. rosetta*

In metazoans, fascin functions as a filament-bundling protein that localizes to filopodia, and in some cell types also to microvilli (DeRosier and Tilney 2000; Kureishy et al. 2002; Tilney et al. 2004). Given that we identified clear fascin homologs in *S. rosetta*, *M. brevicollis*, and *C. owczarzaki* (fig. 1; supplementary fig. S2B, Supplementary Material online) (see above), we next investigated whether fascin homologs in choanoflagellates might function in filopodia-like structures and microvilli as they do in the metazoans. Western blot analysis showed that a commercially available fascin antibody, which was originally raised against the human protein fascin, recognizes a single band of the expected size when used to probe *S. rosetta* lysate (fig. 3D). The *S. rosetta* genome encodes two fascin paralogs with predicted molecular weights of 54.3 and 54.6 kDa. Thus, we performed immunolocalization studies of fascin in *S. rosetta*. Interestingly, fascin localizes to the




Fig. 4. Expression of filopodial and related genes in unicellular holozoans. (A) Log₂-fold expression (see Materials and Methods) of *Capsaspora owczarzaki* filopodial genes between filopodiated and cystic stages. (B) Log₂-fold expression of *Salpingoeca rosetta* filopodial genes between attached and colonial stages. Red lines highlight 2-fold expression differences. For clarity, negative values indicate overexpression in one stage compared with the other, and vice versa.

Our expression data suggest that this complex complement is, indeed, functionally conserved in unicellular holozoans, as most filopodial genes are overexpressed in *C. owczarzaki* filopodial cells. Moreover the main actin-bundling protein fascin localizes to filopodia and microvilli in *S. rosetta* and in many metazoan cell types, suggesting that fascin functioned as an actin-crosslinking protein in the filopodia and microvilli of the Urmetazoa.

In the common ancestor of choanoflagellates and metazoans, the complexity of the filopodial apparatus was further expanded as filopodial specialization in the form of the microvillar collar evolved. Our findings support the hypothesis that microvilli and filopodia are related, with microvilli reusing part of the filopodial toolkit, while also depending on the function of proteins like ERM and Eps8, which are restricted to choanoflagellates and metazoans. Finally, in




FIG. 5. Evolution of the metazoan filopodial toolkit. Schematic representation of the metazoan filopodial toolkit (left), with colors indicating the inferred evolutionary origin of each gene (white, eukaryotes; blue, Amorphea; yellow, opisthokonts + apusozoans; orange, holozoans; green, choanoflagellates + metazoans; red, metazoans). The cladogram (right) represents gains and losses and the presence of filopodia and microvilli in different groups. Dashed filopodia in other eukaryotes represent the presence of filopodia only in some groups (see main text).

metazoans the toolkit further expanded, particularly with the evolution of new RhoGTPases that are known to act instead of Cdc42 in specific cell-types.

Our analyses suggest that other non-metazoan eukaryotic lineages evolved their specific filopodial toolkits based on an ancient molecular machinery that included core actin linking proteins (profilin, twinfilin, fimbrin, cofilin, and others) and an ancestral filopodia formation mechanism (Arp2/3-WASP-DRFs). We hypothesize this mechanism was deployed under the control of different signaling triggers (e.g., Cdc42 in metazoans and Rac1 in amoebozoans) together with different specific co-factors. The recently reported NET superfamily (Deeks et al. 2012), a plant specific membrane-actin cytoskeleton adaptor protein, exemplifies this idea of convergence (in this case, to mediate the interaction between the membrane and the actin cytoskeleton). Therefore, we infer that metazoan-type filopodia originated at the stem of Holozoa, built upon many ancient proteins and acting with some more recently evolved (i.e., holozoan-specific) molecular components.

In any case, the study of the cell biology and genome content of filopodiated chlorarachniophytes (Rhizaria) (Ota and Vaultot 2012), labyrinthulomycetes, other filopodiated stramenopiles (Tsui et al. 2009; Gómez et al. 2011),

and filopodiated Excavata, such as *N. gruberi* (Preston and King 2005), will be crucial to gain new insights into the question of whether there is a common, functionally homologous, molecular toolkit underlying all eukaryotic filopodia.

Conclusions

Our study reconstructs in detail the evolutionary assembly of the metazoan filopodial and microvillar molecular toolkit. We find that many components of the metazoan filopodial toolkit are paneukaryotic, whereas other elements evolved in stem Amorphea. Finally, a number of metazoan filopodial and microvillar components evolved in stem holozoans. Moreover, some of the components of metazoan microvilli appeared concomitantly with the evolution of the feeding collar at the stem of choanoflagellates and metazoans and likely played a crucial role in the feeding mode of the Urmetazoan, as it does in choanoflagellates.

We further demonstrate that fascin is expressed both in filopodia-like structures and microvilli in the choanoflagellate *S. rosetta* and that filopodial genes are differentially upregulated in *C. owczarzaki*'s filopodial cell-stage. This suggests functional conservation of the metazoan filopodial toolkit in both choanoflagellates and *C. owczarzaki*. Given the

predicted homology of filopodia between metazoans and their unicellular relatives, we hypothesize that the existence of a complex filopodial toolkit in the ancestors of metazoans may have contributed to the origin of metazoans, by being co-opted to function in cell–cell and cell-matrix adhesion functions within a multicellular context.

Materials and Methods

Gene Searches and Phylogenetic Analysis

A primary search was performed using the basic local alignment sequence tool (BLAST: BlastP and TblastN) using *Homo sapiens* and *C. owczarzaki* proteins as queries against Protein and Genome databases with the default BLAST parameters and an *e*-value threshold of e^{-5} at the National Center for Biotechnology Information (NCBI), the Joint Genome Institute (JGI), the Broad Institute (for *S. rosetta* and *S. punctatus*), as well as the *Amphimedon queenslandica* genome database (www.metazome.net; last accessed June 2013). For some proteins, we also performed Hmmer searches using HMMER3.0b2 (Eddy 1998) to confirm that we were retrieving all orthologs.

We performed searches using the following taxon sampling: seven metazoans (*H. sapiens*, *Drosophila melanogaster*, *Daphnia pulex*, *Capitella teleta*, *Lottia gigantea*, *Nematostella vectensis*, and *A. queenslandica*), two choanoflagellates (*M. brevicollis* and *S. rosetta*), one filasterean (*C. owczarzaki*), four fungi (*Laccaria bicolor*, *Saccharomyces cerevisiae*, *A. macrogynus*, and *S. punctatus*), one apusozoan (*T. trahens*), two amoebozoans (*Acanthamoeba castellanii* and *D. discoideum*), three viridiplantae (*Arabidopsis thaliana*, *Ostreococcus taurii*, and *Chlamydomonas reinhardtii*), two excavates (*Trichomonas vaginalis* and *N. gruberi*), and three chromalveolates (*Thalassiosira pseudonana*, *Tetrahymena thermophila*, and *Toxoplasma gondii*).

Alignments were constructed using the MAFFT v.6 online server (Kato et al. 2002) and then manually inspected and edited using Geneious software. Only those species and those positions that were unambiguously aligned were included in the final analyses. Maximum likelihood (ML) phylogenetic trees were estimated by RaxML (Stamatakis 2006) using the PROTGAMMAWAG + Γ + I model, which uses the WAG amino acid exchangeabilities and accounts for among-site rate variation with a four category discrete gamma approximation and a proportion of invariable sites. Statistical support for bipartitions was estimated by performing 100-bootstrap replicates using RaxML with the same model. Bayesian analyses were performed with MrBayes3.2 (Huelsenbeck and Ronquist 2001), using the LG + Γ + I model of evolution, with four chains, a subsampling frequency of 100 and two parallel runs. Runs were stopped when the average standard deviation of split frequencies of the two parallel runs was <0.01 , usually at around 18,000,000 generations. The two LnL graphs were checked and an appropriate burn-in length established. Bayesian posterior probabilities were used to assess the confidence values of each bipartition.

Cell Culture and Microscopy

Salpingoeca rosetta cultures enriched for attached cells were maintained in artificial sea water and split 1:5 every 3 days. For immunofluorescence, the cells were grown to a density of 10^6 cells/mL and carefully scraped off from the surface of the culture flasks. Cells were then pelleted by spinning for 10 min at $4,000 \times g$ and resuspended in a small volume of artificial seawater. Approximately 0.4 mL of the cells were applied to poly-L-lysine coated coverslips, left to attach for 30 min.

Capsaspora owczarzaki cells were grown on coverslips in ATCC medium 1034 (modified PYNFH medium) for two days and directly fixed.

For both *S. rosetta* and *C. owczarzaki*, cells were fixed for 5 min with 6% acetone and for 15 min with 4% formaldehyde. The coverslips were washed gently four times with 100 mM Pipes at pH 6.9, 1 mM EGTA, and 0.1 mM $MgSO_4$ (PEM), incubated for 30 min in blocking solution (PEM + : 1% BSA, 0.3% Triton X-100), 1 h in primary antibodies solution (in PEM +), and after further washes (PEM +), 1 h in the dark with fluorescent secondary antibodies (1:100 in PEM + , Alexa Fluor 488 goat anti-mouse, and Alexa Fluor 568 goat anti-rabbit; Invitrogen) and washed again four times (PEM). To visualize F-actin coverslips were incubated for 15 min in the dark with rhodamine phalloidin (6 U/ml in PEM; Molecular Probes). After 3 washes (PEM), coverslips were mounted onto slides with Fluorescent Mounting Media (4 μ L; Prolong Gold Antifade, Invitrogen). The following primary antibodies have been used: mouse monoclonal antibody against β -tubulin (E7, 1:400; Developmental Studies Hybridoma Bank); mouse monoclonal antibody against Fascin (ab78487, 1:100; Abcam). Images were taken with a 100 \times oil immersion objective on an inverted Leica microscope.

For SEM, *C. owczarzaki* cells were fixed for 1 h with 2.5% glutaraldehyde and 1 h with 1% osmium tetroxide, followed by sequential dehydration with ethanol. Next, drying critical point was performed and samples were coated with carbon. Samples were observed in a Hitachi S-4100 microscope.

For TEM, choanoflagellate cells were concentrated by gentle centrifugation, loaded into 100- μ m deep specimen carriers and high pressure frozen in a Bal-Tec HPM 010 high pressure freezer (Bal-Tec AG, Liechtenstein). Freeze-substitution was performed over 2 h by the SQFS method of McDonald and Webb (2011), then infiltrated with Eponate 12 resin and polymerized in a Pelco Biowave research microwave oven (Ted Pella, Inc., Redding, CA) over a period of 2 h. Sections were cut at 70-nm thickness, poststained with uranyl acetate and lead citrate, and viewed in a Tecnai 12 transmission EM (FEI Inc., Hillsboro, OR) operating at 120 kV. Images were recorded on a Gatan Ultrascan 1000 CCD camera.

Gene Expression Analyses

Total RNA from *C. owczarzaki*'s described life stages was extracted using Trizol reagent. Libraries were sequenced with 76 base pair reads on an Illumina HiSeq instrument (Illumina). mRNA was isolated from *S. rosetta* cultures enriched for

colonial and attached cells using the RNAeasy (Qiagen) and Oligotex (Qiagen) kits. Libraries were sequenced with 68 base paired-end reads on an Illumina GAI instrument (Illumina) following manufacturer's recommendations. In both cases, fragments per kilobase per million reads mapped per CDS was calculated and colonial and attached values averaged and \log_2 transformed (resulting in negative or positive values corresponding to overexpression in one particular cell stage or the other, this relationship is arbitrary and only for visualization purposes).

Supplementary Material

Supplementary figures S1–S9 and table S1 are available at *Molecular Biology and Evolution* online (<http://www.mbe.oxfordjournals.org/>).

Acknowledgments

The authors thank Matt Welch for providing critical feedback on the manuscript. They also thank the Joint Genome Institute and Broad Institute for making data publicly available. N.K. is a Senior Fellow in the Integrated Microbial Biodiversity Program of the Canadian Institute for Advanced Research. This work was supported by an Institució Catalana de Recerca i Estudis Avançats contract, a European Research Council Starting Grant (ERC-2007-StG-206883), a grant (BFU2011-23434) from Ministerio de Economía y Competitividad (MINECO) to I.R.-T., funding from NIH NIGMS GM089977 to N.K., pregraduate Formacion Profesorado Universitario grant from MICINN to A.S.-P., a Deutsche Forschungsgemeinschaft (DFG) postdoctoral fellowship to P.B., and the Canadian Research Chair program grant to B.F.L.

References

Adl SM, Simpson AGB, Lane CE, et al. (25 co-authors). 2012. The revised classification of eukaryotes. *J Eukaryot Microbiol.* 59:429–514.

Antón IM, Jones GE, Wandosell F, Geha R, Ramesh N. 2007. WASP-interacting protein (WIP): working in polymerisation and much more. *Trends Cell Biol.* 17:555–562.

Berg JS, Cheney RE. 2002. Myosin-X is an unconventional myosin that undergoes intrafilopodial motility. *Nat Cell Biol.* 4:246–250.

Bohil AB, Robertson BW, Cheney RE. 2006. Myosin-X is a molecular motor that functions in filopodia formation. *Proc Natl Acad Sci U S A.* 103:12411.

Boureaux A, Vignal E, Faure S, Fort P. 2007. Evolution of the Rho family of ras-like GTPases in eukaryotes. *Mol Biol Evol.* 24:203–216.

Breshears LM, Wessels D, Soll DR, Titus MA. 2010. An unconventional myosin required for cell polarization and chemotaxis. *Proc Natl Acad Sci U S A.* 107:6918–6923.

Bretscher A, Edwards K, Fehon RG. 2002. ERM proteins and merlin: integrators at the cell cortex. *Nat Rev Mol Cell Biol.* 3:586–599.

Brown MW, Kolisko M, Silberman JD, Roger AJ. 2012. Aggregative multicellularity evolved independently in the eukaryotic supergroup Rhizaria. *Curr Biol.* 22:1–5.

Cavalier-Smith T. 2003. Phylogeny of Choanozoa, Apusozoa, and other Protozoa and early eukaryote megaevolution. *J Mol Evol.* 56:540–563.

Cavalier-Smith T. 2013. Early evolution of eukaryote feeding modes, cell structural diversity, and classification of the protozoan phyla Loukozoa, Sulcozoa, and Choanozoa. *Eur J Protistol.* 49:115–178.

Cavalier-Smith T, Chao EE. 2010. Phylogeny and evolution of Apusomonadida (Protozoa: Apusozoa): new genera and species. *Protist* 161:549–576.

Chalkia D, Nikolaidis N, Makalowski W, Klein J, Nei M. 2008. Origins and evolution of the formin multigene family that is involved in the formation of actin filaments. *Mol Biol Evol.* 25:2717–2733.

Dayel MJ, Alegado RA, Fairclough SR, Levin TC, Nichols SA, McDonald K, King N. 2011. Cell differentiation and morphogenesis in the colony-forming choanoflagellate *Salpingoeca rosetta*. *Dev Biol.* 357:73–82.

Deeks MJ, Calcutt JR, Ingle EKS, et al. (12 co-authors). 2012. A superfamily of actin-binding proteins at the actin-membrane nexus of higher plants. *Curr Biol.* 22:1–6.

Derelle R, Lang BF. 2012. Rooting the eukaryotic tree with mitochondrial and bacterial proteins. *Mol Biol Evol.* 29:1277–1289.

DeRosier DJ, Tilney LG. 2000. F-actin bundles are derivatives of microvilli: what does this tell us about how bundles might form? *J Cell Biol.* 148:1–6.

Dumontier M, Höcht P, Mintert U, Faix J. 2000. Rac1 GTPases control filopodia formation, cell motility, endocytosis, cytokinesis and development in *Dictyostelium*. *J Cell Sci.* 113:2253–2265.

Eddy SR. 1998. Profile hidden Markov models. *Bioinformatics* 14:755.

Fairclough SR, Chen Z, Kramer E, et al. (15 co-authors). 2013. Premetazoan genome evolution and the regulation of cell differentiation in the choanoflagellate *Salpingoeca rosetta*. *Genome Biol.* 14:R15.

Faix J, Breitsprecher D, Stradal TE, Rottner K. 2009. Filopodia: complex models for simple rods. *Int J Biochem Cell Biol.* 41:1656–1664.

Faix J, Rottner K. 2006. The making of filopodia. *Curr Opin Cell Biol.* 18:18–25.

Fletcher DA, Mullins RD. 2010. Cell mechanics and the cytoskeleton. *Nature* 463:485–492.

Goley ED, Welch MD. 2006. The ARP2/3 complex: an actin nucleator comes of age. *Nat Rev Mol Cell Biol.* 7:713–726.

Gómez F, Moreira D, Benzerara K, López-García P. 2011. *Solenicola setigera* is the first characterized member of the abundant and cosmopolitan uncultured marine stramenopile group MAST-3. *Environ Microbiol.* 13:193–202.

Gonobobleva E, Maldonado M. 2009. Choanocyte ultrastructure in *Halisarca dujardini* (Demospongiae, Halisarcida). *J Morphol.* 270:615–627.

Gupton SL, Gertler FB. 2007. Filopodia: the fingers that do the walking. *Sci Signal.* 2007:re5.

Hoeflich KP, Ikura M. 2004. Radixin: cytoskeletal adopter and signaling protein. *Int J Biochem Cell Biol.* 36:2131–2136.

Huelsenbeck JPP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754–755.

Karpov S, Leadbeater BSC. 1998. Cytoskeleton structure and composition in choanoflagellates. *J Eukaryot Microbiol.* 45:361–367.

Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 30:3059.

Khurana S, George SP. 2008. Regulation of cell structure and function by actin-binding proteins: villin's perspective. *FEBS Lett.* 582:2128–2139.

King N, Westbrook M, Young S, Kuo A. 2008. The genome of the choanoflagellate *Monosiga brevicollis* and the origin of metazoans. *Nature* 21:4300–4305.

Kinley AW, Weed SA, Weaver AM, Karginov AV, Bissonette E, Cooper JA, Parsons JT. 2003. Cortactin interacts with WIP in regulating Arp2/3 activation and membrane protrusion. *Curr Biol.* 13:384–393.

Kollmar M, Lbik D, Enge S. 2012. Evolution of the eukaryotic ARP2/3 activators of the WASP family: WASP, WAVE, WASH, and WHAMM, and the proposed new family members WAWH and WAML. *BMC Res Notes.* 5:1–23.

Kureishy N, Sapountzi V, Prag S, Anilkumar N, Adams JC. 2002. Fascins, and their roles in cell structure and function. *Bioessays* 24:350–361.

Ladwein M, Rottner K. 2008. On the Rho'd: the regulation of membrane protrusions by Rho-GTPases. *FEBS Lett.* 582:2066–2074.

Leadbeater BSC. 1979. Developmental and ultrastructural observations on two stalked marine Choanoflagellates, *Acanthoecopsis spiculifera* Norris and *Acanthoeca spectabilis* Ellis. *Proc R Soc Lond B Biol Sci.* 204:57–66.

- Leadbeater BSC, Morton C. 1974. A microscopical study of a marine species of *Codosiga* James-Clark (Choanoflagellata) with special reference to the ingestion of bacteria. *Biol J Linn Soc Lond.* 6:337–347.
- Lundquist EA. 2009. The finer points of filopodia. *PLoS Biol.* 7:e1000142.
- Magie CR, Daly M, Martindale MQ. 2007. Gastrulation in the cnidarian *Nematostella vectensis* occurs via invagination not ingression. *Dev Biol.* 305:483–497.
- Mattila PK, Lappalainen P. 2008. Filopodia: molecular architecture and cellular functions. *Nat Rev Mol Cell Biol.* 9:446–454.
- McClatchey AI. 2012. ERM proteins. *Curr Biol.* 22:R784–R785.
- McDonald KL, Webb RI. 2011. Freeze substitution in 3 hours or less. *J Microsc.* 243:227–233.
- Mellor H. 2010. The role of formins in filopodia formation. *Biochim Biophys Acta.* 1803:191–200.
- Mikrjukov KA, Mylnikov AP. 2001. A study of the fine structure and the mitosis of a lamellicristate amoeba, *Micronuclearia podoventralis* gen. et sp. nov. (Nucleariidae, Rotosphaerida). *Eur J Protistol.* 37: 15–24.
- Nagy S, Ricca BL, Norstrom MF, Courson DS, Brawley CM, Smithback PA, Rock RS. 2008. A myosin motor that selects bundled actin for motility. *Proc Natl Acad Sci U S A.* 105:9616–9620.
- Nambiar R, McConnell RE, Tyska MJ. 2010. Myosin motor function: the ins and outs of actin-based membrane protrusions. *Cell Mol Life Sci.* 67:1239–1254.
- Niggli V, Rossy J. 2008. Ezrin/radixin/moesin: versatile controllers of signaling molecules and of the cortical cytoskeleton. *Int J Biochem Cell Biol.* 40:344–349.
- Odrionitz F, Kollmar M. 2007. Drawing the tree of eukaryotic life based on the analysis of 2,269 manually annotated myosins from 328 species. *Genome Biol.* 8:R196.
- Ota S, Eikrem W, Edvardsen B. 2011. Ultrastructure and molecular phylogeny of Thaumatomonads (Cercozoa) with emphasis on *Thaumatomastix salina* from Oslofjorden, Norway. *Protist* 163: 560–573.
- Ota S, Vaulot D. 2012. *Lotharella reticulosa* sp. nov.: a highly reticulated network forming chlorarachniophyte from the Mediterranean Sea. *Protist* 163:91–104.
- Pawlowski J. 2008. The twilight of Sarcodina: a molecular perspective on the polyphyletic origin of amoeboid protists. *Protistology* 5:281–302.
- Pollard TD. 2007. Regulation of actin filament assembly by Arp2/3 complex and formins. *Annu Rev Biophys Biomol Struct.* 36:451–477.
- Preston T, King C. 2005. Locomotion and phenotypic transformation of the Amoeboflagellate *Naegleria gruberi* at the water/air interface. *J Eukaryot Microbiol.* 50:245–251.
- Ren G, Crampton MS, Yap AS. 2009. Cortactin: coordinating adhesion and the actin cytoskeleton at cellular protrusions. *Cell Motil Cytoskeleton.* 66:865–873.
- Revenu C, Athman R, Robine S, Louvard D. 2004. The co-workers of actin filaments: from cell structures to signals. *Nat Rev Mol Cell Biol.* 5:635–646.
- Richards TA, Cavalier-Smith T. 2005. Myosin domain evolution and the primary divergence of eukaryotes. *Nature* 436:1113–1118.
- Ridley AJ. 2006. Rho GTPases and actin dynamics in membrane protrusions and vesicle trafficking. *Trends Cell Biol.* 16:522–529.
- Ruiz-Trillo I, Burger G, Holland PWH, King N, Lang BF, Roger AJ, Gray MW. 2007. The origins of multicellularity: a multi-taxon genome initiative. *Trends Genet.* 23:113–118.
- Ruiz-Trillo I, Roger AJ, Burger G, Gray MWMW, Lang BF. 2008. A phylogenomic investigation into the origin of metazoa. *Mol Biol Evol.* 25: 664–672.
- Schäfer C, Born S, Möhl C, Houben S, Kirchgeßner N, Merkel R. 2010. Dependence of adhesion, actin bundles, force generation and transmission on filopodia. *Cell Adh Migr.* 4:2:215–225.
- Schirenbeck A, Arasada R, Bretschneider T, Stradal TEB, Schleicher M, Faix J. 2006. The bundling activity of vasodilator-stimulated phosphoprotein is required for filopodium formation. *Proc Natl Acad Sci U S A.* 103:7694–7699.
- Sebé-Pedrós A, Roger A, Lang BF, King N, Ruiz-Trillo I. 2010. Ancient origin of the integrin-mediated adhesion and signaling machinery. *Proc Natl Acad Sci U S A.* 107:10142–10147.
- Silacci P, Mazzolai L, Gauci C, Stergiopoulos N, Yin HL, Hayoz D. 2004. Gelsolin superfamily proteins: key regulators of cellular functions. *Cell Mol Life Sci.* 61:2614–2623.
- Small J, Auinger S, Nemethova M, Koestler S, Goldie K, Hoenger A, Resch G. 2008. Unravelling the structure of the lamellipodium. *J Microsc.* 231:479–485.
- Small JV, Stradal T, Vignal E, Rottner K. 2002. The lamellipodium: where motility begins. *Trends Cell Biol.* 12:112–120.
- Soldati D, Meissner M. 2004. Toxoplasma as a novel system for motility. *Curr Opin Cell Biol.* 16:32–40.
- Sousa AD, Cheney RE. 2005. Myosin-X: a molecular motor at the cell's fingertips. *Trends Cell Biol.* 15:533–539.
- Stamatakis A. 2006. RAXML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22:2688.
- Suga H, Dacre M, De Mendoza A, Shalchian-Tabrizi K, Manning G, Ruiz-Trillo I. 2012. Genomic survey of premetazoans shows deep conservation of cytoplasmic tyrosine kinases and multiple radiations of receptor tyrosine kinases. *Sci Signal.* 5:ra35–ra35.
- Takenawa T, Miki H. 2001. WASP and WAVE family proteins: key molecules for rapid rearrangement of cortical actin filaments and cell movement. *J Cell Sci.* 114:1801–1809.
- Tilney LG, Connelly PS, Guild GM. 2004. Microvilli appear to represent the first step in actin bundle formation in *Drosophila bristles*. *J Cell Sci.* 117:3531–3538.
- Tsui CKM, Marshall W, Yokoyama R, Honda D, Lippmeier JC, Craven KD, Peterson PD, Berbee ML. 2009. Labyrinthulomycetes phylogeny and its implications for the evolutionary loss of chloroplasts and gain of ectoplasmic gliding. *Mol Phylogenet Evol.* 50:129–140.
- Tuxworth RJ, Weber I, Wessels D, Addicks GC, Soll DR, Gerisch G, Titus MA. 2001. A role for myosin VII in dynamic cell adhesion. *Curr Biol.* 11:318–329.
- Vallotton P, Small JV. 2009. Shifting views on the leading role of the lamellipodium in cell migration: speckle tracking revisited. *J Biosci.* 122:1955–1958.
- Veltman DM, Insall RH. 2010. WASP family proteins: their evolution and its physiological implications. *Mol Biol Cell.* 21:2880.
- Vlahou G, Rivero F. 2006. Rho GTPase signaling in *Dictyostelium discoideum*: insights from the genome. *Eur J Cell Biol.* 85:947–959.
- Weaver AM, Karginov AV, Kinley AW, Weed SA, Li Y, Parsons JT, Cooper JA. 2001. Cortactin promotes and stabilizes Arp2/3-induced actin filament network formation. *Curr Biol.* 11:370–374.
- Weaver AM, Young ME, Lee W-L, Cooper JA. 2003. Integration of signals to the Arp2/3 complex. *Curr Opin Cell Biol.* 15:23–30.
- Yang C, Svitkina T. 2011. Filopodia initiation—focus on the Arp2/43 complex and formins. *Cell Adh Migr.* 5:402–408.
- Zettler LA, Nerad TA, O'Kelly CJ, Sogin ML. 2001. The nuclearioid amoebae: more protists at the animal-fungal boundary. *J Eukaryot Microbiol.* 48:293–297.
- Zhang H, Berg JS, Li Z, Wang Y, Lang P, Sousa AD, Bhaskar A, Cheney RE, Stromblad S. 2004. Myosin-X provides a motor-based link between integrins and the cytoskeleton. *Nat Cell Biol.* 6:523–531.
- Zwaenepoel I, Naba A, Da Cunha MML, Del Maestro L, Formstecher E, Louvard D, Arpin M. 2012. Ezrin regulates microvillus morphogenesis by promoting distinct activities of Eps8 proteins. *Mol Biol Cell.* 23:1080–1094.