


Role of the FtsA C Terminus as a Switch for Polymerization and Membrane Association

Marcin Krupka,^a Elisa J. Cabré,^b Mercedes Jiménez,^b Germán Rivas,^b  Ana Isabel Rico,^a Miguel Vicente^a

Centro Nacional de Biotecnología, Consejo Superior de Investigaciones Científicas (CNB-CSIC), Madrid, Spain^a; Centro de Investigaciones Biológicas, Consejo Superior de Investigaciones Científicas (CIB-CSIC), Madrid, Spain^b

ABSTRACT Together with ATP, the C-terminal region of the essential streptococcal FtsA protein acts as an intramolecular switch to promote its polymerization and attachment to the membrane. During septation, FtsA is known to anchor the constricting FtsZ ring and, subsequently, the divisome to the membrane. Truncation of the C terminus of the streptococcal FtsA (FtsAΔCt) facilitates a more rapid ATP-dependent polymerization in solution than is seen with the full-length protein (FtsA⁺). The FtsAΔCt polymers are more organized and compact than those formed in solution by FtsA⁺, resembling the shape of the membrane-associated FtsA⁺ polymers. We find that ATP, besides being needed for polymerization, is required for the attachment of FtsA⁺ to lipid monolayers and to vesicle membranes. We propose a model in which the binding of ATP activates a switch favoring the polymerization of FtsA and at the same time driving the amphipathic helix at its C terminus to become attached to the membrane. Conversely, when FtsA is in the cytoplasm, the C terminus is not engaged in the attachment to the membrane, and it obstructs polymerization. ATP-dependent polymerization of FtsA inside membrane vesicles causes vesicle shrinkage, suggesting that, besides providing a membrane attachment for FtsZ, the FtsA C terminus may also introduce local alterations in the membrane to facilitate septation.

IMPORTANCE FtsA is a protein needed in many bacteria to construct a septum that divides one fully grown cell, producing two daughters. We show that the region located at the C-terminal end of the *Streptococcus pneumoniae* FtsA protein works as a switch triggered by ATP, a molecule that stores energy. This region contains an amphipathic helix that obstructs the assembly of FtsA into polymers in the cytoplasm. In the presence of ATP, the obstruction is removed by switching the position of the helix. The switch directs the helix to the membrane and simultaneously facilitates the polymerization of the protein. The accumulation of FtsA molecules at the membrane causes distortions, an effect produced also by proteins such as MinD, MreB, and SepF that also contain amphipathic helices as membrane attachment devices. In the case of FtsA, these distortions may also facilitate the initial events that lead to the division of bacteria.

Received 30 October 2014 Accepted 3 November 2014 Published 25 November 2014

Citation Krupka M, Cabré EJ, Jiménez M, Rivas G, Rico AI, Vicente M. 2014. Role of the FtsA C terminus as a switch for polymerization and membrane association. *mBio* 5(6): e02221-14. doi:10.1128/mBio.02221-14.

Editor Philippe J. Sansonetti, Pasteur Institute

Copyright © 2014 Krupka et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Noncommercial-ShareAlike 3.0 Unported license](https://creativecommons.org/licenses/by-nc-sa/4.0/), which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Address correspondence to Ana Isabel Rico, airico@cnb.csic.es, or Miguel Vicente, mvicente@cnb.csic.es.

This article is a direct contribution from a Fellow of the American Academy of Microbiology.

At the earliest stage of bacterial division, the FtsA protein connects FtsZ, the principal component of the division machinery, to the cell membrane and forms a structure called the proto-ring at the division site (reviewed in reference 1). Correct assembly of proto-ring components enables progression of bacterial division by recruiting in successive steps the remainder of the essential divisome proteins involved in septum synthesis and cell pole formation. FtsA is a multiple connector protein with various regions able to interact independently with several of the early- and late-assembling division proteins, including itself.

FtsA belongs structurally to the actin/Hsp70/hexokinase superfamily (2–4). In the presence of ATP, FtsA monomers polymerize through a head-to-tail assembly mechanism that involves its 1C domain and the S12 and S13 β-strands (5–8). These regions are essential for cell division in *Escherichia coli* (9), and the residues involved in the self-interaction have been mapped in the crystal

model from the *Thermotoga maritima* FtsA dimer (8). The 1C domain is also the main factor in recruitment of late-assembling division proteins (9, 10); dissociation of the FtsA oligomer to release this domain is thus proposed to help to recruit the late-assembling division proteins (11). In addition to the head-to-tail interactions between monomers, the polymers also establish lateral interactions to form bundles. The H1 alpha helix in domain 1A is involved in these interactions, and it is needed for the integrity of the Z ring (11). Consistent with their negligible ATP hydrolytic activity and the apparently insignificant differences between the ATP-bound and unbound crystal forms (4), FtsA polymers remain stable *in vitro* (6–8).

At its C terminus, FtsA has a flexible, disordered hydrophilic linker that connects the core of the protein to a membrane-targeting sequence (MTS). In *E. coli* FtsA, the MTS is an essential, conserved amphipathic helix which participates in anchoring the

protein to the cell membrane (12) and in regulating FtsA self-interaction (13).

In this work, we have investigated the role of the streptococcal FtsA C-terminal region in its ATP-dependent polymerization and explored the relationship between the two activities previously assigned to this region, i.e., FtsA self-interaction and attachment to the cytoplasmic membrane. We used FtsA from *Streptococcus pneumoniae* produced heterologously in *E. coli* (5), since this protein is more stable than the *E. coli* FtsA and shows ATP-dependent polymerization activity (6). Our results suggest that, upon ATP binding, the position of the MTS acts as a switch to modify simultaneously the polymerization activity and the interaction of the protein with the membrane.

RESULTS

The FtsA C-terminal region is involved in FtsA polymerization.

The C terminus of *E. coli* FtsA, containing the MTS, has been described to have a role in the interaction between FtsA molecules (13). To study the role of the FtsA C-terminal region in the protein self-interaction, we took advantage of the ability of *S. pneumoniae* FtsA to polymerize in the presence of ATP (5, 6). Purification of FtsA from *S. pneumoniae* yields two distinct bands of different sizes in analysis by SDS-PAGE (6). The mobility of the larger band corresponds to the molecular weight of the full gene product, whereas the smaller one fits with a deletion comprising the last 52 residues at the C terminus of the protein as confirmed by amino acid sequencing (6). We purified a homogeneous variant of the streptococcal FtsA lacking the 52 C-terminal amino acids (FtsA Δ Ct) fused to a His tag after expressing a suitably deleted genetic construction (see Materials and Methods).

Measurement of polymerization by light-scattering assay showed that at a 2.5 or 5 μ M concentration, the FtsA Δ Ct protein assembled into large structures in the presence of 2 mM ATP (Fig. 1A). The C-terminal-truncated protein assembled more rapidly than the purified full-length protein (containing a six-His tag at its N-terminal end; simplified here as FtsA⁺) and presented a 20% reduced lag time to initiation of polymerization. Consistently with the absence of measurable ATPase activity (6), the polymers are stable and their amount depends only on the protein concentration. Nonetheless, our results suggest that all the FtsA⁺ and FtsA Δ Ct monomers are finally incorporated into stable polymers in solution, as polymerization reaches the maximal scatter intensity level in reaction mixtures containing FtsA⁺ or FtsA Δ Ct or an equimolar mixture of FtsA⁺ and FtsA Δ Ct (Fig. 1A). From these results, we infer that the MTS causes a temporal delay of the FtsA polymerization in solution but does not affect the final amount of polymer.

Images of the polymers formed by FtsA Δ Ct showed bundles of helical protofilaments morphologically similar to those formed by FtsA⁺ (Fig. 1B and C). This suggests that the two proteins may establish similar head-to-tail interactions and lateral contacts when polymerizing. FtsA Δ Ct polymers nevertheless had a more regular and compact organization than FtsA⁺ polymers, frequently forming tubular structures (18 ± 2 nm in diameter), suggesting a role of the C terminus as a steric hindrance for polymerization.

Since FtsA⁺ and FtsA Δ Ct assembled in bundles of helical protofilaments when assayed separately, we reasoned that similar bundles would result from the simultaneous polymerization of

the two proteins in a mixture at equimolar concentrations. The lag time and the molar mass values obtained for a polymerization mixture containing FtsA⁺ and FtsA Δ Ct were within the ranges individually shown by each species (Fig. 1A). However, transmission electron microscopy (EM) images nonetheless showed that the polymers formed by the mixture of the two proteins were a disordered network of short filaments and protein clumps (Fig. 1D). In contrast to the results seen with the FtsA⁺ or FtsA Δ Ct polymers, the filaments formed when the two proteins were assayed together barely bundled, suggesting that the lateral contacts among protofilaments were impaired. Given the tendency to self-associate shown by the hydrophobic faces within amphipathic helices, we suggest that MTS-MTS interactions may partially counteract the negative effect of this region, as the FtsA⁺ polymers were more ordered than those formed by the mixture.

These results indicate that the C-terminal region might interfere with FtsA polymerization in solution, probably by disturbing the initial nucleation event proposed for the FtsA⁺ polymerization (6) and/or the subsequent addition of the monomers to the protofilaments.

The C-terminal region of FtsA affects polymerization but not the head-to-tail interactions.

The interference of the C-terminal ends in polymerization could be exerted through the distortion of the surfaces involved in the head-to-tail interaction or, alternatively, by masking other, unknown interaction sites in the FtsA⁺ molecule. To discriminate between the two possibilities, we measured the ability of FtsA Δ Ct to polymerize in the presence of truncated FtsA proteins from *S. pneumoniae* that lacked S12-to-S13 β strands or the 1C domain (at the top or bottom of the crystallographic model, respectively). These proteins inhibit FtsA⁺ head-to-tail bidirectional polymerization by capping the opposite ends and cannot polymerize (5). We designed FtsA Δ Ct constructs in which these regions were additionally deleted. Deletion of the 1C domain from FtsA Δ Ct resulted in a protein (FtsA Δ Ct Δ 1C) with a high tendency to aggregate, making analysis in solution unreliable (not shown).

We have measured whether the double-deletion FtsA Δ Ct Δ S protein, in which both the C-terminal end and the S12-to-S13 region were lacking, affected the polymerization and morphology of the FtsA Δ Ct polymers. FtsA Δ Ct Δ S itself did not polymerize, and, when added to FtsA Δ Ct, it reduced both the maximum level and rate of polymerization (Fig. 1E). In addition, the polymers obtained from a mixture containing both proteins (Fig. 1F) were shorter, less abundant, less compact, and less defined than polymers formed by FtsA Δ Ct alone (Fig. 1C). This effect is similar to that exerted by FtsA Δ S on FtsA⁺ polymerization where both proteins contain their C termini (5). These results indicate that, in addition to the head-to-tail interaction, the C terminus has a distinct role in FtsA polymerization.

Association of the C-terminal region of *Streptococcus pneumoniae* FtsA (SnFtsA) with the *E. coli* cytoplasmic membrane prevents the initiation of constrictions and distorts the membrane.

The FtsA proteins from *E. coli*, *T. maritima*, and *Bacillus subtilis*, which do not form polymers in solution, can polymerize inside the cytoplasm of *E. coli* cells (8, 12). Given that the streptococcal FtsA⁺ and FtsA Δ Ct proteins polymerize *in vitro*, we tested whether they could also form polymers inside *E. coli*. For this purpose, these two streptococcal proteins were overproduced separately in *E. coli* as fusions to the fluorescent mCherry protein. Data in Fig. 2A and B show that mCherry FtsA⁺ appeared in close




FIG 1 *In vitro* polymerization of FtsA proteins. (A) Light-scattering measurements of 5 μM FtsA⁺ (○), 5 μM FtsA Δ Ct (●), 2.5 μM FtsA⁺ (□), 2.5 μM FtsA Δ Ct (■), and a mixture of 2.5 μM FtsA⁺ with 2.5 μM FtsA Δ Ct (△) after the addition of 2 mM ATP. A.U., arbitrary units. (B to D) Electron microscopy of polymers formed by 5 μM FtsA⁺ (B), 5 μM FtsA Δ Ct (C), and a mixture of 2.5 μM FtsA⁺ and 2.5 μM FtsA Δ Ct (D). Bars, 100 nm. (E) Light-scattering measurements of 5 μM FtsA Δ Ct (●), 5 μM FtsA Δ Ct Δ S (□), or 5 μM FtsA Δ Ct in the presence of 1.25, 2.5, 3.75, and 5 μM FtsA Δ Ct Δ S (■, ▲, ◆, and ○, respectively). (F) Electron microscopy images of polymers formed by a mixture of 5 μM FtsA Δ Ct and 2.5 μM FtsA Δ Ct Δ S. Bar, 100 nm.

proximity to the membrane and inhibited septation. This effect may have resulted from an excess of C-terminal amphipathic FtsA helices attached to the membrane. Indeed, our results show a high density of labeled proteins in the vicinity of the cytoplasmic membrane that appears distorted (Fig. 2A to C).

As expected, the mCherry FtsA Δ Ct fusion, in which the membrane-targeting helix was absent, failed to localize near the cytoplasmic membrane. Its overproduction did not prevent the initiation of constrictions and caused no apparent distortions in the membrane (Fig. 2E to G). These constrictions failed to complete division, perhaps because the bulk of the mCherry FtsA Δ Ct observed at the constriction may block them.

To examine if the accumulated FtsA⁺ proteins could be arranged as polymers on the cytoplasmic membrane, we further

analyzed the *in vitro* polymerization of FtsA⁺ on a lipid monolayer. Purified FtsA⁺ formed long, tubular polymers densely covering a lipid monolayer 40 min after the addition of ATP. They appear as nested tubes similar to those found for FtsA Δ Ct when polymerized in solution (Fig. 1B and 2D). In contrast to FtsA⁺, only occasional FtsA Δ Ct polymers were found on the lipid monolayer in the presence of ATP, as exemplified by the one shown in Fig. 2H. As FtsA Δ Ct lacks the membrane-targeting sequence, its binding to the lipid monolayer after washing was likely unspecific.

Fusions of mCherry to the amino terminus of the nonpolymerizing FtsA Δ S or FtsA Δ 1C protein resulted in proteins that, as expected, did not polymerize in solution. These proteins nevertheless contained an intact C terminus (5), and they consequently localized in the proximity of the cytoplasmic membrane similarly




FIG 2 Localization of streptococcal FtsA⁺ and FtsA Δ Ct overproduced in *E. coli* and on lipid monolayers. FtsA (A to C) and FtsA Δ Ct (E to G) were fused with mCherry (red) at the N terminus and overproduced in *E. coli* (3 h). Membranes were stained with FM1-43FX (green). Samples were analyzed by confocal microscopy (A and E) and electron microscopy (B, C, F, and G). Panels B and F show the gold immunolocalization of FtsA and FtsA Δ Ct, respectively. For monolayer assay, a lipid-coated electron microscopy grid was incubated with FtsA⁺ or FtsA Δ Ct, followed by the addition of 2 mM ATP (D or H, respectively). Samples were washed extensively with polymerization buffer and fixed with 2% uranyl acetate (see Materials and Methods). (H) Note that polymers formed by FtsA Δ Ct occurred very occasionally on the lipid monolayer. Scale bars are as indicated.

to the FtsA⁺ protein (see Fig. S1 in the supplemental material). However, these nonpolymerizing proteins failed to distort the membrane as FtsA⁺ does. On the other hand, a similar mCherry fusion to the double-deletion FtsA Δ Ct Δ S protein was distributed all over the cytoplasm, as would be expected given the absence of a membrane-attaching amphipathic helix and its inability to polymerize. When overproduced, the products of this FtsA Δ Ct Δ 1C fusion localized as inclusion bodies, as would be expected from the tendency of the double mutant to form aggregates (see Fig. S1). Together, these results suggest that the FtsA C terminus, besides mediating the association of the protein with the membrane,

might produce an alteration in its structure when attached in an orderly manner in large amounts, as would occur in the FtsA polymers.

The attachment of FtsA⁺ polymers to the inner surface of GUVs induces shrinkage. FtsA⁺ associates with cell membranes *in vivo*, and, with the addition of ATP, it forms lipid-anchored polymers *in vitro*. We have analyzed the localization of FtsA⁺ inside giant unilamellar vesicles (GUVs) and the effects of its polymerization. The use of GUVs provides a membrane-confined environment, intermediate between the cell and the lipid monolayer, in which the effects of polymerization can be followed over time.




FIG 3 Spatial distribution of Alexa 488-labeled FtsA⁺, FtsA Δ Ct, and FtsA Δ S inside GUVs. The indicated proteins (green) were encapsulated in GUVs (see Materials and Methods). Representative equatorial cross-sectional images for each protein taken shortly after addition of 2 mM ATP (3 to 7 min) and after prolonged (>10-min) incubation are shown. The percentages of vesicles showing similar localization patterns are indicated in the upper right corner of the respective images. The total number of vesicles observed in each case is indicated in brackets. Bar, 10 μ m.

Purified FtsA⁺ and its variants were labeled with Alexa 488 and incorporated separately into vesicles. In the absence of ATP, all the FtsA variants were distributed homogeneously in the vesicle lumen (Fig. 3). After addition of ATP, the bulk of FtsA⁺ localized at the vesicle membrane and only a residual amount could be observed in the lumen. This protein attached more efficiently to lipids than the nonpolymerizing FtsA Δ S variant (Fig. 3), which conserves the C terminus and the ATP-binding pocket (5), suggesting that a weak membrane affinity of FtsA monomers should be compensated for by the protein multimerization. The FtsA Δ S could be detected on the membrane only in the presence of ATP (Fig. 3), suggesting that the ATP may shift the protein from a state in which it hinders membrane attachment to a membrane-bound state even as a monomer. On the other hand, the majority of the FtsA Δ Ct protein, lacking an MTS, failed to localize at the vesicle membrane upon addition of ATP (Fig. 3) even if it had a higher tendency to polymerize (Fig. 1). Therefore, we propose that the shift promoted by ATP involves the movement of the C-terminal region and, consequently, of the MTS. However, we then tested the ability of a double-deletion FtsA Δ Ct Δ S protein to interact with vesicles and found that it was not totally abolished (not shown). This result suggests that FtsA may contain secondary membrane-interacting regions additional to the C terminus that could be exposed upon the ATP addition.

These results suggest that, to efficiently associate to the membrane, FtsA needs both the presence of the MTS and the formation of polymers triggered by ATP. The fact that polymerization and membrane attachment are synergistically enhanced in the presence of ATP would also explain why *E. coli* and *T. maritima* FtsA

polymers cannot be detected *in vitro* unless they are membrane associated (7, 8).

After 10 min of incubation with ATP, over 70% of the vesicles containing FtsA⁺ shrank, while those containing any of the truncated FtsA variants showed no apparent deformation. After further incubation periods (over 45 min) or at higher protein concentrations (over 13 μ M), the truncated proteins were also able to promote deformation in 70% of the observed vesicles (not shown). We postulate that vesicle deformation may result from the progression of polymerization, leading to the formation of large assemblies near the membrane and thus to the accumulation of membrane-anchored amphipathic helices. In the case of FtsA Δ Ct, we observed brilliant protein clusters in the lumen of vesicles, in accordance with its ability to polymerize. Consistently with our *in vivo* results with the mCherry-tagged FtsA Δ Ct, these structures were not attached to the vesicle membrane, and therefore the vesicles showed no deformation after prolonged observation times (Fig. 2E to G and 3).

When FtsA⁺ was added to the buffer outside the vesicles, it formed polymers that could be found decorating their outer surface in part upon the addition of ATP (see Fig. S2 in the supplemental material), which confirmed the affinity of this protein for lipid membranes. We conclude that the recruitment of FtsA⁺ to the membrane is predominantly mediated by the C terminus and that it requires the polymerization induced by ATP.

DISCUSSION

We propose that an intramolecular switch, driven by ATP binding and involving the displacement of its C terminus, operates in the




FIG 4 The role of the C terminus in FtsA ATP-dependent polymerization and attachment to the membrane. (A) FtsA monomers (blue) are not anchored to the membrane in the absence of nucleotide, and free C-terminal ends temporarily disturb the polymerization process in the cytoplasm (bottom). The FtsA molecule domains (1A, 1C, 2A, and 2B), the MTS (blue rectangles), and the flexible linker (blue lines) that connects the MTS to the rest of the protein are represented. ATP (green triangles) triggers the FtsA polymerization and membrane attachment mediated by a shift of the position of the MTS (top). Local perturbations produced by the abundance of MTS in the membrane (gray) are represented in red. (B) The lack of C termini in FtsA Δ Ct facilitates the ATP-dependent polymerization of the protein in solution, but it does not interact specifically with the membrane.

essential FtsA cell division protein from *S. pneumoniae* to allow its polymerization and its attachment to the membrane (Fig. 4).

Our proposal is based on data showing the different behavior of the FtsA Δ Ct protein, in which the C terminus is absent, relative to that of FtsA⁺. Differences can be observed in three properties: the ability to polymerize in solution, the shape of the polymers, and the localization in the cytoplasmic membrane or in lipid surfaces. In the presence of ATP, FtsA Δ Ct polymerizes in solution faster than FtsA⁺. Whereas polymers of FtsA⁺ in solution appear loose, the polymers formed both by FtsA Δ Ct in solution and by FtsA⁺ when associated to lipid surfaces are more compact. Finally, FtsA Δ Ct cannot localize, even in the presence of ATP, either in the cytoplasmic membrane of *E. coli* cells when expressed heterologously or in the membranes of GUVs when artificially included in them. Moreover, FtsA Δ Ct forms organized structures in the cytoplasm or in the lumen of the vesicles, whereas under similar conditions the FtsA⁺ structures are readily associated to the membrane both in cells and vesicles.

A similar increased polymerization when associated to the membrane has been reported for other proteins such as MinD, MreB, and SepF in which their attachment to the membrane is mediated by an amphipathic helix (MTS) (14–16). In the case of MinD, ATP binding causes the displacement of the MTS, allowing both the interaction between monomers and association to the membrane (17).

A possible mechanism to facilitate FtsA polymerization in the presence of ATP may be based on the different position occupied by the 1C domain in the nucleotide-free dimer structure relative

to the one adopted in the ATP-bound one. Upon overlaying the structure of the ATP-free monomer on the top monomer of the ATP-bound form, it became evident that the 1C domain in the top monomer did not contact the 1A domain of the bottom one (see Fig. S3 in the supplemental material). It is possible, then, that the binding of ATP works by bringing into closer contact the 1C and 1A domains of monomers and therefore promoting polymerization.

Our results favoring the operation of an intramolecular switch acting on both polymerization and membrane attachment do not offer specific clues on which of the two processes occurs earlier during the assembly of the proto-ring. Pichoff and Lutkenhaus (12, 18) have suggested that the interaction between FtsA and FtsZ is likely preceded by both ATP binding and attachment of FtsA to the membrane, an interpretation compatible with our results. In *E. coli*, FtsA initially works together with ZipA to assemble FtsZ into the proto-ring; in addition, a role for the MTS FtsA in the recruitment of the late *E. coli* divisome proteins (FtsN) has been suggested (19, 20). It is possible as well that the local enrichment in amphipathic helices created at the membrane regions in which FtsA becomes associated may weaken the membrane at the site of insertion (21). This effect has been observed to happen in the attachment of SepF (14), MreB (16), and MinD (17, 22) as well. Our results favor this possibility, as the attachment of FtsA⁺ to the membrane of vesicles upon ATP addition results in their shrinking. Besides promoting polymerization and association to the membrane, the ATP-driven switch would initiate the differentia-

tion of a potential septation site in which first a proto-ring and later a divisome are assembled.

Note that, while the *E. coli* proto-ring contains an additional protein, ZipA, to provide attachment to the membrane, the proto-ring of *S. pneumoniae* does not contain ZipA. In this case, the role of ZipA may be replaced by those of other membrane proteins recruiting FtsZ, such as EzrA or SepF (reviewed in reference 23), or by the higher number of FtsA molecules, such as the nearly 2,200 in *S. pneumoniae* (6), instead of the approximately 740 present in *E. coli* (24). Consistent with the role of both proteins being membrane anchors for FtsZ, a large amount of ZipA produces invaginations in the cytoplasmic membrane of *E. coli* (25), a behavior reminiscent of the deformations caused in vesicles by the presence of the ATP-bound streptococcal FtsA protein.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *E. coli* strain DH5 α (26) was used as a host for gene cloning and *E. coli* strain C41 (27) for overproduction and purification of streptococcal proteins. *E. coli* strains were routinely grown at 37°C in Luria-Bertani (LB) agar or broth supplemented with ampicillin (100 $\mu\text{g ml}^{-1}$) when necessary.

Plasmid constructions and DNA manipulation. Standard protocols for molecular cloning, transformation, and DNA analysis were performed as previously described (28). Oligonucleotides were purchased from Sigma-Aldrich and restriction enzymes and DNA polymerase from Roche Diagnostics.

Primers MK1 (5' CGGGATCCATGGCTAGAGAAGGCTT 3') and MK3 (5' CGGAATTC TTAAGCTGTTTTTGCAG 3') were used to amplify the pRSETA_ftsA sequence (6), except the sequence encoding the last 52 amino acids of the FtsA C-terminal region from Gln406 to Glu457. The plasmid named pMKV1 bears the desired deletion to produce truncated FtsA lacking the C terminus, termed FtsA Δ Ct. A similar procedure was used to amplify the ftsA Δ Ct genes additionally lacking the sequence encoding domain 1C (ftsA Δ Ct Δ 1C) or the S12 and S13 β sheets (ftsA Δ Ct Δ S) from the pARV48 and pARV49 plasmids encoding FtsA Δ 1C and FtsA Δ S (5) to yield the plasmids pMKV12 and pMKV2, respectively.

Plasmids from pMKV29 to pMKV34 encoding FtsA and its deletion mutants (FtsA Δ 1C, FtsA Δ S, FtsA Δ Ct, FtsA Δ Ct Δ S, and FtsA Δ Ct Δ 1C, respectively), fused with mCherry at the N-terminal end, were obtained by inserting the mCherry gene into the pRSETA_ftsA, pARV48, pARV49, pMKV1, pMKV2 and pMKV12 plasmids, respectively, to span the His tag-encoding sequence derived from the pRSETA vector (Invitrogen). All mutations were confirmed by DNA sequencing.

Purification of streptococcal His-tagged proteins. The FtsA protein and its variants were overproduced in the C41 strain and purified from the inclusion bodies by affinity chromatography in denaturing conditions, as previously described (5). Samples were renaturalized by sequential dialysis in polymerization buffer (20 mM Tris-HCl [pH 7.5], 50 mM NaCl, 5 mM MgCl₂, 10% glycerol) supplemented with decreasing concentrations (2.5, 1.25, 0.5, and 0 M) of guanidine chloride and stored at -80°C.

FtsA polymerization assays in solution and visualization of polymer morphology. For standard polymerization assays, thawed purified proteins were centrifuged (13,000 \times g, 15 min) to remove protein aggregates and then diluted to a 5 μM final concentration in polymerization buffer. After purified protein was tempered in the reaction mix (room temperature, 5 min), 2 mM ATP (buffered at pH 8) was added to initiate polymerization. The polymerization process was followed by 90° light-scattering measurements using a Hitachi F-2500 fluorescence spectrophotometer. Excitation and emission wavelengths were set to 320 nm with 5-nm slit widths (29).

The morphology of polymers containing FtsA or its variants was observed using transmission electron microscopy 20 min after ATP addition. Samples were applied to a 400-mesh collodion-coated glow-discharged copper grid, negatively stained with 2% uranyl acetate,

inspected with a Jeol JEM1011 transmission electron microscope, and photographed with an 11-megapixel charge-coupled Gatan Erlanghens ES1000W camera. The software used for image capture was Gatan Digital Micrograph 1.8.0, and the software used for processing was Adobe Photoshop CS5.

FtsA overproduction in vivo. Overnight cultures of *E. coli* C41 cells transformed with plasmids from pMKV29 to pMKV34 encoding mCherry-tagged FtsA or its deletion proteins (FtsA Δ 1C, FtsA Δ S, FtsA Δ Ct, FtsA Δ Ct Δ S, and FtsA Δ Ct Δ 1C, respectively) were diluted 1:100 and grown at 37°C in ampicillin-supplemented LB medium. Cultures were maintained by sequential dilution under exponentially balanced growth conditions at below an optical density at 600 nm (OD₆₀₀) of 0.2 for at least 3 h; they were induced with 1 mM isopropyl- β -D-thiogalactopyranoside (IPTG) (3 h), and samples were collected for visualization.

For confocal microscopy imaging, cells were subjected to membrane staining with FM1-43FX (Molecular Probes) and fixed according to methods described in reference 30. Images were collected with a Leica TCS SP5 AOBS inverted confocal microscope with a 63 \times (numerical aperture [NA], 1.4 to 0.6; oil HCX PL APO lambda-blue) immersion objective (Leica, Mannheim, Germany). Argon ion lasers were used for laser lines at 488 nm, which excited FM1-43FX; a diode-pumped solid-state (DPSS) laser was used to excite mCherry at 561 nm.

To examine the organization of these C41 cells, cryo-transmission electron microscopy was performed as previously described (25). For FtsA and FtsA Δ Ct immunolabeling, ultrathin sections of the samples were first blocked with TBST (30 mM Tris-HCl [pH 8], 150 mM NaCl, 1% bovine serum albumin [BSA], 0.02% Tween 20) followed by a 1-h incubation with MVA1 anti-FtsA antibody (6). Samples were incubated with EM goat F(ab')₂ anti-rabbit immunogold conjugate (BBInternational, Cardiff, United Kingdom). All images were processed with Adobe Photoshop CS5.

Lipid monolayer assay. Monolayer assays were performed as previously described (8) with some modifications. Based on the previously estimated streptococcal polar lipid membrane composition (31, 32), 0.2 μg phosphatidylglycerol and cardiolipin (Avanti) at a 1:1 ratio were dissolved in chloroform and floated on polymerization buffer in Teflon wells. After the chloroform evaporated, electron microscopy grids were placed onto the lipid monolayer and FtsA⁺ or FtsA Δ Ct proteins were injected underneath, into the polymerization buffer, to achieve a final concentration of 2 μM and incubated (40 min), and 2 mM ATP was added to trigger polymerization. The reaction was allowed to proceed for 20 min, followed by standard grid staining with 2% uranyl acetate, as described for the polymer visualization.

Polymerization of proteins encapsulated in giant unilamellar vesicles. FtsA⁺ and its truncation mutants (0.75 mg/ml) were labeled with Alexa 488 (Molecular Probes) (1:5 M ratio)–20 mM HEPES-HCl [pH 8]–50 mM NaCl–5 mM MgCl₂ at 4°C for 30 min. The reaction was terminated by addition of Tris-HCl (pH 8) to achieve a final concentration of 100 mM. Excess free probe was removed by gel filtration (HiTrap Desalting; GE Healthcare), and the protein was divided into aliquots and stored at -80°C. Labeling efficiency was 0.1 to 0.2 mol Alexa 488 per mol of protein. Polymerization of labeled proteins was confirmed by 90° light-scattering assays and electron microscopy.

Giant unilamellar vesicles (GUVs) containing FtsA⁺ or its deletion forms (7 μM) were prepared by the droplet-transfer method (25), except that the lipid mixture contained 40% egg phosphatidylcholine (ePC), 10% dimyristoylphosphatidylcholine (DMPC), 25% phosphatidylglycerol (PG), and 25% cardiolipin (CL). This composition allows vesicle formation and maintains the 1 PG/1 CL lipid ratio (31, 32). DMPC increases membrane fluidity, allowing diffusion of external ATP into GUVs (33). Membranes were stained with 8% rhodamine (Avanti). A high concentration of inert macromolecules (50 mg/ml Ficoll) was added to mimic the conditions of the cytosol (34). ATP (2 mM) was added to the vesicles to trigger FtsA polymerization. Images were collected with a Leica TCS SP5 AOBS inverted confocal microscope with a 63 \times immersion objective

(NA, 1.4 to 0.6; oil HCX PL APO lambda-blue). Argon ion lasers were used for laser lines at 488 nm to excite Alexa 488. Images were processed with Adobe Photoshop CS5.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <http://mbio.asm.org/lookup/suppl/doi:10.1128/mBio.02221-14/-/DCSupplemental>.

Figure S1, TIF file, 2.3 MB.

Figure S2, TIF file, 1.6 MB.

Figure S3, TIF file, 2.6 MB.

ACKNOWLEDGMENTS

This work was funded by grant BIO2008-04478-C03 from Spanish Ministerio de Educación y Ciencia and by contract HEALTH-F3-2009-223431 (DIVINOCELL) from the European Commission (to both M.V. and G.R.). M.K. was supported by La Caixa Foundation International Fellowship Program (La Caixa/CNB).

We thank Orietta Massidda for critical reading of the manuscript. We also thank the Confocal Microscopy Facility of CNB-CSIC and CIB-CSIC and the Electron Microscopy Facility of CNB-CSIC for their technical support.

REFERENCES

- Rico AI, Krupka M, Vicente M. 2013. In the beginning, *Escherichia coli* assembled the proto-ring: an initial phase of division. *J. Biol. Chem.* **288**: 20830–20836. <http://dx.doi.org/10.1074/jbc.R113.479519>.
- Bork P, Sander C, Valencia A. 1992. An ATPase domain common to prokaryotic cell cycle proteins, sugar kinases, actin, and hsp70 heat shock proteins. *Proc. Natl. Acad. Sci. U. S. A.* **89**:7290–7294. <http://dx.doi.org/10.1073/pnas.89.16.7290>.
- Sánchez M, Valencia A, Ferrándiz MJ, Sander C, Vicente M. 1994. Correlation between the structure and biochemical activities of FtsA, an essential cell division protein of the actin family. *EMBO J.* **13**:4919–4925.
- van den Ent F, Löwe J. 2000. Crystal structure of the cell division protein FtsA from *Thermotoga maritima*. *EMBO J.* **19**:5300–5307. <http://dx.doi.org/10.1093/emboj/19.20.5300>.
- Krupka M, Rivas G, Rico AI, Vicente M. 2012. Key role of two terminal domains in the bidirectional polymerization of FtsA protein. *J. Biol. Chem.* **287**:7756–7765. <http://dx.doi.org/10.1074/jbc.M111.311563>.
- Lara B, Rico AI, Petruzzelli S, Santana A, Dumas J, Biton J, Vicente M, Mingorance J, Massidda O. 2005. Cell division in cocci: localization and properties of the *Streptococcus pneumoniae* FtsA protein. *Mol. Microbiol.* **55**:699–711. <http://dx.doi.org/10.1111/j.1365-2958.2004.04432.x>.
- Martos A, Monterroso B, Zorrilla S, Reija B, Alfonso C, Mingorance J, Rivas G, Jiménez M. 2012. Isolation, characterization and lipid-binding properties of the recalcitrant FtsA division protein from *Escherichia coli*. *PLoS One* **7**:e39829. <http://dx.doi.org/10.1371/journal.pone.0039829>.
- Szwedziak P, Wang Q, Freund SM, Löwe J. 2012. FtsA forms actin-like protofilaments. *EMBO J.* **31**:2249–2260. <http://dx.doi.org/10.1038/emboj.2012.76>.
- Rico AI, García-Ovalle M, Mingorance J, Vicente M. 2004. Role of two essential domains of *Escherichia coli* FtsA in localization and progression of the division ring. *Mol. Microbiol.* **53**:1359–1371. <http://dx.doi.org/10.1111/j.1365-2958.2004.04245.x>.
- Corbin BD, Geissler B, Sadasivam M, Margolin W. 2004. Z-ring-independent interaction between a subdomain of FtsA and late septation proteins as revealed by a polar recruitment assay. *J. Bacteriol.* **186**: 7736–7744. <http://dx.doi.org/10.1128/JB.186.22.7736-7744.2004>.
- Pichoff S, Shen B, Sullivan B, Lutkenhaus J. 2012. FtsA mutants impaired for self-interaction bypass ZipA suggesting a model in which FtsA's self-interaction competes with its ability to recruit downstream division proteins. *Mol. Microbiol.* **83**:151–167. <http://dx.doi.org/10.1111/j.1365-2958.2011.07923.x>.
- Pichoff S, Lutkenhaus J. 2005. Tethering the Z ring to the membrane through a conserved membrane targeting sequence in FtsA. *Mol. Microbiol.* **55**:1722–1734. <http://dx.doi.org/10.1111/j.1365-2958.2005.04522.x>.
- Yim L, Vandenbussche G, Mingorance J, Rueda S, Casanova M, Ruyschaert JM, Vicente M. 2000. Role of the carboxy terminus of *Escherichia coli* FtsA in self-interaction and cell division. *J. Bacteriol.* **182**:6366–6373. <http://dx.doi.org/10.1128/JB.182.22.6366-6373.2000>.
- Duman R, Ishikawa S, Celik I, Strahl H, Ogasawara N, Troc P, Löwe J, Hamoen LW. 2013. Structural and genetic analyses reveal the protein SepF as a new membrane anchor for the Z ring. *Proc. Natl. Acad. Sci. U. S. A.* **110**:E4601–E4610. <http://dx.doi.org/10.1073/pnas.1313978110>.
- Hu Z, Lutkenhaus J. 2003. A conserved sequence at the C-terminus of MinD is required for binding to the membrane and targeting MinC to the septum. *Mol. Microbiol.* **47**:345–355. <http://dx.doi.org/10.1046/j.1365-2958.2003.03321.x>.
- Salje J, van den Ent F, de Boer P, Löwe J. 2011. Direct membrane binding by bacterial actin MreB. *Mol. Cell* **43**:478–487. <http://dx.doi.org/10.1016/j.molcel.2011.07.008>.
- Hu Z, Gogol EP, Lutkenhaus J. 2002. Dynamic assembly of MinD on phospholipid vesicles regulated by ATP and MinE. *Proc. Natl. Acad. Sci. U. S. A.* **99**:6761–6766. <http://dx.doi.org/10.1073/pnas.102059099>.
- Pichoff S, Lutkenhaus J. 2007. Identification of a region of FtsA required for interaction with FtsZ. *Mol. Microbiol.* **64**:1129–1138. <http://dx.doi.org/10.1111/j.1365-2958.2007.05735.x>.
- Shiomi D, Margolin W. 2007. Dimerization or oligomerization of the actin-like FtsA protein enhances the integrity of the cytokinetic Z ring. *Mol. Microbiol.* **66**:1396–1415. <http://dx.doi.org/10.1111/j.1365-2958.2007.05998.x>.
- Shiomi D, Margolin W. 2008. Compensation for the loss of the conserved membrane targeting sequence of FtsA provides new insights into its function. *Mol. Microbiol.* **67**:558–569. <http://dx.doi.org/10.1111/j.1365-2958.2007.06085.x>.
- Zimmerberg J, Kozlov MM. 2006. How proteins produce cellular membrane curvature. *Nat. Rev. Mol. Cell Biol.* **7**:9–19. <http://dx.doi.org/10.1038/nrm1784>.
- Suefuji K, Valluzzi R, RayChaudhuri D. 2002. Dynamic assembly of MinD into filament bundles modulated by ATP, phospholipids, and MinE. *Proc. Natl. Acad. Sci. U. S. A.* **99**:16776–16781. <http://dx.doi.org/10.1073/pnas.262671699>.
- Massidda O, Nováková L, Vollmer W. 2013. From models to pathogens: how much have we learned about *Streptococcus pneumoniae* cell division? *Environ. Microbiol.* **15**:3133–3157. <http://dx.doi.org/10.1111/1462-2920.12189>.
- Rueda S, Vicente M, Mingorance J. 2003. Concentration and assembly of the division ring proteins FtsZ, FtsA, and ZipA during the *Escherichia coli* cell cycle. *J. Bacteriol.* **185**:3344–3351. <http://dx.doi.org/10.1128/JB.185.11.3344-3351.2003>.
- Cabré EJ, Sánchez-Gorostiaga A, Carrara P, Roperio N, Casanova M, Palacios P, Stano P, Jiménez M, Rivas G, Vicente M. 2013. Bacterial division proteins FtsZ and ZipA induce vesicle shrinkage and cell membrane invagination. *J. Biol. Chem.* **288**:26625–26634. <http://dx.doi.org/10.1074/jbc.M113.491688>.
- Hanahan D. 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**:557–580. [http://dx.doi.org/10.1016/S0022-2836\(83\)80284-8](http://dx.doi.org/10.1016/S0022-2836(83)80284-8).
- Miroux B, Walker JE. 1996. Over-production of proteins in *Escherichia coli*: mutant hosts that allow synthesis of some membrane proteins and globular proteins at high levels. *J. Mol. Biol.* **260**:289–298. <http://dx.doi.org/10.1006/jmbi.1996.0399>.
- Sambrook J, Fritsch EF, Maniatis T. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, New York, NY.
- Mingorance J, Rueda S, Gómez-Puertas P, Valencia A, Vicente M. 2001. *Escherichia coli* FtsZ polymers contain mostly GTP and have a high nucleotide turnover. *Mol. Microbiol.* **41**:83–91. <http://dx.doi.org/10.1046/j.1365-2958.2001.02498.x>.
- Uehara T, Dinh T, Bernhardt TG. 2009. LytM-domain factors are required for daughter cell separation and rapid ampicillin-induced lysis in *Escherichia coli*. *J. Bacteriol.* **191**:5094–5107. <http://dx.doi.org/10.1128/JB.00505-09>.
- Epanand RF, Savage PB, Epanand RM. 2007. Bacterial lipid composition and the antimicrobial efficacy of cationic steroid compounds (ceragenins). *Biochim. Biophys. Acta* **1768**:2500–2509. <http://dx.doi.org/10.1016/j.bbmem.2007.05.023>.
- Trombe MC, Lanéelle MA, Lanéelle G. 1979. Lipid composition of aminopterin-resistant and sensitive strains of *Streptococcus pneumoniae*. Effect of aminopterin inhibition. *Biochim. Biophys. Acta* **574**:290–300. [http://dx.doi.org/10.1016/0005-2760\(79\)90010-9](http://dx.doi.org/10.1016/0005-2760(79)90010-9).

33. Bessonov K, Harauz G. 2010. Molecular dynamics investigation of myelin basic protein stability on lipid membranes. *Studies by Undergraduate Researchers at Guelph* 4:79–86.
34. Jiménez M, Martos A, Vicente M, Rivas G. 2011. Reconstitution and organization of *Escherichia coli* proto-ring elements (FtsZ and FtsA) inside giant unilamellar vesicles obtained from bacterial inner membranes. *J. Biol. Chem.* 286:11236–11241. <http://dx.doi.org/10.1074/jbc.M110.194365>.
35. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera—a visualization system for exploratory research and analysis. *J. Comput. Chem.* 25:1605–1612. <http://dx.doi.org/10.1002/jcc.20084>.