

UDK 37

ISSN 2545 - 4439

ISSN 1857 - 923X

# INTERNATIONAL JOURNAL

Institute of Knowledge Management

# KNOWLEDGE



Scientific Papers

MEDICAL SCIENCES AND HEALTH

Vol. 38.4

View metadata, citation and similar papers at CORE.ac.uk

Indexed by UGD Academic Repository

rought to you by COBE

KIJ

Vol. 38

No. 4

pp. 639 - 1010

Skopje 2020

Global Impact & Quality Factor 1.822 (2017) <http://globalimpactfactor.com/knowledge-international-journal/>

KNOWLEDGE – International Journal  
Vol.38.4

---

**KNOWLEDGE**



**INTERNATIONAL JOURNAL**

**SCIENTIFIC PAPERS  
VOL. 38.4**

*March, 2020*



# KNOWLEDGE

## International Journal Scientific Papers Vol. 38.4

### ADVISORY BOARD

Vlado Kambovski PhD, Robert Dimitrovski PhD, Siniša Zarić PhD, Maria Kavdanska PhD, Venelin Terziev PhD, Mirjana Borota – Popovska PhD, Cezar Birzea PhD, Veselin Videv PhD, Ivo Zupanovic, PhD, Savo Ashtalkoski PhD, Zivota Radosavljević PhD, Laste Spasovski PhD, Mersad Mujevic PhD, Milka Zdravkovska PhD, Drago Cvijanovic PhD, Predrag Trajković PhD, Lazar Stosic PhD, Krasimira Staneva PhD, Nebojsa Pavlović PhD, Daniela Todorova PhD, Baki Koleci PhD, Lisen Bashkurti PhD, Zoran Srzentić PhD, Sinisa Opic PhD, Marija Kostic PhD

**Print:** GRAFOPROM – Bitola

**Editor:** IKM – Skopje

**Editor in chief**

Robert Dimitrovski, PhD

**KNOWLEDGE - International Journal Scientific Papers Vol. 38.4**

**ISSN 1857-923X** (for e-version)

**ISSN 2545 – 4439** (for printed version)

CONTENTS

XANTHOGRANULOMATOUS EPIDIDYMO-ORCHITIS AS A UNCOMMON AND RARE PSEUDONEOPLASTIC CONDITION .....	657
Ilbert Ademi.....	657
Adnan Vrajnko .....	657
Majlinda Ademi.....	657
CORRELATION BETWEEN THE POST 5ARI SYNDROME AND DEHIDROEPIANDROSTERONE LEVELS IN PATIENTS WITH BPH .....	663
Maja Sofronievska Glavinov .....	663
IMPACT OF COMPLEX PHARMACOLOGICAL AND NON-PHARMACOLOGICAL TREATMENT IN PATIENTS WITH CHRONIC PROSTATITIS/CHRONIC PELVIC PAIN SYNDROME .....	671
Galina Mratskova .....	671
Rumen Deliev .....	671
PREVENTION OF THROMBOPHILIA AND THE FOLLOWING PREGNANCY COMPLICATIONS .....	679
Anna Mihaylova .....	679
Petar Uchikov .....	679
Ekaterina Uchikova .....	679
Nikoleta Parahuleva.....	679
THE CORELATION BETWEEN FETO-PLACENTAL UNIT, FETAL BIOMETRY AND AMNIOTIC FLUID IN PREGNANT WOMEN WITH PREECLAMPSIA IN SECOND TRIMESTER .....	685
Pranvera Izairi.....	685
Kristina Skeparovska.....	685
Nevenka Velickova.....	685
LABORATORY DIAGNOSIS OF INFECTIONS CAUSED BY CHLAMYDIA TRACHOMATIS, EXPERIENCES FROM THE PHF CLINICAL HOSPITAL “M.D. TRIFUN PANOVSKI” – BITOLA FOR THE PERIOD FROM 2013 TO 2019 .....	691
Blagica Krsteska .....	691
Dino Karpicarov .....	691
Biljana Gjorgjeska .....	691
PROLONGED NEONATAL JAUNDICE – FREQUENCY AND CAUSE OF ITS OCCURRENCE...697	697
Tatyana Itova .....	697
Victorya Atanasova .....	697
ANATOMICAL CHARACTERISTICS OF THE VERTEBRAL ARTERY USING CT ANGIOGRAPHY .....	701
Svetlana Jovevska.....	701
Merima Idrizovikj.....	701
APPLICATION OF SPECIALIZED METHODS OF PFYSIOTHERAPY ON THE FLEXIBLE MUSCLE GROUP IN PATIENTS WITH COXARTROSIS.....	705
Zhulieta Gerenova .....	705
ANKYLOSING SPONDYLITIS (MORBUS BECHTEREW).....	711
Merima Idrizovikj.....	711
Svetlana Jovevska.....	711
METHODS FOR NEUROMUSCULAR REEDUCATION FOR REDUCING MOTOR DEFICIT AND MUSCLE SPASM IN PEOPLE WITH CEREBRAL PALSY.....	717
Lence Nikolovska.....	717
Zana Markova.....	717
KINESITHERAPY EFFECT IN PATIENTS WITH SHOULDER BURSITIS .....	723
Danche Vasileva .....	723
Suat Berisha.....	723

---

EPIDEMIOLOGICAL CHARACTERISTICS OF ACUTE VIRAL HEPATITIS TYPE C IN MACEDONIA .....	729
Velo Markovski .....	729
Vlatko Cvetanovski .....	729
EPIDEMIOLOGICAL CHARACTERISTICS OF MEASLES IN MACEDONIA DURING THE PERIOD 2010-2019.....	735
Kolevska Natasha .....	735
Bosevska Golubinka .....	735
Zdravkovska Milka.....	735
Markovski Velo .....	735
NURSING PROCESS IN TRAUMATIC CHEST INJURY OF THE HEART - A CLINICAL CASE..	741
Lilyana Stancheva.....	741
Ivanka Stambolova .....	741
Tsvetelina Evtimova .....	741
HEALTH-RELATED QUALITY OF LIFE TO PEOPLE WITH CHRONIC RESPIRATORY AND CARDIOVASCULAR DISEASES .....	747
Katya Popova.....	747
Galina Terzieva.....	747
Monika Obreykova .....	747
SELF-CONTROL OF DISEASE IN PATIENTS WITH ARTERIAL HYPERTENSION .....	753
Lyubomir Marinov .....	753
Pavlinka Dobrilova .....	753
PATIENTS WITH CHRONIC DISEASE AWARENESS ABOUT THE OPPORTUNITIES FOR PREVENTIVE INTERVENTIONS .....	759
Pavlinka Dobrilova .....	759
PHARMACOTHERAPY APPROACH IN KIDNEY CANCER THERAPY .....	767
Sanela Andova .....	767
Tijana Serafimovska .....	767
Milka Zdravkovska.....	767
Trajan Balkanov .....	767
Marija Darkovska Serafimovska .....	767
ANTHROPOLOGY IN ORTHODONTICS .....	773
Natasa Toseska-Spasova.....	773
Natasha Stavreva .....	773
Biljana Dzipunova .....	773
DISCOLORATION OF NANO COMPOSITE RESIN BY EVERYDAY DRINKS: IN VITRO STUDY .....	779
Aleksandar Andreevski.....	779
Ivona Kovacevska.....	779
METHODS AND TECHNIQUES FOR BLEACHING VITAL TEETH .....	785
Natasa Longurova.....	785
Ivona Kovacevska.....	785
PROSTHODONTIC SOLUTION OF FLABBY RIDGES USING WINDOW IMPRESSION TECHNIQUE.....	791
Natasha Stavreva .....	791
Natasha Tosheska Spasova .....	791
THE DIGITAL IMPRESSION AS AN INITIAL UNIT OF MODERN DENTAL MEDICINE - LITERATURE REVIEW .....	797
Dobromira Shopova.....	797
Miroslava Yordanova .....	797
Svetlana Yordanova.....	797

PROSTHETIC TREATMENT WITH PLATE IN PATIENT WITH MAXILLARY RESECTION .....	803
Ivan Gerdzhikov .....	803
RISKS DURING ORTHODONTIC TREATMENT .....	807
Ana Radeska-Panovska .....	807
DRUG SUPPLY CHAIN SAFETY - FIGHT AGAINST COUNTERFEIT DRUGS .....	813
Elena Drakalska Sersemova .....	813
Bistra Angelovska.....	813
Dijana Miceva.....	813
Elmedina Sakir .....	813
MONITORING OF THE RENAL FUNCTION IN PIROXICAM TREATED PATIENTS WITH HEADACHE.....	819
Drita Yzeiri Havziu .....	819
Biljana Gjorgjeska .....	819
Edita Alili Idrizi.....	819
Gjylaj Alija .....	819
Sihana Lika .....	819
Merita Dauti.....	819
SESAME ALLERGY: A GROWING FOOD ALLERGY OF GLOBAL PROPORTIONS? .....	825
Verica Jakjimoska.....	825
Biljana Gjorgjeska .....	825
REHABILITATION IN PATIENTS WITH TYPE 2 DIABETES .....	831
Alexandra Yankova Maleva .....	831
DEPRESSION AND DIABETES .....	837
Aneta Spasovska Trajanovska .....	837
Danijela Janicevic Ivanovska .....	837
Zora Mitic.....	837
FINAL ANALYSIS OF THE RESULTST IN MAN WITH MULTIPLE SCULPROSIS AFTER A PHYSIOTHERAPEUTIC PROGRAM .....	843
Inna Ivanova .....	843
PSYCHOSOMATIC DISORDERS IN ADOLESCENTS .....	849
Lubomira Dimitrova .....	849
CHEMICAL CORROSIVE DISEASES OF THE EYE .....	853
Gazepov Strahil .....	853
Viktorija Micevska .....	853
Alen Gorgiev .....	853
Slavena Stojkova .....	853
HEALTH BEHAVIOR AND PARENTS' AWARENESS HOW TO PROTECT THE CHILDREN'S EYE HEALTH.....	859
Krassimira Dimitrova .....	859
Christina Grupcheva .....	859
CONTROL MATERIALS-MEDICAL DEVICES FOR INCREASING THE QUALITY OF CLINICAL LABORATORY ANALYSES .....	865
Denitsa Trancheva .....	865
LABORATORY QUALITY MANAGEMENT.....	871
Katerina Tosheska-Trajkovska .....	871
MEDICAL SUPPLIES FOR CHEMICAL ACCIDENTS AND TERRORIST ATTACKS WITH CHLORINE IN THE PHARMACY NETWORK IN VARNA .....	877
Maria Panteleeva .....	877
Nikolina Radeva .....	877
Rostislav Kostadinov .....	877

AWARENESS AND ATTITUDES OF NURSING STAFF REGARDING TO EVALUATION IN HOSPITAL .....	883
Daniela Stankeva .....	883
Galina Tchaneva .....	883
Milena Stoycheva .....	883
HEALTH CARE CENTERS IN BULGARIA – NECESSITY AND CHALLENGES .....	889
Ivanka Stambolova .....	889
Stefan Stambolov .....	889
NURSING CARE ASSESSMENT AND NURSES' PROFESSIONAL ATTITUDES .....	895
Galya Yankova .....	895
CREATION, DEVELOPMENT AND PERSPECTIVES OF NURSING EDUCATION IN BULGARIA .....	903
Silviya Mladenova .....	903
THE ROLE OF THE FAMILY NURSE IN THE IMPLEMENTATION OF IMMUNIZATION AND EDUCATION OF THE POPULATION IN THE MUNICIPALITY OF KOCANI FOR THE PERIOD 2015-2019 .....	909
Gordana Panova .....	909
Christina Janeva .....	909
STRESS AND CRISIS IN THE FAMILY IN CHILDREN WITH AUTISM AND DOWN SYNDROME .....	915
Plamen Tzokov .....	915
TRAINING AND ADDITIONAL QUALIFICATIONS OF SPECIALISTS WORKING IN THE FIELD OF CHILD REHABILITATION IN ORDER TO IMPROVE THE QUALITY OF THE SERVICES OFFERED .....	923
Stela Marinkova .....	923
Antoaneta Terzieva .....	923
ROLE OF THE PHYSICAL ACTIVITY AND REHABILITATION IN THE PROCESS OF AGING .....	929
Vanina Mihaylova .....	929
Ina Ivanova .....	929
Adolf Alakidi .....	929
Kristina Kilova .....	929
Mariana Liochkova .....	929
TREATMENT OF PATIENTS WITH FIBROMYALGIA WITH METHODS OF TRADITIONAL CHINESE MEDICINE .....	937
Lence Nikolovska .....	937
Bojan Radinski .....	937
THE ROLE OF THE PATRONAGE NURSE IN CONDUCTING AND EDUCATING THE ELDERLY .....	943
Hristina Trajkovska .....	943
Gordana Panova .....	943
THE CHALLENGES OF THE ELECTROMAGNETIC RADIATION- ARE WE READY FOR THEM - A MINI REVIEW .....	949
Vassilka Ilieva .....	949
TELEMEDICINE, TELEHEALTHCARE AND TELEHEALTH – LITERATURE REVIEW .....	957
Stanka Uzunova .....	957
Kristina Kilova .....	957
MEDICAL DEVICES AND ADAPTED MEANS IN OCCUPATIONAL THERAPY PROFESSION .....	965
Yoana Negrova .....	965
SOCIAL SIGNIFICANT ASPECTS OF COMBUSTION INJURY .....	971
Anushka Dimitrova .....	971

ROLE OF NON-PROFIT ORGANIZATIONS FOR PROTECTION OF THE QUALITY OF LIFE AMONG PEOPLE WITH ONCOLOGICAL DISEASES .....	979
Petya Stefanova .....	979
Zlatina Lecheva .....	979
APPLICATION OF THE PARETO PRINCIPLE IN THE CONTRACTAL PROCESS FOR ONCOLOGICAL DISEASE .....	985
Darina Mineva .....	985
ADMINISTRATIVE CONTROL FOR THE GUARANTEE OF MEDICAL DEVICES AT THEIR PURCHASE AS CONSUMER GOODS IN THE REPUBLIC OF BULGARIA.....	991
Mikael Chakaldziyan.....	991
LEGAL MEANS FOR PROTECTION FROM MEDICAL ERRORS .....	997
Momchil Mavrov.....	997
COMMUNICATION PROBLEMS IN A MEDICAL ORGANIZATION .....	1005
Svetlana Radeva .....	1005



---

## DEPRESSION AND DIABETES

**Aneta Spasovska Trajanovska**

Psychiatric Hospital, Skopje, N. Macedonia

**Danijela Janicevic Ivanovska**

University clinic of Clinical biochemistry, Skopje, N. Macedonia, University “Goce Delchev”, Faculty of  
Medical Sciences, Stip, N. Macedonia, djanicevic@yahoo.com,

**Zora Mitic**

Private Practice in Psychiatry D-r Zora Mitik, Skopje, N. Macedonia

**Abstract: Introduction:** Some researchers suggest that people with depression are more likely to develop diabetes. The **AIM** of this study is determine the effects of SSRIs antidepressant in depressive patients with diabetes mellitus. **Methods:** The study was designed as observation cross-section including 80 depressive patients with diabetes, observed as two groups. The first group was presented patients without antidepressant medication in them (the patients who starting in the treatment), and the second group were patients with antidepressant therapy (sertraline). Depression status was assessed using BDI. Quantitative determination of glucose level in serum was performed with enzymatic reference method with hexokinase. **Results :** There was significant differences in distribution on according the level of blood glucose in observed groups. . In the first group 58% of patients have high level of blood glucose but in second group only 20% of them.. Data obtained in the first groups show significant positive correlation between score of BDI and blood glucose level.  $P=0,003$ . The results in examination groups show positive correlation between blood glucose level and socio-demographic and clinical parameters. **Conclusion :** The results obtained in our study showed high blood glucose levels in depressant patients but treatment with sertraline was diminished this level. So screening this condition is important, because that patients should be treated differently.

**Keywords:** patients, depression, diabetes mellitus, therapy

### 1. INTRODUCTION

According to the International Diabetes Federation diabetes is one of the largest global health emergencies of the 21st century (1). Also depression is a common and very serious disease with lifetime prevalence ranging from approximately 11% and low-income countries to 15% in high-income countries (2). There is evidence that prevalence of depression is moderately increased in pre diabetic patients and in undiagnosed diabetic patients, and markedly increased in the previously diagnosed diabetic patients compared to normal glucose metabolism individuals (3). So some studies show that depression occurrence is two to three times higher in people with diabetes mellitus, the majority of the case remaining under-diagnosed. Some researchers suggest that this could be due to diabetes metabolic effects on brain as well as the toll day to day management can take. It is also possible that people with depression are more likely to develop diabetes (4). Although the psychological burden of diabetes may contribute to depression, this explanation does not fully explain the relationship between these 2 conditions. Both conditions may be driven by shared underlying biological and behavioral mechanisms, such as hypothalamic-pituitary-adrenal axis activation, inflammation, sleep disturbance, inactive lifestyle, poor dietary habits and environmental and cultural risk factors. (5). Also chronic stress activates the hypothalamus-pituitary-adrenal axis and the sympathetic nervous system, increasing the production of cortisol in the adrenal cortex and the production of adrenalin and noradrenalin in the adrenal medulla (6). Chronic hypercortisolemia and prolonged SNS activation promote insulin resistance, visceral obesity and lead to metabolic syndrome and diabetes mellitus (7). On the other hand it is thought that alteration in brain chemistry tied to diabetes may be related to the development of depression. For example, damage resulting from diabetic neuropathy or blocked blood vessels in the brain may contribute to the development of depression in people with diabetes (4). Conversely, changes in the brain due to depression may cause an increased risk for diabetes complication. Some study found that people with diabetes and depression often have higher blood sugar levels (2). Additionally the results of another study suggested that people who have both conditions are 82 percentage more likely to experience a heart attack (5). On the other hand poor diabetes management can also prompt symptoms similar to those of depression. If the blood sugar is too high or too low, people may experience increased feeling of anxiety, restlessness, or low energy. Low blood sugar levels can also cause people to feel shaky and sweaty, which are symptoms similar to anxiety (3). It is also possible that people with depression are more likely to develop diabetes. Because of this, it is recommended that people who have a history of depression be screened for diabetes(7).

## 2. AIM

The AIM of this study is determine the effects of SSRIs- antidepressant in depressive patients with diabetes mellitus

## 3. MATERIALS AND METHODS

The patients of this cross-section study were examination in the Private Psychiatric Ambulance "Zora Mitic" and Psychiatric Hospital Skopje during 2019 years. The study was approved by the regional ethics committee for research. All participants gave written consent. We evaluated prevalence of diabetes mellitus in 80 patients with depression. The patients were observed in two groups. The first groups was presented with patients without antidepressant medication in them only detection the depression and diabetes mellitus (the patients who starting in the treatment). The second group consisted patients with antidepressant therapy (SSRIs\_\_sertraline) and diabetes mellitus. For the collection of data we use medical documentation (history of illness and outpatient diary) The socio-demographic data were collected by structural questionnaire . All the patients followed criteria for depression. Criteria for elimination was: chronic liver or renal diseases, HIV diseases or active infection .All the patients there followed by routine medical examination, with included blood chemistry , urinoanalysis to determine the serology for hepatitis and HIV infection. . Blood samples of patients were obtained the morning between 9,00-11,00 H in the University Institute of Clinical biochemistry Skopje. The glucose plasma levels were assayed using the Enzymatic reference method with hexokinase. Hexokinase catalyzes the phosphorylation of glucose to glucose-6-phosphate by ATP. Glucose-6-phosphate dehydrogenase oxidizes glucose-6- phosphate in the presence of NADP to gluconate-6-phosphate. No other carbohydrate is oxidized. The rate of NADPH formation during the reaction is directly proportional to the glucose concentration and is measured photometrically. The normal range of glucose level was 3,5-6,1 mmol/l. To determine exactly results of plasma glucose I levels we use QS of the BIORAD . Depression was assays with Beck Depression Inventory (BDI) considerate with 21 items. The score BDI  $\geq 10$  determine depressive symptoms. The results were an analyzed statistically using: descriptive methods, t-test for independent simplex and the Pearson coefficient of linear correlation The statistical test were considered significant at the level  $p \leq 0,05$ .

## 4. RESULTS

Our study included two groups of patients with depression (BDI  $\geq 10$ ). There was significant differences in distribution on according the level of blood glucose in observed groups. (Table 1). In the first group 58% of patients have high level of blood glucose but in second group only 20% of them.

**Table 1. Distribution of depressive patients in examination groups according their blood glucose levels**

Level of blood glucose	Group 1 (N=41)		Group 2 (N=39)		P level
	N	%	N	%	
normal	18	42	30	80	sig
high	23	58	9	20	
low	0	0	0	0	

Data obtained in the first groups show significant positive correlation between score of BDI and blood glucose level. (Table 2)

**Table 2. Correlation between score of BDI and blood glucose level in examination groups.**

Parameters	p-spearmen koeficient	P valvule
Score BDI/ Blood glucose level		
Group A	P=0,365	0,003
Group B	P= 0,290	0,098

The results in examination groups show positive correlation between blood glucose level and socio-demographic and clinical parameters. (Table3,4).

**Table 3. Correlation of the socio-demographic and clinical parameters with blood glucose level in the first examination group.**

Parameters	p-Spearman coefficient	p valvule
Gender (female)	0,350	0,002
Age	0,070	0,082
Smoking	0,274	0,062
Physical activity	0,211	0,003
Stress	0,075	0,081
BMI	0,325	0,009
Marital status (single)	0,071	0,089
Education	0,285	0,095

**Table 4. Correlation of the socio-demographic and clinical parameters with blood glucose level in depressant patients after three mount of SSRI (sertraline) therapy (second group).**

Parameters	p-Spearman coefficient	p valvule
Gender(female)	0,356	0,032
Age	0,295	0,096
Smoking	0,552	0,004
Physical activity	0,301	0,061
Stress	0,215	0,095
BMI	0,604	0,002
Marital status (single)	0,065	0,074
Education	0,360	0,002

## 5. DISCUSSION

The association between mental illness and diabetes has recognized for many years (8). In the 17<sup>th</sup> century, Thomas Willis, the famous anatomic and founding member of the Royal Society, described how " diabetes is a consequence of prolonged sorrow" (9) . In our study we got that 58% of depressant patients in the first group and only 20 % in second group have high level of blood glucose . This results correlate with another study who reported that twenty-nine percent of male and 30,5% of female participants with depression had diabetes mellitus (10) . This results also correlate with another studies who suggested that rates of depression are also elevated in either type 1 or type 2 diabetes with prevalence rates ranging from 9%-26% (11). Significant depressive symptoms affect approximately 1 in 4 adults with type 1 and type 2 diabetes, where a formal diagnosis of depressive disorders is made in approximately 10%-15% of people with diabetes (12). As observed by Thomas Willis, epidemiologic studies have demonstrated that the association between depression and diabetes is be-directional (13,14). A meta-analysis of 9 cohort studies found that adults with depression had 37% increased risk of developing type 2 diabetes (15) after according for factors common to both disorders including sex, body mass index, and poverty. Also both disorders some researches reported that common especially in women (10). This findings correlate with results in our study we got that 65 % in the first groups and 55% in the second groups of patients were women.

In both examination groups we got positive correlation between blood glucose level and socio-demographic and clinical parameters (smoking, physical activity, stress, BMI, marital satus and education (table3,4). This results correlate with another studies who reported that in additional , there a number of diabetes specific risk factors associated with depression.General population risk factor for depression, including female sex, marital status, childhood, adversity and social deprivation also apply to people with diabetes. (16,17). Lifestyle factor are hypothesized to play a role in priming or reinforcing the co morbidity of depression and diabetes. For example, people with depression are more likely to be sedentary and eat diets that are rich in saturated fats and refined sugar while avoiding fruit and vegetables , which may contribute to the risk of developing diabetes (18,19).Both conditions (depression and diabetes ) is also associated with premature mortality through a range of physical conditions (20) blood pressure lipids, body mass index, tobacco, living alone and education level (21).This finding correlate with our results. In our study we also got positive correlation between diabetess mellitus(blood glucose levels) and marital status, smoking status, fizical activity, body mass index.,stressful lifestyle. Lifestyle change such as increased physical activity or exercise , dietary modification, adequate relaxation, sleep and social interaction, use of mindfulness-based meditation techniques, and the reduction of recreational substances such as nicotine, drugs and alcohol already proved their benefits in the improvement of depression as well as diabetes

Both psychological therapies and antidepressant medication should be determined on an individual basis. As poor

metabolic control, low rates of blood glucose self-monitoring, and diabetes complications all predict inadequate response to depression treatment, an equal emphasis on both diabetes and depression is needed (18). Until recently, people with diabetes were specifically excluded from trials of the depression treatment and so consequently, there are relatively few studies examining antidepressant and psychotherapy treatment of depression in this population. Nevertheless studies published over the last decade have clearly indicated that treating both conditions is effective (22). There is preliminary evidence to suggest the web-based psychological therapies may also be effective in treating depression in people with diabetes but may have limited effects on glycemic outcomes (15,16). The best response to psychological interventions occurs when these are combined with diabetes education, to provide diabetes self-management skills as well as the psychological support to use these affectively (22). Some antidepressants, including mirtazapine, paroxetine and some tricyclic antidepressants, are associated with significant weight gain and are less suitable because of worsening insulin resistance and potentially glycemic control (21). Furthermore, unlike SSRIs, it does not appear to worsen sexual function and therefore may have advantages for people with diabetes (22). Sertraline may have specific advantages for glycemic control (20). So in our study also we got results which show that treatment of diabetes in depressed patients with antidepressant sertraline diminished the percentage of depressed patients with high blood glucose level. Only 20% of patients in second group (treated with sertraline) have high blood glucose level. Psychiatric intervention may be necessary in addition to lifestyle change to prevent the exponential increase in the occurrence of diabetes. In addition a common approach including psychiatric treatment in diabetes care may be necessary to achieve improved glycemic control in this population. (2).

## 6. CONCLUSION

The results obtained in our study showed high blood glucose levels in depressed patients but treatment with sertraline was diminished this level. So screening this condition is important, because that patients should be treated differently than those who have normal blood glucose level. Psychiatric interventions (psychotherapy, social support) and treatment with SSRIs- sertraline may prevent the exponential increase in the occurrence of diabetes mellitus in those patients.

## REFERENCES

- International Diabetes Federation .IDF Diabetes. 7 ed. Brussels, Belgium: International Diabetes Federation; 2015. <http://www.diabetesatlas.org>. [Google Scholar]
- Bromet, E., Andrade, L.H., Hwang, I., Sampson, N.A., Alonso, J., de G, de GR, Demyttenaere, K., Hu, C., Iwata N, Karam AN, Kaur J, Kostyuchenko S, Lepine JP, Levinson D, Matschinger H, Mora ME, Browne MO, Posada-Villa J, Viana MC, Williams DR, Kessler RC. (2011). Cross-national epidemiology of DSM-IV major depressive episode. *BMC Med.* 9:90.
- Grigsby AB, Anderson RJ, Freedland KE, Clouse RE, Lustman PJ. (2002). Prevalence of anxiety in adults with diabetes: a systematic review. *J Psychosom Res.* 53:1053–1060.
- Gonzalez JS, Peyrot M, McCarl LA, Collins EM, Serpa L, Mimiaga MJ, Safren SA. (2008). Depression and diabetes treatment nonadherence: a meta-analysis. *Diabetes Care.* 31:2398–2403.
- Baumeister H, Hutter N, Bengel J, Harter M. (2011). Quality of life in medically ill persons with comorbid mental disorders: a systematic review and meta-analysis. *Psychother Psychosom.* 80:275–286.
- Egede LE, Nietert PJ, Zheng D. (2005). Depression and all-cause and coronary heart disease mortality among adults with and without diabetes. *Diabetes Care.* 28:1339–1345.
- Mezuk B, Eaton WW, Albrecht S, Golden SH. (2008). Depression and type 2 diabetes over the lifespan: a meta-analysis. *Diabetes Care.* 31:2383–2390.
- Rubin RR, Ma Y, Marrero DG, Peyrot M, Barrett-Connor EL, Kahn SE, Haffner SM, Price DW, Knowler WC. (2008). Elevated depression symptoms, antidepressant medicine use, and risk of developing diabetes during the diabetes prevention program. *Diabetes Care.* 31:420–426.
- Kyrou I, Tsigos C. (2009). Stress hormones: physiological stress and regulation of metabolism. *Curr Opin Pharmacol.* 9:787–793.
- Holt RI. (2011). Undoing Descartes: integrating diabetes care for those with mental illness. *Pract Diabetes.* 28:270–5.
- Sartorius N, Cimino L. (2012). The Dialogue on Diabetes and Depression (DDD): origins and achievements. *J Affect Disord.* 142(Suppl):S4–7.
- Reynolds KA, Helgeson VS. (2011). Children with diabetes compared with peers: Depressed? Distressed? A meta-analytic review. *Ann Behav Med.* 42:29–41.

- Mezuk B, Eaton WW, Albrecht S, Golden SH. (2008). Depression and type 2 diabetes over the lifespan: a meta-analysis. *Diabetes Care*. 31:2383–90.
- Golden SH, Lazo M, Carnethon M, Bertoni AG, Schreiner PJ, Diez Roux AV, et al. (2008). Examining a bidirectional association between depressive symptoms and diabetes. *JAMA*. 299:2751–9.
- Knol MJ, Twisk JW, Beekman AT, Heine RJ, Snoek FJ, Pouwer F. (2006). Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. *Diabetologia*. 49:837–45]
- Hermanns N, Kulzer B, Krichbaum M, Kubiak T, Haak T. (2005). Affective and anxiety disorders in a German sample of diabetic patients: prevalence, comorbidity and risk factors. *Diabetes Med*. 22:293–300. ]
- Li C, Ford ES, Strine TW, Mokdad AH. (2008). Prevalence of depression among U.S. adults with diabetes: findings from the 2006 behavioral risk factor surveillance system. *Diabetes Care*. 31:105–7.
- McMartin SE, Jacka FN, Colman I. (2013). The association between fruit and vegetable consumption and mental health disorders: evidence from five waves of a national survey of Canadians. *Prev Med*. 56:225–30.
- Weyerer S. (1992). Physical inactivity and depression in the community. Evidence from the Upper Bavarian Field Study. *Int J Sports Med*. 13:492–6.
- Aikens JE, Perkins DW, Piette JD, Lipton B. (2008). Association between depression and concurrent Type 2 diabetes outcomes varies by diabetes regimen. *Diabetes Med*. 25:1324–9.
- O'Connor PJ, Narayan KM, Anderson R, Feeney P, Fine L, Ali MK, et al. (2012). Effect of intensive vs standard blood pressure control on depression and health-related quality of life in type 2 diabetes: the ACCORD trial. *Diabetes Care*. 35:1479–81. [
- van der Feltz-Cornelis CM, Nuyen J, Stoop C, Chan J, Jacobson AM, Katon W, et al. (2010). Effect of interventions for major depressive disorder and significant depressive symptoms in patients with diabetes mellitus: a systematic review and meta-analysis. *Gen Hosp Psychiatry*. 32:380–95.
- Serretti A, Mandelli L. (2010). Antidepressants and body weight: a comprehensive review and meta-analysis. *J Clin Psychiatry*. 71:1259–72.
- Sayuk GS, Gott BM, Nix BD, Lustman PJ. (2011). Improvement in sexual functioning in patients with type 2 diabetes and depression treated with bupropion. *Diabetes Care*. 34:332–4.