



Optimal boundary control for steady motions of a self-propelled body in a Navier-Stokes liquid

Toshiaki Hishida, Ana Leonor Silvestre, Takéo Takahashi

► To cite this version:

Toshiaki Hishida, Ana Leonor Silvestre, Takéo Takahashi. Optimal boundary control for steady motions of a self-propelled body in a Navier-Stokes liquid. *ESAIM: Control, Optimisation and Calculus of Variations*, EDP Sciences, 2020, 26 (92), 10.1051/cocv/2020073 . hal-02502289

HAL Id: hal-02502289

<https://hal.archives-ouvertes.fr/hal-02502289>

Submitted on 9 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Optimal boundary control for steady motions of a self-propelled body in a Navier-Stokes liquid

Toshiaki Hishida^{*1}, Ana Leonor Silvestre^{†2}, and Takéo Takahashi^{‡3}

¹Graduate School of Mathematics, Nagoya University, Nagoya 464-8602 Japan
`hishida@math.nagoya-u.ac.jp`

²CEMAT and Department of Mathematics, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal
`Ana.Silvestre@math.tecnico.ulisboa.pt`

³Université de Lorraine, CNRS, Inria, IECL, F-54000 Nancy, France
`takeo.takahashi@inria.fr`

March 9, 2020

Abstract

Consider a rigid body $\mathcal{S} \subset \mathbb{R}^3$ immersed in an infinitely extended Navier-Stokes liquid and the motion of the body-fluid interaction system described from a reference frame attached to \mathcal{S} . We are interested in steady motions of this coupled system, where the region occupied by the fluid is the exterior domain $\Omega = \mathbb{R}^3 \setminus \mathcal{S}$. This paper deals with the problem of using boundary controls v_* , acting on the whole $\partial\Omega$ or just on a portion Γ of $\partial\Omega$, to generate a self-propelled motion of \mathcal{S} with a target velocity $V(x) := \xi + \omega \times x$ and to minimize the drag about \mathcal{S} . Firstly, an appropriate drag functional is derived from the energy equation of the fluid and the problem is formulated as an optimal boundary control problem. Then the minimization problem is solved for localized controls, such that $\text{supp } v_* \subset \Gamma$, and for tangential controls, i.e. $v_* \cdot n|_{\partial\Omega} = 0$, where n is the outward unit normal to $\partial\Omega$. We prove the existence of optimal solutions, justify the Gâteaux derivative of the control-to-state map, establish the well-posedness of the corresponding adjoint equations and, finally, derive the first order optimality conditions. The results are obtained under smallness restrictions on the objectives $|\xi|$ and $|\omega|$ and on the boundary controls.

Keywords: 3-D Navier-Stokes equations; Exterior domain; Rotating body; Self-propelled motion; Boundary control; Drag reduction.

2010 Mathematics Subject Classification. 76D05, 49K21, 76D55, 49J21

Contents

1	Introduction	2
2	Notation and auxiliary results	3

^{*}T. Hishida is partially supported by Grant-in-Aid for Scientific Research, 18K03363, from JSPS.

[†]A. L. Silvestre acknowledges the financial support of the Portuguese FCT - Fundação para a Ciência e a Tecnologia, through the projects UIDB/04621/2020 and UIDP/04621/2020 of CEMAT/IST-ID.

[‡]T. Takahashi is partially supported by the project IFSMACS ANR-15-CE40-0010, financed by the French Agence Nationale de la Recherche.

3	L^2-estimate of the solution to a linearized problem	6
4	The state system and the cost functional	13
5	Existence of optimal controls	17
6	Regularity of the control-to-state mapping	19
7	Necessary first order conditions for an optimal control	23
7.1	Introduction of the Lagrangian	23
7.2	Well-posedness of the adjoint system	27
7.3	Optimality condition	29

1 Introduction

Consider a rigid body \mathcal{S} moving by self-propulsion through an infinitely extended incompressible viscous fluid \mathcal{F} . This means that the total net force and torque, external to the system $\{\mathcal{S}, \mathcal{F}\}$, acting on \mathcal{S} , are identically zero. Since the shape of the body is constant, the self-propelled motion of \mathcal{S} through \mathcal{F} is due to the boundary values for the velocity of the system at the boundary of \mathcal{S} , which are prescribed relative to \mathcal{S} . For instance, the propulsion may be produced by drawing fluid inwards across portions of the boundary and by expelling it from others, or by the tangential motion of certain portions of the boundary, as by belts. The propulsion of jet planes and submarines or of minute organisms like ciliates and flagellates can be considered of this nature.

In this context, the set of equations describing the motion of $\{\mathcal{S}, \mathcal{F}\}$, in a reference frame attached to \mathcal{S} , is

$$-\operatorname{div} \sigma(v, p) + (v - V) \cdot \nabla v + \omega \times v = 0 \quad \text{in } \Omega \tag{1.1}$$

$$\operatorname{div} v = 0 \quad \text{in } \Omega \tag{1.2}$$

$$v = V + v_* \quad \text{on } \partial\Omega \tag{1.3}$$

$$\lim_{|x| \rightarrow \infty} v(x) = 0 \tag{1.4}$$

$$m\xi \times \omega + \int_{\partial\Omega} [-\sigma(v, p)n + (v_* \cdot n)(v_* + V + \omega \times x)] \, d\gamma = 0 \tag{1.5}$$

$$(I\omega) \times \omega + \int_{\partial\Omega} x \times [-\sigma(v, p)n + (v_* \cdot n)(v_* + V + \omega \times x)] \, d\gamma = 0 \tag{1.6}$$

where the quantities $v = v(x)$ and $p = p(x)$ represent, respectively, the velocity field and the pressure of the liquid and

$$V(x) = \xi + \omega \times x, \quad x \in \mathbb{R}^3, \tag{1.7}$$

represents the velocity of the solid, as seen by an observer attached to \mathcal{S} . We assume that the density of the fluid is constant and equal to 1. Moreover, we denote by $\sigma(v, p)$ the Cauchy stress tensor defined by

$$\sigma(v, p) := 2D(v) - p\mathbb{I}_3,$$

where the viscosity of the fluid is also assumed equal to 1, \mathbb{I}_3 is the 3×3 identity matrix and $D(v)$ is the symmetric part of the gradient,

$$D(v) := \frac{1}{2} \left(\nabla v + (\nabla v)^\top \right).$$

Due to the incompressibility condition (1.2), we can use the relation $\operatorname{div} \sigma(v, p) = \Delta v - \nabla p$ in equation (1.1). Moreover, in equation (1.1) and in what follows, for sufficiently regular vector fields u and v , $u \cdot \nabla v$ is the vector field with components $(u \cdot \nabla v)_i = \sum_{j=1}^3 u_j \frac{\partial v_i}{\partial x_j}$. The set \mathcal{S} representing the rigid body is a compact simply connected set, with non empty interior, so that $\Omega = \mathbb{R}^3 \setminus \mathcal{S}$ is a three-dimensional exterior domain. In (1.5)–(1.6) and in what follows, n denotes the outward unit normal to $\partial\Omega$, which is well defined provided $\partial\Omega$ is locally Lipschitzian. Throughout this paper, nevertheless, $\partial\Omega$ is assumed to be of class C^3 (which is

in fact needed in Theorem 4.1.) We will assume that the center of gravity of the rigid body is located at the origin, that is $\int_{\mathcal{S}} x \, dx = 0$, and define the mass and inertia matrix by

$$m := \int_{\mathcal{S}} dx, \quad I := \int_{\mathcal{S}} (|x|^2 \mathbb{I}_3 - x \otimes x) \, dx,$$

where we are assuming that the density of the rigid body is constant and equal to 1. The boundary values v_* represent the thrust velocity, responsible for the motion of \mathcal{S} . The model (1.1)–(1.6) is inspired by Galdi [7, 8, 9], the equation (1.5) having been obtained as the net force exerted by the fluid on the solid and the equation (1.6) being the corresponding balance of torques.

While moving through the fluid, \mathcal{S} will experience a drag, i.e., a net force in the direction of flow, due to the pressure and shear stress on the surface of \mathcal{S} , which tends to slow down its motion. In this paper, our aim is to use the Dirichlet boundary data v_* not only to self-propell \mathcal{S} , but also to minimize the work needed to overcome the drag exerted by the fluid on \mathcal{S}

$$W(v, p) := \int_{\partial\Omega} v \cdot (\sigma(v, p)n) \, d\gamma = \int_{\partial\Omega} (v_* + V) \cdot (\sigma(v, p)n) \, d\gamma \quad (1.8)$$

when it performs a motion with the objective velocity V .

In our previous work [12], by the present authors, we have solved the control problem (1.1)–(1.6) for v_* in finite-dimensional control spaces: \mathcal{C}_χ for localized controls and \mathcal{C}_τ for tangential controls (see the definition of those spaces in (4.5) and (4.4)).

In this paper, our aim is to consider controls v_* in infinite-dimensional spaces and to show the existence of an optimal control which minimizes the drag functional given by (1.8). Similar problems have been solved in [5, 6] in the unsteady case without spin of the body by taking for the state equations only the classical Navier-Stokes system. A control problem for the swimming of microscopic organisms was solved in [15]. Here, we also want to characterize the minima by means of the adjoint system. The optimality systems can serve as the basis for computing approximations to optimal solutions numerically as in [3]. The previous results of [12] will be used to construct a corrector of the control v_* which ensures that the self-propelling conditions are satisfied (see Theorem 4.1).

The main difficulties we have to overcome are the usual ones when dealing with exterior domains [10, 7], namely the need of knowing the asymptotic behavior of solutions as well as the presence of terms associated with the rotation of the solid. For example, a good knowledge of the rate of decay of the velocity field is crucial to establish the energy equation for a flow (1.1)–(1.4), which, in turn, permits to write the cost functional (1.8) in a more convenient form, where, in particular, the pressure is not present. The definition of the Lagrangian is another delicate issue in the analysis of the control problems, which benefits from the extra regularity at infinity provided by the self-propelling condition (1.5), as justified in detail in [12]. In particular, we use the fact that $v \in L^2(\Omega)$, which in general does not hold in exterior domains. We develop even L^2 -estimate of the velocity v , which seems to be of independent interest, and apply it to justify the Gâteaux differentiability of the control-to-state map with values in $W^{2,2}(\Omega)$ (especially in $L^2(\Omega)$), which plays a role to deduce the optimal condition at the final stage. For technical reasons, as in [6], we need to impose some restrictions on the size of the control v_* and on the size of the objective velocity components ξ and ω .

The plan of the paper is the following. In Section 2, we gather some notation and an auxiliary result. The new L^2 -estimates for the linearized problem associated with (1.1)–(1.4), which are consequence of a condition on the net force exerted by the fluid to the rigid body similar to (1.5), are deduced in Section 3. In Section 4, we precisely state our optimal control problems associated with system (1.1)–(1.6). Existence of minima to those control problems is shown in Section 5. Then in order to characterize minima, we study, in Section 6, the regularity of the control-to-state mapping, more specifically, its Gâteaux differentiability. Finally, in Section 7, we obtain optimality conditions of order 1 for our problems. These conditions require the well-posedness of the adjoint system.

2 Notation and auxiliary results

Throughout the paper we shall use the same font style to denote scalar, vector and tensor-valued functions and corresponding function spaces. We use the usual notation to denote Lebesgue and Sobolev spaces on a domain \mathcal{A} , namely, $L^q(\mathcal{A})$ and $W^{m,q}(\mathcal{A})$, with norms $\|\cdot\|_{q,\mathcal{A}}$ and $\|\cdot\|_{m,q,\mathcal{A}}$, respectively. By $W^{m-\frac{1}{q},q}(\partial\mathcal{A})$

we indicate the trace space on the (sufficiently) smooth boundary $\partial\mathcal{A}$ of \mathcal{A} , for functions from $W^{m,q}(\mathcal{A})$, equipped with the usual norm $\|\cdot\|_{m-\frac{1}{q},q,\partial\mathcal{A}}$. The homogeneous Sobolev space of order $(k, q) \in \mathbb{N} \times (1, \infty)$ is defined by

$$D^{k,q}(\mathcal{A}) := \{u \in L^1_{loc}(\mathcal{A}) ; D^\alpha u \in L^q(\mathcal{A}) \text{ for any multi-index } \alpha \text{ with } |\alpha| = k\}$$

with associated seminorm $|u|_{k,q,\mathcal{A}} = \sum_{|\alpha|=k} \|D^\alpha u\|_{q,\mathcal{A}}$. The space $D_0^{1,q}(\mathcal{A})$ is the closure of $C_0^\infty(\mathcal{A})$ in $D^{1,q}(\mathcal{A})$. In particular, the dual space of $D_0^{1,2}(\mathcal{A})$ defined for a domain \mathcal{A} , $D^{-1,2}(\mathcal{A})$ with norm $|\cdot|_{-1,2,\mathcal{A}}$, will be used in this work.

In what follows, we can assume that Ω is the exterior domain $\mathbb{R}^3 \setminus \mathcal{S}$. By $\mathcal{D}(\Omega)$ we denote the space of $C_0^\infty(\Omega)^3$ -functions which are divergence free. For a vector or second-order tensor field G , $\alpha \geq 0$ and a positive function ϖ defined on Ω , we use the notation

$$[G]_{\alpha,\varpi,\Omega} := \sup_{x \in \Omega} [\varpi(x)^\alpha |G(x)|], \quad (2.1)$$

and denote by $L_{\alpha,\varpi}^\infty(\Omega)^3$ and $L_{\alpha,\varpi}^\infty(\Omega)^{3 \times 3}$ the spaces of those $G \in L^\infty(\Omega)^3$ or $G \in L^\infty(\Omega)^{3 \times 3}$ such that the norm $[G]_{\alpha,\varpi,\Omega}$ is finite. To alleviate the notation, we will write only $L_{\alpha,\varpi}^\infty(\Omega)$ instead of $L_{\alpha,\varpi}^\infty(\Omega)^3$ and $L_{\alpha,\varpi}^\infty(\Omega)^{3 \times 3}$.

For $R > 0$, we denote by B_R the open ball $B_R := \{x \in \mathbb{R}^3; |x| < R\}$ and by A_{R_1,R_2} ($R_2 > R_1$) the spherical-annulus domain $A_{R_1,R_2} := \{x \in \mathbb{R}^3; R_1 < |x| < R_2\}$. If $\mathbb{R}^3 \setminus \Omega \subset B_\varrho$ for some $\varrho > 0$, we set $\Omega_R := \Omega \cap B_R$ and $\Omega^R := \Omega \setminus \overline{B_R}$ for $R > \varrho$.

Now we collect a number of useful results concerning the generalized Oseen system

$$-\operatorname{div} \sigma(u, q) - V \cdot \nabla u + \omega \times u = f \quad \text{in } \Omega \quad (2.2)$$

$$\operatorname{div} u = 0 \quad \text{in } \Omega \quad (2.3)$$

$$u = u_* \quad \text{on } \partial\Omega \quad (2.4)$$

$$\lim_{|x| \rightarrow \infty} u(x) = 0 \quad (2.5)$$

where V is given by (1.7). For the case of external force $f = \operatorname{div} F$ with F satisfying some anisotropic pointwise estimate, see [12, Proposition 2.1]. The following proposition provides a solution with several properties when less assumptions are imposed on f .

Proposition 2.1. *If Ω is of class C^2 , $f \in L^2(\Omega) \cap D^{-1,2}(\Omega)$ and $u_* \in W^{3/2,2}(\partial\Omega)$, then there is a unique solution (u, q) to (2.2)–(2.5) such that*

$$|u|_{1,2,\Omega} + |u|_{2,2,\Omega} + \|u\|_{\infty,\Omega} + \|q\|_{1,2,\Omega} \leq C(B, \Omega) (\|f\|_{2,\Omega} + |f|_{-1,2,\Omega} + \|u_*\|_{3/2,2,\partial\Omega}),$$

where $B > 0$ is such that $|\xi|, |\omega| \leq B$.

Proof. For the sake of the readers, we give the main steps of the proof although the results are more or less known. First, existence of a weak solution $u \in D^{1,2}(\Omega)$ of (2.2)–(2.5) is obtained by applying [10, Theorem VIII.1.2, p. 501]. Note that for this step, it is sufficient that Ω is a locally Lipschitz domain, $f \in D^{-1,2}(\Omega)$ and $u_* \in W^{1/2,2}(\partial\Omega)$. The pressure $q \in L^2_{loc}(\overline{\Omega})$ is recovered by [10, Lemma VIII.1.1, p. 500]. This weak solution satisfies the following estimates:

$$\|u\|_{2,\Omega_R} + \|\nabla u\|_{2,\Omega} + \|u\|_{6,\Omega} \leq C(R, \Omega) [|f|_{-1,2,\Omega} + (1 + |\xi| + |\omega|) \|u_*\|_{1/2,2,\partial\Omega}], \quad (2.6)$$

$$\|q - \bar{q}_R\|_{2,\Omega_R} \leq C(R, \Omega) [|f|_{-1,2,\Omega} + (1 + |\xi| + |\omega|) \|\nabla u\|_{2,\Omega}], \quad (2.7)$$

for R sufficiently large, where $\bar{q}_R := \frac{1}{|\Omega_R|} \int_{\Omega_R} q$. Uniqueness of the weak solution (up to constants for the pressure, however, we will actually single out it later) is given by [10, Lemma VIII.2.3, p. 512] (see also Theorem VIII.2.1).

Then, we apply [10, Lemma VIII.2.1, p. 505] to obtain

$$\|D^2 u\|_{2,\Omega^r} \leq C(r, \rho, B) (\|f\|_{2,\Omega^\rho} + \|\nabla u\|_{2,\Omega}) \quad (2.8)$$

with $r > \rho$ provided $f \in L^2(\Omega^\rho)$ for some $\rho > 0$. Moreover, [10, Lemma VIII.6.1, p. 554] yields estimates for the second derivatives of u and first derivatives of q in a bounded domain Ω_r , provided $f \in L^2(\Omega_R)$ for some $R > r$, $u_* \in W^{3/2,2}(\partial\Omega)$ and Ω is of class C^2 :

$$\|u\|_{2,2,\Omega_r} + \|\nabla q\|_{2,\Omega_r} \leq C(r, R, \Omega, B) (\|f\|_{2,\Omega_R} + \|u_*\|_{3/2,2,\partial\Omega} + \|u\|_{2,\Omega_R} + \|q - \bar{q}_R\|_{2,\Omega_R}). \quad (2.9)$$

From the equation (2.2) together with (2.6) and (2.8) it follows that

$$\frac{|\nabla q(x)|}{1+|x|} \leq |\Delta u(x)| + |f(x)| + C(\xi, \omega) \left(|\nabla u(x)| + \frac{|u(x)|}{1+|x|} \right) \text{ a.e. } \Omega,$$

and since the right-hand side of this inequality is square-summable near infinity (Hardy inequality is used for the last term), we have

$$\frac{\nabla q}{1+|x|} \in L^2(\Omega^r) \quad (2.10)$$

for every sufficiently large $r > 0$.

Given $r > 0$ large enough such that $\mathbb{R}^3 \setminus \Omega \subset B_{r/2}$, we fix a cut-off function $\psi \in C_0^\infty(B_r; [0, 1])$ satisfying $\psi(x) = 1$ for $x \in B_{r/2}$. Given a pressure q , let us consider the pair $(\tilde{u}, \tilde{q}) = ((1-\psi)u, (1-\psi)(q-\bar{q}_r))$, that obeys

$$-\operatorname{div} \sigma(\tilde{u}, \tilde{q}) - V \cdot \nabla \tilde{u} + \omega \times \tilde{u} = \tilde{f}, \quad \operatorname{div} \tilde{u} = g \quad \text{in } \mathbb{R}^3$$

with

$$\tilde{f} := (1-\psi)f + 2\nabla\psi \cdot \nabla u + (\Delta\psi + V \cdot \nabla\psi)u - (\nabla\psi)(q-\bar{q}_r), \quad g := -u \cdot \nabla\psi.$$

Then

$$\Delta \tilde{q} = \operatorname{div}(\tilde{f} + \nabla g + gV) \quad \text{in } \mathbb{R}^3 \quad (2.11)$$

and from (2.10) we deduce $\nabla \tilde{q} \in \mathcal{S}'(\mathbb{R}^3)$ and thereby, $\tilde{q} \in \mathcal{S}'(\mathbb{R}^3)$. Every solution to (2.11) within $\mathcal{S}'(\mathbb{R}^3)$ is represented as

$$\tilde{q} = Q + P, \quad Q = -\mathcal{F}^{-1} \left(\frac{i\zeta \cdot \hat{h}}{|\zeta|^2} \right) \quad (2.12)$$

where P is a harmonic polynomial, \mathcal{F}^{-1} denotes the Fourier inverse transform and

$$h := \tilde{f} + \nabla g + gV \in L^2(\mathbb{R}^3).$$

Since $\nabla Q \in L^2(\mathbb{R}^3)$, which follows from $h \in L^2(\mathbb{R}^3)$, one needs $(\nabla P)/(1+|x|) \in L^2(\Omega^r)$ to accomplish (2.10) and this, in turn, is possible only if P is a constant. We thus obtain

$$\nabla \tilde{q} = \nabla Q \quad (2.13)$$

together with the estimate

$$\|\nabla q\|_{2,\Omega^r} \leq \|\nabla \tilde{q}\|_{2,\mathbb{R}^3} \leq \|\tilde{f} + \nabla g + gV\|_{2,\mathbb{R}^3} \leq C(r, B) (\|f\|_{2,\Omega} + \|u\|_{1,2,\Omega_r} + \|q - \bar{q}_r\|_{2,\Omega_r}). \quad (2.14)$$

To deduce this estimate, we have used Plancherel Theorem. Combining (2.14) with (2.9) leads to $\nabla q \in L^2(\Omega)$ and, therefore, there is a constant $a \in \mathbb{R}$ such that $\|q - a\|_{6,\Omega} \leq C\|\nabla q\|_{2,\Omega}$. We now single out the pressure q with $a = 0$, that is, $q \in L^6(\Omega)$. Then we have

$$\|q\|_{2,\Omega_r} \leq C(r)\|q\|_{6,\Omega_r} \leq C(r)\|\nabla q\|_{2,\Omega}. \quad (2.15)$$

With this pressure at hand, let us go back to the cut-off procedure above and (2.11).

Considering the pairing $\langle h, \varphi \rangle_{\mathbb{R}^3}$ with $\varphi \in C_0^\infty(\mathbb{R}^3)$ and noting $\|\varphi\|_{2,\Omega_r} \leq C\|\nabla \varphi\|_{2,\mathbb{R}^3}$ by the same reasoning as above, we find that $h \in D^{-1,2}(\mathbb{R}^3)$ by duality. By [13, Lemma 2.2] (see also [10, Theorem II.8.2, p. 112]), there is a tensor field $H \in L^2(\mathbb{R}^3)$ such that $\langle h, \varphi \rangle_{\mathbb{R}^3}$ is given by

$$\langle h, \varphi \rangle_{\mathbb{R}^3} = \int_{\mathbb{R}^3} H : \nabla \varphi dx = -\langle \operatorname{div} H, \varphi \rangle_{(C_0^\infty(\mathbb{R}^3))' \times C_0^\infty(\mathbb{R}^3)}, \quad \forall \varphi \in C_0^\infty(\mathbb{R}^3),$$

and the following estimate holds:

$$\|H\|_{2,\mathbb{R}^3} = \|h\|_{-1,2,\mathbb{R}^3} = \|\tilde{f} + \nabla g + gV\|_{-1,2,\mathbb{R}^3} \leq C(r, B) (\|f\|_{-1,2,\Omega} + \|u\|_{2,\Omega_r} + \|q - \bar{q}_r\|_{2,\Omega_r}). \quad (2.16)$$

From (2.12) and Plancherel Theorem, we get

$$Q = -\mathcal{F}^{-1} \left(\frac{(\zeta \otimes \zeta) : \hat{H}}{|\zeta|^2} \right), \quad \|Q\|_{2,\mathbb{R}^3} \leq \|H\|_{2,\mathbb{R}^3}. \quad (2.17)$$

By (2.13) there is a constant $b \in \mathbb{R}$ such that $\tilde{q} + b = Q \in L^2(\mathbb{R}^3)$ and, therefore, $q - \bar{q}_r + b \in L^2(\Omega^r)$, where $r > 0$ is fixed at the outset of the cut-off procedure. However, it follows from $q \in L^6(\Omega)$ that $b = \bar{q}_r$. As a consequence, we obtain $q \in L^2(\Omega^r)$ and, furthermore, by (2.16)–(2.17)

$$\|q\|_{2,\Omega^r} \leq \|\tilde{q} + \bar{q}_r\|_{2,\mathbb{R}^3} = \|Q\|_{2,\mathbb{R}^3} \leq C(r, B)(\|f\|_{-1,2,\Omega} + \|u\|_{2,\Omega^r} + \|q - \bar{q}_r\|_{2,\Omega^r}), \quad (2.18)$$

which together with (2.15) implies that $q \in L^2(\Omega)$. We collect (2.6)–(2.9), (2.14), (2.15) and (2.18) to conclude the desired estimates except the L^∞ -norm.

Finally, using [2, Lemma 4.1] and Sobolev inequalities, we conclude that $u, \nabla u \in L^6(\Omega)$, $u \in L^\infty(\Omega)$ and

$$\|u\|_{\infty,\Omega} \leq C(\Omega, B)(\|f\|_{2,\Omega} + \|f\|_{-1,2,\Omega} + \|u_*\|_{3/2,2,\partial\Omega}).$$

□

3 L^2 -estimate of the solution to a linearized problem

Given a rigid motion V as in (1.7), in this section, we consider better asymptotic behavior at infinity of the solution to the generalized Oseen system

$$-\operatorname{div} \sigma(u, q) - V \cdot \nabla u + \omega \times u = \operatorname{div} F, \quad \operatorname{div} u = 0 \quad (3.1)$$

in an exterior domain Ω without explicitly specifying any boundary condition at $\partial\Omega$ when

$$\begin{aligned} N &= 0, & \text{if } \omega &= 0, \\ \omega \cdot N &= 0, & \text{if } \omega &\in \mathbb{R}^3 \setminus \{0\}, \end{aligned} \quad (3.2)$$

where N stands for the net force exerted by the fluid to the rigid body, that is,

$$N = N_{u,q} = \int_{\partial\Omega} [\sigma(u, q) + u \otimes V - (\omega \times x) \otimes u + F] n \, d\gamma, \quad (3.3)$$

which is well-defined as long as (u, q) and F are of class (3.4)–(3.5) below (yielding

$$[\sigma(u, q) + u \otimes V - (\omega \times x) \otimes u + F] n \in W^{-1/2,2}(\partial\Omega)$$

as the normal trace). Note that (1.5) is equivalent to $N_{v,p} = 0$ with $F = -v \otimes v$. Under the condition (3.2), we know from asymptotic structure of the flow at infinity that $u \in L^2(\Omega)$, see [4] and [14] especially for the case $\omega \in \mathbb{R}^3 \setminus \{0\}$, however, to our knowledge useful estimates are not available so far in the literature. For later use, we are aiming at deduction of the following L^2 -estimate.

Proposition 3.1. *Suppose*

$$[x \mapsto (1 + |x|)F(x)] \in L^2(\Omega). \quad (3.4)$$

Let (u, q) be a solution to (3.1) of class

$$\nabla u, q \in L^2(\Omega), \quad u \in L^6(\Omega). \quad (3.5)$$

Let us set

$$\Phi = \int_{\partial\Omega} n \cdot u \, d\gamma.$$

1. Let $\omega = 0$. If $N = 0$, then we have $u \in L^2(\Omega)$ subject to

$$\|u\|_{2,\Omega} \leq C[(1 + |\xi|)(\|\nabla u\|_{2,\Omega} + |\Phi|) + \|q\|_{2,\Omega} + \|F\|_{2,\Omega}] + C'\| |x|F \|_{2,\Omega} \quad (3.6)$$

with some constants $C, C' > 0$ which are independent of u, q, F, ξ and Φ .

2. Let $\omega \in \mathbb{R}^3 \setminus \{0\}$. If $\omega \cdot N = 0$, then we have $u \in L^2(\Omega)$ subject to

$$\|u\|_{2,\Omega} \leq CK(u, q, F, \xi, \omega, \Phi) + C'\| |x|F \|_{2,\Omega} \quad (3.7)$$

with some constants $C, C' > 0$ which are independent of u, q, F, ξ, ω , where

$$\begin{aligned} K(u, q, F, \xi, \omega, \Phi) &= \left(1 + |\omega|^{-1/4} + \frac{|\omega \cdot \xi|^{1/2}}{|\omega|} + \frac{|\omega \times \xi|}{|\omega|^2}\right) \\ &\quad \cdot [(1 + |\xi| + |\omega|)(\|\nabla u\|_{2,\Omega} + |\Phi|) + \|q\|_{2,\Omega} + \|F\|_{2,\Omega}]. \end{aligned} \quad (3.8)$$

For the latter case $\omega \in \mathbb{R}^3 \setminus \{0\}$, it is also possible to deduce a bit different estimate from (3.7):

$$\begin{aligned} \|u\|_{2,\Omega} &\leq C \left(|\omega|^{-1/4} + \frac{|\omega \cdot \xi|^{1/2}}{|\omega|} \right) (|N| + |\omega| |\Phi|) + C' \| |x| F \|_{2,\Omega} \\ &\quad + C \left(1 + \frac{|\omega \times \xi|}{|\omega|^2} \right) [(1 + |\xi| + |\omega|) (\|\nabla u\|_{2,\Omega} + |\Phi|) + \|q\|_{2,\Omega} + \|F\|_{2,\Omega}]. \end{aligned} \quad (3.9)$$

As usual, by a cut-off procedure, the problem in exterior domains will be reduced to the one in the whole space. When $\omega \in \mathbb{R}^3 \setminus \{0\}$, we then use the Moggi-Chasles transform, see [10] and [11], to modify the resulting problem in the following way

$$\begin{aligned} y &= M \left(x - \frac{\omega \times \xi}{|\omega|^2} \right), \\ \mathfrak{s}(y) &= s \left(M^\top y + \frac{\omega \times \xi}{|\omega|^2} \right) \text{ for a scalar field,} \\ \mathfrak{v}(y) &= Mv \left(M^\top y + \frac{\omega \times \xi}{|\omega|^2} \right) \text{ for a vector field,} \\ \mathfrak{T}(y) &= (MTM^\top) \left(M^\top y + \frac{\omega \times \xi}{|\omega|^2} \right) \text{ for a tensor field,} \end{aligned} \quad (3.10)$$

where $M \in \mathbb{R}^{3 \times 3}$ being an orthogonal matrix that fulfills $M \frac{\omega}{|\omega|} = e_1$, to obtain the generalized Oseen system in which the direction of the translation is parallel to the axis of rotation that becomes the e_1 -direction. Thus, let us consider the system

$$-\Delta v + \nabla p - \mathcal{R} \partial_1 v - |\omega| [(e_1 \times y) \cdot \nabla v - e_1 \times v] = f, \quad \operatorname{div} v = 0 \quad \text{in } \mathbb{R}_y^3 \quad (3.11)$$

within the class of tempered distributions, where

$$\mathcal{R} = \frac{\omega \cdot \xi}{|\omega|}. \quad (3.12)$$

For the external force f of a suitable class, we know that:

1. when $\omega = 0$, to the classical Oseen system

$$-\Delta v + \nabla p - \xi \cdot \nabla v = f, \quad \operatorname{div} v = 0 \text{ in } \mathbb{R}^3,$$

we see that

$$\widehat{v}(\zeta) = \frac{1}{|\zeta|^2 - i\xi \cdot \zeta} \left(\mathbb{I}_3 - \frac{\zeta \otimes \zeta}{|\zeta|^2} \right) \widehat{f}(\zeta) \quad (3.13)$$

is a solution on the Fourier side.

2. when $\omega \in \mathbb{R}^3 \setminus \{0\}$

$$\widehat{v}(\zeta) = \int_0^\infty e^{-(|\zeta|^2 - i\mathcal{R}\zeta_1)t} O_\omega(t)^\top \left(\mathbb{I}_3 - \frac{(O_\omega(t)\zeta) \otimes (O_\omega(t)\zeta)}{|\zeta|^2} \right) \widehat{f}(O_\omega(t)\zeta) dt \quad (3.14)$$

is a solution for (3.11) on the Fourier side, where

$$O_\omega(t) = O(|\omega|t), \quad O(t) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos t & -\sin t \\ 0 & \sin t & \cos t \end{bmatrix}.$$

Relation (3.14) is classical but we recall here the idea to obtain it: first we notice that the Fourier transform of (v, p) satisfies

$$\begin{aligned} (|\zeta|^2 - i\mathcal{R}\zeta_1) \widehat{v}(\zeta) - |\omega| [(e_1 \times \zeta) \cdot \nabla_\zeta \widehat{v}(\zeta) - e_1 \times \widehat{v}(\zeta)] - i\zeta \widehat{p}(\zeta) &= \widehat{f}(\zeta), \\ i\zeta \cdot \widehat{v}(\zeta) &= 0, \quad \zeta \in \mathbb{R}^3. \end{aligned}$$

Eliminating the pressure, we find

$$(|\zeta|^2 - i\mathcal{R}\zeta_1)\widehat{v}(\zeta) - |\omega| [(e_1 \times \zeta) \cdot \nabla_\zeta \widehat{v}(\zeta) - e_1 \times \widehat{v}(\zeta)] = \left(\mathbb{I}_3 - \frac{\zeta \otimes \zeta}{|\zeta|^2} \right) \widehat{f}(\zeta), \quad \zeta \in \mathbb{R}^3.$$

Then we define

$$\mathcal{V}(t, \zeta) = O_\omega(t) \widehat{v}(O_\omega(t)^\top \zeta) = O_\omega(t) \widehat{v}(O_\omega(-t)\zeta) \quad (3.15)$$

and some standard computation yields that $\mathcal{V}(t, \zeta)$ is time-periodic and satisfies

$$\frac{\partial}{\partial t} \mathcal{V}(t, \zeta) + (|\zeta|^2 - i\mathcal{R}\zeta_1) \mathcal{V}(t, \zeta) = O_\omega(t) \left(\mathbb{I}_3 - \frac{(O_\omega(t)^\top \zeta) \otimes (O_\omega(t)^\top \zeta)}{|\zeta|^2} \right) \widehat{f}(O_\omega(t)^\top \zeta).$$

By Duhamel's principle, we deduce

$$\mathcal{V}(t, \zeta) = \int_{-\infty}^t e^{(s-t)(|\zeta|^2 - i\mathcal{R}\zeta_1)} O_\omega(s) \left(\mathbb{I}_3 - \frac{(O_\omega(s)^\top \zeta) \otimes (O_\omega(s)^\top \zeta)}{|\zeta|^2} \right) \widehat{f}(O_\omega(s)^\top \zeta) ds$$

and using (3.15), we recover (3.14).

Let $(v_0, p_0) \in \mathcal{S}'(\mathbb{R}^3)$ be a solution to (3.11) with $f = 0$, then we see that $\text{supp } \widehat{v}_0 \subset \{0\}$. In fact, since

$$|\zeta|^2 \widehat{v}_0 + i\zeta \widehat{p}_0 - i\mathcal{R}\zeta_1 \widehat{v}_0 - |\omega| [(e_1 \times \zeta) \cdot \nabla_\zeta \widehat{v}_0 - e_1 \times \widehat{v}_0] = 0, \quad i\zeta \cdot \widehat{v}_0 = 0,$$

we have $|\zeta|^2 \widehat{p}_0 = 0$, which implies that $\text{supp } \widehat{p}_0 \subset \{0\}$ and that

$$|\zeta|^2 [|\zeta|^2 \widehat{v}_0 - i\mathcal{R}\zeta_1 \widehat{v}_0 - |\omega| [(e_1 \times \zeta) \cdot \nabla_\zeta \widehat{v}_0 - e_1 \times \widehat{v}_0]] = 0.$$

Given arbitrary vector field $\varphi \in C_0^\infty(\mathbb{R}^3 \setminus \{0\})$, we set

$$\tau(\zeta) = \int_0^\infty e^{-(|\zeta|^2 + i\mathcal{R}\zeta_1)t} O_\omega(t) \varphi(O_\omega(t)^\top \zeta) dt \in C_0^\infty(\mathbb{R}^3 \setminus \{0\}),$$

which solves the adjoint system

$$|\zeta|^2 \tau + i\mathcal{R}\zeta_1 \tau + |\omega| [(e_1 \times \zeta) \cdot \nabla_\zeta \tau - e_1 \times \tau] = \varphi.$$

We thus obtain

$$\begin{aligned} \langle \widehat{v}_0, \varphi \rangle &= \langle \widehat{v}_0, |\zeta|^2 \tau + i\mathcal{R}\zeta_1 \tau + |\omega| [(e_1 \times \zeta) \cdot \nabla_\zeta \tau - e_1 \times \tau] \rangle \\ &= \left\langle |\zeta|^2 \{ |\zeta|^2 \widehat{v}_0 - i\mathcal{R}\zeta_1 \widehat{v}_0 - |\omega| [(e_1 \times \zeta) \cdot \nabla_\zeta \widehat{v}_0 - e_1 \times \widehat{v}_0] \}, \frac{\tau}{|\zeta|^2} \right\rangle = 0, \end{aligned}$$

yielding $\text{supp } \widehat{v}_0 \subset \{0\}$. Therefore, $v = \mathcal{F}^{-1} \widehat{v}$ with (3.14) is the only solution to (3.11) up to (specific) polynomials within $\mathcal{S}'(\mathbb{R}^3)$. It is actually the only solution when the polynomials are excluded on account of the asymptotic behavior at infinity. The same thing for the case $\omega = 0$ is shown even more straightforward. Thus the following L^2 -estimate for (3.13)–(3.14) plays an important role.

Lemma 3.2. *Suppose that f is of the form $f = g + \text{div } G$ with*

$$[y \mapsto (1 + |y|)G(y)] \in L^2(\mathbb{R}^3), \quad g \in L^1(\mathbb{R}^3), \quad [y \mapsto |y|g(y)] \in L^s(\mathbb{R}^3) \quad (3.16)$$

for some $s \in [1, 6/5)$.

1. Let $\omega = 0$, and let \widehat{v} be as in (3.13). If

$$\int_{\mathbb{R}^3} g(y) dy = 0, \quad (3.17)$$

then we have $v = \mathcal{F}^{-1} \widehat{v} \in L^2(\mathbb{R}^3)$ subject to

$$\|v\|_{2, \mathbb{R}^3} \leq C (\|g\|_{1, \mathbb{R}^3} + \|G\|_{2, \mathbb{R}^3}) + C' \| |y|G \|_{2, \mathbb{R}^3} + C'' \| |y|g \|_{s, \mathbb{R}^3} \quad (3.18)$$

with some constants $C, C', C'' > 0$ which are independent of g, G and ξ .

2. Let $\omega \in \mathbb{R}^3 \setminus \{0\}$, and let \widehat{v} be as in (3.14). If

$$e_1 \cdot \int_{\mathbb{R}^3} g(y) dy = 0, \quad (3.19)$$

then we have $v = \mathcal{F}^{-1}\widehat{v} \in L^2(\mathbb{R}^3)$ subject to

$$\begin{aligned} \|v\|_{2,\mathbb{R}^3} &\leq C \left[\left(|\omega|^{-1/4} + |\mathcal{R}|^{1/2} |\omega|^{-1/2} \right) \left| \int_{\mathbb{R}^3} g(y) dy \right| + \|g\|_{1,\mathbb{R}^3} + \|G\|_{2,\mathbb{R}^3} \right] \\ &\quad + C' \| |y|G \|_{2,\mathbb{R}^3} + C'' \| |y|g \|_{s,\mathbb{R}^3} \end{aligned} \quad (3.20)$$

with some constants $C, C', C'' > 0$ which are independent of g, G, ω and \mathcal{R} .

Proof. Let us discuss the case $\omega \in \mathbb{R}^3 \setminus \{0\}$ and consider

$$\int_{\mathbb{R}^3} |\widehat{v}(\zeta)|^2 d\zeta = \int_{|\zeta| \geq 1} |\widehat{v}(\zeta)|^2 d\zeta + \int_{|\zeta| < 1} |\widehat{v}(\zeta)|^2 d\zeta.$$

Using the Schwarz inequality in the integral with respect to t and then the Fubini theorem followed by the change of variable $\zeta \mapsto O_\omega(t)^\top \zeta$ together with the decomposition $\widehat{f}(\zeta) = \widehat{g}(\zeta) + i\zeta \cdot \widehat{G}(\zeta)$, we see that the high frequency part of (3.14) is estimated as

$$\begin{aligned} \int_{|\zeta| \geq 1} |\widehat{v}(\zeta)|^2 d\zeta &\leq C \int_{|\zeta| \geq 1} \left(\int_0^\infty e^{-|\zeta|^2 t} |\widehat{f}(O_\omega(t)\zeta)| dt \right)^2 d\zeta \\ &\leq C \int_{|\zeta| \geq 1} \int_0^\infty e^{-|\zeta|^2 t} |\widehat{f}(O_\omega(t)\zeta)|^2 dt \frac{d\zeta}{|\zeta|^2} \\ &= C \int_0^\infty \int_{|\zeta| \geq 1} e^{-|\zeta|^2 t} |\widehat{f}(\zeta)|^2 \frac{d\zeta}{|\zeta|^2} dt \\ &= C \int_{|\zeta| \geq 1} \frac{|\widehat{f}(\zeta)|^2}{|\zeta|^4} d\zeta \\ &\leq C \|\widehat{g}\|_{\infty,\mathbb{R}^3}^2 \int_{|\zeta| \geq 1} \frac{d\zeta}{|\zeta|^4} + C \|\widehat{G}\|_{2,\mathbb{R}^3}^2 \\ &\leq C (\|g\|_{1,\mathbb{R}^3}^2 + \|G\|_{2,\mathbb{R}^3}^2). \end{aligned} \quad (3.21)$$

Our main task is thus to study the low frequency part, which will be based on the decomposition

$$\widehat{f}(O_\omega(t)\zeta) = \widehat{g}(0) + \int_0^1 (O_\omega(t)\zeta) \cdot (\nabla \widehat{g})(\sigma O_\omega(t)\zeta) d\sigma + i(O_\omega(t)\zeta) \cdot \widehat{G}(O_\omega(t)\zeta).$$

Note that the function \widehat{g} is uniformly continuous by $g \in L^1(\mathbb{R}^3)$ and thus $\widehat{g}(0)$ makes sense. The above decomposition of $\widehat{f}(O_\omega(t)\zeta)$ splits (3.14) into three parts:

$$\begin{aligned} &\int_{|\zeta| < 1} |\widehat{v}(\zeta)|^2 d\zeta \\ &= \int_{|\zeta| < 1} \left| \int_0^\infty e^{-(|\zeta|^2 - i\mathcal{R}\zeta_1)t} O_\omega(t)^\top \left(\mathbb{I}_3 - \frac{(O_\omega(t)\zeta) \otimes (O_\omega(t)\zeta)}{|\zeta|^2} \right) dt \widehat{g}(0) \right. \\ &\quad \left. + \int_0^\infty e^{-(|\zeta|^2 - i\mathcal{R}\zeta_1)t} O_\omega(t)^\top \left(\mathbb{I}_3 - \frac{(O_\omega(t)\zeta) \otimes (O_\omega(t)\zeta)}{|\zeta|^2} \right) \int_0^1 (O_\omega(t)\zeta) \cdot (\nabla \widehat{g})(\sigma O_\omega(t)\zeta) d\sigma dt \right. \\ &\quad \left. + i \int_0^\infty e^{-(|\zeta|^2 - i\mathcal{R}\zeta_1)t} O_\omega(t)^\top \left(\mathbb{I}_3 - \frac{(O_\omega(t)\zeta) \otimes (O_\omega(t)\zeta)}{|\zeta|^2} \right) (O_\omega(t)\zeta) \cdot \widehat{G}(O_\omega(t)\zeta) dt \right|^2 d\zeta \\ &\leq (I_1 + I_2 + I_3)^2. \end{aligned}$$

For the last two integrals I_2 and I_3 , one may ignore the oscillation. In fact, we see from the Hardy inequality that

$$\begin{aligned} I_3^2 &\leq C \int_{|\zeta| < 1} \int_0^\infty e^{-|\zeta|^2 t} |\widehat{G}(O_\omega(t)\zeta)|^2 dt d\zeta = C \int_{|\zeta| < 1} |\widehat{G}(\zeta)|^2 \frac{d\zeta}{|\zeta|^2} \\ &\leq C \int_{\mathbb{R}^3} \frac{|\widehat{G}(\zeta)|^2}{|\zeta|^2} d\zeta \leq C \|\nabla \widehat{G}\|_{2,\mathbb{R}^3}^2 = C \| |y|G \|_{2,\mathbb{R}^3}^2 \end{aligned} \quad (3.22)$$

and that

$$\begin{aligned}
I_2^2 &\leq C \int_{|\zeta|<1} \int_0^\infty e^{-|\zeta|^2 t} \int_0^1 |(\nabla \widehat{g})(\sigma O_\omega(t)\zeta)|^2 d\sigma dt d\zeta \\
&= C \int_{|\zeta|<1} \int_0^1 |(\nabla \widehat{g})(\sigma\zeta)|^2 d\sigma \frac{d\zeta}{|\zeta|^2} = C \int_0^1 \int_{|\zeta|<\sigma} |(\nabla \widehat{g})(\zeta)|^2 \frac{d\zeta}{|\zeta|^2} \frac{d\sigma}{\sigma} \\
&\leq C \int_0^1 \left(\int_{|\zeta|<\sigma} |(\nabla \widehat{g})(\zeta)|^{s/(s-1)} d\zeta \right)^{2(1-1/s)} \sigma^{-6(1-1/s)} d\sigma \\
&\leq C \|\nabla \widehat{g}\|_{s/(s-1), \mathbb{R}^3}^2 \leq C \|y|g|\|_{s, \mathbb{R}^3}^2
\end{aligned} \tag{3.23}$$

where (3.16) with $s \in [1, 6/5]$ is employed. We thus obtain

$$I_2 \leq C \|y|g|\|_{s, \mathbb{R}^3}, \quad I_3 \leq C \|y|G|\|_{2, \mathbb{R}^3}. \tag{3.24}$$

For the crucial part I_1 we do need the assumption (3.19), that is, $e_1 \cdot \widehat{g}(0) = 0$, as well as the oscillation caused by the rotation. By the relations

$$\widehat{g}(0) = (e_1 \cdot \widehat{g}(0))e_1 + (e_1 \times \widehat{g}(0)) \times e_1 = (e_1 \times \widehat{g}(0)) \times e_1$$

and

$$O_\omega(t)^\top \left(\mathbb{I}_3 - \frac{(O_\omega(t)\zeta) \otimes (O_\omega(t)\zeta)}{|\zeta|^2} \right) O_\omega(t) = \mathbb{I}_3 - \frac{\zeta \otimes \zeta}{|\zeta|^2}$$

we may write

$$I_1^2 = \int_{|\zeta|<1} |w(\zeta)|^2 d\zeta$$

with

$$\begin{aligned}
w(\zeta) &:= \int_0^\infty e^{-(|\zeta|^2 - i\mathcal{R}\zeta_1)t} \left(\mathbb{I}_3 - \frac{\zeta \otimes \zeta}{|\zeta|^2} \right) O_\omega(t)^\top dt [(e_1 \times \widehat{g}(0)) \times e_1] \\
&= \left(\mathbb{I}_3 - \frac{\zeta \otimes \zeta}{|\zeta|^2} \right) \phi(\zeta),
\end{aligned}$$

where

$$\phi(\zeta) := \int_0^\infty e^{-(|\zeta|^2 - i\mathcal{R}\zeta_1)t} O_\omega(t)^\top dt [(e_1 \times \widehat{g}(0)) \times e_1].$$

An elementary computation yields

$$\phi(\zeta) = \frac{1}{2} \begin{bmatrix} 0 \\ \alpha_- J_+ + \alpha_+ J_- \\ i\alpha_- J_+ - i\alpha_+ J_- \end{bmatrix}$$

with

$$\alpha_\pm := \widehat{g}_2(0) \pm i\widehat{g}_3(0), \quad J_\pm(\zeta) := \frac{1}{|\zeta|^2 - i(\mathcal{R}\zeta_1 \pm |\omega|)}.$$

We then find

$$\int_{|\zeta|<1} |J_\pm(\zeta)|^2 d\zeta \leq \int_{\mathbb{R}^3} \frac{d\zeta}{|\zeta|^4 + (\mathcal{R}\zeta_1 \pm |\omega|)^2} \leq C \left(\frac{1}{\sqrt{|\omega|}} + \frac{|\mathcal{R}|}{|\omega|} \right) \tag{3.25}$$

for all $|\omega| > 0$ and $\mathcal{R} \in \mathbb{R}$, see (3.12). In fact, we immediately see that

$$\int_{\mathbb{R}^3} \frac{d\zeta}{|\zeta|^4 + |\omega|^2} = \frac{1}{\sqrt{|\omega|}} \int_{\mathbb{R}^3} \frac{d\zeta}{|\zeta|^4 + 1}$$

for the case $\mathcal{R} = 0$, while the integral above for the other case $\mathcal{R} \neq 0$ is rewritten as

$$\frac{2\pi}{\sqrt{|\omega|}} \int_0^\pi \sin \theta \int_0^\infty \frac{\rho^2 d\rho}{\rho^4 + \left(\frac{\mathcal{R}}{\sqrt{|\omega|}} \rho \cos \theta \pm 1 \right)^2} d\theta.$$

For the latter case, it is reasonable to split the integral with respect to ρ into two parts:

$$\int_0^\infty = \int_0^{\sqrt{|\omega|}/(2|\mathcal{R}|)} + \int_{\sqrt{|\omega|}/(2|\mathcal{R}|)}^\infty.$$

Then we have

$$\int_0^{\sqrt{|\omega|}/(2|\mathcal{R}|)} \frac{\rho^2 d\rho}{\rho^4 + \left(\frac{\mathcal{R}}{\sqrt{|\omega|}} \rho \cos \theta \pm 1\right)^2} \leq \int_0^\infty \frac{\rho^2}{\rho^4 + \frac{1}{4}} d\rho$$

as well as

$$\int_{\sqrt{|\omega|}/(2|\mathcal{R}|)}^\infty \frac{\rho^2 d\rho}{\rho^4 + \left(\frac{\mathcal{R}}{\sqrt{|\omega|}} \rho \cos \theta \pm 1\right)^2} \leq \int_{\sqrt{|\omega|}/(2|\mathcal{R}|)}^\infty \frac{d\rho}{\rho^2} \leq \frac{2|\mathcal{R}|}{\sqrt{|\omega|}}.$$

We thus obtain (3.25), which leads us to

$$I_1 \leq C \left(\frac{1}{\sqrt{|\omega|}} + \frac{|\mathcal{R}|}{|\omega|} \right)^{1/2} |\widehat{g}(0)|. \quad (3.26)$$

We collect (3.21), (3.24) and (3.26) to conclude (3.20).

For the other case $\omega = 0$, the high frequency part is the same as in (3.21)₃ and the low frequency part can be treated as in (3.22)–(3.23) by use of $\widehat{g}(0) = 0$, so that we obtain (3.18). The proof is complete. \square

Proof of Proposition 3.1. We first discuss the flow with vanishing flux condition. Let us fix $R > 0$ such that $\mathbb{R}^3 \setminus \Omega \subset B_R$, and take a cut-off function $\psi \in C_0^\infty(B_{3R}; [0, 1])$ such that $\psi(x) = 1$ for $x \in B_{2R}$. Given (u, q) which is of class (3.5) together with $\int_{\partial\Omega} n \cdot u d\gamma = 0$ and satisfies (3.1), we set

$$\tilde{u} = (1 - \psi)u + \mathbb{B}[u \cdot \nabla\psi], \quad \tilde{q} = (1 - \psi)q,$$

where \mathbb{B} denotes the Bogovskii operator in the domain $A_{R,3R}$, see [1] and [10, Theorem III.3.3, p.179]. Note that $u \cdot \nabla\psi \in W_0^{1,2}(A_{R,3R})$ with

$$\int_{A_{R,3R}} u \cdot \nabla\psi dx = \int_{\partial B_R} u \cdot \frac{-x}{R} d\gamma = \int_{\partial\Omega} u \cdot n d\gamma = 0$$

so that $\mathbb{B}[u \cdot \nabla\psi] \in W_0^{2,2}(A_{R,3R})$ is well-defined with the relation $\operatorname{div} \mathbb{B}[u \cdot \nabla\psi] = u \cdot \nabla\psi$ and

$$\|\mathbb{B}[u \cdot \nabla\psi]\|_{2,2,A_{R,3R}} \leq C \|u \cdot \nabla\psi\|_{1,2,A_{R,3R}} \leq C \|u\|_{1,2,A_{R,3R}} \leq C (\|u\|_{6,A_{R,3R}} + \|\nabla u\|_{2,A_{R,3R}}) \leq C \|\nabla u\|_{2,\Omega}. \quad (3.27)$$

Then the pair (\tilde{u}, \tilde{q}) obeys

$$-\Delta \tilde{u} + \nabla \tilde{q} - V \cdot \nabla \tilde{u} + \omega \times \tilde{u} = h + \operatorname{div} [(1 - \psi)F], \quad \operatorname{div} \tilde{u} = 0 \quad \text{in } \mathbb{R}^3, \quad (3.28)$$

where

$$h = 2\nabla\psi \cdot \nabla u + (\Delta\psi + V \cdot \nabla\psi)u - \Delta\mathbb{B}[u \cdot \nabla\psi] - V \cdot \nabla\mathbb{B}[u \cdot \nabla\psi] + \omega \times \mathbb{B}[u \cdot \nabla\psi] - (\nabla\psi)q + F(\nabla\psi)$$

which satisfies

$$\int_{\mathbb{R}^3} h(x) dx = N, \quad (3.29)$$

and N denotes the net force (3.3). In fact,

$$\begin{aligned} \int_{\mathbb{R}^3} h(x) dx &= - \int_{A_{R,3R}} \operatorname{div} [\sigma(\tilde{u}, \tilde{q}) + \tilde{u} \otimes V - (\omega \times x) \otimes \tilde{u} + (1 - \psi)F] dx \\ &= - \int_{|x|=3R} [\sigma(u, q) + u \otimes V - (\omega \times x) \otimes u + F] \frac{x}{3R} d\gamma \end{aligned}$$

yielding (3.29) by use of (3.1).

By the transformations (3.10), we consider

$$\begin{aligned} v(y) &= M\tilde{u} \left(M^\top y + \frac{\omega \times \xi}{|\omega|^2} \right), \\ p(y) &= \tilde{q} \left(M^\top y + \frac{\omega \times \xi}{|\omega|^2} \right), \\ g(y) &= Mh \left(M^\top y + \frac{\omega \times \xi}{|\omega|^2} \right), \\ G(y) &= \left(M[(1-\psi)F]M^\top \right) \left(M^\top y + \frac{\omega \times \xi}{|\omega|^2} \right), \end{aligned}$$

and it can be shown that (v, p) obeys (3.11) with $f = g + \operatorname{div} G$ which satisfies (3.16)–(3.19) subject to

$$\begin{aligned} & \|g\|_{1, \mathbb{R}^3} + \| |y|g \|_{1, \mathbb{R}^3} \\ &= \|h\|_{1, \mathbb{R}^3} + \left\| \left| x - \frac{\omega \times \xi}{|\omega|^2} \right| h \right\|_{1, \mathbb{R}^3} \\ &\leq C_R \left(1 + \frac{|\omega \times \xi|}{|\omega|^2} \right) [(1 + |\xi| + |\omega|) \|\nabla u\|_{2, \Omega} + \|q\|_{2, \Omega} + \|F\|_{2, \Omega}], \end{aligned} \quad (3.30)$$

as well as

$$\|G\|_{2, \mathbb{R}^3} + \| |y|G \|_{2, \mathbb{R}^3} \leq \left(1 + \frac{|\omega \times \xi|}{|\omega|^2} \right) \|F\|_{2, \Omega} + \| |x|F \|_{2, \Omega}. \quad (3.31)$$

The condition (3.19) is in fact verified as

$$e_1 \cdot \int_{\mathbb{R}^3} g(y) dy = \frac{\omega}{|\omega|} \cdot \int_{\mathbb{R}^3} h(x) dx = 0 \quad (3.32)$$

by use of (3.29) and the assumption $\omega \cdot N = 0$.

Since $u \in L^6(\Omega)$ implies that $v \in L^6(\mathbb{R}^3) \subset \mathcal{S}'(\mathbb{R}^3)$, v coincides with (3.14) on the Fourier side by the reasoning mentioned just before Lemma 3.2. By taking (3.30)–(3.32) into account, we obtain from Lemma 3.2 that there are constants $C, C' > 0$ satisfying

$$\|u\|_{2, \Omega_{3R}} \leq \|\tilde{u}\|_{2, \mathbb{R}^3} = \|v\|_{2, \mathbb{R}^3} \leq CK(u, q, F, \xi, \omega, 0) + C' \| |x|F \|_{2, \Omega},$$

where $K(u, q, F, \xi, \omega, 0)$ is given by (3.8) with $\Phi = 0$. Here, $|\int_{\mathbb{R}^3} g(y) dy|$ has been just replaced by $\|g\|_{1, \mathbb{R}^3}$ in (3.20). On the other hand, we have

$$\|u\|_{2, \Omega_{3R}} \leq C \|u\|_{6, \Omega} \leq C \|\nabla u\|_{2, \Omega}.$$

Combining the estimates above implies (3.7) when $\Phi = 0$. If we prefer to keep

$$\left| \int_{\mathbb{R}^3} g(y) dy \right| = \left| \int_{\mathbb{R}^3} Mh(x) dx \right| = |N|$$

in (3.20) as it is, we obtain (3.9) with $\Phi = 0$ as well.

For general case without any condition at the boundary $\partial\Omega$, let us reduce the problem to the case discussed above by lifting the flux $\Phi = \int_{\partial\Omega} n \cdot u d\gamma$. We fix $x_0 \in \operatorname{int}(\mathbb{R}^3 \setminus \Omega)$ and take the flux carrier

$$\mathcal{W}(x) = \Phi \nabla \frac{1}{4\pi|x-x_0|}.$$

Note that one cannot always choose $x_0 = 0$ (center of mass of the rigid body). Then the pair

$$U = u - \mathcal{W}, \quad Q = q - (\xi + \omega \times x_0) \cdot \mathcal{W}$$

obeys

$$-\Delta U + \nabla Q - V \cdot \nabla U + \omega \times U = \operatorname{div} F, \quad \operatorname{div} U = 0$$

in Ω subject to $\int_{\partial\Omega} n \cdot U \, d\gamma = 0$. Concerning the net force (3.3), we observe

$$N_{U,Q} = N_{u,q} + \Phi(\omega \times x_0)$$

as verified in [12, Section 6] (in which the nonlinear momentum flux is discussed, however, all the computations for the linear part are included there), so that the condition $\omega \cdot N_{u,q} = 0$ implies $\omega \cdot N_{U,Q} = 0$. Hence, we already know that

$$\|U\|_{2,\Omega} \leq CK(U, Q, F, \xi, \omega, 0) + C' \|x|F\|_{2,\Omega}.$$

Since

$$\|\nabla U\|_{2,\Omega} \leq \|\nabla u\|_{2,\Omega} + C|\Phi|, \quad \|Q\|_{2,\Omega} \leq \|q\|_{2,\Omega} + C(|\xi| + |\omega|)|\Phi|$$

as well as

$$\|\mathcal{W}\|_{2,\Omega} \leq C|\Phi|,$$

one concludes (3.7). The other case $\omega = 0$ is also discussed in the same way as above. \square

4 The state system and the cost functional

As explained in the introduction, our aim is to find a control v_* for which

$$\inf W(v, p) = \inf_{\partial\Omega} \int_{\partial\Omega} v \cdot \sigma(v, p) n \, d\gamma, \quad (4.1)$$

is attained (see (1.8)). In (4.1) the infimum is taken over the set of all possible states (v, p) satisfying (1.1)–(1.6) for v_* either in

$$\mathcal{V}_\tau := \left\{ v_* \in W^{3/2,2}(\partial\Omega) ; v_* \cdot n = 0 \text{ on } \partial\Omega \right\} \quad (4.2)$$

or in

$$\mathcal{V}_\Gamma := \left\{ v_* \in W^{3/2,2}(\partial\Omega) ; v_* = 0 \text{ on } \partial\Omega \setminus \Gamma \right\}, \quad (4.3)$$

where Γ is a nonempty open subset of $\partial\Omega$.

We recall that in [12] we studied the case of subspaces of \mathcal{V}_τ and of \mathcal{V}_Γ of finite dimension:

$$\mathcal{C}_\tau := \text{span} \left\{ (g^{(i)} \times n) \times n, (G^{(i)} \times n) \times n ; i = 1, 2, 3 \right\}, \quad (4.4)$$

and

$$\mathcal{C}_\chi := \text{span} \left\{ \chi g^{(i)}, \chi G^{(i)} ; i = 1, 2, 3 \right\}, \quad (4.5)$$

where $\chi \not\equiv 0$ is a non-negative smooth function such that $\text{supp}(\chi) \subset \Gamma$ and where $g^{(i)}, G^{(i)}$ are defined as follows.

First, we introduce a set of generalized Oseen systems associated with the basic rigid motions: for each $i \in \{1, 2, 3\}$, $(v^{(i)}, q^{(i)})$ and $(V^{(i)}, Q^{(i)})$ are the solutions of

$$\begin{aligned} -\text{div} \sigma(v^{(i)}, q^{(i)}) + (\xi + \omega \times x) \cdot \nabla v^{(i)} - \omega \times v^{(i)} &= 0 \quad \text{in } \Omega \\ \text{div} v^{(i)} &= 0 \quad \text{in } \Omega \\ v^{(i)} &= e_i \quad \text{on } \partial\Omega \\ \lim_{|x| \rightarrow \infty} v^{(i)} &= 0 \end{aligned} \quad (4.6)$$

and

$$\begin{aligned} -\text{div} \sigma(V^{(i)}, Q^{(i)}) + (\xi + \omega \times x) \cdot \nabla V^{(i)} - \omega \times V^{(i)} &= 0 \quad \text{in } \Omega \\ \text{div} V^{(i)} &= 0 \quad \text{in } \Omega \\ V^{(i)} &= e_i \times x \quad \text{on } \partial\Omega \\ \lim_{|x| \rightarrow \infty} V^{(i)} &= 0 \end{aligned} \quad (4.7)$$

where (e_1, e_2, e_3) is the canonical basis of \mathbb{R}^3 . Then the fields $g^{(i)}$ and $G^{(i)}$ are given by

$$g^{(i)} := \sigma(v^{(i)}, q^{(i)})n \quad \text{on } \partial\Omega, \quad (4.8)$$

$$G^{(i)} := \sigma(V^{(i)}, Q^{(i)})n \quad \text{on } \partial\Omega, \quad (4.9)$$

being in $W^{3/2,2}(\partial\Omega)$ provided $\partial\Omega \in C^3$, see [12, Lemma 3.2]. If $|\xi|$ and $|\omega|$ are small enough, then \mathcal{C}_τ and \mathcal{C}_χ are of dimension 6 (see [12, Theorem 1.1]). We proved in [12] that, for $|\xi|$ and $|\omega|$ small enough, there exists only one $v_* \in \mathcal{C}_\tau$ (resp. $v_* \in \mathcal{C}_\chi$) such that there exists a solution (v, p) of (1.1)–(1.6).

Here, we want to consider controls v_* in \mathcal{V}_τ or \mathcal{V}_Γ and characterize the optimal controls that minimize (1.8). However, before starting the analysis of the minimization problem, we must notice that for an arbitrary $v_* \in \mathcal{V}_\tau$ or $v_* \in \mathcal{V}_\Gamma$, system (1.1)–(1.6) has no solution in general since the unique solution (v, p) of (1.1)–(1.4) may not verify the self-propelled conditions (1.5), (1.6). In order to handle this difficulty, the solution to our problem consists of a boundary velocity v_* which can be decomposed into two parts, one part which effectively acts as the infinite dimensional control (to alleviate the presentation, we keep denoting it by v_*) and another part, say v_*^c , that “corrects” the control in order to enforce the self-propelled conditions and belongs to the finite dimensional spaces \mathcal{C}_τ and \mathcal{C}_χ .

Then, assuming that the rigid body velocity V is given and having in mind the boundary control problems for $v_* \in \mathcal{V}_\tau$ or $v_* \in \mathcal{V}_\Gamma$, we write the state system (1.1)–(1.6) in the form

$$-\operatorname{div} \sigma(v, p) + (v - V) \cdot \nabla v + \omega \times v = 0 \quad \text{in } \Omega \quad (4.10)$$

$$\operatorname{div} v = 0 \quad \text{in } \Omega \quad (4.11)$$

$$v = V + v_* + v_*^c \quad \text{on } \partial\Omega \quad (4.12)$$

$$\lim_{|x| \rightarrow \infty} v(x) = 0 \quad (4.13)$$

$$m\xi \times \omega + \int_{\partial\Omega} \left[-\sigma(v, p)n + \left((v_* + v_*^c) \cdot n \right) (v_* + v_*^c + V + \omega \times x) \right] d\gamma = 0 \quad (4.14)$$

$$(I\omega) \times \omega + \int_{\partial\Omega} x \times \left[-\sigma(v, p)n + \left((v_* + v_*^c) \cdot n \right) (v_* + v_*^c + V + \omega \times x) \right] d\gamma = 0 \quad (4.15)$$

$$v_*^c \in \mathcal{C}_\tau \quad \text{or} \quad v_*^c \in \mathcal{C}_\chi. \quad (4.16)$$

In what follows, we use the notation

$$\varpi(x) := \begin{cases} \left(1 + \left| x - \frac{\omega \times \xi}{|\omega|^2} \right| \right) \left(1 + 2 \frac{|\omega \cdot \xi|}{|\omega|} s(x) \right), & \omega \neq 0, \\ (1 + |x|) (1 + 2(|\xi||x| + \xi \cdot x)), & \omega = 0, \end{cases} \quad (4.17)$$

where

$$s(x) := \left| x - \frac{\omega \times \xi}{|\omega|^2} \right| + \frac{\operatorname{sign}(\omega \cdot \xi)}{|\omega|} \omega \cdot x, \quad \omega \neq 0.$$

Extending the results of [12, Theorem 1.1 and Theorem 1.2], we obtain the following for our state system.

Theorem 4.1. *Let Ω be of class C^3 . There exist constants $c_0, C_1, C_2 > 0$, which depend on Ω , such that if $\xi, \omega \in \mathbb{R}^3$ and $v_* \in \mathcal{V}_\tau$ (resp. $v_* \in \mathcal{V}_\Gamma$) satisfy*

$$|\xi| \leq c_0, \quad |\omega| \leq c_0, \quad \|v_*\|_{3/2,2,\partial\Omega} \leq c_0, \quad (4.18)$$

then the following assertions hold.

1. A solution (v, p, v_*^c) of the problem (4.10)–(4.16) can be found within the class

$$\varpi v \in L^\infty(\Omega), \quad (v, p) \in W^{2,2}(\Omega) \times W^{1,2}(\Omega), \quad v_*^c \in \mathcal{C}_\tau \quad (\text{resp. } v_*^c \in \mathcal{C}_\chi) \quad (4.19)$$

along with estimates

$$\|v\|_{1,\varpi,\Omega} + \|\nabla v\|_{2,\Omega} + \|v_*^c\|_{3/2,2,\partial\Omega} \leq C_1 (|\xi, \omega| + \|v_*\|_{3/2,2,\partial\Omega}), \quad (4.20)$$

$$\|\nabla^2 v\|_{2,\Omega} + \|p\|_{1,2,\Omega} \leq C_2 (|\xi, \omega| + \|v_*\|_{3/2,2,\partial\Omega}), \quad (4.21)$$

and the energy equation

$$\int_{\partial\Omega} (\sigma(v, p)n) \cdot v d\gamma = 2 \int_{\Omega} |D(v)|^2 dx + \frac{1}{2} \int_{\partial\Omega} (v_* + v_*^c) \cdot n |V + v_* + v_*^c|^2 d\gamma. \quad (4.22)$$

2. The solution of the problem (4.10)–(4.16) is unique (up to constants for the pressure) within the class of functions satisfying (4.20) as well as $v_*^c \in \mathcal{C}_\tau$ (resp. $v_*^c \in \mathcal{C}_\chi$). The pressure is singled out under the additional condition $p \in L^2(\Omega)$.

The emphasis is finite kinetic energy $v \in L^2(\Omega)$, see (4.19), as a consequence of the self-propelling condition (4.14), and this helps us to justify the energy relation (4.22). As in [12], our notion of solution to (4.10)–(4.16) is the standard weak solution $\nabla v \in L^2(\Omega)$, such that

$$2 \int_{\Omega} D(v) : D(\varphi) dx + \int_{\Omega} (v - V) \cdot \nabla v \cdot \varphi dx + \int_{\Omega} (\omega \times v) \cdot \varphi dx = 0, \quad \forall \varphi \in \mathcal{D}(\Omega), \quad (4.23)$$

which, due to the extra regularity obtained in Theorem 4.1, will satisfy the equations (1.5)–(1.6) in the strong form. We will only sketch the proof of Theorem 4.1 in the case of localized controls \mathcal{C}_χ since it is completely similar to the proof of [12, Theorem 1.1].

Proof of Theorem 4.1. Let

$$\mathcal{X} := \{(v, \alpha, \beta) \in D^{1,2}(\Omega) \times \mathbb{R}^3 \times \mathbb{R}^3 ; [v]_{1,\varpi,\Omega} < \infty\}$$

endowed with the norm

$$\|(v, \alpha, \beta)\|_{\mathcal{X}} := |v|_{1,2,\Omega} + [v]_{1,\varpi,\Omega} + |(\alpha, \beta)|.$$

Assume $(\bar{v}, \bar{\alpha}, \bar{\beta}) \in \mathcal{X}$, and let

$$\bar{v}_*^c := \sum_{i=1}^3 \left(\bar{\alpha}_i \chi g^{(i)} + \bar{\beta}_i \chi G^{(i)} \right),$$

$$f(\bar{v}) := -\bar{v} \cdot \nabla \bar{v} = \operatorname{div} F(\bar{v}), \quad F(\bar{v}) := -\bar{v} \otimes \bar{v},$$

$$\xi_f(\bar{v}_*^c) := - \int_{\partial\Omega} ((v_* + \bar{v}_*^c) \cdot n) ((v_* + \bar{v}_*^c + V + \omega \times x) d\gamma - m\xi \times \omega - \int_{\partial\Omega} (V \cdot n)(V + v_* + \bar{v}_*^c) d\gamma, \quad (4.24)$$

$$\omega_f(\bar{v}_*^c) := - \int_{\partial\Omega} x \times (v_* + \bar{v}_*^c + V + \omega \times x) ((v_* + \bar{v}_*^c) \cdot n) d\gamma - (I\omega) \times \omega - \int_{\partial\Omega} x \times (V + v_* + \bar{v}_*^c) (V \cdot n) d\gamma. \quad (4.25)$$

Following [12, Lemma 4.1], we introduce the following auxiliary linear systems

$$\left\{ \begin{array}{l} -\operatorname{div} \sigma(u^{(j)}, p^{(j)}) - V \cdot \nabla u^{(j)} + \omega \times u^{(j)} = 0 \\ \operatorname{div} u^{(j)} = 0 \\ u^{(j)} = \chi g^{(j)} \text{ or } (g^{(j)} \times n) \times n \text{ on } \partial\Omega \\ \lim_{|x| \rightarrow \infty} u^{(j)} = 0, \end{array} \right. \quad (4.26)$$

$$\left\{ \begin{array}{l} -\operatorname{div} \sigma(U^{(j)}, P^{(j)}) - V \cdot \nabla U^{(j)} + \omega \times U^{(j)} = 0 \\ \operatorname{div} U^{(j)} = 0 \\ U^{(j)} = \chi G^{(j)} \text{ or } (G^{(j)} \times n) \times n \text{ on } \partial\Omega \\ \lim_{|x| \rightarrow \infty} U^{(j)} = 0. \end{array} \right. \quad (4.27)$$

$$\left\{ \begin{array}{l} -\operatorname{div} \sigma(u_f, p_f) - V \cdot \nabla u_f + \omega \times u_f = f(\bar{v}) \\ \operatorname{div} u_f = 0 \\ u_f = V + v_* \text{ on } \partial\Omega \\ \lim_{|x| \rightarrow \infty} u_f = 0. \end{array} \right. \quad (4.28)$$

Using systems (4.26)–(4.28) and [12, Proposition 4.5], we can solve the following problem: for any $v_* \in \mathcal{V}_\Gamma$, and for ξ and ω satisfying (4.18) with some constant c_0 small enough, there exists a unique (v, v_*^c, p) such

that

$$-\operatorname{div} \sigma(v, p) - V \cdot \nabla v + \omega \times v = f(\bar{v}) \quad \text{in } \Omega \quad (4.29)$$

$$\operatorname{div} v = 0 \quad \text{in } \Omega \quad (4.30)$$

$$v = V + v_* + v_*^c \quad \text{on } \partial\Omega \quad (4.31)$$

$$\lim_{|x| \rightarrow \infty} v(x) = 0 \quad (4.32)$$

$$-\int_{\partial\Omega} [\sigma(v, p)n + (V \cdot n)v] d\gamma = \xi_f(\bar{v}_*^c) \quad (4.33)$$

$$-\int_{\partial\Omega} x \times [\sigma(v, p)n + (V \cdot n)v] d\gamma = \omega_f(\bar{v}_*^c) \quad (4.34)$$

$$v_*^c = \sum_{i=1}^3 \left(\alpha_i \chi g^{(i)} + \beta_i \chi G^{(i)} \right) \in \mathcal{C}_\chi. \quad (4.35)$$

Moreover, we have

$$\begin{aligned} & \sup_{x \in \Omega} [\varpi(x)|v(x)|] + \|\nabla v\|_{1,2,\Omega} + \|p\|_{1,2,\Omega} + |(\alpha, \beta)| \\ & \leq C [|(\xi_f(\bar{v}_*^c), \omega_f(\bar{v}_*^c))| + \|f(\bar{v})\|_{2,\Omega} + [F(\bar{v})]_{2,\varpi,\Omega} + |(\xi, \omega)| + \|v_*\|_{3/2,2,\partial\Omega} + |(\xi, \omega)| (|(\xi, \omega)| + \|v_*\|_{3/2,2,\partial\Omega})]. \end{aligned} \quad (4.36)$$

This allows us to define the mapping

$$\mathcal{Z} : \mathcal{X} \ni (\bar{v}, \bar{\alpha}, \bar{\beta}) \mapsto (v, \alpha, \beta) \in \mathcal{X},$$

where (v, v_*^c) is the solution of (4.29)–(4.35).

Following the proof of [12, Theorem 1.1], based on [12, Proposition 4.5], we obtain

$$\|\mathcal{Z}(\bar{v}, \bar{\alpha}, \bar{\beta})\|_{\mathcal{X}} \leq C (|(\xi, \omega)| + \|v_*\|_{3/2,2,\partial\Omega} + |(\xi, \omega)|^2 + \|v_*\|_{3/2,2,\partial\Omega}^2) + C \|(\bar{v}, \bar{\alpha}, \bar{\beta})\|_{\mathcal{X}}^2.$$

Taking $c_0 \in (0, 1)$ small enough in (4.18), we see that a ball

$$\mathcal{X}_{R_0} := \{(v, \alpha, \beta) \in \mathcal{X} ; \|(v, \alpha, \beta)\|_{\mathcal{X}} \leq R_0\}, \quad R_0 = C_1 (|(\xi, \omega)| + \|v_*\|_{3/2,2,\partial\Omega}) \quad (4.37)$$

is invariant by \mathcal{Z} . In a similar way, we also obtain that \mathcal{Z} is a strict contraction on \mathcal{X}_{R_0} for c_0 small enough. This gives us the existence of a solution to problem (4.10)–(4.16). The corresponding estimates (4.20)–(4.21) follow from (4.36).

To prove that $v \in L^2(\Omega)$, we apply [12, Theorem 1.2] where we only have to replace v_* by $v_* + v_*^c$ (see the definition (1.13) of N in [12]).

In order to obtain the energy equation, for $R > 0$ large enough such that $\mathbb{R}^3 \setminus \Omega \subset B_R$, we use a radially symmetric cut-off function $\psi_R(x) = \tilde{\psi}(|x|/R)$ with $\tilde{\psi} \in C_0^1((-2, 2); [0, 1])$ which fulfills $\tilde{\psi} \equiv 1$ on $[-1, 1]$. Then we have

$$\|\nabla \psi_R\|_{q, \mathbb{R}^3} = CR^{-1+3/q} \quad (3 \leq q \leq \infty) \quad (4.38)$$

as well as $(\omega \times x) \cdot \nabla \psi_R = 0$. From those properties it follows that

$$\|V \cdot \nabla \psi_R\|_{\infty, \mathbb{R}^3} = \|\xi \cdot \nabla \psi_R\|_{\infty, \mathbb{R}^3} = C/R.$$

Taking the inner product of both sides of the equation (4.10) with $\psi_R v$ and integrating by parts over Ω yield

$$\begin{aligned} & -\int_{\partial\Omega} (\sigma(v, p)n) \cdot v d\gamma + \int_{\Omega} \sigma(v, p) : (v \otimes \nabla \psi_R) dx + 2 \int_{\Omega} |D(v)|^2 \psi_R dx \\ & \quad + \int_{\partial\Omega} \frac{|v|^2}{2} (v - V) \cdot n d\gamma - \int_{\Omega} \frac{|v|^2}{2} (v - V) \cdot \nabla \psi_R dx = 0 \end{aligned}$$

Since $v \in W^{1,2}(\Omega)$ and $p \in L^2(\Omega)$, we have

$$\left| -\int_{\Omega} \frac{|v|^2}{2} (v - V) \cdot \nabla \psi_R dx + \int_{\Omega} \sigma(v, p) : (v \otimes \nabla \psi_R) dx \right| \leq \frac{C}{R} \int_{\Omega} (|v|^3 + |v|^2 + |D(v)|^2 + |p|^2) dx.$$

Using the dominated convergence Theorem and (4.19), we can pass to the limit $R \rightarrow \infty$ and we deduce the result. \square

Using the energy equation (4.22), we can rewrite the drag functional (1.8) in the following way:

Corollary 4.2. *Let (v, p, v_*^c) be the solution of (4.10)–(4.16) obtained in Theorem 4.1. Then*

$$W(v, p) = 2\|D(v)\|_{2,\Omega}^2 + \frac{1}{2} \int_{\partial\Omega} (v_* + v_*^c) \cdot n |V + v_* + v_*^c|^2 d\gamma.$$

From now on, we assume

$$|\xi| \leq c_0, \quad |\omega| \leq c_0 \quad (4.39)$$

and, given $\kappa \in (0, c_0]$, we define

$$\mathcal{V}_\tau^\kappa := \left\{ v_* \in \mathcal{V}_\tau ; \|v_*\|_{W^{3/2,2}(\partial\Omega)} \leq \kappa \right\}, \quad (4.40)$$

$$\mathcal{V}_\Gamma^\kappa := \left\{ v_* \in \mathcal{V}_\Gamma ; \|v_*\|_{W^{3/2,2}(\partial\Omega)} \leq \kappa \right\}, \quad (4.41)$$

where \mathcal{V}_τ and \mathcal{V}_Γ are defined by (4.2) and (4.3), while c_0 is the constant in Theorem 4.1.

Using Corollary 4.2, we deduce that the problem (4.1) reduces here to minimize

$$J(v_*) := 2\|D(v)\|_{2,\Omega}^2 + \frac{1}{2} \int_{\partial\Omega} (v_* + v_*^c) \cdot n |V + v_* + v_*^c|^2 d\gamma, \quad (4.42)$$

where (v, p, v_*^c) is the solution of (4.10)–(4.16) associated with either $v_* \in \mathcal{V}_\tau^\kappa$ or $v_* \in \mathcal{V}_\Gamma^\kappa$. This functional J is well-defined since Theorem 4.1 allows us to define the control-to-state mapping $v_* \mapsto (v, p, v_*^c)$. In the following sections, we study this mapping and consider the optimal control problems:

$$\inf_{v_* \in \mathcal{V}_\tau^\kappa} J(v_*) \quad (4.43)$$

or

$$\inf_{v_* \in \mathcal{V}_\Gamma^\kappa} J(v_*). \quad (4.44)$$

Under the condition (4.39), in the next section, it turns out that (4.43) and (4.44) respectively admit solutions for every $\kappa \in (0, c_0]$, however, the radius κ of the admissible closed balls (4.40)–(4.41) as well as (ξ, ω) should be taken still smaller (see Theorems 6.1 and 7.4) in order to characterize the optimal solution in Theorem 7.5.

5 Existence of optimal controls

Here we show that problems (4.43) and (4.44) have a solution, so that the infima are actually minima.

Theorem 5.1. *Assume that $\xi, \omega \in \mathbb{R}^3$ satisfy (4.39). Let $\kappa \in (0, c_0]$. Then each of the optimal control problems (4.43) and (4.44) admits a solution.*

Proof. We only consider problem (4.44), the case of tangential controls can be treated with exactly the same arguments. By the embedding $W^{3/2,2}(\partial\Omega) \subset L^3(\partial\Omega)$ and (4.41), we have

$$\|v_*\|_{L^3(\partial\Omega)} \leq C\kappa$$

and from (4.20) it follows that

$$\|v_*^c\|_{L^3(\partial\Omega)} \leq C(\kappa + c_0).$$

Therefore, the functional J is bounded from below on $\mathcal{V}_\Gamma^\kappa$ since

$$J(v_*) \geq -C \int_{\partial\Omega} \left(|v_* + v_*^c|^3 + |v_* + v_*^c| \right) d\gamma.$$

Thus, there exists a sequence

$$\{v_{*k}\}_{k \in \mathbb{N}} \subset \mathcal{V}_\Gamma^\kappa \text{ such that } J(v_{*k}) \rightarrow \inf_{v_* \in \mathcal{V}_\Gamma^\kappa} J(v_*) \quad (k \rightarrow \infty).$$

We will denote by $\{(v_k, p_k, v_{*k}^c)\}_{k \in \mathbb{N}}$ the corresponding sequence of states, that is, (v_k, p_k, v_{*k}^c) is the solution of problem (4.10)–(4.16) for v_{*k} . Since the admissible set \mathcal{V}_Γ^c is weakly sequentially compact, there exist $\widehat{v}_* \in \mathcal{V}_\Gamma^c$ and a subsequence of $\{v_{*k}\}_{k \in \mathbb{N}}$, still denoted by $\{v_{*k}\}_{k \in \mathbb{N}}$, such that

$$v_{*k} \rightharpoonup \widehat{v}_* \text{ weakly in } W^{3/2,2}(\partial\Omega) \quad (5.1)$$

In what follows we take suitable subsequences in order although they are always denoted by the same symbol $\{v_k\}$ etc. From (4.20)–(4.21), we also deduce the existence of $(\widehat{v}, \widehat{p}, \widehat{v}_*^c)$ such that

$$(\nabla v_k, p_k) \rightharpoonup (\nabla \widehat{v}, \widehat{p}) \text{ weakly in } W^{1,2}(\Omega), \quad (5.2)$$

$$v_{*k}^c \rightarrow \widehat{v}_*^c \text{ strongly in } W^{3/2,2}(\partial\Omega). \quad (5.3)$$

Concerning (5.3), what we see at once is that \mathcal{C}_τ (resp. \mathcal{C}_Γ) $\ni v_{*k}^c$ tends to \widehat{v}_*^c weakly in $W^{3/2,2}(\partial\Omega)$ along a subsequence as $k \rightarrow \infty$, which yields $\widehat{v}_*^c \in \mathcal{C}_\tau$ (resp. \mathcal{C}_Γ) since the subspace is weakly closed; then, we eventually obtain the strong convergence above because it is a finite dimensional space. Since $v_k(x)$ tends to zero as $|x| \rightarrow \infty$, from a classical embedding inequality we also deduce (see, for instance, [10, Theorem II.6.1])

$$\|v_k\|_{6,\Omega} \leq C$$

and thus

$$v_k \rightharpoonup \widehat{v} \text{ weakly in } L^6(\Omega). \quad (5.4)$$

We also have

$$\varpi v_k \rightharpoonup \varpi \widehat{v} \text{ weakly } * \text{ in } L^\infty(\Omega).$$

Indeed, there exists $\vartheta \in L^\infty(\Omega)$ such that

$$\int_\Omega \varpi(x) v_k(x) u(x) dx \rightarrow \int_\Omega \vartheta(x) u(x) dx, \quad \forall u \in L^1(\Omega).$$

and, in particular,

$$\int_\Omega \varpi(x) v_k(x) \varphi(x) dx \rightarrow \int_\Omega \vartheta(x) \varphi(x) dx, \quad \forall \varphi \in C_0^\infty(\overline{\Omega}). \quad (5.5)$$

On the other hand, due to (5.4),

$$\int_\Omega \varpi(x) v_k(x) \varphi(x) dx \rightarrow \int_\Omega \varpi(x) v(x) \varphi(x) dx, \quad \forall \varphi \in C_0^\infty(\overline{\Omega}). \quad (5.6)$$

From (5.5) and (5.6) it follows that

$$\int_\Omega \vartheta(x) \varphi(x) dx = \int_\Omega \varpi(x) v(x) \varphi(x) dx, \quad \forall \varphi \in C_0^\infty(\overline{\Omega})$$

and therefore $\vartheta = \varpi v$ a.e. in Ω .

Finally, using classical compactness results, we also deduce from (4.20) that

$$(\nabla v_k, p_k) \rightarrow (\nabla \widehat{v}, \widehat{p}) \text{ strongly in } L_{loc}^2(\overline{\Omega}),$$

$$v_k \rightarrow \widehat{v} \text{ strongly in } L_{loc}^2(\overline{\Omega}),$$

and

$$v_{*k} \rightarrow \widehat{v}_* \text{ strongly in } L^3(\partial\Omega). \quad (5.7)$$

Using the above convergences, we can pass to the limit in (4.23) where v is replaced by v_k . We also know that \widehat{v} satisfies (4.11) and (4.13). We have

$$\|v_k - \widehat{v}\|_{2,\partial\Omega} \leq C \|v_k - \widehat{v}\|_{2,\Omega_R}^{1/2} \|v_k - \widehat{v}\|_{1,2,\Omega_R}^{1/2} \rightarrow 0 \quad (k \rightarrow \infty),$$

and using (5.3) and (5.1), we deduce (4.12).

Similarly,

$$\begin{aligned} & \|\sigma(v_k, p_k)n - \sigma(\widehat{v}, \widehat{p})n\|_{2,\partial\Omega} \\ & \leq C \|\nabla v_k - \nabla \widehat{v}\|_{2,\Omega_R}^{1/2} \|\nabla v_k - \nabla \widehat{v}\|_{1,2,\Omega_R}^{1/2} + C \|p_k - \widehat{p}\|_{2,\Omega_R}^{1/2} \|p_k - \widehat{p}\|_{1,2,\Omega_R}^{1/2} \rightarrow 0 \quad (k \rightarrow \infty), \end{aligned}$$

so that, using (5.7) and (5.3), we deduce (4.14) and (4.15).

The pair $(\widehat{v}, \widehat{v}_*^c)$ is in the class satisfying (4.20) and thus, by the assumption (4.39) and the uniqueness result in Theorem 4.1, $(\widehat{v}, \widehat{p}, \widehat{v}_*^c)$ is the state associated with \widehat{v}_* . Now using (5.2), (5.7) and (5.3) we deduce

$$J(\widehat{v}_*) \leq \liminf_{k \rightarrow \infty} J(v_{*k})$$

and therefore \widehat{v}_* is actually a minimizer:

$$J(\widehat{v}_*) = \inf_{v_* \in \mathcal{V}_\Gamma^c} J(v_*).$$

□

6 Regularity of the control-to-state mapping

Let

$$\mathcal{W} := \left\{ (v, v_*^c) ; v \in D^{1,2}(\Omega) \cap L_{1,\varpi}^\infty(\Omega), \nabla \cdot v = 0 \text{ in } \Omega, v_*^c \in W^{3/2,2}(\partial\Omega) \right\}$$

endowed with the norm

$$\|(v, v_*^c)\|_{\mathcal{W}} := |v|_{1,2,\Omega} + [v]_{1,\varpi,\Omega} + \|v_*^c\|_{3/2,2,\partial\Omega} = \|\nabla v\|_{2,\Omega} + [v]_{1,\varpi,\Omega} + \|v_*^c\|_{3/2,2,\partial\Omega},$$

and consider the subset

$$\mathcal{W}_{R_0} := \left\{ (v, v_*^c) \in \mathcal{W} ; \|(v, v_*^c)\|_{\mathcal{W}} \leq R_0 \right\}, \quad (6.1)$$

where R_0 is the same as in (4.37), that is the right-hand side of (4.20). Suppose (4.39). Given $\kappa \in (0, c_0]$, Theorem 4.1 allows us to define the following control-to-state mappings (recall (4.40) and (4.41))

$$\Lambda_\tau : \mathcal{V}_\tau^\kappa \rightarrow \mathcal{W}_{R_0} \cap [(D^{2,2}(\Omega) \cap L^2(\Omega)) \times \mathcal{C}_\tau], \quad v_* \mapsto (v, v_*^c) \quad (6.2)$$

$$\Lambda_\chi : \mathcal{V}_\Gamma^\kappa \rightarrow \mathcal{W}_{R_0} \cap [(D^{2,2}(\Omega) \cap L^2(\Omega)) \times \mathcal{C}_\chi], \quad v_* \mapsto (v, v_*^c) \quad (6.3)$$

where (v, v_*^c) together with the pressure $p \in W^{1,2}(\Omega)$ is the solution of problem (4.10)–(4.16) associated with v_* given in the admissible set \mathcal{V}_τ^κ or $\mathcal{V}_\Gamma^\kappa$.

Our aim is to show that the maps Λ_τ and Λ_χ are Gâteaux differentiable. We analyze this problem in detail for Λ_τ , the idea being similar for the mapping Λ_χ .

In order to compute the Gâteaux derivative of Λ_τ at $v_* \in \mathcal{V}_\tau^\kappa$ in the direction $\overline{v_*} \in \mathcal{V}_\tau$, that is denoted by

$$D\Lambda_\tau(v_*)\overline{v_*} = (z, z_*^c),$$

we suppose that $v_* + h\overline{v_*}$ is also in \mathcal{V}_τ^κ for $0 < h < h_0$ with sufficiently small h_0 ; in fact, this is accomplished as long as $\langle v_*, \overline{v_*} \rangle_{W^{3/2,2}(\partial\Omega)} < 0$ even if $\|v_*\|_{3/2,2,\partial\Omega} = \kappa$.

We consider

$$(v_h, v_{h_*}^c, p_h) \in \left(\mathcal{W}_{R_0} \cap [(D^{2,2}(\Omega) \cap L^2(\Omega)) \times \mathcal{C}_\tau] \right) \times W^{1,2}(\Omega)$$

which is the solution to (4.10)–(4.16) associated with $v_* + h\overline{v_*}$, that is, $(v_h, v_{h_*}^c) := \Lambda_\tau(v_* + h\overline{v_*})$. Then

$$\begin{aligned} (z_h, z_{h_*}^c, r_h) &:= \frac{(v_h, v_{h_*}^c, p_h) - (v, v_*^c, p)}{h} \\ &= \frac{(\Lambda_\tau(v_* + h\overline{v_*}), p_h) - (\Lambda_\tau(v_*), p)}{h} \in \left(\mathcal{W} \cap [(D^{2,2}(\Omega) \cap L^2(\Omega)) \times \mathcal{C}_\tau] \right) \times W^{1,2}(\Omega) \end{aligned}$$

satisfies the following system

$$-\operatorname{div} \sigma(z_h, r_h) + v \cdot \nabla z_h + z_h \cdot \nabla v_h - V \cdot \nabla z_h + \omega \times z_h = 0 \quad \text{in } \Omega \quad (6.4)$$

$$\operatorname{div} z_h = 0 \quad \text{in } \Omega \quad (6.5)$$

$$z_h = z_{h_*}^c + \overline{v_*} \quad \text{on } \partial\Omega \quad (6.6)$$

$$\lim_{|x| \rightarrow \infty} z_h(x) = 0 \quad (6.7)$$

$$\int_{\partial\Omega} \sigma(z_h, r_h) n \, d\gamma = 0 \quad (6.8)$$

$$\int_{\partial\Omega} x \times \sigma(z_h, r_h) n \, d\gamma = 0, \quad (6.9)$$

$$z_{h_*}^c \in \mathcal{C}_\tau. \quad (6.10)$$

Our aim is to show that, when $h \rightarrow 0$, (z_h, z_{h*}^C, r_h) converges to (z, z_*^C, r) , which solves the linearized state equations

$$-\operatorname{div} \sigma(z, r) + v \cdot \nabla z + z \cdot \nabla v - V \cdot \nabla z + \omega \times z = 0 \quad \text{in } \Omega \quad (6.11)$$

$$\operatorname{div} z = 0 \quad \text{in } \Omega \quad (6.12)$$

$$z = z_*^C + \bar{v}_* \quad \text{on } \partial\Omega \quad (6.13)$$

$$\lim_{|x| \rightarrow \infty} z(x) = 0 \quad (6.14)$$

$$\int_{\partial\Omega} \sigma(z, r) n \, d\gamma = 0 \quad (6.15)$$

$$\int_{\partial\Omega} x \times \sigma(z, r) n \, d\gamma = 0 \quad (6.16)$$

$$z_*^C \in \mathcal{C}_\tau. \quad (6.17)$$

In the case of localized controls, instead of the equations (6.8)–(6.10), we have

$$\int_{\partial\Omega} [-\sigma(z_h, r_h) + (v_* + v_*^C) \cdot n(\bar{v}_* + z_{h*}^C) + (\bar{v}_* + z_{h*}^C) \cdot n(v_* + h\bar{v}_* + v_{h*}^C + V + \omega \times x)] \, d\gamma = 0 \quad (6.18)$$

$$\int_{\partial\Omega} x \times [-\sigma(z_h, r_h) + (v_* + v_*^C) \cdot n(\bar{v}_* + z_{h*}^C) + (\bar{v}_* + z_{h*}^C) \cdot n(v_* + h\bar{v}_* + v_{h*}^C + V + \omega \times x)] \, d\gamma = 0 \quad (6.19)$$

$$z_{h*}^C \in \mathcal{C}_\chi \quad (6.20)$$

and in the corresponding linearized state equations conditions (6.15)–(6.17) are replaced by

$$\int_{\partial\Omega} [-\sigma(z, r) n + (v_* + v_*^C) \cdot n(\bar{v}_* + z_*^C) + (\bar{v}_* + z_*^C) \cdot n(v_* + v_*^C + V + \omega \times x)] \, d\gamma = 0 \quad (6.21)$$

$$\int_{\partial\Omega} x \times [-\sigma(z, r) n + (v_* + v_*^C) \cdot n(\bar{v}_* + z_*^C) + (\bar{v}_* + z_*^C) \cdot n(v_* + v_*^C + V + \omega \times x)] \, d\gamma = 0 \quad (6.22)$$

$$z_*^C \in \mathcal{C}_\chi. \quad (6.23)$$

More specifically, we have

Theorem 6.1. *There exists a constant $\kappa_1 \in (0, c_0]$ depending on Ω such that if $\xi, \omega \in \mathbb{R}^3$ and $v_* \in \mathcal{V}_\tau$ (resp. $v_* \in \mathcal{V}_\Gamma$) satisfy*

$$|\xi| \leq \kappa_1, \quad |\omega| \leq \kappa_1, \quad \|v_*\|_{3/2, 2, \partial\Omega} \leq \kappa_1, \quad (6.24)$$

then the following assertion holds, where c_0 is the constant in Theorem 4.1:

Let (v, v_*^C, p) be the solution to the state equations (4.10)–(4.16) associated with $v_* \in \mathcal{V}_\tau^{\kappa_1}$ (resp. $v_* \in \mathcal{V}_\Gamma^{\kappa_1}$) obtained in Theorem 4.1. Then the mapping Λ_τ (resp. Λ_χ) defined by (6.2) (resp. (6.3)) with $\kappa = \kappa_1$ is Gâteaux differentiable with values in $[W^{2,2}(\Omega) \times L_{1,\varpi}^\infty(\Omega)] \times W^{3/2,2}(\partial\Omega)$ at $v_* \in \mathcal{V}_\tau^{\kappa_1}$ (resp. $v_* \in \mathcal{V}_\Gamma^{\kappa_1}$) in the direction $\bar{v}_* \in \mathcal{V}_\tau$ (resp. $\bar{v}_* \in \mathcal{V}_\Gamma$), where \bar{v}_* must be taken such that $\langle v_*, \bar{v}_* \rangle_{W^{3/2,2}(\partial\Omega)} < 0$ if $\|v_*\|_{3/2, 2, \partial\Omega} = \kappa_1$, whereas it can be arbitrary if $\|v_*\|_{3/2, 2, \partial\Omega} < \kappa_1$, and its derivative is given by $D\Lambda_\tau(v_*)\bar{v}_* = (z, z_*^C)$, (resp. $D\Lambda_\chi(v_*)\bar{v}_* = (z, z_*^C)$) with (z, z_*^C, r) being the solution to the problem (6.11)–(6.17) (resp. (6.11)–(6.14), (6.21)–(6.23)). Namely, we have

$$\|z_h - z\|_{2,2,\Omega} + \|z_h - z\|_{1,\varpi,\Omega} + \|z_{h*}^C - z_*^C\|_{3/2, 2, \partial\Omega} \rightarrow 0,$$

where z_h satisfies (6.4)–(6.10) (resp. (6.4)–(6.7), (6.18)–(6.20)).

Proof. We write a detailed proof for Λ_τ but point out the main differences for the mapping Λ_χ . Equation (6.4) can be written in the form

$$-\operatorname{div} \sigma(z_h, r_h) - V \cdot \nabla z_h + \omega \times z_h = f_h$$

with

$$f_h := -(v \cdot \nabla z_h + z_h \cdot \nabla v) = -\operatorname{div} (z_h \otimes v + v_h \otimes z_h) = \operatorname{div} F_h, \\ F_h := -(z_h \otimes v + v_h \otimes z_h).$$

If $v, v_h \in \mathcal{W}_{R_0}$, then the following estimates hold for f_h and F_h :

$$\|f_h\|_{2,\Omega} \leq C([\![v]\!]_{1,\varpi,\Omega} \|\nabla z_h\|_{2,\Omega} + [\![z_h]\!]_{1,\varpi,\Omega} \|\nabla v_h\|_{2,\Omega}) \leq CR_0([\![z_h]\!]_{1,2,\Omega} + [\![z_h]\!]_{1,\varpi,\Omega}),$$

$$[\![F_h]\!]_{2,\varpi,\Omega} = [\![z_h \otimes v + v_h \otimes z_h]\!]_{2,\varpi,\Omega} \leq CR_0[\![z_h]\!]_{1,\varpi,\Omega},$$

where R_0 is given by (4.37), see (6.1). In order to apply [12, Proposition 4.5], which is still valid even though V is replaced by \bar{v}_* , we rewrite the conditions (6.8) and (6.9) as

$$\int_{\partial\Omega} [\sigma(z_h, r_h)n + (V \cdot n)z_h] d\gamma = \int_{\partial\Omega} (V \cdot n)z_h d\gamma \quad (6.25)$$

$$\int_{\partial\Omega} x \times [\sigma(z_h, r_h)n + (V \cdot n)z_h] d\gamma = \int_{\partial\Omega} x \times [(V \cdot n)z_h] d\gamma, \quad (6.26)$$

to find

$$\begin{aligned} & |z_h|_{1,2,\Omega} + |z_h|_{2,2,\Omega} + [\![z_h]\!]_{1,\varpi,\Omega} + \|r_h\|_{1,2,\Omega} + \|z_{h*}^c\|_{3/2,2,\partial\Omega} \\ & \leq C \left[R_0(|z_h|_{1,2,\Omega} + [\![z_h]\!]_{1,\varpi,\Omega}) + \|\bar{v}_*\|_{3/2,2,\partial\Omega} + (|\xi| + |\omega|)(\|\bar{v}_*\|_{3/2,2,\partial\Omega} + \|z_{h*}^c\|_{3/2,2,\partial\Omega}) \right]. \end{aligned}$$

For R_0 and (ξ, ω) small enough, we deduce that (z_h, r_h, z_{h*}^c) is uniformly bounded in

$$[D^{1,2}(\Omega) \cap D^{2,2}(\Omega) \cap L_{1,\varpi}^\infty(\Omega)] \times W^{1,2}(\Omega) \times W^{3/2,2}(\partial\Omega).$$

Now using (3.6) or (3.9), depending on $\omega = 0$ or $\omega \neq 0$, we get

$$\|z_h\|_{2,\Omega} \leq C[(1 + |\xi|)(\|\nabla z_h\|_{2,\Omega} + |\Phi_h|) + \|r_h\|_{2,\Omega} + \|F_h\|_{2,\Omega}] + C'\|x\|F_h\|_{2,\Omega}$$

or

$$\begin{aligned} \|z_h\|_{2,\Omega} & \leq C \left(|\omega|^{-1/4} + \frac{|\omega \cdot \xi|^{1/2}}{|\omega|} \right) (|N_h| + |\omega| |\Phi_h|) + C'\|x\|F_h\|_{2,\Omega} \\ & \quad + C \left(1 + \frac{|\omega \times \xi|}{|\omega|^2} \right) [(1 + |\xi| + |\omega|)(\|\nabla z_h\|_{2,\Omega} + |\Phi_h|) + \|r_h\|_{2,\Omega} + \|F_h\|_{2,\Omega}] \end{aligned}$$

where

$$\begin{aligned} N_h & := \int_{\partial\Omega} [\sigma(z_h, r_h) + z_h \otimes V - (\omega \times x) \otimes z_h + F_h] n d\gamma \\ & = \int_{\partial\Omega} [\sigma(z_h, r_h)n + z_h(V - v) \cdot n - (v_h + \omega \times x)z_h \cdot n] d\gamma \\ & = \int_{\partial\Omega} [\sigma(z_h, r_h)n - (v_* + v_*^c) \cdot n(\bar{v}_* + z_{h*}^c) - (\bar{v}_* + z_{h*}^c) \cdot n(v_* + h\bar{v}_* + v_{h*}^c + V + \omega \times x)] d\gamma = 0 \end{aligned} \quad (6.27)$$

on account of the conditions (6.8) for tangential controls (then it is $(v_* + v_*^c) \cdot n = (\bar{v}_* + z_{h*}^c) \cdot n = 0$) or (6.18) if we are considering localized controls; in fact, since v and v_h satisfy (1.5), we should have $N_h = 0$ for z_h , no matter which kind of controls we would adopt. Moreover,

$$\Phi_h := \int_{\partial\Omega} n \cdot z_h d\gamma = \int_{\partial\Omega} n \cdot (z_{h*}^c + \bar{v}_*) d\gamma$$

which is zero if $z_{h*}^c \in \mathcal{C}_\tau$, $\bar{v}_* \in \mathcal{V}_\tau$. For localized controls, we use the estimate

$$|\Phi_h| = \left| \int_{\partial\Omega} n \cdot (z_{h*}^c + \bar{v}_*) d\gamma \right| \leq C(\|z_{h*}^c\|_{3/2,2,\partial\Omega} + \|\bar{v}_*\|_{3/2,2,\partial\Omega}).$$

Now we can use the estimate

$$\|(1 + |x|)F_h\|_{2,\Omega} = \|(1 + |x|)(z_h \otimes v + v \otimes z_h + h z_h \otimes z_h)\|_{2,\Omega} \leq C[\![z_h]\!]_{1,\varpi,\Omega}(\|v\|_{2,\Omega} + h\|z_h\|_{2,\Omega})$$

together with the uniform boundedness of z_h , r_h and z_{h*}^c in $D^{1,2}(\Omega) \cap D^{2,2}(\Omega) \cap L_{1,\varpi}^\infty(\Omega)$, $W^{1,2}(\Omega)$ and $W^{3/2,2}(\partial\Omega)$, respectively, to conclude that z_h is uniformly bounded in $L^2(\Omega)$ when h is close to zero.

Therefore, there exists

$$(z, r, z_*^C) \in [W^{2,2}(\Omega) \cap L_{1,\varpi}^\infty(\Omega)] \times W^{1,2}(\Omega) \times \mathcal{C}_\tau$$

such that

$$\varpi z_h \rightharpoonup \varpi z \quad \text{weakly } * \text{ in } L^\infty(\Omega), \quad (6.28)$$

$$(z_h, r_h) \rightharpoonup (z, r) \quad \text{weakly in } W^{2,2}(\Omega) \times W^{1,2}(\Omega), \quad (6.29)$$

$$z_{h*}^C \rightarrow z_*^C \quad \text{strongly in } W^{3/2,2}(\partial\Omega), \quad (6.30)$$

along a subsequence as $h \rightarrow 0$, where the strong convergence (6.30) follows from the same reasoning as in (5.3).

By classical compactness results, we also have

$$z_h \rightarrow z \quad \text{strongly in } W_{loc}^{1,2}(\overline{\Omega}),$$

and since $v_h = v + h z_h$,

$$v_h \rightarrow v \quad \text{strongly in } W_{loc}^{1,2}(\overline{\Omega}).$$

Proceeding as in the proof in Theorem 5.1, we can pass to the limit $h \rightarrow 0$ in (6.4)-(6.10) and show that (z, r, z_*^C) satisfies (6.11)-(6.17).

Now we prove the convergence of $z_h - z$ in the norm $\|\cdot\|_{2,2,\Omega} + [\cdot]_{1,\varpi,\Omega}$. Consider the problem

$$\begin{aligned} & -\operatorname{div} \sigma(z_h - z, r_h - r) - V \cdot \nabla(z_h - z) + \omega \times (z_h - z) \\ & = -[(v \cdot \nabla z_h + z_h \cdot \nabla v_h) - (v \cdot \nabla z + z \cdot \nabla v)] =: g_h \quad \text{in } \Omega \\ & \operatorname{div}(z_h - z) = 0 \quad \text{in } \Omega \\ & z_h - z = z_{h*}^C - z_*^C \quad \text{on } \partial\Omega \\ & \lim_{|x| \rightarrow \infty} (z_h(x) - z(x)) = 0 \\ & \int_{\partial\Omega} \sigma(z_h - z, r_h - r) n d\gamma = 0 \\ & \int_{\partial\Omega} x \times \sigma(z_h - z, r_h - r) n d\gamma = 0, \\ & z_{h*}^C - z_*^C \in \mathcal{C}_\tau. \end{aligned}$$

and notice that, since $v_h - v = h z_h$, we can write

$$\begin{aligned} -g_h & := (v \cdot \nabla z_h + z_h \cdot \nabla v_h) - (v \cdot \nabla z + z \cdot \nabla v) \\ & = v \cdot \nabla(z_h - z) + (z_h - z) \cdot \nabla v_h + z \cdot \nabla(v_h - v) \\ & = v \cdot \nabla(z_h - z) + (z_h - z) \cdot \nabla v + h z_h \cdot \nabla z_h \end{aligned}$$

and

$$-g_h = \operatorname{div}((z_h - z) \otimes v + v \otimes (z_h - z) + h z_h \otimes z_h) =: -\operatorname{div} G_h.$$

We use the fact that z_h is uniformly bounded in $W^{2,2}(\Omega) \cap L_{1,\varpi}^\infty(\Omega)$ to get the following estimates

$$\begin{aligned} \|g_h\|_{2,\Omega} & = \|v \cdot \nabla(z_h - z) + (z_h - z) \cdot \nabla v + h z_h \cdot \nabla z_h\|_{2,\Omega} \\ & \leq C [R_0(|z_h - z|_{1,2,\Omega} + [z_h - z]_{1,\varpi,\Omega}) + h [z_h]_{1,\varpi,\Omega} |z_h|_{1,2,\Omega}], \end{aligned}$$

$$[G_h]_{2,\varpi,\Omega} = [(z_h - z) \otimes v + v \otimes (z_h - z) + h z_h \otimes z_h]_{2,\varpi,\Omega} \leq C (R_0 [z_h - z]_{1,\varpi,\Omega} + h [z_h]_{1,\varpi,\Omega}^2),$$

with R_0 as specified above. Then using [12, Proposition 4.5], we first deduce that

$$\begin{aligned} & |z_h - z|_{1,2,\Omega} + |z_h - z|_{2,2,\Omega} + [z_h - z]_{1,\varpi,\Omega} + \|r_h - r\|_{1,2,\Omega} + \|z_{h*}^C - z_*^C\|_{3/2,2,\partial\Omega} \\ & \leq C \left[R_0 ([z_h - z]_{1,\varpi,\Omega} + |z_h - z|_{1,2,\Omega}) + (|\xi| + |\omega|) \|z_{h*}^C - z_*^C\|_{3/2,2,\partial\Omega} + h [z_h]_{1,\varpi,\Omega} |z_h|_{1,2,\Omega} + h [z_h]_{1,\varpi,\Omega}^2 \right]. \end{aligned}$$

This estimate yields a first convergence result for $z_h - z$:

$$|z_h - z|_{1,2,\Omega} + |z_h - z|_{2,2,\Omega} + [z_h - z]_{1,\varpi,\Omega} + \|r_h - r\|_{1,2,\Omega} + \|z_{h*}^c - z_*^c\|_{3/2,2,\partial\Omega} \rightarrow 0 \quad (h \rightarrow 0) \quad (6.31)$$

under suitable smallness assumptions on R_0 as well as (ξ, ω) and the uniform boundedness of z_h already established.

Let

$$N := \int_{\partial\Omega} [\sigma(z, r) + z \otimes V - (\omega \times x) \otimes z + F] n \, d\gamma, \quad (6.32)$$

where $F = -(z \otimes v + v \otimes z)$. Then we have

$$|N_h - N| \leq C \left\| (\nabla(z_h - z), r_h - r) \right\|_{2,\partial\Omega} + C(1 + \|v\|_{2,\partial\Omega}) \|z_{h*}^c - z_*^c\|_{2,\partial\Omega} + Ch \|z_{h*}^c + \bar{v}_*\|_{2,\partial\Omega}^2$$

which goes to zero as $h \rightarrow 0$ by (6.31), yielding $N = 0$ because of $N_h = 0$, see (6.27).

By (3.9), in the case $\omega \neq 0$ (the case $\omega = 0$ is even simpler), we have

$$\begin{aligned} \|z_h - z\|_{2,\Omega} \leq C \left(|\omega|^{-1/4} + \frac{|\omega \cdot \xi|^{1/2}}{|\omega|} \right) (|M_h| + |\omega| |\Psi_h|) + C' \| |x| G_h \|_{2,\Omega} \\ + C \left(1 + \frac{|\omega \times \xi|}{|\omega|^2} \right) [(1 + |\xi| + |\omega|) (\|\nabla(z_h - z)\|_{2,\Omega} + |\Psi_h|) + \|r_h - r\|_{2,\Omega} + \|G_h\|_{2,\Omega}], \end{aligned}$$

where

$$M_h := \int_{\partial\Omega} [\sigma(z_h - z, r_h - r) + (z_h - z) \otimes V - (\omega \times x) \otimes (z_h - z) + G_h] n \, d\gamma = N_h - N = 0 \quad (6.33)$$

on account of $N_h = N = 0$, see (6.27) and (6.32), no matter which kind of controls we would adopt, while we have

$$\Psi_h := \int_{\partial\Omega} n \cdot (z_h - z) \, d\gamma \rightarrow 0$$

as $h \rightarrow 0$ since $|\Psi_h| \leq C \|z_{h*}^c - z_*^c\|_{2,\partial\Omega}$. Hence, the estimate

$$\|(1 + |x|) G_h\|_{2,\Omega} \leq C (\|v\|_{2,\Omega} [z_h - z]_{1,\varpi,\Omega} + h [z_h]_{1,\varpi,\Omega} \|z_h\|_2)$$

and the previous convergence results (6.31) yield $\|z_h - z\|_{2,\Omega} \rightarrow 0$ when $h \rightarrow 0$. We have completed the proof provided that R_0 given by (4.37) is small enough as we have mentioned twice, which is accomplished through (6.24) with some $\kappa_1 \in (0, c_0]$. \square

7 Necessary first order conditions for an optimal control

In this section, we introduce the Lagrangian associated with problems (4.43) and (4.44), analyze the adjoint system and obtain a characterization of the optimal controls.

7.1 Introduction of the Lagrangian

Let us define

$$\mathcal{Y} := \{v \in W^{2,2}(\Omega) \cap L_{1,\varpi}^\infty(\Omega) ; \nabla \cdot v = 0 \text{ in } \Omega\}, \quad (7.1)$$

$$\begin{aligned} \mathcal{U} := \{u \in L^6(\Omega) \cap D^{1,2}(\Omega) \cap D^{2,2}(\Omega) ; V \cdot \nabla u - \omega \times u \in L^2(\Omega), \nabla \cdot u = 0 \text{ in } \Omega, \\ \exists \ell_u, k_u \in \mathbb{R}^3 \quad u = \ell_u + k_u \times x \quad \text{on } \partial\Omega\}, \quad (7.2) \end{aligned}$$

$$\mathcal{Z} := L^2(\partial\Omega). \quad (7.3)$$

Using these spaces, we can obtain a weak formulation for our problems (4.10)–(4.16).

Proposition 7.1. *Assume*

$$(v, p, v_*^c) \in \mathcal{Y} \times W^{1,2}(\Omega) \times \mathcal{C}_\chi \quad (7.4)$$

is a solution of (4.10)–(4.16) associated with $v_* \in \mathcal{V}_\Gamma$. Then

$$\begin{aligned} & 2 \int_{\Omega} Dv : Du \, dx - \int_{\Omega} (v \cdot \nabla u) \cdot v \, dx + \int_{\Omega} (V \cdot \nabla u - \omega \times u) \cdot v \, dx \\ & - m(\xi \times \omega) \cdot \ell_u - ((I\omega) \times \omega) \cdot k_u - \int_{\partial\Omega} \left((v_* + v_*^c) \cdot n \right) (\omega \times x) \cdot (\ell_u + k_u \times x) \, d\gamma = 0, \quad \forall u \in \mathcal{U}, \end{aligned} \quad (7.5)$$

$$\langle v - V - v_* - v_*^c, \zeta \rangle_{\partial\Omega} = 0, \quad \forall \zeta \in \mathcal{Z}. \quad (7.6)$$

If $(v, p, v_*^c) \in \mathcal{Y} \times W^{1,2}(\Omega) \times \mathcal{C}_\tau$ is a solution of (4.10)–(4.16) associated with $v_* \in \mathcal{V}_\tau$ then, a similar result holds true, but instead of (7.5), we have

$$\begin{aligned} & 2 \int_{\Omega} Dv : Du \, dx - \int_{\Omega} (v \cdot \nabla u) \cdot v \, dx + \int_{\Omega} (V \cdot \nabla u - \omega \times u) \cdot v \, dx \\ & - m(\xi \times \omega) \cdot \ell_u - ((I\omega) \times \omega) \cdot k_u = 0, \quad \forall u \in \mathcal{U}. \end{aligned} \quad (7.7)$$

Conversely, if $(v, v_*^c) \in \mathcal{Y} \times \mathcal{C}_\chi$ (resp. $\mathcal{Y} \times \mathcal{C}_\tau$) satisfies (7.5) (resp. (7.7)) together with (7.6), then there exists $p \in W^{1,2}(\Omega)$ such that (4.10)–(4.16) hold.

Proof. We use the same cut-off function ψ_R as in the final stage of the proof of Theorem 4.1. Recalling that $(\omega \times x) \cdot \nabla \psi_R = 0$ and (4.38), from which we obtain

$$\|V \cdot \nabla \psi_R\|_{3, \mathbb{R}^3} = \|\xi \cdot \nabla \psi_R\|_{3, \mathbb{R}^3} = C|\xi|$$

with a constant $C > 0$ independent of R . Assume $u \in \mathcal{U}$, multiply (4.10) by $\psi_R u$ and integrate by parts:

$$\begin{aligned} 0 &= - \int_{\Omega} \operatorname{div} \sigma(v, p) \cdot (\psi_R u) \, dx + \int_{\Omega} ((v - V) \cdot \nabla v) \cdot (\psi_R u) \, dx + \int_{\Omega} (\omega \times v) \cdot (\psi_R u) \, dx \\ &= - \int_{\partial\Omega} \sigma(v, p) n \cdot u \, d\gamma + 2 \int_{\Omega} \psi_R D(v) : D(u) \, dx - \int_{\Omega} \psi_R ((v - V) \cdot \nabla u) \cdot v \, dx \\ &\quad + \int_{\partial\Omega} (v - V) \cdot n (v \cdot u) \, dx - \int_{\Omega} \psi_R (\omega \times u) \cdot v \, dx \\ &\quad + \int_{\Omega} (2D(v) - p\mathbb{I}_3) : (u \otimes \nabla \psi_R) \, dx - \int_{\Omega} \nabla \psi_R \cdot (v - V)(u \cdot v) \, dx. \end{aligned} \quad (7.8)$$

On the other hand, using (4.14) and (4.15), we have

$$m(\xi \times \omega) \cdot \ell_u + ((I\omega) \times \omega) \cdot k_u + \int_{\partial\Omega} \left[-\sigma(v, p) n + \left((v_* + v_*^c) \cdot n \right) (v_* + v_*^c + V + \omega \times x) \right] \cdot u \, d\gamma = 0.$$

Combining the above relation with (7.8) yields

$$\begin{aligned} 0 &= 2 \int_{\Omega} \psi_R D(v) : D(u) \, dx - \int_{\Omega} \psi_R v \cdot \nabla u \cdot v \, dx + \int_{\Omega} \psi_R (V \cdot \nabla u - \omega \times u) \cdot v \, dx \\ &\quad + \int_{\Omega} (2D(v) - p\mathbb{I}_3) : (u \otimes \nabla \psi_R) \, dx - \int_{\Omega} \nabla \psi_R \cdot (v - V)(u \cdot v) \, dx \\ &\quad - m(\xi \times \omega) \cdot \ell_u - ((I\omega) \times \omega) \cdot k_u - \int_{\partial\Omega} \left((v_* + v_*^c) \cdot n \right) (\omega \times x) \cdot u \, d\gamma. \end{aligned} \quad (7.9)$$

Recalling (4.38) together with the summability properties given in (7.1) and (7.2), we get

$$\left| \int_{\Omega} (2D(v) - p\mathbb{I}_3) : (u \otimes \nabla \psi_R) \, dx \right| \leq \|\nabla \psi_R\|_{3, A_{R, 2R}} (\|D(v)\|_{2, \Omega} + \|p\|_{2, \Omega}) \|u\|_{6, A_{R, 2R}} \rightarrow 0 \quad \text{as } R \rightarrow \infty \quad (7.10)$$

and

$$\left| \int_{\Omega} \nabla \psi_R \cdot (v - V)(u \cdot v) \, dx \right| \leq C \left(\frac{1}{R} \|v\|_{2,\Omega}^2 \|u\|_{\infty,\Omega} + \|\nabla \psi_R\|_{3,A_R,2R} \|v\|_{2,\Omega} \|u\|_{6,A_R,2R} \right). \quad (7.11)$$

Hence, letting $R \rightarrow \infty$ in (7.9), yields (7.5).

Conversely, by taking $u \in \mathcal{D}(\Omega)$ in (7.5), we find that there exists $p \in L^2_{\text{loc}}(\overline{\Omega})$ such that (4.10) holds. Then applying [12, Proposition 2.1] with $v \otimes v \in L^2_{\infty,\varpi}(\Omega)$ and $f = -\text{div}(v \otimes v) \in L^2(\Omega)$, we deduce that $p \in W^{1,2}(\Omega)$. Finally, multiplying (4.10) by $\psi_R u \in \mathcal{U}$ as above, taking $R \rightarrow \infty$ and comparing with (7.5), we obtain (4.14) and (4.15). \square

Remark 7.2. *The summability properties assumed for v and u imply that the weak formulation (7.5) is meaningful. In particular, $v \in L^2(\Omega) \cap L^\infty(\Omega)$ and $\nabla u \in L^2(\Omega)$ guarantee that the integral $\int_{\Omega} (v \cdot \nabla u) \cdot v \, dx$ is finite. Moreover, the integral $\int_{\Omega} [(\omega \times x) \cdot \nabla u - \omega \times u] \cdot v \, dx$ is finite because $(\omega \times x) \cdot \nabla u - \omega \times u \in L^2(\Omega)$.*

Proposition 7.1 and the above remarks lead to the following definition of the Lagrangians:

$$\begin{aligned} \mathcal{L}_{\Gamma}(v, v_*^c, v_*, u, \zeta) &:= \int_{\Omega} |D(v)|^2 \, dx + \frac{1}{4} \int_{\partial\Omega} (v_* + v_*^c) \cdot n |V + v_* + v_*^c|^2 \, d\gamma \\ &\quad - 2 \int_{\Omega} D(v) : D(u) \, dx + \int_{\Omega} (v \cdot \nabla u) \cdot v \, dx \\ &\quad - \int_{\Omega} (V \cdot \nabla u - \omega \times u) \cdot v \, dx + m(\xi \times \omega) \cdot \ell_u + ((I\omega) \times \omega) \cdot k_u \\ &\quad + \int_{\partial\Omega} \left((v_* + v_*^c) \cdot n \right) (\omega \times x) \cdot (\ell_u + k_u \times x) \, d\gamma - \langle v - V - v_* - v_*^c, \zeta \rangle_{\partial\Omega} \end{aligned} \quad (7.12)$$

for $(v, v_*^c, v_*, u, \zeta) \in \mathcal{Y} \times \mathcal{C}_{\chi} \times \mathcal{V}_{\Gamma} \times \mathcal{U} \times \mathcal{Z}$, and

$$\begin{aligned} \mathcal{L}_{\tau}(v, v_*^c, v_*, u, \zeta) &:= \int_{\Omega} |D(v)|^2 \, dx - 2 \int_{\Omega} D(v) : D(u) \, dx + \int_{\Omega} (v \cdot \nabla u) \cdot v \, dx \\ &\quad - \int_{\Omega} (V \cdot \nabla u - \omega \times u) \cdot v \, dx + m(\xi \times \omega) \cdot \ell_u + ((I\omega) \times \omega) \cdot k_u \\ &\quad - \langle v - V - v_* - v_*^c, \zeta \rangle_{\partial\Omega} \end{aligned} \quad (7.13)$$

for $(v, v_*^c, v_*, u, \zeta) \in \mathcal{Y} \times \mathcal{C}_{\tau} \times \mathcal{V}_{\tau} \times \mathcal{U} \times \mathcal{Z}$.

Let \widehat{v}_* be a solution of the problem (4.44) and denote by

$$(\widehat{v}, \widehat{v}_*^c, \widehat{p}) \in \mathcal{Y} \times \mathcal{C}_{\chi} \times W^{1,2}(\Omega)$$

the corresponding solution of (4.10)–(4.16) given by Theorem 4.1. We now obtain the adjoint system by considering the equations

$$D_v \mathcal{L}_{\Gamma}(\widehat{v}, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta})v = 0 \quad \forall v \in \mathcal{Y}, \quad (7.14)$$

$$D_{v_*^c} \mathcal{L}_{\Gamma}(\widehat{v}, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta})v_*^c = 0 \quad \forall v_*^c \in \mathcal{C}_{\chi} \quad (7.15)$$

for the unknowns $\widehat{u}, \widehat{\zeta}$. In the case of tangential controls, equations (7.14) and (7.15) are replaced by

$$D_v \mathcal{L}_{\tau}(\widehat{v}, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta})v = 0 \quad \forall v \in \mathcal{Y}, \quad (7.16)$$

$$D_{v_*^c} \mathcal{L}_{\tau}(\widehat{v}, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta})v_*^c = 0 \quad \forall v_*^c \in \mathcal{C}_{\tau}. \quad (7.17)$$

By computing the Gâteaux derivatives of \mathcal{L}_{Γ} or \mathcal{L}_{τ} , we can rewrite the above equations. Such a calculation is standard but for sake of completeness, we give it in the case of (7.16): we have to pass to the limit $h \rightarrow 0$ in

$$\frac{\mathcal{L}_{\tau}(\widehat{v} + hv, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}_{\tau}(\widehat{v}, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta})}{h}$$

where

$$\begin{aligned} \mathcal{L}_\tau(\widehat{v} + hv, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta}) &= \int_\Omega |D(\widehat{v} + hv)|^2 dx - 2 \int_\Omega D(\widehat{v} + hv) : D(\widehat{u}) dx + \int_\Omega ((\widehat{v} + hv) \cdot \nabla \widehat{u}) \cdot (\widehat{v} + hv) dx \\ &\quad - \int_\Omega (V \cdot \nabla \widehat{u} - \omega \times \widehat{u}) \cdot (\widehat{v} + hv) dx + m(\xi \times \omega) \cdot \ell_{\widehat{u}} + ((I\omega) \times \omega) \cdot k_{\widehat{u}} \\ &\quad - \langle \widehat{v} + hv - V - \widehat{v}_* - \widehat{v}_*^c, \widehat{\zeta} \rangle_{\partial\Omega} \end{aligned}$$

and

$$\begin{aligned} \mathcal{L}_\tau(\widehat{v}, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta}) &= \int_\Omega |D(\widehat{v})|^2 dx - 2 \int_\Omega D(\widehat{v}) : D(\widehat{u}) dx + \int_\Omega (\widehat{v} \cdot \nabla \widehat{u}) \cdot \widehat{v} dx \\ &\quad - \int_\Omega (V \cdot \nabla \widehat{u} - \omega \times \widehat{u}) \cdot \widehat{v} dx + m(\xi \times \omega) \cdot \ell_{\widehat{u}} + ((I\omega) \times \omega) \cdot k_{\widehat{u}} \\ &\quad - \langle \widehat{v} - V - \widehat{v}_* - \widehat{v}_*^c, \widehat{\zeta} \rangle_{\partial\Omega}. \end{aligned}$$

Simplifying the above expressions and letting $h \rightarrow 0$ in

$$\begin{aligned} &\frac{\mathcal{L}_\tau(\widehat{v} + hv, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}_\tau(\widehat{v}, \widehat{v}_*^c, \widehat{v}_*, \widehat{u}, \widehat{\zeta})}{h} \\ &= 2 \int_\Omega D(\widehat{v}) : D(v) dx + h \int_\Omega |D(v)|^2 \\ &\quad + \int_\Omega (\widehat{v} \cdot \nabla \widehat{u}) \cdot v dx + \int_\Omega (v \cdot \nabla \widehat{u}) \cdot \widehat{v} dx + h \int_\Omega (v \cdot \nabla \widehat{u}) \cdot v dx \\ &\quad - 2 \int_\Omega D(v) : D(\widehat{u}) dx - \int_\Omega [(V \cdot \nabla) \widehat{u} - \omega \times \widehat{u}] \cdot v dx - \langle v, \widehat{\zeta} \rangle_{\partial\Omega} \end{aligned}$$

yields that (7.16) is equivalent to

$$\begin{aligned} 2 \int_\Omega D(\widehat{v} - \widehat{u}) : D(v) dx + \int_\Omega (\widehat{v} \cdot \nabla \widehat{u}) \cdot v dx + \int_\Omega (v \cdot \nabla \widehat{u}) \cdot \widehat{v} dx \\ - \int_\Omega (V \cdot \nabla \widehat{u} - \omega \times \widehat{u}) \cdot v dx - \langle v, \widehat{\zeta} \rangle_{\partial\Omega} = 0, \quad \forall v \in \mathcal{Y}. \end{aligned} \quad (7.18)$$

By similar calculations, we see that the relation (7.14) is also equivalent to (7.18), whereas (7.15) and (7.17) can be respectively written as follows:

$$\begin{aligned} \frac{1}{4} \int_{\partial\Omega} v_*^c \cdot n |V + \widehat{v}_* + \widehat{v}_*^c|^2 d\gamma + \frac{1}{2} \int_{\partial\Omega} (\widehat{v}_* + \widehat{v}_*^c) \cdot n (V + \widehat{v}_* + \widehat{v}_*^c) \cdot v_*^c d\gamma \\ + \int_{\partial\Omega} (v_*^c \cdot n) (\omega \times x) \cdot \widehat{u} d\gamma + \langle v_*^c, \widehat{\zeta} \rangle_{\partial\Omega} = 0, \quad \forall v_*^c \in \mathcal{C}_\chi, \end{aligned} \quad (7.19)$$

and

$$\langle v_*^c, \widehat{\zeta} \rangle_{\partial\Omega} = 0, \quad \forall v_*^c \in \mathcal{C}_\tau. \quad (7.20)$$

Once we have a solution $(\widehat{u}, \widehat{\zeta}) \in \mathcal{U} \times \mathcal{Z}$ to (7.18), see (7.2)–(7.3), we deduce that there exists a pressure $\widehat{q} \in L_{\text{loc}}^2(\overline{\Omega})$ which together with \widehat{u} obeys

$$-\operatorname{div} \sigma(\widehat{u} - \widehat{v}, \widehat{q} - \widehat{p}) - \widehat{v} \cdot \nabla \widehat{u} - (\nabla \widehat{u})^\top \widehat{v} + V \cdot \nabla \widehat{u} - \omega \times \widehat{u} = 0 \quad \text{in } \Omega \quad (7.21)$$

$$\operatorname{div} \widehat{u} = 0 \quad \text{in } \Omega \quad (7.22)$$

$$\widehat{u} = \ell_{\widehat{u}} + k_{\widehat{u}} \times x \quad \text{on } \partial\Omega \quad (7.23)$$

$$\lim_{|x| \rightarrow \infty} \widehat{u}(x) = 0. \quad (7.24)$$

Since $\widehat{v} \cdot \nabla \widehat{u} + (\nabla \widehat{u})^\top \widehat{v} - \Delta \widehat{v} + \nabla \widehat{p} \in L^2(\Omega) \cap D^{-1,2}(\Omega)$, see Lemma 7.3 below, we employ Proposition 2.1 to see that the pressure $\widehat{q} \in W^{1,2}(\Omega)$ can be singled out.

Taking the scalar product of (7.21) with $\psi_R v$, where $v \in \mathcal{Y}$ is arbitrary and ψ_R is the same cut-off function as in the final stage of the proof of Theorem 4.1, integrating by parts and letting $R \rightarrow \infty$ (where $\widehat{q} \in L^2(\Omega)$ is used), we deduce from (7.18) that

$$\widehat{\zeta} = \sigma(\widehat{v} - \widehat{u}, \widehat{p} - \widehat{q})n. \quad (7.25)$$

Recalling that $\mathcal{Z} = L^2(\partial\Omega)$ and replacing (7.25) in (7.19), yields

$$\begin{aligned} & \int_{\partial\Omega} (\sigma(\widehat{v} - \widehat{u}, \widehat{p} - \widehat{q})n) \cdot v_*^c \, d\gamma + \frac{1}{4} \int_{\partial\Omega} v_*^c \cdot n |V + \widehat{v}_* + \widehat{v}_*^c|^2 \, d\gamma \\ & + \frac{1}{2} \int_{\partial\Omega} (\widehat{v}_* + \widehat{v}_*^c) \cdot n (V + \widehat{v}_* + \widehat{v}_*^c) \cdot v_*^c \, d\gamma + \int_{\partial\Omega} (v_*^c \cdot n) (\omega \times x) \cdot \widehat{u} \, d\gamma = 0, \quad \forall v_*^c \in \mathcal{C}_\chi. \end{aligned} \quad (7.26)$$

In the case of tangential controls, (7.20) takes the form

$$\int_{\partial\Omega} (\sigma(\widehat{v} - \widehat{u}, \widehat{p} - \widehat{q})n) \cdot v_*^c \, d\gamma = 0, \quad \forall v_*^c \in \mathcal{C}_\tau. \quad (7.27)$$

7.2 Well-posedness of the adjoint system

Now we show that the adjoint system (7.21)–(7.24) subject to (7.26)/(7.27) is well-posed. With $\widehat{v} \in \mathcal{Y}$ given, we can define the following mapping

$$\mathfrak{F} = \mathfrak{F}_{\widehat{v}} : D^{1,2}(\Omega) \rightarrow L^2(\Omega) \cap D^{-1,2}(\Omega), \quad u \mapsto \widehat{v} \cdot \nabla u + (\nabla u)^\top \widehat{v}. \quad (7.28)$$

Lemma 7.3. *Assume that $\widehat{v} \in \mathcal{Y}$. Then the mapping \mathfrak{F} is well-defined and continuous:*

$$\|\mathfrak{F}(u)\|_{L^2(\Omega) \cap D^{-1,2}(\Omega)} \leq C[\widehat{v}]_{1,\varpi,\Omega} |u|_{1,2,\Omega}.$$

Proof. Both terms in the formula of $\mathfrak{F}_{\widehat{v}}(u)$ can be handled in the same way. First, it is immediate to obtain

$$\|\widehat{v} \cdot \nabla u + (\nabla u)^\top \widehat{v}\|_{2,\Omega} \leq 2\|\widehat{v}\|_{\infty,\Omega} \|\nabla u\|_{2,\Omega} \leq 2[\widehat{v}]_{1,\varpi,\Omega} |u|_{1,2,\Omega}.$$

Now we notice that, for arbitrary $\varphi \in D_0^{1,2}(\Omega)$, by the Hardy inequality,

$$\langle \widehat{v} \cdot \nabla u + (\nabla u)^\top \widehat{v}, \varphi \rangle_{D^{-1,2}(\Omega), D_0^{1,2}(\Omega)} \leq C[\widehat{v}]_{1,\varpi,\Omega} \|\nabla u\|_{2,\Omega} \left\| \frac{\varphi}{\varpi} \right\|_{2,\Omega} \leq C[\widehat{v}]_{1,\varpi,\Omega} \|\nabla u\|_{2,\Omega} \|\nabla \varphi\|_{2,\Omega},$$

where $\varpi(x)$ is given by (4.17) and therefore

$$|\widehat{v} \cdot \nabla u + (\nabla u)^\top \widehat{v}|_{-1,2,\Omega} \leq C[\widehat{v}]_{1,\varpi,\Omega} |u|_{1,2,\Omega}.$$

□

Theorem 7.4. *There exists a constant $\kappa_2 \in (0, c_0]$ depending on Ω such that if $\xi, \omega \in \mathbb{R}^3$ and $v_* \in \mathcal{V}_\Gamma$ (resp. $v_* \in \mathcal{V}_\tau$) satisfy*

$$|\xi| \leq \kappa_2, \quad |\omega| \leq \kappa_2, \quad \|v_*\|_{3/2,2,\partial\Omega} \leq \kappa_2, \quad (7.29)$$

then the adjoint system (7.21)–(7.24) subject to (7.26) (resp. (7.27)) admits a unique solution $(\widehat{u}, \widehat{q}, \ell_{\widehat{u}}, k_{\widehat{u}}) \in \mathcal{U} \times W^{1,2}(\Omega) \times \mathbb{R}^3 \times \mathbb{R}^3$.

Proof. We show only the case of localized boundary values since the tangential case is similar. Consider the space

$$\mathcal{B} := \{u \in L^6(\Omega) \cap D^{1,2}(\Omega) \cap D^{2,2}(\Omega) ; \nabla \cdot u = 0 \text{ in } \Omega\},$$

with the norm $\|u\|_{\mathcal{B}} := |u|_{1,2,\Omega} + |u|_{2,2,\Omega}$.

Assume $\bar{u} \in \mathcal{B}$ and consider the unique solution of the linear problem

$$-\operatorname{div} \sigma(u, q) + V \cdot \nabla u - \omega \times u = \mathfrak{F}(\bar{u}) - \operatorname{div} \sigma(\widehat{v}, \widehat{p}) \quad \text{in } \Omega \quad (7.30)$$

$$\operatorname{div} u = 0 \quad \text{in } \Omega \quad (7.31)$$

$$u = a + b \times x \quad \text{on } \partial\Omega \quad (7.32)$$

$$\lim_{|x| \rightarrow \infty} u(x) = 0, \quad (7.33)$$

$$\int_{\partial\Omega} (\sigma(u, q)n) \cdot v_*^c \, d\gamma = \int_{\partial\Omega} (\mathfrak{G}(\bar{u}) + \mathfrak{H}) \cdot v_*^c \, d\gamma, \quad \forall v_*^c \in \mathcal{C}_\chi, \quad (7.34)$$

where

$$\begin{aligned}\mathfrak{G}(\bar{u}) &:= [(\omega \times x) \cdot \bar{u}]n, \\ \mathfrak{H} &:= \sigma(\hat{v}, \hat{p})n + \frac{1}{4}|V + \hat{v}_* + \hat{v}_*^c|^2 n + \frac{1}{2}(\hat{v}_* + \hat{v}_*^c) \cdot n(V + \hat{v}_* + \hat{v}_*^c).\end{aligned}$$

and $\mathfrak{F}(\bar{u})$ is given by (7.28).

More precisely, the above linear system is solved by decomposing u and q as follows

$$u = \sum_{i=1}^3 a_i v^{(i)} + \sum_{i=1}^3 b_i V^{(i)} + u_f, \quad q = \sum_{i=1}^3 a_i q^{(i)} + \sum_{i=1}^3 b_i Q^{(i)} + q_f \quad (7.35)$$

where $(v^{(i)}, q^{(i)})$ and $(V^{(i)}, Q^{(i)})$ are given by (4.6) and (4.7), whereas (u_f, q_f) is the solution of

$$\begin{aligned}-\operatorname{div} \sigma(u_f, q_f) + V \cdot \nabla u_f - \omega \times u_f &= \mathfrak{F}(\bar{u}) - \operatorname{div} \sigma(\hat{v}, \hat{p}) \quad \text{in } \Omega \\ \operatorname{div} u_f &= 0 \quad \text{in } \Omega \\ u_f &= 0 \quad \text{on } \partial\Omega \\ \lim_{|x| \rightarrow \infty} u_f(x) &= 0,\end{aligned}$$

obtained in Proposition 2.1. Then (u, p) automatically satisfies (7.30)–(7.33). It only remains to choose (a, b) such that (7.34) holds. Using the basis (4.5) of \mathcal{C}_χ , we reduce (7.34) to

$$\int_{\partial\Omega} (\sigma(u, q)n) \cdot \chi g^{(i)} d\gamma = \int_{\partial\Omega} (\mathfrak{G}(\bar{u}) + \mathfrak{H}) \cdot \chi g^{(i)} d\gamma \quad (i = 1, 2, 3), \quad (7.36)$$

$$\int_{\partial\Omega} (\sigma(u, q)n) \cdot \chi G^{(i)} d\gamma = \int_{\partial\Omega} (\mathfrak{G}(\bar{u}) + \mathfrak{H}) \cdot \chi G^{(i)} d\gamma \quad (i = 1, 2, 3), \quad (7.37)$$

which, in view of (7.35) together with (4.8)–(4.9), can be written as

$$A \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} c \\ d \end{bmatrix} \quad (7.38)$$

where

$$A_{i,j} := \int_{\partial\Omega} \chi g^{(i)} \cdot g^{(j)} d\gamma \quad (i, j \leq 3), \quad A_{i,j} := \int_{\partial\Omega} \chi g^{(i)} \cdot G^{(j-3)} d\gamma \quad (i \leq 3, j \geq 4), \quad (7.39)$$

$$A_{i,j} := \int_{\partial\Omega} \chi G^{(i-3)} \cdot g^{(j)} d\gamma \quad (i \geq 4, j \leq 3), \quad A_{i,j} := \int_{\partial\Omega} \chi G^{(i-3)} \cdot G^{(j-3)} d\gamma \quad (i, j \geq 4), \quad (7.40)$$

$$c_i := \int_{\partial\Omega} (\mathfrak{G}(\bar{u}) + \mathfrak{H} - \sigma(u_f, q_f)n) \cdot \chi g^{(i)} d\gamma \quad (i = 1, 2, 3), \quad (7.41)$$

$$d_i := \int_{\partial\Omega} (\mathfrak{G}(\bar{u}) + \mathfrak{H} - \sigma(u_f, q_f)n) \cdot \chi G^{(i)} d\gamma \quad (i = 1, 2, 3). \quad (7.42)$$

By [12, Lemma 4.3] we know that, for ξ and ω satisfying (4.39), the matrix A is invertible; in fact, this is needed and even crucial in Theorem 4.1 although it is hidden in [12, Proposition 4.5]. Thus, we obtain the existence and uniqueness of (a, b) satisfying (7.38)–(7.42) and deduce that for any $\bar{u} \in \mathcal{B}$ there exists a unique solution $(u, q, a, b) \in \mathcal{B} \times W^{1,2}(\Omega) \times \mathbb{R}^3 \times \mathbb{R}^3$ of (7.30)–(7.34). By the equation (7.30) we have $V \cdot \nabla u - \omega \times u \in L^2(\Omega)$ as well, so that $u \in \mathcal{U}$.

To obtain the existence of a solution of the adjoint system (7.21)–(7.24) and (7.26), we only need to show that the mapping

$$\Xi : \mathcal{B} \rightarrow \mathcal{B}, \quad \bar{u} \mapsto u$$

is contractive as long as (ξ, ω) and v_* are small enough. Using the linearity of the adjoint system, it is sufficient to consider the system

$$\begin{aligned}-\operatorname{div} \sigma(u, q) + V \cdot \nabla u - \omega \times u &= \mathfrak{F}(\bar{u}) \quad \text{in } \Omega \\ \operatorname{div} u &= 0 \quad \text{in } \Omega \\ u &= a + b \times x \quad \text{on } \partial\Omega \\ \lim_{|x| \rightarrow \infty} u(x) &= 0,\end{aligned}$$

with

$$\begin{bmatrix} a \\ b \end{bmatrix} = A^{-1} \begin{bmatrix} c \\ d \end{bmatrix} \quad (7.43)$$

where

$$c_i := \int_{\partial\Omega} (\mathfrak{G}(\bar{u}) - \sigma(u_f, q_f)n) \cdot \chi g^{(i)} d\gamma \quad (i = 1, 2, 3), \quad (7.44)$$

$$d_i := \int_{\partial\Omega} (\mathfrak{G}(\bar{u}) - \sigma(u_f, q_f)n) \cdot \chi G^{(i)} d\gamma \quad (i = 1, 2, 3) \quad (7.45)$$

and

$$\begin{aligned} -\operatorname{div} \sigma(u_f, q_f) + (V \cdot \nabla)u_f - \omega \times u_f &= \mathfrak{F}(\bar{u}) \quad \text{in } \Omega \\ \operatorname{div} u_f &= 0 \quad \text{in } \Omega \\ u_f &= 0 \quad \text{on } \partial\Omega \\ \lim_{|x| \rightarrow \infty} u_f(x) &= 0. \end{aligned}$$

Using the trace theorem and Proposition 2.1, we have

$$|(a, b)| \leq C (\|\mathfrak{F}(\bar{u})\|_{L^2(\Omega) \cap D^{-1,2}(\Omega)} + |\omega| |\bar{u}|_{1,2,\Omega})$$

and thus, using again Proposition 2.1 and Lemma 7.3 we deduce

$$\|\Xi(\bar{u})\|_{\mathcal{B}} = \|u\|_{\mathcal{B}} \leq C (\|\mathfrak{F}(\bar{u})\|_{L^2(\Omega) \cap D^{-1,2}(\Omega)} + |(a, b)|) \leq C (\lceil \widehat{v} \rceil_{1,\varpi,\Omega} + |\omega|) |\bar{u}|_{1,2,\Omega} \leq C(R_0 + |\omega|) \|\bar{u}\|_{\mathcal{B}},$$

where R_0 is given by (4.37). This yields the existence and uniqueness of a solution of the adjoint system (7.21)–(7.24) subject to (7.26) under the condition (7.29) with $\kappa_2 \in (0, c_0]$ small enough. \square

7.3 Optimality condition

Recall the mappings $\Lambda_\tau : v_* \mapsto (v, v_*^c)$ and $\Lambda_\chi : v_* \mapsto (v, v_*^c)$ defined by (6.2) and (6.3), respectively. For both cases, we abbreviate them to Λ and, similarly, we write \mathcal{L} instead of \mathcal{L}_τ or \mathcal{L}_χ . Because of Proposition 7.1, the functional (4.42) can be written as

$$J(v_*) = 2 \mathcal{L}(\Lambda(v_*), v_*, u, \zeta), \quad (7.46)$$

no matter which $(u, \zeta) \in \mathcal{U} \times \mathcal{Z}$ may be. Assume (7.29) and let us take, in particular, the solution $\widehat{u} \in \mathcal{U}$ to the adjoint system (7.21)–(7.24) subject to (7.26)/(7.27) obtained in Theorem 7.4 together with $\widehat{\zeta} \in \mathcal{Z}$ given by (7.25) so that (7.14)–(7.15) or (7.16)–(7.17) are satisfied.

We are now in a position to provide the optimality conditions for problems (4.43) and (4.44):

Theorem 7.5. *Let Ω be of class C^3 . Set $\kappa_0 := \min\{\kappa_1, \kappa_2\}$, where κ_1 and κ_2 are the constants in Theorem 6.1 and Theorem 7.4, respectively. Suppose that $\xi, \omega \in \mathbb{R}^3$ and $v_* \in \mathcal{V}_\Gamma$ (resp. \mathcal{V}_τ) satisfy*

$$|\xi| \leq \kappa_0, \quad |\omega| \leq \kappa_0, \quad \|v_*\|_{3/2,2,\partial\Omega} \leq \kappa_0.$$

Let \widehat{v}_ be a solution of the optimal control problem (4.44) (resp. (4.43)) with $\kappa = \kappa_0$, $(\widehat{v}, \widehat{v}_*^c, \widehat{p})$ the corresponding state obtained in Theorem 4.1 and $(\widehat{u}, \widehat{q})$ the solution of the adjoint system (7.21)–(7.24) subject to (7.26) (resp. (7.27)) obtained in Theorem 7.4. Then we have*

$$\begin{aligned} &\int_{\partial\Omega} (\sigma(\widehat{v} - \widehat{u}, \widehat{p} - \widehat{q})n) \cdot (v_* - \widehat{v}_*) d\gamma + \frac{1}{4} \int_{\partial\Omega} (v_* - \widehat{v}_*) \cdot n |V + \widehat{v}_* + \widehat{v}_*^c|^2 d\gamma \\ &+ \frac{1}{2} \int_{\partial\Omega} (\widehat{v}_* + \widehat{v}_*^c) \cdot n (V + \widehat{v}_* + \widehat{v}_*^c) \cdot (v_* - \widehat{v}_*) d\gamma + \int_{\partial\Omega} ((v_* - \widehat{v}_*) \cdot n) (\omega \times x) \cdot \widehat{u} d\gamma \geq 0, \quad \forall v_* \in \mathcal{V}_\Gamma^{\kappa_0}, \end{aligned} \quad (7.47)$$

in the case of localized controls, while

$$\int_{\partial\Omega} (\sigma(\widehat{v} - \widehat{u}, \widehat{p} - \widehat{q})n) \cdot (v_* - \widehat{v}_*) d\gamma \geq 0, \quad \forall v_* \in \mathcal{V}_\tau^{\kappa_0}, \quad (7.48)$$

in the case of tangential controls.

Proof. We begin by showing that

$$\frac{1}{2} D_{v_*} J(\widehat{v}_*) v_* = D_{(v, v_*^c)} \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta}) D_{v_*} \Lambda(\widehat{v}_*) v_* + D_{v_*} \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta}) v_*, \quad (7.49)$$

where the direction $v_* \in \mathcal{V}_\Gamma$ must be taken such that $\langle \widehat{v}_*, v_* \rangle_{W^{3/2,2}(\partial\Omega)} < 0$ if $\|\widehat{v}_*\|_{3/2,2,\partial\Omega} = \kappa_0$, whereas it can be arbitrary if $\|\widehat{v}_*\|_{3/2,2,\partial\Omega} < \kappa_0$.

In order to compute

$$D_{v_*} J(\widehat{v}_*) v_* = \lim_{h \rightarrow 0} \frac{J(\widehat{v}_* + hv_*) - J(\widehat{v}_*)}{h}$$

we write

$$\begin{aligned} \frac{1}{2} \frac{J(\widehat{v}_* + hv_*) - J(\widehat{v}_*)}{h} &= \frac{\mathcal{L}(\Lambda(\widehat{v}_* + hv_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta})}{h} \\ &= \frac{\mathcal{L}(\Lambda(\widehat{v}_* + hv_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta})}{h} \\ &\quad + \frac{\mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta})}{h} \end{aligned}$$

and, by the definition of Gateaux derivative, it is clear that

$$\lim_{h \rightarrow 0} \frac{\mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta})}{h} = D_{v_*} \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta}) v_*.$$

The relation

$$\Lambda(\widehat{v}_* + hv_*) = \Lambda(\widehat{v}_*) + h D_{v_*} \Lambda(\widehat{v}_*) v_* + o(h) = \Lambda(\widehat{v}_*) + h(\widehat{z}_h, \widehat{z}_{h*}^c),$$

with $(\widehat{z}_h, \widehat{z}_{h*}^c) := D_{v_*} \Lambda(\widehat{v}_*) v_* + o(h)/h$, yields

$$\begin{aligned} &\frac{\mathcal{L}(\Lambda(\widehat{v}_* + hv_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta})}{h} \\ &= \frac{\mathcal{L}(\Lambda(\widehat{v}_*) + h(\widehat{z}_h, \widehat{z}_{h*}^c), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta})}{h} \\ &= D_{(v, v_*^c)} \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta})(\widehat{z}_h, \widehat{z}_{h*}^c) + \frac{o(h)}{h}. \end{aligned}$$

Therefore

$$\lim_{h \rightarrow 0} \frac{\mathcal{L}(\Lambda(\widehat{v}_* + hv_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta})}{h} = \lim_{h \rightarrow 0} D_{(v, v_*^c)} \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta})(\widehat{z}_h, \widehat{z}_{h*}^c).$$

Note that, by Theorem 6.1, the convergence

$$(\widehat{z}_h, \widehat{z}_{h*}^c) = \frac{\Lambda(\widehat{v}_* + hv_*) - \Lambda(\widehat{v}_*)}{h} \rightarrow D_{v_*} \Lambda(\widehat{v}_*) v_* =: (\widehat{z}, \widehat{z}^c) \quad (h \rightarrow 0)$$

means that

$$\|\widehat{z}_h - \widehat{z}\|_{2,2,\Omega} + [\widehat{z}_h - \widehat{z}]_{1,\varpi,\Omega} + \|\widehat{z}_{h*}^c - \widehat{z}^c\|_{3/2,2,\partial\Omega} \rightarrow 0. \quad (7.50)$$

In the case of localized controls, we have

$$\begin{aligned} D_v \mathcal{L}_\Gamma(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) \widehat{z}_h &= 2 \int_\Omega D(\widehat{v} - \widehat{u}) : D(\widehat{z}_h) dx + \int_\Omega (\widehat{v} \cdot \nabla \widehat{u}) \cdot \widehat{z}_h dx + \int_\Omega (\widehat{z}_h \cdot \nabla \widehat{u}) \cdot \widehat{v} dx \\ &\quad - \int_\Omega [(V \cdot \nabla) \widehat{u} - \omega \times \widehat{u}] \cdot \widehat{z}_h dx - \int_{\partial\Omega} \widehat{z}_h \cdot \widehat{\zeta} ds \quad (\widehat{z}_h \in \mathcal{Y}), \end{aligned} \quad (7.51)$$

$$\begin{aligned} D_{v_*^c} \mathcal{L}_\Gamma(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) \widehat{z}_{h*}^c &= \frac{1}{4} \int_{\partial\Omega} \widehat{z}_{h*}^c \cdot n |V + \widehat{v}_* + hv_* + \widehat{v}_*^c|^2 d\gamma \\ &\quad + \frac{1}{2} \int_{\partial\Omega} (\widehat{v}_* + hv_* + \widehat{v}_*^c) \cdot n (V + \widehat{v}_* + hv_* + \widehat{v}_*^c) \cdot \widehat{z}_{h*}^c d\gamma \\ &\quad + \int_{\partial\Omega} (\widehat{z}_{h*}^c \cdot n) (\omega \times x) \cdot \widehat{u} d\gamma + \langle \widehat{z}_{h*}^c, \widehat{\zeta} \rangle_{\partial\Omega} \quad (\widehat{z}_{h*}^c \in \mathcal{C}_\chi), \end{aligned} \quad (7.52)$$

and analogously for \mathcal{L}_τ , so that by taking the limit $h \rightarrow 0$ in (7.51) and (7.52), and using (7.50), we get

$$\lim_{h \rightarrow 0} \frac{\mathcal{L}(\Lambda(\widehat{v}_* + hv_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta}) - \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_* + hv_*, \widehat{u}, \widehat{\zeta})}{h} = D_{(v, v_*^c)} \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta}) D_{v_*} \Lambda(\widehat{v}_*) v_*.$$

Having shown (7.49) and using the definition of the adjoint system, specifically (7.14)–(7.15) or (7.16)–(7.17) we get

$$\frac{1}{2} D_{v_*} J(\widehat{v}_*) v_* = D_{v_*} \mathcal{L}(\Lambda(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta}) v_*. \quad (7.53)$$

Now, using that $\mathcal{V}_\tau^{\kappa_0}$ and $\mathcal{V}_\Gamma^{\kappa_0}$ are convex sets, we deduce from $J(\widehat{v}_* + h(v_* - \widehat{v}_*)) \geq J(\widehat{v}_*)$ for every $v_* \in \mathcal{V}_\tau^{\kappa_0}$ or $\mathcal{V}_\Gamma^{\kappa_0}$ and $h \in (0, 1)$ that

$$\frac{1}{2} D_{v_*} J(\widehat{v}_*)(v_* - \widehat{v}_*) = D_{v_*} \mathcal{L}_\tau(\Lambda_\tau(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta})(v_* - \widehat{v}_*) \geq 0, \quad \forall v_* \in \mathcal{V}_\tau^{\kappa_0}, \quad (7.54)$$

and that

$$\frac{1}{2} D_{v_*} J(\widehat{v}_*)(v_* - \widehat{v}_*) = D_{v_*} \mathcal{L}_\Gamma(\Lambda_\chi(\widehat{v}_*), \widehat{v}_*, \widehat{u}, \widehat{\zeta})(v_* - \widehat{v}_*) \geq 0, \quad \forall v_* \in \mathcal{V}_\Gamma^{\kappa_0}. \quad (7.55)$$

Using (7.12), (7.13) and (7.25), we deduce (7.47) and (7.48). \square

Note that in Theorem 7.5, if in particular $\|\widehat{v}_*\|_{3/2, 2, \partial\Omega} < \kappa_0$, then conditions (7.47) and (7.48) become respectively

$$\left[\sigma(\widehat{v} - \widehat{u}, \widehat{p} - \widehat{q})n + \frac{1}{4}|V + \widehat{v}_* + \widehat{v}_*^c|^2 n + \frac{1}{2}((\widehat{v}_* + \widehat{v}_*^c) \cdot n)(V + \widehat{v}_* + \widehat{v}_*^c) + ((\omega \times x) \cdot \widehat{u})n \right] \perp \mathcal{V}_\Gamma \quad (7.56)$$

and

$$\sigma(\widehat{v} - \widehat{u}, \widehat{p} - \widehat{q})n \perp \mathcal{V}_\tau \quad (7.57)$$

in $L^2(\partial\Omega)$.

References

- [1] M.E. Bogovskii, Solution of the first boundary value problem for the equation of continuity of an incompressible medium, *Sov. Math. Dokl.* **20** (1979), 1094–1098.
- [2] F. Crispo and P. Maremonti, An interpolation inequality in exterior domains. *Rend. Sem. Mat. Univ. Padova*, **112** (2004), 11–39.
- [3] L. Dedè, Optimal flow control for Navier-Stokes equations: drag minimization, *Int. J. Numer. Meth. Fl.*, **55**:4 (2007), 347–366.
- [4] R. Farwig and T. Hishida, Asymptotic profile of steady Stokes flow around a rotating obstacle, *Manuscripta Math.* **136** (2011), 315–338.
- [5] A. V. Fursikov, M. D. Gunzburger and L. S. Hou, Boundary value problems and optimal boundary control for the Navier-Stokes system: the two-dimensional case, *SIAM J. Control Optim.*, **36** (1998), No. 3, 852–894.
- [6] A. V. Fursikov, M. D. Gunzburger and L. S. Hou, Optimal boundary control for the evolutionary Navier-Stokes system: the three-dimensional case, *SIAM J. Control Optim.* **43** (2005), No. 6, 2191–2232.
- [7] G.P. Galdi, On the Steady, Translational Self-Propelled Motion of a Symmetric Body in a Navier-Stokes Fluid, *Quaderni di Matematica della II Università di Napoli*, Vol. 1 (1997), 97–169.
- [8] G.P. Galdi, On the steady self-propelled motion of a body in a viscous incompressible fluid. *Arch. Rational Mech. Anal.* **148** (1999), 53–88.
- [9] G. P. Galdi, On the Motion of a Rigid Body in a Viscous Liquid: A Mathematical Analysis with Applications, *Handbook of Mathematical Fluid Dynamics*, Vol. 1, (2002) 655–679.
- [10] G.P. Galdi, *An Introduction to the Mathematical Theory of the Navier-Stokes Equations, Steady-State Problem*, Second Edition, Springer, 2011.
- [11] G.P. Galdi and A.L. Silvestre, The steady motion of a Navier-Stokes liquid around a rigid body, *Arch. Rational Mech. Anal.* **184** (2007), 371–400.

- [12] T. Hishida, A.L. Silvestre and T. Takahashi, A boundary control problem for the steady self-propelled motion of a rigid body in a Navier-Stokes fluid, *Ann. I.H.Poincaré, Analyse Non Linéaire* **34** (2017), 1507–1541.
- [13] H. Kozono and H. Sohr, New a priori estimates for the Stokes equations in exterior domains, *Indiana U. Math. J.*, **40** (1991), No.1, 1–27.
- [14] M. Kyed, Asymptotic profile of a linearized Navier-Stokes flow past a rotating body, *Q. Appl. Math.* **71** (2013), 489–500.
- [15] J. San Martín, T. Takahashi, M. Tucsnak, A control theoretic approach to the swimming of microscopic organisms, *Q. Appl. Math.*, **65** (2007), 405–424.