

H<sub>3</sub>Rs and spinal cord

## **Histamine H<sub>3</sub> receptors expressed in ventral horns modulate spinal motor output**

Tamara Coslovich<sup>1,2\*</sup>, Alberto Della Mora<sup>3\*</sup>, Giuseppe D'Angelo<sup>1,2</sup>, Fulvia Ortolani<sup>3</sup>, and Giuliano Taccola<sup>1,2</sup>

<sup>1</sup>Neuroscience department, International School for Advanced Studies (SISSA), via Bonomea 265, Trieste, (TS) Italy; <sup>2</sup>SPINAL (Spinal Person Injury Neurorehabilitation Applied Laboratory), Istituto di Medicina Fisica e Riabilitazione (IMFR), via Gervasutta 48, Udine (UD) Italy; <sup>3</sup>Department of Experimental Clinical Medicine, University of Udine, Piazzale Kolbe 3, Udine, Italy;

\* T.C. and A.DM. contributed equally to this work

*Corresponding author:*

Giuliano Taccola

Department of Neuroscience

SISSA - Scuola Internazionale Superiore di Studi Avanzati

Via Bonomea 265

34136 Trieste ITALY

Phone: (+39) 0403787111

Email: [taccola@sissa.it](mailto:taccola@sissa.it)

**Abstract (135 words)**

Motoneuron activity is modulated by histamine receptors. While H<sub>1</sub> and H<sub>2</sub> receptors have been widely explored, H<sub>3</sub> histamine receptors (H<sub>3</sub>Rs) have not been sufficiently characterized. This paper targets the effects of the selective activation of H<sub>3</sub>Rs and their expression on the membranes of large ventral horn cells. The application of selective pharmacological agents to spinal cords isolated from neonatal rats was used to identify the presence of functional H<sub>3</sub>Rs on the membrane of physiologically identified lumbar motoneurons. Intra and extracellular recordings revealed that H<sub>3</sub>R agonist,  $\alpha$ -methylhistamine, depolarized both single motoneurons and VRs, even in the presence of tetrodotoxin (TTX), an effect prevented by H<sub>3</sub>R antagonist, thioperamide. Finally, immunohistochemistry located the expression of H<sub>3</sub>Rs on a subpopulation of large cells in lamina IX. This study identifies H<sub>3</sub>Rs as a new exploitable pharmacological target against motor disturbances.

**Keywords:** motoneurons, motor pools, motor reflexes, spontaneous activity

**Abbreviations:** CV, cresyl violet; DRG, dorsal root ganglion; H<sub>3</sub>Rs, H<sub>3</sub> histamine receptors; P, postnatal; VR, ventral root

## **Histamine H<sub>3</sub> receptors expressed in ventral horns modulate spinal motor output**

**(2005 words)**

Histamine in the spinal cord is released by fibers descending from the tuberomammillary nucleus of the posterior hypothalamus, the exclusive location for histaminergic neurons (Haas et al., 2008). Histamine-immunoreactive spinal fibers are located around the central canal (Inagaki et al., 1988) and scattered in the anterior horn of the lumbar cord, mainly condensed in lamina X (Seybold, 1985).

In the mammalian spinal cord, the rhythmic patterns generated by neural circuits are modulated by histamine (Coslovich et al., 2018). Four metabotropic histamine receptors have been reported in the CNS (H<sub>1-4</sub>; Haas et al., 2008) and are also identified in the spinal cord (Taylor et al., 1982 for H<sub>1</sub> subtype; Murakami et al., 1999 for H<sub>2</sub>; Cannon et al., 2007 for H<sub>3</sub>; Strakhova et al., 2009 for H<sub>4</sub>). So far, the effects of histamine on spinal motoneurons are thought to mainly rely on H<sub>1</sub> and H<sub>2</sub> subtypes (Constanti and Nistri, 1976; Taylor et al., 1982; Saito et al., 1984; Wu et al., 2012). Nevertheless, H<sub>3</sub> histamine receptors (H<sub>3</sub>Rs) also play a functional role in the spinal cord (Harasawa, 2000; Hough and Rice, 2011) and may thus modulate ventral spinal neuron activity, as well. We explored this hypothesis using *in vitro* neonatal rat spinal cords, which allow to examine spinal and locomotor functions at both circuit and cellular levels (Brumley et al., 2017). H<sub>3</sub>Rs were studied on functionally-identified motoneurons using the selective pharmacological agonist for H<sub>3</sub>Rs,  $\alpha$ -methylhistamine, at a concentration similar to what previously reported on *in vitro* CNS preparations (Brown et al., 1996; Takei et al., 2012; 2017). Histological tools and selective staining assessed the presence of H<sub>3</sub>Rs on large-diameter (>15  $\mu$ m) cells in the ventral horns, while electrophysiological recordings from lumbar ventral roots (VRs) explored the involvement of H<sub>3</sub>Rs in modulating the output of lumbo-sacral networks.

Experiments were performed on 39 isolated spinal cords of postnatal day (P) 0 - 4 rats, as previously reported (Dose et al., 2014; 2016). Procedures were approved by the International School for Advanced Studies (SISSA) ethics committee and are in accordance with the guidelines of the Italian Animal Welfare Act 24/3/2014 n. 26 implementing the European Union directive on animal experimentation (2010/63/EU). All efforts were made to minimize number and suffering of animals used. Cords were isolated from the midthoracic region to the *cauda equina* and placed in a small recording chamber at room temperature to be continuously superfused (5 mL/min) with a Krebs solution, composed as follows (in mM): 113 NaCl, 4.5 KCl, 1 MgCl<sub>2</sub>·7H<sub>2</sub>O, 2 CaCl<sub>2</sub>, 1 NaH<sub>2</sub>PO<sub>4</sub>, 25 NaHCO<sub>3</sub>, and 11 glucose, gassed with 95 % O<sub>2</sub>-

5 % CO<sub>2</sub>, pH 7.4. Tight-fitting suction electrodes allowed DC-coupled extracellular recordings from lumbar ventral roots (VRs; Dose et al., 2016). Intracellular recordings with sharp electrodes (electrode resistance =  $38.82 \pm 9.98$  M $\Omega$ ) were obtained from 34 motoneurons total, impaled from both left (l) and right (r) L3 - L5 segments (Dose et al., 2014). In control conditions, cells' overall average resting potential was  $-66.63 \pm 9.22$  mV, with membrane resistance of  $32.03 \pm 14.38$  M $\Omega$  and antidromic spike amplitude of  $65.50 \pm 9.32$  mV. Selective histaminergic agents were added to the superfusing medium as needed. Histamine dihydrochloride (Murakoshi et al., 1985) and R- $\alpha$ -methylhistamine dihydrobromide (Dai et al., 2006) were purchased from Tocris (Bristol, UK). Thioperamide maleate (Dai et al., 2006), and tetrodotoxin (TTX) were bought from Ascent Scientific (Bristol, UK).

Histology and immunohistochemistry were performed on freshly isolated spinal cords fixed in 4% paraformaldehyde (Fluka, Buchs, Switzerland) for 3 days. Samples were soaked overnight in cryoprotecting 30% sucrose (Fluka) water solution and embedded in OCT (Kalttek, Padova, Italy). From L4 level, serial 15  $\mu$ m thick cross cryosections were cut rostrocaudally with a 2800 Frigocut N cryostat microtome (Reichert-Jung GmbH, Nussloch, Germany) and mounted on polylysine (Sigma-Aldrich, St. Louis, MO, USA) precoated slices. Serial sections were processed alternatively for staining with 0.1% cresyl violet acetate (Sigma) water solution or H<sub>3</sub>R selective immunostaining (Suppl. Fig. 1). For immunohistochemical analysis, slices were rinsed in PBS 0.1 M and then sections were incubated in 0.3% H<sub>2</sub>O<sub>2</sub> (BDH Laboratory Supplies, Poole, UK) solution in PBS, at room temperature, in order to block endogenous peroxidase activity. After washing, the sections were blocked with 3% normal goat serum (Vector Laboratories, Burlingame, CA, USA) and 0.25 % Triton X-100 (Bio-Optica, Milano, Italy) PBS solution for 2 hours and then incubated overnight with polyclonal antibody NLS476, RRID:AB\_2264153 (Cricco et al., 2008; Novus Biologicals, Littleton, CO, USA) at a dilution of 1:200, in a humidified chamber, at room temperature. After washing, sections were incubated 2 hours in biotinylated secondary antibody (Vector Laboratories) at a dilution of 1:200, and then rinsed. The slices were incubated in ABC solution (Vector Laboratories) for 1 hour at room temperature. Bound antibodies were detected using DAB solution (Vector Laboratories). After immunohistochemical labelling, slices were then counterstained with haematoxylin to visualize cell nuclei (Suppl. Fig. 2). Control experiments included omission of the primary antibody. All slices were subjected to ethanol/xylene (Carlo Erba, Milan, Italy) cleansing and then examined via photomicroscopy (Zeiss, Göttingen, Germany). Large cells in lamina IX (putatively motoneurons) were morphologically identified as cells with a diameter greater than 15  $\mu$ m. The number of H<sub>3</sub>R-positive large cells in ventral horns was calculated

## H<sub>3</sub>Rs and spinal cord

based on immunostained sections ( $n = 84$ ), whereas total number of large cells in ventral horns was calculated based on cresyl violet stained sections ( $n = 84$ ), in a  $350 \times 400 \mu\text{m}$  area, using ImageJ software.

Mean values from each spinal cord (P2-P4;  $n = 5$ ) were then obtained. Since no differences were found between right and left spinal cord sides, mean values of those data were calculated and normality was assessed by means of Shapiro-Wilk test ( $p > 0.05$ ).

As for statistical analysis, all data are indicated as a mean  $\pm$  SD, while  $n$  indicates the number of cells or spinal cords analyzed. Data were distinguished between parametric and non-parametric, using a normality test. Then, all parametric values were analysed with either Student's t-test (paired or unpaired) to compare two groups of data, or with ANOVA when groups were more than two. Non-parametric data was analysed with a Mann-Whitney test for two groups or, for multiple comparisons, with one-way ANOVA on ranks (Kruskal-Wallis) first, followed by a post hoc test (Dunnett's Method, Tukey Test). Results were considered significant when  $p < 0.05$ .

To verify the presence of functional H<sub>3</sub>Rs in the spinal cord, we added the endogenous agonist, histamine, to the superfusing medium. An immediate depolarization (about 15 mV) was observed, associated with the superimposed firing activity (Figure 1A, left) previously reported (Coslovich et al., 2018). After an extensive washout from histamine (at least 15 min), the application of  $\alpha$ -methylhistamine (20  $\mu\text{M}$ ) induced an appreciable depolarization (about 4 mV), accompanied by action potentials (Figure 1A, right). Serial applications of histamine and  $\alpha$ -methylhistamine revealed that depolarizations induced by the first were significantly higher than the ones elicited by  $\alpha$ -methylhistamine on the same cells (paired t-test,  $p = 0.03$ ,  $n = 3$ ).

A depolarization of  $4.40 \pm 1.34$  mV was recorded in 10 out of 34 recorded motoneurons, after  $154.03 \pm 91.31$  s from  $\alpha$ -methylhistamine application. As data were collected from animals very close in age (P1-P3), the observed sensitivity to the agent in the 29% of recorded motoneurons did not appear to depend upon developmental stage, but rather upon different cell properties. In control conditions, the two groups of cells (sensitive and non-sensitive) showed similar membrane resistance ( $33.34 \pm 16.91$  m $\Omega$ , sensitive vs.  $30.70 \pm 13.85$  m $\Omega$  non-sensitive), but different resting membrane potentials ( $-59.73 \pm 7.53$ , sensitive vs.  $-68.68 \pm 9.38$ , non-sensitive; t-test;  $p = 0.010$ ).

To verify the presence of functional H<sub>3</sub>Rs on the motoneuron membrane, we blocked the action potential-mediated transmission with TTX (0.5-1  $\mu\text{M}$ ) before and during  $\alpha$ -methylhistamine application (20  $\mu\text{M}$ ). In control conditions, TTX suppressed spontaneous tonic activity, abolished action potentials, and hyperpolarized cells. Addition of  $\alpha$ -methylhistamine (20  $\mu\text{M}$ ;

## H<sub>3</sub>Rs and spinal cord

Figure 1B) significantly depolarized 60% of motoneurons (mean depolarization  $1.23 \pm 0.79$  mV, paired t-test,  $p = 0.012$ ,  $n = 6$ ). In the presence of TTX, the extent of depolarization induced by  $\alpha$ -methylhistamine was significantly smaller than the one induced by histamine (Figure 1C, paired t-test,  $p = 0.019$ ,  $n = 5, 6$ ).

Since only a subgroup of cells showed to possess H<sub>3</sub>Rs, we examined their selective contribution to the overall output of motoneuronal pools. Thus, VR activity was extracellularly recorded during bath-application of H<sub>3</sub>R agonist,  $\alpha$ -methylhistamine. In a sample cord in Fig. 1D, VRs were depolarized by 367  $\mu$ V, in accordance with the mean depolarization of  $415 \pm 64$   $\mu$ V obtained from 11 experiments, where the selective activation of H<sub>3</sub>Rs did not elicit any alternating activity from VRs.

In the presence of TTX, histamine ( $n = 8$ ) and  $\alpha$ -methylhistamine ( $n = 6$ ) were always able to depolarize VRs (Figure 1E). This provides evidence that H<sub>3</sub>Rs, although functionally expressed only by a third of cells, once activated, play an important role in modulating overall spinal motor output. In additional five preparations, co-application of histamine, TTX and the selective antagonist for the H<sub>3</sub>R subtype, thioperamide, statistically reduced the depolarization induced by histamine (Figure 1E; one-way ANOVA followed by all pair-wise multiple comparison procedures with Tukey test;  $p = 0.005$ ). The observation showed that H<sub>3</sub>Rs mediate part of the effects of histamine application.

In summary, data indicate that, although H<sub>3</sub>Rs were functionally involved in modulating the 29% of intracellularly-recorded motoneurons, their role is predominant as the overall VR motor output was indeed depolarized by H<sub>3</sub>R selective agents.

Because electrophysiological experiments with TTX strongly indicated the presence of H<sub>3</sub>Rs on motoneuron membranes, we performed immunohistochemical labelling to visualize the distribution of H<sub>3</sub>Rs in the ventral spinal cord. Interestingly, H<sub>3</sub>Rs were expressed on the membrane of 35% of the large diameter ( $>15$   $\mu$ m) cells in lamina IX, as morphologically confirmed using cresyl violet staining (Figure 2,  $n = 84$  cryosections from 5 spinal cords).

Indeed,  $\alpha$ -methylhistamine in TTX affected the activity of only a portion of functionally-identified motoneurons, in line with the expression of H<sub>3</sub>Rs on the membranes of one third of large cells in the ventral horn, immunohistochemically labelled.

The identification of H<sub>3</sub>Rs on motoneurons explains part of the motor behavior following pharmacological manipulation with selective histamine agents (Chiavegatto et al., 1998; Toyota et al., 2002) and opens a new path in exploring therapeutics to alleviate spinal cord pathologies. Histamine is well known to be involved in nociceptive and antinociceptive processing (Wei et al., 2016) and itch sensations (Lee et al., 2016; Luo et al., 2015). In

particular, H<sub>3</sub>Rs, acting on sensory afferents and sensory-responsive neurons in the spinal cord (Cannon et al., 2007), attenuate mechanically-induced nociception (Cannon et al., 2003). Moreover, H<sub>3</sub>R antagonism has been suggested as a potential mechanism for the treatment of neuropathic pain in preclinical rat models (Coward et al., 2012; Hsieh et al., 2010). Our findings suggest that H<sub>3</sub>R agents also act on motoneurons in the ventral spinal cord to influence motor behavior and can represent promising pharmacological targets in the CNS, because of their characteristic of being constitutively active *in vivo* (Passani and Blandina, 2011). A direct effect of H<sub>3</sub>Rs on interneuronal spinal networks for locomotion should be excluded, since no locomotor-like alternating patterns (Taccola and Nistri, 2005) were elicited by  $\alpha$ -methylhistamine alone. The inverse agonism of these receptors unveils their ability to modulate neuronal excitability, even in the absence of any endogenous histamine, as shown by innovative molecules that underwent preclinical testing for the treatment of cognitive disorders and feeding behavior (Arrang et al., 2007).

Recently, histamine has been reported to improve long-term recovery in a rat model of spinal injury, by decreasing the lesion area, inhibiting astrogliosis and glial scar formation, and improving locomotor recovery (Zhao et al., 2015). Although many of the effects were reversed by a treatment with a H<sub>1</sub> receptor antagonist, the authors suggested that the overall protective role of histamine may be due to different histamine receptors in different cell types. Then, the modulation of H<sub>3</sub>Rs expressed in a subpopulation of large ventral horn cells could represent a potential neuroprotective strategy to rescue spinal tissue after damage (Kukko-Lukjanov et al., 2006).


### **Compliance with Ethical Standards**

**Ethical Approval:** Procedures were approved by the International School for Advanced Studies (SISSA) ethics committee and are in accordance with the guidelines of the Italian Animal Welfare Act 24/3/2014 n. 26 implementing the European Union directive on animal experimentation (2010/63/EU). All procedures performed in studies involving animals were in accordance with the ethical standards of the SISSA Laboratory Animal Center and the protocol was approved by the local Animal Committee (Organismo preposto al benessere degli animali, OPBA) of SISSA, Italy.

**Conflict of interest:** no conflicts of interest to declare

**Acknowledgments:** We thank Ms. Elvita Titolo and Mr Walter Zangrando for their generous support. We are grateful to Elisa Ius for the excellent assistance in preparing the manuscript.

**Authors' contributions:** All authors had full access to the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Conceptualization, G.T.; Methodology, G.T.; Investigation, T.C., G.D'A., A.DM. and G.T.; Formal Analysis, T.C., G.D'A., A.DM. and G.T.; Writing - Original Draft, T.C., G.D'A., A.DM. and G.T.; Writing - Review & Editing, A.DM. and G.T.; Visualization, G.D'A., A.DdM. and G.T.; Supervision, F.O. and G.T.; Funding Acquisition, G.T.









**Figure 1.  $\alpha$ -methylhistamine directly depolarizes a subset of recorded motoneurons and affects the motor output recorded from VRs.**

A.) After 84 s from the application of histamine (upper gray bar; 20  $\mu$ M), a sustained depolarization (15.32 mV) with superimposed intense firing (3.97 Hz) is recorded from a rL5 single motoneuron (resting potential = -65 mV). After 15 min of washout, the baseline returns to control level and only sporadic action potentials can be derived (trace break = 2 min). Application of  $\alpha$ -methylhistamine ( $\alpha$ -MH) slightly depolarizes the same cell (4.15 mV) with the occurrence of faster spiking activity. Based on the extent of depolarization induced by 20  $\mu$ M  $\alpha$ -methylhistamine ( $\alpha$ -MH), single motoneurons are *a posteriori* divided in sensitive and non-sensitive. B.) A different rL5 motoneuron (- 63 mV initial resting potential,  $V_m$ ) is hyperpolarized (- 4.94 mV) by TTX (1 and then 0.5  $\mu$ M) with the suppression of both spontaneous firing and tonic activity, as well.  $\alpha$ -methylhistamine ( $\alpha$ -MH, 20  $\mu$ M) slightly depolarizes the motoneuron (2.56 mV) as indicated at a higher magnification in the insert

## H<sub>3</sub>Rs and spinal cord


above. C.) Analysis of pooled data from different experiments during TTX perfusion indicates that the mean depolarization elicited by histamine (20  $\mu$ M; dark gray bar) is significantly higher than the one induced by  $\alpha$ -methylhistamine (20  $\mu$ M; light gray bar; \*, t test;  $p = 0.019$ ;  $n = 5, 6$ ). D.) Application of  $\alpha$ -methylhistamine (20  $\mu$ M, 15 min application) induces comparable stable depolarizations with sporadic synchronous events among all VRs (VRrL2 = 349  $\mu$ V; VRrL5 = 452  $\mu$ V; VRIL5 = 321  $\mu$ V; VRIL2 = 347  $\mu$ V). After 5 min of the following washout, traces return to baseline and discharges are progressively attenuated, until mostly suppressed. E.) Histamine (20  $\mu$ M) and  $\alpha$ -methylhistamine ( $\alpha$ -MH; 20  $\mu$ M) depolarize VRs in the presence of TTX, while the selective H<sub>3</sub>R antagonist thioperamide (20  $\mu$ M) abolishes the depolarization induced by histamine (\*, one-way ANOVA followed by all pair-wise multiple comparison procedures with Tukey test;  $p = 0.005$ ;  $n = 8, 6, 5$ ). Note that A and B traces come from different motoneurons.




**Figure 2. Immunostaining confirms that a subgroup of large cells in lamina IX contains H<sub>3</sub>Rs.** A.) Cresyl violet staining of 15 µm thick cross cryosection of L4 spinal cord. Dotted rectangle delimiting lamina IX, in which large diameter (>15 µm) ventral horn cells are morphologically identified, is shown in C.) and E.) at a higher magnification. Scale bars: 250 µm A.), 200 µm C.) and 100 µm E.), respectively.

D.) Immunohistochemical detection by H<sub>3</sub>R marker on a serial slice from the same cord reveals diffuse labelling concentrated around a set of large ventral horn cells. Dotted rectangle delimiting lamina IX, is shown in F.) at a higher magnification. Scale bars: 200 µm D.) and 100 µm. F.)

B.) Quantification of mean number of H<sub>3</sub>R-positive cells among the total number of large ventral horn cells in lamina IX (motor neurons). Large cells in lamina IX were morphologically identified as cells with a diameter greater than 15 µm. Number of H<sub>3</sub>R-positive cells is significantly different from the total number of cells, morphologically identified in cresyl violet stained (CV+) sections (\*; Student's t-test,  $p < 0.001$ ,  $n = 84$  sections from five different cords).



**Supplementary Figure 1.** The cartoon schematizes the histological processing. Spinal cords, fixed in PFA, were cryosected rostrocaudally from L4 level. Serial 15 µm thick sections were processed alternatively for staining with cresyl violet or H<sub>3</sub>R selective immunostaining. After immunohistochemical labelling, slices were then counterstained with haematoxylin to visualize cell nuclei.



**Supplementary Figure 2.** Double staining by H<sub>3</sub>R antibody and haematoxylin shows the expression of H<sub>3</sub>R on four large lamina IX cells (black arrows). Scale bar: 50  $\mu$ m.

## References (36)

Arrang JM, Morisset S, Gbahou F (2007). Constitutive activity of the histamine H<sub>3</sub> receptor. *Trends Pharmacol Sci* 28, 350-357.

Brown RE, Reymann KG (1996). Histamine H<sub>3</sub> receptor-mediated depression of synaptic transmission in the dentate gyrus of the rat in vitro. *J Physiol* 496, 175-184.

Brumley MR, Guertin PA, Taccola G. (2017). Multilevel analysis of locomotion in immature preparations suggests innovative strategies to reactivate stepping after spinal cord injury. *Curr Pharm Des. In press*

Cannon KE, Nalwalk JW, Stadel R, Ge P, Lawson D, Silos-Santiago I, Hough LB (2003). Activation of spinal histamine H<sub>3</sub> receptors inhibits mechanical nociception. *Eur J Pharmacol* 470, 139-147.

Cannon KE, Chazot PL, Hann V, Shenton F, Hough LB, Rice FL (2007). Immunohistochemical localization of histamine H<sub>3</sub> receptors in rodent skin, dorsal root ganglia, superior cervical ganglia, and spinal cord: potential antinociceptive targets. *Pain* 129, 76-92.

Chiavegatto S, Nasello AG, Bernardi MM (1998). Histamine and spontaneous motor activity: biphasic changes, receptors involved and participation of the striatal dopamine system. *Life Sci.* 62, 1875-1888.

Constanti A, Nistri A (1976). Antagonism by some antihistamines of the amino acid-evoked responses recorded from the lobster muscle fibre and the frog spinal cord. *Br J Pharmacol* 58, 583-592.

Coslovich T, Brumley MR, D'Angelo G, Della Mora A, Swann HE, Ortolani F, Taccola G (2018). Histamine modulates spinal motoneurons and locomotor circuits. *J Neurosci Res* 96, 889-900.

Cowart M, Hsieh G, Black LA, Zhan C, Gomez EJ, Pai M, Strakhova M, Manelli A, Carr T, Wetter J, Lee A, Diaz G, Garrison T, Brioni JD (2012). Pharmacological characterization of A-960656, a histamine H<sub>3</sub> receptor antagonist with efficacy in animal models of osteoarthritis and neuropathic pain. *Eur J Pharmacol* 684, 87-94.

## H<sub>3</sub>Rs and spinal cord

Cricco GP, Mohamad NA, Sambuco LA, Genre F, Croci M, Gutiérrez AS, Medina VA, Bergoc RM, Rivera ES, Martín GA (2008). Histamine regulates pancreatic carcinoma cell growth through H<sub>3</sub> and H<sub>4</sub> receptors. *Inflamm Res* 57:S23-4. 5

Dai H, Zhang Z, Zhu Y, Shen Y, Hu W, Huang Y, Luo J, Timmerman H, Leurs R, Chen Z (2006). Histamine protects against NMDA-induced necrosis in cultured cortical neurons through H receptor/cyclic AMP/protein kinase A and H receptor/GABA release pathways. *J Neurochem* 96, 1390-1400.

Dose F, Zanon P, Coslovich T, Taccola G (2014). Nanomolar oxytocin synergizes with weak electrical afferent stimulation to activate the locomotor CpG of the rat spinal cord in vitro. *PLoS One* 9(3):e92967.

Dose F, Deumens R, Forget P, Taccola G (2016). Staggered multi-site low-frequency electrostimulation effectively induces locomotor patterns in the isolated rat spinal cord. *Spinal Cord* 54, 93-101.

Haas HL, Sergeeva OA, Selbach O (2008). Histamine in the nervous system. *Physiol Rev* 88, 1183-11241.

Harasawa K (2000). Does the histaminergic system play a role in spinal nociception? *Hokkaido Igaku Zasshi* 75, 275-281.

Hough LB, Rice FL (2011). H<sub>3</sub> receptors and pain modulation: peripheral, spinal, and brain interactions. *J Pharmacol Exp Ther* 336, 30-37.

Hsieh GC, Honore P, Pai M, Wensink EJ, Chandran P, Salyers AK, Wetter JM, Zhao C, Liu H, Decker MW, Esbenshade TA, Cowart MD, Brioni JD (2010). Antinociceptive effects of histamine H<sub>3</sub> receptor antagonist in the preclinical models of pain in rats and the involvement of central noradrenergic systems. *Brain Res* 1354, 74-84.

Inagaki N, Yamatodani A, Ando-Yamamoto M, Tohyama M, Watanabe T, Wada H (1988). Organization of histaminergic fibers in the rat brain. *J Comp Neurol* 273, 283-300.

Kukko-Lukjanov TK, Soini S, Taira T, Michelsen KA, Panula P, Holopainen IE (2006). Histaminergic neurons protect the developing hippocampus from kainic acid-induced neuronal damage in an organotypic coculture system. *J Neurosci* 26, 1088-1097.

## H<sub>3</sub>Rs and spinal cord

Lee JS, Han JS, Lee K, Bang J, Lee H (2016). The peripheral and central mechanisms underlying itch. *BMB Rep* 49, 474-487.

Luo J, Feng J, Liu S, Walters ET, Hu H (2015). Molecular and cellular mechanisms that initiate pain and itch. *Cell Mol Life Sci* 72, 3587-3588.

Murakami H, Sun-Wada GH, Matsumoto M, Wada Y, Futai M (1999). Human histamine H<sub>2</sub> receptor gene: multiple transcription initiation and tissue-specific expression. *FEBS Lett* 451, 327-331.

Murakoshi T, Suzue T, Tamai S (1985). A pharmacological study on the respiratory rhythm in the isolated brainstem-spinal cord preparation of the newborn rat. *Br J Pharmacol* 86, 95-104.

Passani MB, Blandina P (2011). Histamine receptors in the CNS as targets for therapeutic intervention. *Trend Pharmacol Sci* 32, 242-249.

Saito T, Tagami H, Kamei C, Tasaka K (1984). Excitatory and inhibitory effects of H<sub>1</sub>-blockers on isolated bullfrog spinal cords. *Arch Int Pharmacodyn Ther* 271, 127-134.

Seybold VS (1985). Distribution of histaminergic, muscarinic and serotonergic binding sites in cat spinal cord with emphasis on the region surrounding the central canal. *Brain Res* 342, 291-296.

Strakhova MI, Nikkel AL, Manelli AM, Hsieh GC, Esbenshade TA, Brioni, Bitner RS (2009). Localization of histamine H<sub>4</sub> receptors in the central nervous system of human and rat. *Brain Res* 1250, 41-48.

Taccola G, Nistri A (2005). Characteristics of the electrical oscillations evoked by 4-aminopyridine on dorsal root fibers and their relation to fictive locomotor patterns in the rat spinal cord in vitro. *Neuroscience* 132, 1187-1197.

Takei H, Song L, Ebihara K, Shirakawa T, Koshikawa N, Kobayashi M (2012). Histaminergic effects on the frequency of repetitive spike firing in rat insular cortex. *Neurosci Lett* 518, 55-59.

Takei H, Yamamoto K, Bae YC, Shirakawa T, Kobayashi M (2017). Histamine H<sub>3</sub> heteroreceptors suppress glutamatergic and gabaergic synaptic transmission in the rat insular cortex. *Front Neural Circuits* 11:85.

### H<sub>3</sub>Rs and spinal cord

Taylor JE, Yaksh TL, Richelson E (1982). Histamine H1 receptors in the brain and spinal cord of the cat. *Brain Res* 243, 391-394.

Toyota H, Dugovic C, Koehl M, Laposky AD, Weber C, Ngo K, Wu Y, Lee DH, Yanai K, Sakurai E, Watanabe T, Liu C, Chen J, Barbier AJ, Turek FW, Fung-Leung WP, Lovenberg TW (2002). Behavioral characterization of mice lacking histamine H(3) receptors. *Mol Pharmacol* 62, 389-397.

van Neerven SG, Pannaye P, Bozkurt A, Van Nieuwenhoven F, Joosten E, Hermans E, Taccola G, Deumens R. (2013). Schwann cell migration and neurite outgrowth are influenced by media conditioned by epineurial fibroblasts. *Neuroscience* 252, 144-153.

Wei H, Vilasanen H, You HJ, Pertovaara A (2016). Spinal histamine in attenuation of mechanical hypersensitivity in the spinal nerve ligation-induced model of experimental neuropathy. *Eur J Pharmacol* 772, 162-171.

Wu GY, Han XH, Zhuang J, Yung WH, Chan YS, Zhu JN, Wang JJ (2012). Excitatory effect of histamine on rat spinal motoneurons by activation of both H1 and H2 receptors in vitro. *J Neurosci Res* 90, 132-142.

Zhao Y-Y, Yuan Y, Chen, Y, Jiang L, Liao R-J, Wang L, Zhang X-N, Ohtsu H, Hu W-W, Chen Z (2015). Histamine promotes locomotion recovery after spinal cord hemisection via inhibiting astrocytic scar formation. *CNS Neurosci Ther* 21, 454-462.