CERN-PH-EP/2012-103
2013/01/25

CMS-EXO-11-024

Search for leptonic decays of W' bosons in pp collisions at $\sqrt{s} = 7$ TeV

The CMS Collaboration*

Abstract

A search for a new heavy gauge boson W' decaying to an electron or muon, plus a low mass neutrino, is presented. This study uses data corresponding to an integrated luminosity of 5.0 fb^{-1} , collected using the CMS detector in pp collisions at a centre-of-mass energy of 7 TeV at the LHC. Events containing a single electron or muon and missing transverse momentum are analyzed. No significant excess of events above the standard model expectation is found in the transverse mass distribution of the lepton-neutrino system, and upper limits for cross sections above different transverse mass thresholds are presented. Mass exclusion limits at 95% CL for a range of W' models are determined, including a limit of 2.5 TeV for right-handed W' bosons with standard-model-like couplings and limits of 2.43–2.63 TeV for left-handed W' bosons, taking into account their interference with the standard model W boson. Exclusion limits have also been set on Kaluza–Klein W_{KK} states in the framework of split universal extra dimensions.

Submitted to the Journal of High Energy Physics

*See Appendix A for the list of collaboration members

1 Introduction

This Letter describes a search for a new heavy gauge boson W' , using proton-proton collision data collected during 2011 using the Compact Muon Solenoid (CMS) detector [1] at the Large Hadron Collider (LHC) at a centre-of-mass energy of 7 TeV. The dataset corresponds to an integrated luminosity of $5.0 \pm 0.1 \text{ fb}^{-1}$ [2]. The search attempts to identify an excess of events with a charged lepton (an electron or muon) and a neutrino in the final state, and an interpretation of the results is provided in the context of several theoretical models.

2 Physics models

New heavy gauge bosons such as the W' and Z' are predicted by various extensions of the standard model (SM). In the sequential standard model (SSM) [3], the W' boson is considered to be a left-handed heavy analogue of the W . It is assumed to be a narrow s -channel resonance with decay modes and branching fractions similar to those of the W , with the addition of the $t\bar{b}$ channel that becomes relevant for W' masses above 180 GeV. Interference between the W' and W is assumed to be negligible. If the W' is heavy enough to decay to top and bottom quarks, the predicted branching fraction is about 8.5% for each of the two leptonic channels studied in the present analysis. Under these assumptions, the width of a 1 TeV W' is about 33 GeV. Decays of the W' into WZ dibosons are usually suppressed in this model.

The assumptions of the SSM were used in previous searches in leptonic channels at the Tevatron [4, 5] and the LHC [6–9]. The signature of a charged high-momentum lepton and a neutrino would also be observed in the decays of a right-handed W'_R , predicted by left-right symmetric models [10–13]. This particle is typically predicted to decay to a heavy right-handed neutrino [14–16].

However, the mass of the right-handed neutrino is not constrained, and it could be light as long as it does not couple to SM weak bosons. This would result in the same W'_R decay signature as for the W .

If the W' is right-handed it will not interfere with the W . However, if it is left-handed (W'_L), interference with the W is expected [17–19]. Constructive (destructive) interference occurs in the mass range between W and W' if the coupling of the W' boson to quarks and leptons has opposite sign to (same sign as) the coupling of the W boson to left-handed fermions (g_L). While constructive interference increases the W' production cross section, and therefore allows experimental sensitivity at higher masses, destructive interference would yield a lower cross section, rendering previously published LHC mass exclusion limits [7, 9] slightly optimistic. Interference has previously been considered in searches for the decay to top and bottom quarks [19, 20], but never for leptonic decays.

Figure 1 shows the transverse mass distribution for a W' of 2.5 TeV mass for the cases of constructive, destructive and non-interference, along with the background due to the SM W . In the absence of interference the cross sections and transverse mass spectrum of left- and right-handed W' are identical. The W' manifests itself as a Jacobian peak with its width almost independent of the presence and type of interference. However, the intermediate region around $M_T \sim 1 \text{ TeV}$ shows a clear variation of the shape. Destructive interference of a W'_L boson with mass $\geq 2 \text{ TeV}$ modulates the W transverse mass tail, resulting in a faster fall-off. The modulation strength and the resulting effect on the cross section both increase with the W' mass and width. Given sufficient detector resolution, the constructive and destructive interference scenarios may be distinguishable.

The leptonic final states under study may also be interpreted in the framework of universal extra dimensions (UED) with bulk mass fermions, or split-UED [21, 22]. This is a model based on an extended space-time with an additional compact fifth dimension of radius R . All SM fermions and gauge bosons have Kaluza–Klein (KK) states, for instance W_{KK}^n , where n denotes the n -th KK excitation mode, and

$$m_{W_{\text{KK}}^n}^2 \equiv m_n^2 = m_W^2 + \left(\frac{n}{R}\right)^2, \quad (1)$$

$$g_n = g^{\text{SM}} \mathcal{F}_n(\pi\mu R), \quad (2)$$

$$\mathcal{F}_n(x) = \begin{cases} 0 & \text{if } n = 2m + 1 \\ \frac{x^2[-1+(-1)^m e^{2x}](\coth x - 1)}{\sqrt{2(1+\delta_{m0})(x^2+m^2\pi^2/4)}} & \text{if } n = 2m. \end{cases} \quad (3)$$

Here μ is the bulk mass parameter in five dimensions of the fermion field, with $[1/R, \mu]$ defining the UED parameter space. The coupling of the W_{KK}^n to SM fermions is denoted g_n and defined as a modification of the SM coupling g^{SM} of the W . The function $\mathcal{F}_{2m}(x)$ tends to approach $(-1)^m \sqrt{2}$ as $x \rightarrow \infty$. In minimal UED models, the parameter μ is assumed to be zero [23]. Following [21, 22], we assume a non-zero value for μ , thus increasing the cross sections sufficiently to allow observation by LHC experiments.

KK-odd modes of W_{KK}^n do not couple to SM fermions, owing to KK-parity conservation. Moreover, there is no expected sensitivity for $n \geq 4$ modes at the LHC centre-of-mass energy and luminosity used in this analysis. W_{KK}^2 is therefore the only mode considered. Under this assumption, the decay to leptons is kinematically identical to the sequential SM-like W' decay, and the observed limits obtained from the $W' \rightarrow e\nu$ and $W' \rightarrow \mu\nu$ searches can directly be reinterpreted in terms of the W_{KK}^n mass considering the different widths. The width of a W_{KK}^n is \mathcal{F}_n^2 times the SSM-like W' width:

$$\Gamma_{W_{\text{KK}}^n} = \mathcal{F}_n^2 \frac{4}{3} \frac{m_{W_{\text{KK}}^n}}{m_W} \Gamma_W. \quad (4)$$

3 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid, of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the field volume are the silicon pixel and strip tracker, the crystal electromagnetic calorimeter and the brass/scintillator hadron calorimeter. The electromagnetic calorimeter consists of nearly 76 000 lead tungstate crystals. The energy resolution for electrons with the very high transverse momentum used in this analysis, which are predominantly in the central pseudorapidity region, is about 1%. In the forward region the resolution is about 2%. Muons are measured in gas-ionization detectors embedded in the steel return yoke. Central and forward regions are instrumented with four muon stations combining high precision tracking detectors (drift tubes in the central region and forward cathode strip chambers) with resistive plate chambers, which contribute to the trigger as well as the track measurement. The muon transverse momentum, p_{T}^μ , is determined from the curvature of its track, measured as it traverses the magnetized return yoke. Each muon track is matched to a track measured in the silicon tracker, resulting in a muon p_{T} resolution of 1 to 10% for p_{T} of




Figure 1: MADGRAPH and COMPHEP predictions of the transverse mass distribution for the SM W background and various W' models for $m(W')=2.5$ TeV. In the absence of interference, W'_R and W'_L cross sections are identical. A W'_L could experience constructive or destructive interference with the SM W , yielding the shown modulation of the M_T spectrum.

up to 1 TeV. CMS uses a two-level trigger system comprising custom hardware processors and a High-Level Trigger processor farm. Together, these systems select around 300 Hz of the most interesting recorded bunch-crossings for permanent storage. A detailed description of CMS can be found in Ref. [1].

A cylindrical coordinate system about the beam axis is used, in which the polar angle θ is measured with respect to the counterclockwise beam direction and the azimuthal angle ϕ is measured in the xy plane, where the x axis points towards the center of the LHC ring. The quantity η is the pseudo-rapidity, defined as $\eta = -\ln[\tan \theta/2]$.

4 Event selection

Candidate events with at least one high-transverse-momentum (p_T) lepton were selected using single-muon and single-electron triggers. The trigger thresholds were raised as the LHC luminosity increased during the data-taking period, the highest values being $p_T > 80$ GeV for electrons and $p_T > 40$ GeV for muons. Offline, electrons and muons were required to have p_T at least 5 GeV higher than the online threshold, which does not impair the search in the high mass region.

Muons were reconstructed by combining tracks from the inner tracker and the outer muon system. Well-reconstructed muons were selected by requiring at least one pixel hit, hits in eight tracker layers and segments in two muon stations. Since the segments have multiple hits and are typically found in different muon detectors separated by thick layers of iron, the latter requirement significantly reduces the amount of hadronic punch-through. The transverse impact parameter $|d_0|$ of a muon track with respect to the beam spot is required to be less

than 0.02 cm, in order to reduce the cosmic ray muon background. Furthermore, the muon is required to be isolated within a $\Delta R \equiv \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} < 0.3$ cone around its direction. Muon isolation requires that the scalar sum of the transverse momenta of all tracks originating at the interaction vertex, excluding the muon, is less than 15% of its p_T . An additional requirement is that there be no second muon in the event with $p_T > 25$ GeV to reduce the Z, Drell-Yan and cosmic ray muon backgrounds.

Electrons were reconstructed as isolated objects in the electromagnetic calorimeter, with additional requirements on the shower shape and the ratio of hadronic to electromagnetic deposited energies. The electrons were required to have at least one inner hit, a transverse energy greater than 85 GeV, and required to be isolated in a cone of radius $\Delta R < 0.3$ around the electron candidate direction, both in the tracker and in the calorimeter. In the tracker, the sum of the p_T of the tracks, excluding tracks within an inner cone of 0.04, was required to be less than 5 GeV. For the isolation using calorimeters, the total transverse energy in the barrel, excluding deposits associated to the electron, was required to be less than $0.03 \cdot p_T^{\text{ele}} + 2.0$ GeV. The isolation requirements were modified as luminosity increased, owing to the increase in the typical number of additional pp interactions ('pile-up') per LHC bunch crossing. These selections are designed to ensure high efficiency for electrons and a high rejection of misreconstructed electrons from multi-jet backgrounds.

The main observable in this search is the transverse mass M_T of the lepton- E_T^{miss} system, calculated as

$$M_T \equiv \sqrt{2 \cdot p_T^\ell \cdot E_T^{\text{miss}} \cdot (1 - \cos \Delta\phi_{\ell,\nu})} \quad (5)$$

where $\Delta\phi_{\ell,\nu}$ is the azimuthal opening angle between the charged lepton's transverse momentum (p_T^ℓ) and missing transverse energy (E_T^{miss}) direction. The neutrino is not detected directly, but gives rise to experimentally observed E_T^{miss} . This quantity was determined using a particle-flow technique [24], an algorithm designed to reconstruct a complete list of distinct particles using all the subcomponents of the CMS detector. Muons, electrons, photons, and charged and neutral hadrons were all reconstructed individually. The E_T^{miss} for each event was then calculated as the vector opposing the total transverse momentum of all reconstructed particles in each event.

In W' decays, the lepton and E_T^{miss} are expected to be almost back-to-back in the transverse plane, and balanced in transverse energy. Candidate events were therefore selected through a requirement on the ratio of the lepton p_T and the E_T^{miss} , $0.4 < p_T/E_T^{\text{miss}} < 1.5$. A requirement was also imposed on the angular difference in the transverse plane of the lepton and E_T^{miss} direction, $\Delta\phi_{\ell,\nu} > 0.8 \times \pi$. No selection is made on jets. After these selections, the average W' signal efficiency for masses up to 2.5 TeV in simulated events was found to be around 80% in both channels, including the roughly 90% geometrical acceptance corresponding to a requirement of $|\eta_\mu| < 2.1$ for muons, and with $|\eta_e| < 1.442$ or $1.56 < |\eta_e| < 2.5$ for electrons. The transverse mass distributions after these selections are shown in Figure 2.

5 Signal and background simulation

Several large samples of simulated events were used to evaluate signal and background efficiencies. The generated events were processed through a full simulation of the CMS detector based on GEANT4 [25, 26], a trigger emulation, and the event reconstruction chain.

The event samples for the W'_R signal were produced separately from the SM W sample, using the PYTHIA 6.4.9 generator [27]. This is consistent with the case of non-interference assumed




Figure 2: Observed transverse mass distributions for the electron (left) and muon (right) channels. Simulated signal distributions for a (left- or right-handed) W' without interference of 2.3 TeV mass are also shown, including detector resolution effects. The simulated background labelled as ‘diboson’ includes WW , ZZ and WZ contributions. The top background prediction includes single top and top pair production. The total background prediction from a fit to the simulated transverse mass spectrum in each channel is shown by the dashed line.

for the previous ATLAS and CMS studies. In order to include interference of W'_L and W in this analysis, a model of a single new heavy vector boson W' with a SM-like left-handed coupling strength $|g'_L| \approx 0.65$ was implemented in the MADGRAPH event generator [28]. This model includes spin correlations as well as finite-width effects. For such a left-handed scenario with interference, the generation of samples is technically more challenging. Since the scattering amplitude responsible for the lv final state is the sum of W'_L and SM W boson terms, both contributions have to be generated simultaneously. A threshold in M_T was applied to suppress the dominant W contribution around the W -mass, where interference effects are negligible for the W'_L masses considered in this search. The simulation uses MADGRAPH 4.5.1, matched to PYTHIA for showering and hadronisation. For the hadronisation model, the PYTHIA Tune Z2 was used for both the W'_R and W'_L simulations. Both generators simulate at leading order (LO) and use the CTEQ6L1 parton distribution functions (PDF) [29]. Mass-dependent K-factors, varying from 1.14 to 1.36, for the next-to-next-to-leading order (NNLO) correction were calculated with FEWZ [30, 31]. The resulting NNLO $W' \rightarrow lv$ production cross section times branching fraction ranged from 17.7 pb (for $m_{W'} = 0.5$ TeV) to 0.71 fb (for $m_{W'} = 3$ TeV) for a W' without interference (see Table 1 for cross sections). Efficiencies and detector acceptance are then taken into account for estimating the expected number of signal events. The acceptance is nearly maximal since the decay products of such heavy particles tend to populate low pseudo-rapidities. Efficiencies are high because the selections have been optimised. Detailed numbers for both quantities are given in Section 4. The Tevatron $W'_L \rightarrow t\bar{b}$ search used the COMPHEP generator [32, 33] which has the case of destructive interference implemented. The agreement between the model implementations in COMPHEP and MADGRAPH is demonstrated for the case of destructive interference in Figure 1.

The primary source of background is the off-peak, high transverse mass tail of the standard model $W \rightarrow lv$ decays. Other important backgrounds arise from QCD multijet, $t\bar{t}$, and Drell–

Yan events. Dibosons (WW , WZ , ZZ) decaying to electrons, muons, or taus were also considered. The event samples for the electroweak background processes $W \rightarrow \ell\nu$ and $Z \rightarrow \ell\ell$ ($\ell = e, \mu, \tau$) were produced using PYTHIA. NNLO cross sections were accounted for via a single K-factor of 1.32 for the W , and mass-dependent K-factors, ranging from 1.28 to 1.23, for the Z . The PYTHIA generator was also used for QCD multijet events. The $t\bar{t}$ events were generated with MADGRAPH in combination with PYTHIA, and the newly-calculated NNLL (next-to-leading-order including the leading logarithms of NNLO) cross section was applied [34]. All other event samples were normalised to the integrated luminosity of the recorded data, using calculated NNLO cross sections. The only exceptions were the diboson and QCD samples, for which the NLO and LO cross sections were used respectively. We note that multijet background is largely suppressed by the event selection requirements. The simulation of pile-up is included in all event samples by superimposing minimum bias interactions onto the main background processes.

In order to provide a background estimate independent of any interference effects in the W transverse mass tail, the shape of the background was determined from simulation. The full transverse mass spectrum was modelled by a function optimised to best describe the spectrum in either channel up to very high masses. This function, of the form

$$f(M_T) = \frac{a}{(M_T + b)^c} \quad (6)$$

was fitted to the simulation and then normalised to data in the region $200 \text{ GeV} < M_T < 500 \text{ GeV}$, and used to estimate the expected number of SM background events for all transverse mass bins (shown as the dashed lines in Figure 2). A cross check under the assumption of no interference was done by fitting the M_T distribution in data confirming the simulation. To determine the uncertainty introduced by this method, in addition to statistical errors on the fit parameters, two alternative functions were fitted:

$$f(M_T) = \frac{a}{(M_T^2 + b \cdot M_T + c)^d} \quad (7)$$

$$f(M_T) = \frac{a(1 + M_T)^b}{(M_T^{c+d} \cdot \ln M_T)} \quad (8)$$

The largest difference in the background prediction with respect to the original fit was taken as a systematic uncertainty. For M_T larger than 1.4 TeV, this corresponds to an additional uncertainty of 0.14 events with a background expectation of 0.98 events in the muon channel and 0.26 events with a background expectation of 1.28 events in the electron channel.

6 Systematic uncertainties

The expected number of potential signal and background events was evaluated from simulation. In addition to uncertainties due to the fit procedure for the background, systematic uncertainties due to imperfections in the description of the detector performance were included. Uncertainties due to the lepton energy or momentum resolution and scale, ranging between 0.4% and 10% [6, 7] were applied to the transverse mass spectrum. Uncertainties due to momentum scale were evaluated using detailed studies of the $Z \rightarrow \mu\mu$ shape and high p_T muons. The muon p_T resolution has been previously determined with cosmic ray muons to within 10% for high momentum tracks [35]. In order to estimate the uncertainty on the number of

expected events, the muon p_T spectrum was distorted (scaled and smeared) according to the values extracted from comparisons with data. The missing transverse energy was adjusted accordingly, and finally a distorted transverse mass spectrum was obtained and observed to vary by $\sim 1\%$. The electron energy scale uncertainty was around 1% in the ECAL barrel and 3% in the endcaps. Its impact on the number of signal events above the threshold of $M_T > 600$ GeV was ascertained to be less than 1% for all W' masses. We assume an uncertainty of 10% on the hadronic component of the E_T^{miss} resolution (that is, excluding the lepton), and the x and y components of the reconstructed E_T^{miss} in the simulation were smeared accordingly. The impact on the number of signal events was found to be around 2%.

Effects caused by pile-up were modeled by adding to the generated events multiple interactions with a multiplicity distribution matched to the luminosity profile of the collision data. The resulting impact on the signal was studied by varying the mean of the distribution of pile-up interactions by 8%, yielding a variation of the signal efficiency of $\sim 2\%$. Following the recommendations of the PDF4LHC group [36], the signal event samples for W'_R generated with PYTHIA were reweighted using the LHAPDF package [37]. PDF and α_s variations of the MSTW2008 [38], CTEQ6.6 [39] and NNPDF2.0 [40] PDF sets were taken into account and the impact on the signal cross sections was estimated.

7 Results and limits

A $W' \rightarrow e\nu$ or $W' \rightarrow \mu\nu$ signal is expected to manifest itself as an excess over the SM expectation in the tail of the M_T distribution. No significant excess has been observed in the data.

For W' masses well below the centre-of-mass energy of $\sqrt{s} = 7$ TeV the signal events are expected to lie in the Jacobian peak corresponding to the W' mass. For masses above 2.3 TeV, the reduced phase space results in many events below the Jacobian peak, and the acceptance for the M_T^{min} cut drops from about 40% for intermediate masses to 14% at very high W' masses. The expected signal yields given in Table 1 for a range of W'_R masses are largely unaffected when introducing interference effects, owing to the high M_T cut corresponding to the optimum search window, which naturally lies around the Jacobian peak.

We set upper limits on the production cross section times the branching fraction $\sigma_{W'_R} \times \mathcal{B}(W'_R \rightarrow \ell\nu)$, with $\ell = e$ or μ . The observed highest transverse mass events had $M_T = 1.6 \pm 0.1$ TeV in the electron channel, and $M_T = 2.4 \pm 0.1$ TeV in the muon channel. For $M_T > 1.6$ TeV, the background expectation from the fit to simulation is less than one event in each channel. Cross-section limits were derived using a Bayesian method [41] with a uniform prior probability distribution for the signal cross section. The number of data events above an optimised transverse mass threshold M_T^{min} was compared to the expected number of signal and background events. Systematic uncertainties on the signal and background yield were included via nuisance parameters with a log-normal prior distribution. The M_T^{min} threshold was optimised for the best expected exclusion limit, a procedure used in previous analyses [7] which is also appropriate for establishing a W' discovery. The M_T^{min} threshold defining the search window increases with W' mass up to masses around 2.5 TeV, following the Jacobian peak. For larger masses, cross sections become so small that fewer than two events are expected in the recorded data. These events are likely to have lower transverse mass because the production is shifted to the off-peak region, as mentioned above. Both these effects serve to lower the M_T^{min} threshold of the search window for very heavy W' bosons. The expected number of signal and background events listed separately for the two channels are summarized in Table 1. A common theoretical NNLO cross section is assumed.




Figure 3: Upper limits on $\sigma(W'_R) \times B(W'_R \rightarrow \ell\nu)$, with $\ell = e, \mu$, and their combination at 95% confidence level. The one (two) sigma uncertainty bands are shown in green (yellow). The theoretical cross section, with PDF uncertainties, is displayed with and without a mass-dependent NNLO K-factor for the right-handed model without interference. The theoretical cross sections for Kaluza–Klein W_{KK}^2 with $\mu=0.05$ TeV and $\mu=10$ TeV are also shown.

Table 1: M_T^{\min} requirement for different W'_R masses, expected number of signal and background events, number of observed events, theoretical cross section and upper limits on $\sigma(W'_R) \times B(W'_R \rightarrow \ell\nu)$, with $\ell = e, \mu$.

W' mass (GeV)	M_T^{\min} (GeV)	N_{sig} (Events)	N_{bkg} (Events)	N_{obs} (Events)	σ_{theory} (fb)	Exp. Limit (fb)	Obs. Limit (fb)
Electron channel							
500	350	44000 ± 4200	830 ± 85	850	17723	64.15	70.18
700	550	9600 ± 1500	114 ± 15	128	4514	16.94	22.48
900	700	3160 ± 460	37.4 ± 5.7	41	1470	8.38	9.61
1000	800	1730 ± 280	20.0 ± 3.8	22	886	6.77	7.55
1400	1050	294 ± 36	5.4 ± 1.6	6	144	3.56	3.77
1600	1150	128 ± 13	3.4 ± 1.1	5	63.3	3.02	3.80
1800	1200	63.9 ± 5.5	2.79 ± 0.99	3	28.5	2.53	2.57
2100	1350	18.7 ± 1.5	1.55 ± 0.64	2	9.37	2.38	2.61
2400	1450	5.47 ± 0.39	1.08 ± 0.49	2	3.40	2.69	3.39
2700	1450	1.75 ± 0.13	1.08 ± 0.49	2	1.43	3.54	4.46
3000	1400	0.59 ± 0.05	1.29 ± 0.56	2	0.71	5.45	6.42
Muon channel							
500	350	41000 ± 3200	749 ± 47	732	17723	44.65	39.13
700	550	8700 ± 1000	102 ± 10	100	4514	15.42	14.28
900	700	2920 ± 370	32.6 ± 5.0	36	1470	8.24	9.51
1000	750	1840 ± 150	23.3 ± 4.2	26	886	6.62	7.57
1400	1000	313 ± 25	5.6 ± 1.9	6	144	3.37	3.47
1600	1100	136.3 ± 9.2	3.4 ± 1.4	4	63.3	2.83	3.04
1800	1250	56.5 ± 3.7	1.78 ± 0.86	3	28.5	2.48	3.18
2100	1300	18.5 ± 0.9	1.45 ± 0.75	2	9.37	2.35	2.65
2400	1400	5.54 ± 0.26	0.98 ± 0.56	2	3.40	2.59	3.37
2700	1450	1.68 ± 0.08	0.81 ± 0.49	2	1.43	3.45	4.77
3000	1400	0.58 ± 0.03	0.98 ± 0.56	2	0.71	5.17	6.73

The expected and observed upper limits for both channels and their combination, in the right-handed scenario without interference, are shown in Figure 3. Using the central value of the theoretical cross section times the branching fraction, we exclude at 95% confidence level (CL) the existence of a W'_R with SM-like couplings of masses less than 2.5 TeV (compared with an expected limit of 2.6 TeV). Note that the background uncertainty has a negligible impact on the lower limits on W' mass, owing to the lack of observed events in the tail of the M_T distribution.

A similar search procedure was performed including the effect of interference. The theoretical cross sections are approximately 10–30% lower (higher) for destructive (constructive) interference when integrating over the transverse mass spectrum above 500 GeV and hence influence the resulting mass limits [17]. Optimising for the best expected cross section limit resulted in very similar search windows at high M_T , yielding lower limits on the W'_L mass of 2.63 (2.43) TeV for constructive (destructive) interference, based on the same MADGRAPH cross sections and K-factors as the ones used in Figure 3. We note that the interference affects mainly the medium M_T and hardly the Jacobian peak region, with the latter being used to set the limits. The limits shown do not take into account higher order electroweak corrections at high mass, which can be sizable. The effect of these missing corrections would be a reduction of the size of interference effects, leading to limits that are closer to the ones quoted for the no-interference case.

In addition to the model dependent results on W' production, upper limits for the cross section of beyond-the-SM production of charged lepton-neutrino events are given in Table 2 and Figure 4. The results are presented as a function of the transverse mass threshold, M_T^{\min} , and

Table 2: Excluded cross sections times branching fraction in the search window ($M_T > M_T^{\min}$) in the electron and muon channels individually, along with their combination. The number of expected background events was taken from simulation. The expected and observed cross section limits are given for each search window.

M_T^{\min} (GeV)	Electron channel				Muon channel				Combined channels	
	Events		Limit (fb)		Events		Limit (fb)		Limit (fb)	
	N_{bkg}	N_{obs}	Exp.	Obs.	N_{bkg}	N_{obs}	Exp.	Obs.	Exp.	Obs.
500	175 ± 22	192	10.14	13.85	158 ± 14	141	8.20	6.13	6.86	6.04
600	77 ± 10	83	5.99	7.13	67.9 ± 8.1	62	5.12	4.46	4.01	3.95
700	37.4 ± 5.7	41	3.80	4.57	32.6 ± 5.0	36	3.60	4.41	2.65	3.31
800	20.0 ± 3.8	22	3.03	3.24	17.0 ± 3.6	16	2.95	2.54	1.94	1.99
900	11.4 ± 2.6	12	2.10	2.30	9.5 ± 2.6	11	2.01	2.46	1.46	1.68
1000	6.8 ± 1.8	8	1.79	2.02	5.6 ± 1.9	6	1.57	1.80	1.11	1.32
1100	4.3 ± 1.3	6	1.40	1.88	3.4 ± 1.4	4	1.32	1.56	0.94	1.19
1200	2.79 ± 0.98	3	1.32	1.32	2.2 ± 1.0	3	1.18	1.45	0.78	0.92
1300	1.87 ± 0.74	2	1.15	1.15	1.45 ± 0.75	2	0.97	1.26	0.69	0.77
1400	1.29 ± 0.56	2	0.94	1.22	0.98 ± 0.56	2	1.00	1.32	0.59	0.85
1500	0.91 ± 0.43	1	0.97	0.97	0.68 ± 0.43	2	0.72	1.37	0.53	0.76




Figure 4: 95% confidence level upper limits on the cross section times branching fraction for physics beyond the SM (labelled BSM) for the charged lepton-neutrino production with transverse masses exceeding M_T^{\min} . The results for the electron, the muon channel, as well as for both channels combined are presented. The one (two) sigma uncertainty bands are shown in green (yellow).

listed separately for the electron and the muon channels, and their combination. The only assumptions made here are that we are searching for a narrow s -channel produced resonance, using the detector acceptance and selection efficiency outlined in Section 4. Note that the M_T^{\min} threshold is on an experimentally-measured quantity affected by detector resolution.

These exclusion limits on the cross-section can be translated to excluded W' masses within the context of a given model, such as constructive or destructive W'_L, W'_R or something else.




Figure 5: 95% confidence limits on the split-UED parameters μ and R derived from the W' mass limits taking into account the corresponding width of the W_{KK}^2 . The colored areas correspond to the W_{KK}^2 exclusion regions with the same final state as the SM-like W' . Results are shown for the electron and muon channels, as well as for both channels combined. The W_{KK}^2 is the lowest state that can couple to SM fermions. Since it has even parity it can be produced singly.

The observed limits illustrated in Figure 3 can be reinterpreted in terms of the W_{KK}^2 mass, as shown in the same figure for values of the bulk mass parameters $\mu = 0.05$ TeV and $\mu = 10$ TeV. For these parameters the second Kaluza–Klein excitation W_{KK}^2 has been excluded for masses below 1.4 TeV ($\mu = 0.05$ TeV) or 2.9 TeV ($\mu = 10$ TeV), respectively. The corresponding widths (Eq. (4)) are taken into account in the calculation of the cross section times the branching fraction of W_{KK}^2 . These lower limits on the mass can be directly translated to bounds on the split-UED parameter space $[1/R, \mu]$ with μ being the mass parameter for bulk fermions and R the radius of the extra dimension. The results are displayed in Figure 5, using the relations between R, μ and the W_{KK}^2 mass, and the couplings to SM fermions described by expressions (1), (2) and (3). The split-UED model also allows for W - W' interference. When the constructive case is considered, it has a comparable sensitivity to the no-interference case.

8 Summary

A search for an excess of events with a final state consisting of a charged lepton (electron or muon) and significant missing transverse momentum has been performed, using 5.0 fb^{-1} of $\sqrt{s} = 7 \text{ TeV}$ pp collision data. No significant excess over the SM expectation was observed in the distribution of transverse mass. A W'_R in the SSM with a mass of less than 2.5 TeV has been excluded at 95% CL. For the first time in such a study, W - W' interference effects have been taken into account, and mass exclusion limits have been determined as 2.63 TeV and 2.43 TeV for constructive and destructive interference respectively. These are the most stringent limits yet published. An interpretation of the search results has also been made in a specific framework of universal extra dimensions with bulk mass fermions. The second Kaluza–Klein excitation W_{KK}^2 has been excluded for masses below 1.4 TeV, assuming a bulk mass parameter μ of 0.05 TeV or masses below 2.9 TeV for $\mu=10 \text{ TeV}$.

Acknowledgements

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC machine. We thank the technical and administrative staff at CERN and other CMS institutes, and acknowledge support from: FMSR (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MoST, and NSFC (China); COLCIENCIAS (Colombia); MSES (Croatia); RPF (Cyprus); MoER, SF0690030s09 and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); OTKA and NKTH (Hungary); DAE and DST (India); I± (Iran); SFI (Ireland); INFN (Italy); NRF and WCU (Korea); LAS (Lithuania); CINVESTAV, CONACYT, SEP, and UASLP-FAI (Mexico); MSI (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Armenia, Belarus, Georgia, Ukraine, Uzbekistan); MON, RosAtom, RAS and RFBR (Russia); MSTD (Serbia); MICINN and CPAN (Spain); Swiss Funding Agencies (Switzerland); NSC (Taipei); TUBITAK and TAEK (Turkey); STFC (United Kingdom); DOE and NSF (USA). Individuals have received support from the Marie-Curie programme and the European Research Council (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the Council of Science and Industrial Research, India; and the HOMING PLUS programme of Foundation for Polish Science, cofinanced from European Union, Regional Development Fund.

References

- [1] CMS Collaboration, "The CMS experiment at the CERN LHC", *JINST* **03** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [2] CMS Collaboration, "Absolute Calibration of the Luminosity Measurement at CMS: Winter 2012 Update", CMS Physics Analysis Summary CMS-PAS-SMP-12-008, (2012).
- [3] G. Altarelli, B. Mele, and M. Ruiz-Altaba, "Searching for New Heavy Vector Bosons in $p\bar{p}$ Colliders", *Z. Phys. C* **45** (1989) 109, doi:10.1007/BF01556677. [Erratum-ibid. C47 (1990) 676].
- [4] CDF Collaboration, "Search for a New Heavy Gauge Boson W' with Electron + missing ET Event Signature in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV", *Phys. Rev. D* **83** (2011) 031102, doi:10.1103/PhysRevD.83.031102, arXiv:1012.5145.
- [5] D0 Collaboration, "Search for W' bosons decaying to an electron and a neutrino with the D0 detector", *Phys. Rev. Lett.* **100** (2008) 031804, doi:10.1103/PhysRevLett.100.031804, arXiv:0710.2966.
- [6] CMS Collaboration, "Search for a heavy gauge boson W' in the final state with an electron and large missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV", *Phys. Lett. B* **698** (2011) 21, doi:10.1016/j.physletb.2011.02.048, arXiv:1012.5945.
- [7] CMS Collaboration, "Search for a W' boson decaying to a muon and a neutrino in pp collisions at $\sqrt{s} = 7$ TeV", *Phys. Lett. B* **701** (2011) 160, doi:10.1016/j.physletb.2011.05.048, arXiv:1103.0030.

- [8] ATLAS Collaboration, "Search for high-mass states with one lepton plus missing transverse momentum in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector", *Phys. Lett. B* **701** (2011) 50, doi:10.1016/j.physletb.2011.05.043, arXiv:1103.1391.
- [9] ATLAS Collaboration, "Search for a heavy gauge boson decaying to a charged lepton and a neutrino in 1 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector", *Phys. Lett. B* **705** (2011) 28, doi:10.1016/j.physletb.2011.09.093, arXiv:1108.1316.
- [10] J. C. Pati and A. Salam, "Lepton number as the fourth 'color' ", *Phys. Rev. D* **10** (1974), no. 1, 275, doi:10.1103/PhysRevD.10.275. [Erratum-ibid. D **11** (1975) 703].
- [11] R. N. Mohapatra and J. C. Pati, "A Natural Left-Right Symmetry", *Phys. Rev. D* **11** (May, 1975) 2558–2561, doi:10.1103/PhysRevD.11.2558.
- [12] G. Senjanović and R. N. Mohapatra, "Exact Left-Right Symmetry and Spontaneous Violation of Parity", *Phys. Rev. D* **12** (1975), no. 5, 1502, doi:10.1103/PhysRevD.12.1502.
- [13] G. Senjanović, "Spontaneous breakdown of parity in a class of gauge theories", *Nucl. Phys. B* **153** (1979) 334, doi:10.1016/0550-3213(79)90604-7.
- [14] P. Minkowski, " $\mu \rightarrow e\gamma$ at a rate of one out of 1 billion muon decays?", *Phys. Lett. B* **67** (1977) 421, doi:10.1016/0370-2693(77)90435-X.
- [15] R. N. Mohapatra and G. Senjanović, "Neutrino Mass and Spontaneous Parity Nonconservation", *Phys. Rev. Lett.* **44** (1980) 912, doi:10.1103/PhysRevLett.44.912.
- [16] R. N. Mohapatra and G. Senjanović, "Neutrino Masses And Mixings In Gauge Models With Spontaneous Parity Violation", *Phys. Rev. D* **23** (1981) 165, doi:10.1103/PhysRevD.23.165.
- [17] E. Accomando et al., "Interference effects in heavy W' -boson searches at the LHC", (2011). arXiv:1110.0713.
- [18] T. G. Rizzo, "The determination of the helicity of W' boson couplings at the LHC", *JHEP* **05** (2007) 037, doi:10.1088/1126-6708/2007/05/037.
- [19] E. Boos et al., "Interference between W' and W in single-top quark production processes", *Phys. Lett. B* **655** (2007) 245, doi:10.1016/j.physletb.2007.03.064, arXiv:hep-ph/0610080.
- [20] D0 Collaboration, "Search for W' Boson Resonances Decaying to a Top Quark and a Bottom Quark", *Phys. Rev. Lett.* **100** (2008) 211803, doi:10.1103/PhysRevLett.100.211803, arXiv:0803.3256.
- [21] C.-R. Chen et al., "Dark matter and collider phenomenology of split-UED", *JHEP* **09** (2009) 078, doi:10.1088/1126-6708/2009/09/078, arXiv:0903.1971.
- [22] K. Kong, S. C. Park, and T. G. Rizzo, "Collider Phenomenology with Split-UED", *JHEP* **04** (2010) 081, doi:10.1007/JHEP04(2010)081, arXiv:1002.0602.
- [23] T. Appelquist, H.-C. Cheng, and B. A. Dobrescu, "Bounds on universal extra dimensions", *Phys. Rev. D* **64** (2001) 035002, doi:10.1103/PhysRevD.64.035002, arXiv:hep-ph/0012100.

- [24] CMS Collaboration, “Particle–Flow Event Reconstruction in CMS and Performance for Jets, Taus, and E_T^{miss} ”, CMS Physics Analysis Summary CMS-PAS-PFT-09-001, (2009).
- [25] GEANT4 Collaboration, “GEANT4 – a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [26] J. Allison et al., “GEANT4 developments and applications”, *IEEE Trans. Nucl. Sci.* **53** (2006) 270, doi:10.1109/TNS.2006.869826.
- [27] T. Sjöstrand, S. Mrenna, and P. Z. Skands, “PYTHIA 6.4 Physics and Manual”, *JHEP* **05** (2006) 026, doi:10.1088/1126-6708/2006/05/026, arXiv:hep-ph/0603175.
- [28] J. Alwall et al., “MadGraph/MadEvent v4: the new web generation”, *JHEP* **09** (2007) 028, doi:10.1088/1126-6708/2007/09/028, arXiv:0706.2334v1.
- [29] J. Pumplin et al., “New generation of parton distributions with uncertainties from global QCD analysis”, *JHEP* **07** (2002) 012, doi:10.1088/1126-6708/2002/07/012, arXiv:hep-ph/0201195.
- [30] R. Gavin et al., “FEWZ 2.0: A code for hadronic Z production at next-to-next-to-leading order”, *Comput. Phys. Commun.* **182** (2011) 2388, doi:10.1016/j.cpc.2011.06.008, arXiv:1011.3540.
- [31] R. Gavin et al., “W physics at the LHC with FEWZ 2.1”, (2012). arXiv:1201.5896.
- [32] CompHEP Collaboration, “CompHEP 4.4: Automatic computations from Lagrangians to events”, *Nucl. Instrum. Meth. A* **534** (2004) 250, doi:10.1016/j.nima.2004.07.096, arXiv:hep-ph/0403113.
- [33] A. Pukhov et al., “CompHEP: A Package for evaluation of Feynman diagrams and integration over multiparticle phase space”, (1999). arXiv:hep-ph/9908288. User’s manual for version 33.
- [34] N. Kidonakis and R. Vogt, “The Theoretical top quark cross section at the Tevatron and the LHC”, *Phys. Rev. D* **78** (2008) 074005, doi:10.1103/PhysRevD.78.074005, arXiv:0805.3844.
- [35] CMS Collaboration, “Performance of CMS muon reconstruction in cosmic-ray events”, *J. Instrum.* **5** (2009) T03022, doi:10.1088/1748-0221/5/03/T03022.
- [36] M. Botje et al., “The PDF4LHC Working Group Interim Recommendations”, (2011). arXiv:1101.0538.
- [37] M. R. Whalley, D. Bourilkov, and R. C. Group, “The Les Houches Accord PDFs (LHAPDF) and Lhaglué”, (2005). arXiv:hep-ph/0508110.
- [38] A. D. Martin et al., “Parton distributions for the LHC”, *Eur. Phys. J. C* **63** (2009) 189, doi:10.1140/epjc/s10052-009-1072-5, arXiv:0901.0002.
- [39] P. M. Nadolsky et al., “Implications of CTEQ global analysis for collider observables”, *Phys. Rev. D* **78** (2008) 013004, doi:10.1103/PhysRevD.78.013004, arXiv:0802.0007.
- [40] S. Forte et al., “Neural network parametrization of deep inelastic structure functions”, *JHEP* **05** (2002) 062, doi:10.1088/1126-6708/2002/05/062, arXiv:hep-ph/0204232.

- [41] Particle Data Group Collaboration, "Review of Particle Physics", *J. Phys. G* **37** (2010) 075021, doi:10.1088/0954-3899/37/7A/075021. Chap. 33.

A The CMS Collaboration

Yerevan Physics Institute, Yerevan, Armenia

S. Chatrchyan, V. Khachatryan, A.M. Sirunyan, A. Tumasyan

Institut für Hochenergiephysik der OeAW, Wien, Austria

W. Adam, T. Bergauer, M. Dragicevic, J. Erö, C. Fabjan, M. Friedl, R. Frühwirth, V.M. Ghete, J. Hammer¹, N. Hörmann, J. Hrubec, M. Jeitler, W. Kiesenhofer, V. Knünz, M. Krammer, D. Liko, I. Mikulec, M. Pernicka[†], B. Rahbaran, C. Rohringer, H. Rohringer, R. Schöffbeck, J. Strauss, A. Taurok, F. Teischinger, P. Wagner, W. Waltenberger, G. Walzel, E. Widl, C.-E. Wulz

National Centre for Particle and High Energy Physics, Minsk, Belarus

V. Mossolov, N. Shumeiko, J. Suarez Gonzalez

Universiteit Antwerpen, Antwerpen, Belgium

S. Bansal, K. Cerny, T. Cornelis, E.A. De Wolf, X. Janssen, S. Luyckx, T. Maes, L. Mucibello, S. Ochesanu, B. Roland, R. Rougny, M. Selvaggi, H. Van Haeevermaet, P. Van Mechelen, N. Van Remortel, A. Van Spilbeeck

Vrije Universiteit Brussel, Brussel, Belgium

F. Blekman, S. Blyweert, J. D'Hondt, R. Gonzalez Suarez, A. Kalogeropoulos, M. Maes, A. Olbrechts, W. Van Doninck, P. Van Mulders, G.P. Van Onsem, I. Villella

Université Libre de Bruxelles, Bruxelles, Belgium

O. Charaf, B. Clerbaux, G. De Lentdecker, V. Dero, A.P.R. Gay, T. Hreus, A. Léonard, P.E. Marage, T. Reis, L. Thomas, C. Vander Velde, P. Vanlaer

Ghent University, Ghent, Belgium

V. Adler, K. Bernaert, A. Cimmino, S. Costantini, G. Garcia, M. Grunewald, B. Klein, J. Lellouch, A. Marinov, J. McCartin, A.A. Ocampo Rios, D. Ryckbosch, N. Strobbe, F. Thyssen, M. Tytgat, L. Vanelderen, P. Verwilligen, S. Walsh, E. Yazgan, N. Zaganidis

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Basesmez, G. Bruno, L. Ceard, C. Delaere, T. du Pree, D. Favart, L. Forthomme, A. Giammanco², J. Hollar, V. Lemaître, J. Liao, O. Militaru, C. Nuttens, D. Pagano, A. Pin, K. Piotrkowski, N. Schul

Université de Mons, Mons, Belgium

N. Belyi, T. Caebergs, E. Daubie, G.H. Hammad

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves, M. Correa Martins Junior, D. De Jesus Damiao, T. Martins, M.E. Pol, M.H.G. Souza

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior, W. Carvalho, A. Custódio, E.M. Da Costa, C. De Oliveira Martins, S. Fonseca De Souza, D. Matos Figueiredo, L. Mundim, H. Nogima, V. Oguri, W.L. Prado Da Silva, A. Santoro, S.M. Silva Do Amaral, L. Soares Jorge, A. Sznajder

Instituto de Fisica Teorica, Universidade Estadual Paulista, Sao Paulo, Brazil

T.S. Anjos³, C.A. Bernardes³, F.A. Dias⁴, T.R. Fernandez Perez Tomei, E. M. Gregores³, C. Lagana, F. Marinho, P.G. Mercadante³, S.F. Novaes, Sandra S. Padula

Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria

V. Genchev¹, P. Iaydjiev¹, S. Piperov, M. Rodozov, S. Stoykova, G. Sultanov, V. Tcholakov, R. Trayanov, M. Vutova

University of Sofia, Sofia, Bulgaria

A. Dimitrov, R. Hadjiiska, V. Kozhuharov, L. Litov, B. Pavlov, P. Petkov

Institute of High Energy Physics, Beijing, China

J.G. Bian, G.M. Chen, H.S. Chen, C.H. Jiang, D. Liang, S. Liang, X. Meng, J. Tao, J. Wang, J. Wang, X. Wang, Z. Wang, H. Xiao, M. Xu, J. Zang, Z. Zhang

State Key Lab. of Nucl. Phys. and Tech., Peking University, Beijing, China

C. Asawatangtrakuldee, Y. Ban, S. Guo, Y. Guo, W. Li, S. Liu, Y. Mao, S.J. Qian, H. Teng, S. Wang, B. Zhu, W. Zou

Universidad de Los Andes, Bogota, Colombia

C. Avila, B. Gomez Moreno, A.F. Osorio Oliveros, J.C. Sanabria

Technical University of Split, Split, Croatia

N. Godinovic, D. Lelas, R. Plestina⁵, D. Polic, I. Puljak¹

University of Split, Split, Croatia

Z. Antunovic, M. Dzelalija, M. Kovac

Institute Rudjer Boskovic, Zagreb, Croatia

V. Brigljevic, S. Duric, K. Kadija, J. Luetic, S. Morovic

University of Cyprus, Nicosia, Cyprus

A. Attikis, M. Galanti, G. Mavromanolakis, J. Mousa, C. Nicolaou, F. Ptochos, P.A. Razis

Charles University, Prague, Czech Republic

M. Finger, M. Finger Jr.

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

Y. Assran⁶, S. Elgammal, A. Ellithi Kamel⁷, S. Khalil⁸, M.A. Mahmoud⁹, A. Radi^{8,10}

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

M. Kadastik, M. Müntel, M. Raidal, L. Rebane, A. Tiko

Department of Physics, University of Helsinki, Helsinki, Finland

V. Azzolini, P. Eerola, G. Fedi, M. Voutilainen

Helsinki Institute of Physics, Helsinki, Finland

J. Härkönen, A. Heikkinen, V. Karimäki, R. Kinnunen, M.J. Kortelainen, T. Lampén, K. Lassila-Perini, S. Lehti, T. Lindén, P. Luukka, T. Mäenpää, T. Peltola, E. Tuominen, J. Tuominiemi, E. Tuovinen, D. Ungaro, L. Wendland

Lappeenranta University of Technology, Lappeenranta, Finland

K. Banzuzi, A. Korpela, T. Tuuva

DSM/IRFU, CEA/Saclay, Gif-sur-Yvette, France

M. Besancon, S. Choudhury, M. Dejardin, D. Denegri, B. Fabbro, J.L. Faure, F. Ferri, S. Ganjour, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, E. Locci, J. Malcles, L. Millischer, A. Nayak, J. Rander, A. Rosowsky, I. Shreyber, M. Titov

Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France

S. Baffioni, F. Beaudette, L. Benhabib, L. Bianchini, M. Bluj¹¹, C. Broutin, P. Busson, C. Charlot, N. Daci, T. Dahms, L. Dobrzynski, R. Granier de Cassagnac, M. Haguenaer, P. Miné, C. Mironov, C. Ochando, P. Paganini, D. Sabes, R. Salerno, Y. Sirois, C. Veelken, A. Zabi

Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France

J.-L. Agram¹², J. Andrea, D. Bloch, D. Bodin, J.-M. Brom, M. Cardaci, E.C. Chabert, C. Collard, E. Conte¹², F. Drouhin¹², C. Ferro, J.-C. Fontaine¹², D. Gelé, U. Goerlach, P. Juillot, M. Karim¹², A.-C. Le Bihan, P. Van Hove

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules (IN2P3), Villeurbanne, France

F. Fassi, D. Mercier

Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique Nucléaire de Lyon, Villeurbanne, France

S. Beauceron, N. Beaupere, O. Bondu, G. Boudoul, H. Brun, J. Chasserat, R. Chierici¹, D. Contardo, P. Depasse, H. El Mamouni, J. Fay, S. Gascon, M. Gouzevitch, B. Ille, T. Kurca, M. Lethuillier, L. Mirabito, S. Perries, V. Sordini, S. Tosi, Y. Tschudi, P. Verdier, S. Viret

Institute of High Energy Physics and Informatization, Tbilisi State University, Tbilisi, Georgia

Z. Tsamalaidze¹³

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

G. Anagnostou, S. Beranek, M. Edelhoff, L. Feld, N. Heracleous, O. Hindrichs, R. Jussen, K. Klein, J. Merz, A. Ostapchuk, A. Perieanu, F. Raupach, J. Sammet, S. Schael, D. Sprenger, H. Weber, B. Wittmer, V. Zhukov¹⁴

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

M. Ata, J. Caudron, E. Dietz-Laursonn, D. Duchardt, M. Erdmann, A. Güth, T. Hebbeker, C. Heidemann, K. Hoepfner, T. Klimkovich, D. Klingebiel, P. Kreuzer, D. Lanske[†], J. Lingemann, C. Magass, M. Merschmeyer, A. Meyer, M. Olschewski, P. Papacz, H. Pieta, H. Reithler, S.A. Schmitz, J.F. Schulte, L. Sonnenschein, J. Steggemann, D. Teyssier, S. Thüer, M. Weber

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

M. Bontenackels, V. Cherepanov, M. Davids, G. Flügge, H. Geenen, M. Geisler, W. Haj Ahmad, F. Hoehle, B. Kargoll, T. Kress, Y. Kuessel, A. Linn, A. Nowack, L. Perchalla, O. Pooth, J. Rennefeld, P. Sauerland, A. Stahl

Deutsches Elektronen-Synchrotron, Hamburg, Germany

M. Aldaya Martin, J. Behr, W. Behrenhoff, U. Behrens, M. Bergholz¹⁵, A. Bethani, K. Borras, A. Burgmeier, A. Cakir, L. Calligaris, A. Campbell, E. Castro, F. Costanza, D. Dammann, G. Eckerlin, D. Eckstein, D. Fischer, G. Flucke, A. Geiser, I. Glushkov, S. Habib, J. Hauk, H. Jung¹, M. Kasemann, P. Katsas, C. Kleinwort, H. Kluge, A. Knutsson, M. Krämer, D. Krücker, E. Kuznetsova, W. Lange, W. Lohmann¹⁵, B. Lutz, R. Mankel, I. Marfin, M. Marienfeld, I.-A. Melzer-Pellmann, A.B. Meyer, J. Mnich, A. Mussgiller, S. Naumann-Emme, J. Olzem, H. Perrey, A. Petrukhin, D. Pitzl, A. Raspereza, P.M. Ribeiro Cipriano, C. Riedl, M. Rosin, J. Salfeld-Nebgen, R. Schmidt¹⁵, T. Schoerner-Sadenius, N. Sen, A. Spiridonov, M. Stein, R. Walsh, C. Wissing

University of Hamburg, Hamburg, Germany

C. Autermann, V. Blobel, S. Bobrovskiy, J. Draeger, H. Enderle, J. Erfle, U. Gebbert, M. Görner, T. Hermanns, R.S. Höing, K. Kaschube, G. Kaussen, H. Kirschenmann, R. Klanner, J. Lange, B. Mura, F. Nowak, N. Pietsch, D. Rathjens, C. Sander, H. Schettler, P. Schleper, E. Schlieckau, A. Schmidt, M. Schröder, T. Schum, M. Seidel, H. Stadie, G. Steinbrück, J. Thomsen

Institut für Experimentelle Kernphysik, Karlsruhe, Germany

C. Barth, J. Berger, T. Chwalek, W. De Boer, A. Dierlamm, M. Feindt, M. Guthoff¹, C. Hackstein, F. Hartmann, M. Heinrich, H. Held, K.H. Hoffmann, S. Honc, I. Katkov¹⁴, J.R. Komaragiri, D. Martschei, S. Mueller, Th. Müller, M. Niegel, A. Nürnberg, O. Oberst, A. Oehler, J. Ott, T. Peiffer, G. Quast, K. Rabbertz, F. Ratnikov, N. Ratnikova, S. Röcker, C. Saout, A. Scheurer, F.-P. Schilling, M. Schmanau, G. Schott, H.J. Simonis, F.M. Stober, D. Troendle, R. Ulrich, J. Wagner-Kuhr, T. Weiler, M. Zeise, E.B. Ziebarth

Institute of Nuclear Physics "Demokritos", Aghia Paraskevi, Greece

G. Daskalakis, T. Gerasis, S. Kesisoglou, A. Kyriakis, D. Loukas, I. Manolakos, A. Markou, C. Markou, C. Mavrommatis, E. Ntomari

University of Athens, Athens, Greece

L. Gouskos, T.J. Mertzimekis, A. Panagiotou, N. Saoulidou

University of Ioánnina, Ioánnina, Greece

I. Evangelou, C. Foudas¹, P. Kokkas, N. Manthos, I. Papadopoulos, V. Patras

KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary

G. Bencze, C. Hajdu¹, P. Hidas, D. Horvath¹⁶, K. Krajczar¹⁷, B. Radics, F. Sikler¹, V. Veszpremi, G. Vesztergombi¹⁷

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

N. Beni, S. Czellar, J. Molnar, J. Palinkas, Z. Szillasi

University of Debrecen, Debrecen, Hungary

J. Karancsi, P. Raics, Z.L. Trocsanyi, B. Ujvari

Panjab University, Chandigarh, India

S.B. Beri, V. Bhatnagar, N. Dhingra, R. Gupta, M. Jindal, M. Kaur, J.M. Kohli, M.Z. Mehta, N. Nishu, L.K. Saini, A. Sharma, J. Singh, S.P. Singh

University of Delhi, Delhi, India

S. Ahuja, A. Bhardwaj, B.C. Choudhary, A. Kumar, A. Kumar, S. Malhotra, M. Naimuddin, K. Ranjan, V. Sharma, R.K. Shivpuri

Saha Institute of Nuclear Physics, Kolkata, India

S. Banerjee, S. Bhattacharya, S. Dutta, B. Gomber, Sa. Jain, Sh. Jain, R. Khurana, S. Sarkar

Bhabha Atomic Research Centre, Mumbai, India

A. Abdulsalam, R.K. Choudhury, D. Dutta, S. Kailas, V. Kumar, A.K. Mohanty¹, L.M. Pant, P. Shukla

Tata Institute of Fundamental Research - EHEP, Mumbai, India

T. Aziz, S. Ganguly, M. Guchait¹⁸, A. Gurtu¹⁹, M. Maity²⁰, G. Majumder, K. Mazumdar, G.B. Mohanty, B. Parida, K. Sudhakar, N. Wickramage

Tata Institute of Fundamental Research - HECR, Mumbai, India

S. Banerjee, S. Dugad

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

H. Arfaei, H. Bakhshiansohi²¹, S.M. Etesami²², A. Fahim²¹, M. Hashemi, H. Hesari, A. Jafari²¹, M. Khakzad, A. Mohammadi²³, M. Mohammadi Najafabadi, S. Paktinat Mehdiabadi, B. Safarzadeh²⁴, M. Zeinali²²

INFN Sezione di Bari ^a, Università di Bari ^b, Politecnico di Bari ^c, Bari, Italy

M. Abbrescia^{a,b}, L. Barbone^{a,b}, C. Calabria^{a,b,1}, S.S. Chhibra^{a,b}, A. Colaleo^a, D. Creanza^{a,c}, N. De Filippis^{a,c,1}, M. De Palma^{a,b}, L. Fiore^a, G. Iaselli^{a,c}, L. Lusito^{a,b}, G. Maggi^{a,c}, M. Maggi^a, B. Marangelli^{a,b}, S. My^{a,c}, S. Nuzzo^{a,b}, N. Pacifico^{a,b}, A. Pompili^{a,b}, G. Pugliese^{a,c}, G. Selvaggi^{a,b}, L. Silvestris^a, G. Singh^{a,b}, G. Zito^a

INFN Sezione di Bologna ^a, Università di Bologna ^b, Bologna, Italy

G. Abbiendi^a, A.C. Benvenuti^a, D. Bonacorsi^{a,b}, S. Braibant-Giacomelli^{a,b}, L. Brigliadori^{a,b}, P. Capiluppi^{a,b}, A. Castro^{a,b}, F.R. Cavallo^a, M. Cuffiani^{a,b}, G.M. Dallavalle^a, F. Fabbri^a, A. Fanfani^{a,b}, D. Fasanella^{a,b,1}, P. Giacomelli^a, C. Grandi^a, L. Guiducci, S. Marcellini^a, G. Masetti^a, M. Meneghelli^{a,b,1}, A. Montanari^a, F.L. Navarria^{a,b}, F. Odorici^a, A. Perrotta^a, F. Primavera^{a,b}, A.M. Rossi^{a,b}, T. Rovelli^{a,b}, G. Siroli^{a,b}, R. Travaglini^{a,b}

INFN Sezione di Catania ^a, Università di Catania ^b, Catania, Italy

S. Albergo^{a,b}, G. Cappello^{a,b}, M. Chiorboli^{a,b}, S. Costa^{a,b}, R. Potenza^{a,b}, A. Tricomi^{a,b}, C. Tuve^{a,b}

INFN Sezione di Firenze ^a, Università di Firenze ^b, Firenze, Italy

G. Barbagli^a, V. Ciulli^{a,b}, C. Civinini^a, R. D'Alessandro^{a,b}, E. Focardi^{a,b}, S. Frosali^{a,b}, E. Gallo^a, S. Gonzi^{a,b}, M. Meschini^a, S. Paoletti^a, G. Sguazzoni^a, A. Tropiano^{a,1}

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi, S. Bianco, S. Colafranceschi²⁵, F. Fabbri, D. Piccolo

INFN Sezione di Genova, Genova, Italy

P. Fabbriatore, R. Musenich

INFN Sezione di Milano-Bicocca ^a, Università di Milano-Bicocca ^b, Milano, Italy

A. Benaglia^{a,b,1}, F. De Guio^{a,b}, L. Di Matteo^{a,b,1}, S. Fiorendi^{a,b}, S. Gennai^{a,1}, A. Ghezzi^{a,b}, S. Malvezzi^a, R.A. Manzoni^{a,b}, A. Martelli^{a,b}, A. Massironi^{a,b,1}, D. Menasce^a, L. Moroni^a, M. Paganoni^{a,b}, D. Pedrini^a, S. Ragazzi^{a,b}, N. Redaelli^a, S. Sala^a, T. Tabarelli de Fatis^{a,b}

INFN Sezione di Napoli ^a, Università di Napoli "Federico II" ^b, Napoli, Italy

S. Buontempo^a, C.A. Carrillo Montoya^{a,1}, N. Cavallo^{a,26}, A. De Cosa^{a,b}, O. Dogangun^{a,b}, F. Fabozzi^{a,26}, A.O.M. Iorio^{a,1}, L. Lista^a, S. Meola^{a,27}, M. Merola^{a,b}, P. Paolucci^a

INFN Sezione di Padova ^a, Università di Padova ^b, Università di Trento (Trento) ^c, Padova, Italy

P. Azzi^a, N. Bacchetta^{a,1}, P. Bellan^{a,b}, D. Bisello^{a,b}, A. Branca^{a,1}, R. Carlin^{a,b}, P. Checchia^a, T. Dorigo^a, U. Dosselli^a, F. Gasparini^{a,b}, A. Gozzelino^a, K. Kanishchev^{a,c}, S. Lacaprara^a, I. Lazzizzera^{a,c}, M. Margoni^{a,b}, A.T. Meneguzzo^{a,b}, L. Perrozzi^a, N. Pozzobon^{a,b}, P. Ronchese^{a,b}, F. Simonetto^{a,b}, E. Torassa^a, M. Tosi^{a,b,1}, S. Vanini^{a,b}, P. Zotto^{a,b}, G. Zumerle^{a,b}

INFN Sezione di Pavia ^a, Università di Pavia ^b, Pavia, Italy

M. Gabusi^{a,b}, S.P. Ratti^{a,b}, C. Riccardi^{a,b}, P. Torre^{a,b}, P. Vitulo^{a,b}

INFN Sezione di Perugia ^a, Università di Perugia ^b, Perugia, Italy

G.M. Bilei^a, L. Fano^{a,b}, P. Lariccia^{a,b}, A. Lucaroni^{a,b,1}, G. Mantovani^{a,b}, M. Menichelli^a, A. Nappi^{a,b}, F. Romeo^{a,b}, A. Saha, A. Santocchia^{a,b}, S. Taroni^{a,b,1}

INFN Sezione di Pisa ^a, Università di Pisa ^b, Scuola Normale Superiore di Pisa ^c, Pisa, Italy

P. Azzurri^{a,c}, G. Bagliesi^a, T. Boccali^a, G. Broccolo^{a,c}, R. Castaldi^a, R.T. D'Agnolo^{a,c}, R. Dell'Orso^a, F. Fiori^{a,b,1}, L. Foà^{a,c}, A. Giassi^a, A. Kraan^a, F. Ligabue^{a,c}, T. Lomtadze^a, L. Martini^{a,28}, A. Messineo^{a,b}, F. Palla^a, F. Palmonari^a, A. Rizzi^{a,b}, A.T. Serban^{a,29}, P. Spagnolo^a, P. Squillacioti¹, R. Tenchini^a, G. Tonelli^{a,b,1}, A. Venturi^{a,1}, P.G. Verdini^a

INFN Sezione di Roma ^a, Università di Roma "La Sapienza" ^b, Roma, Italy

L. Barone^{a,b}, F. Cavallari^a, D. Del Re^{a,b,1}, M. Diemoz^a, C. Fanelli^{a,b}, M. Grassi^{a,1}, E. Longo^{a,b}, P. Meridiani^{a,1}, F. Micheli^{a,b}, S. Nourbakhsh^a, G. Organtini^{a,b}, F. Pandolfi^{a,b}, R. Paramatti^a, S. Rahatlou^{a,b}, M. Sigamani^a, L. Soffi^{a,b}

INFN Sezione di Torino ^a, Università di Torino ^b, Università del Piemonte Orientale (Novara) ^c, Torino, Italy

N. Amapane^{a,b}, R. Arcidiacono^{a,c}, S. Argiro^{a,b}, M. Arneodo^{a,c}, C. Biino^a, C. Botta^{a,b}, N. Cartiglia^a, R. Castello^{a,b}, M. Costa^{a,b}, N. Demaria^a, A. Graziano^{a,b}, C. Mariotti^{a,1}, S. Maselli^a, E. Migliore^{a,b}, V. Monaco^{a,b}, M. Musich^{a,1}, M.M. Obertino^{a,c}, N. Pastrone^a, M. Pelliccioni^a, A. Potenza^{a,b}, A. Romero^{a,b}, M. Ruspai^{a,c}, R. Sacchi^{a,b}, V. Sola^{a,b}, A. Solano^{a,b}, A. Staiano^a, A. Vilela Pereira^a

INFN Sezione di Trieste ^a, Università di Trieste ^b, Trieste, Italy

S. Belforte^a, F. Cossutti^a, G. Della Ricca^{a,b}, B. Gobbo^a, M. Marone^{a,b,1}, D. Montanino^{a,b,1}, A. Penzo^a, A. Schizzi^{a,b}

Kangwon National University, Chunchon, Korea

S.G. Heo, T.Y. Kim, S.K. Nam

Kyungpook National University, Daegu, Korea

S. Chang, J. Chung, D.H. Kim, G.N. Kim, D.J. Kong, H. Park, S.R. Ro, D.C. Son, T. Son

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

J.Y. Kim, Zero J. Kim, S. Song

Konkuk University, Seoul, Korea

H.Y. Jo

Korea University, Seoul, Korea

S. Choi, D. Gyun, B. Hong, M. Jo, H. Kim, T.J. Kim, K.S. Lee, D.H. Moon, S.K. Park, E. Seo

University of Seoul, Seoul, Korea

M. Choi, S. Kang, H. Kim, J.H. Kim, C. Park, I.C. Park, S. Park, G. Ryu

Sungkyunkwan University, Suwon, Korea

Y. Cho, Y. Choi, Y.K. Choi, J. Goh, M.S. Kim, E. Kwon, B. Lee, J. Lee, S. Lee, H. Seo, I. Yu

Vilnius University, Vilnius, Lithuania

M.J. Bilinskas, I. Grigelionis, M. Janulis, A. Juodagalvis

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

H. Castilla-Valdez, E. De La Cruz-Burelo, I. Heredia-de La Cruz, R. Lopez-Fernandez, R. Magaña Villalba, J. Martínez-Ortega, A. Sánchez-Hernández, L.M. Villasenor-Cendejas

Universidad Iberoamericana, Mexico City, Mexico

S. Carrillo Moreno, F. Vazquez Valencia

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

H.A. Salazar Ibarguen

Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico

E. Casimiro Linares, A. Morelos Pineda, M.A. Reyes-Santos

University of Auckland, Auckland, New Zealand

D. Krofcheck

University of Canterbury, Christchurch, New Zealand

A.J. Bell, P.H. Butler, R. Doesburg, S. Reucroft, H. Silverwood

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

M. Ahmad, M.I. Asghar, H.R. Hoorani, S. Khalid, W.A. Khan, T. Khurshid, S. Qazi, M.A. Shah, M. Shoaib

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

G. Brona, K. Bunkowski, M. Cwiok, W. Dominik, K. Doroba, A. Kalinowski, M. Konecki, J. Krolikowski

Soltan Institute for Nuclear Studies, Warsaw, Poland

H. Bialkowska, B. Boimska, T. Frueboes, R. Gokieli, M. Górski, M. Kazana, K. Nawrocki, K. Romanowska-Rybinska, M. Szleper, G. Wrochna, P. Zalewski

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

N. Almeida, P. Bargassa, A. David, P. Faccioli, P.G. Ferreira Parracho, M. Gallinaro, P. Musella, J. Seixas, J. Varela, P. Vischia

Joint Institute for Nuclear Research, Dubna, Russia

I. Belotelov, M. Gavrilenko, I. Golutvin, I. Gorbunov, A. Kamenev, V. Karjavin, G. Kozlov, A. Lanev, A. Malakhov, P. Moisezenz, V. Palichik, V. Perelygin, M. Savina, S. Shmatov, V. Smirnov, A. Volodko, A. Zarubin

Petersburg Nuclear Physics Institute, Gatchina (St Petersburg), Russia

S. Evstyukhin, V. Golovtsov, Y. Ivanov, V. Kim, P. Levchenko, V. Murzin, V. Oreshkin, I. Smirnov, V. Sulimov, L. Uvarov, S. Vavilov, A. Vorobyev, An. Vorobyev

Institute for Nuclear Research, Moscow, Russia

Yu. Andreev, A. Dermenev, S. Gninenko, N. Golubev, M. Kirsanov, N. Krasnikov, V. Matveev, A. Pashenkov, D. Tlisov, A. Toropin

Institute for Theoretical and Experimental Physics, Moscow, Russia

V. Epshteyn, M. Erofeeva, V. Gavrilov, M. Kossov¹, N. Lychkovskaya, V. Popov, G. Safronov, S. Semenov, V. Stolin, E. Vlasov, A. Zhokin

Moscow State University, Moscow, Russia

A. Belyaev, E. Boos, V. Bunichev, M. Dubinin⁴, L. Dudko, A. Ershov, A. Gribushin, V. Klyukhin, O. Kodolova, I. Lokhtin, A. Markina, S. Obraztsov, M. Perfilov, S. Petrushanko, A. Popov, L. Sarycheva[†], V. Savrin

P.N. Lebedev Physical Institute, Moscow, Russia

V. Andreev, M. Azarkin, I. Dremin, M. Kirakosyan, A. Leonidov, G. Mesyats, S.V. Rusakov, A. Vinogradov

State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, Russia

I. Azhgirey, I. Bayshev, S. Bitioukov, V. Grishin¹, V. Kachanov, D. Konstantinov, A. Korablev, V. Krychkin, V. Petrov, R. Ryutin, A. Sobol, L. Tourtchanovitch, S. Troshin, N. Tyurin, A. Uzunian, A. Volkov

University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia

P. Adzic³⁰, M. Djordjevic, M. Ekmedzic, D. Krpic³⁰, J. Milosevic

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Aguilar-Benitez, J. Alcaraz Maestre, P. Arce, C. Battilana, E. Calvo, M. Cerrada, M. Chamizo Llatas, N. Colino, B. De La Cruz, A. Delgado Peris, C. Diez Pardos, D. Domínguez Vázquez, C. Fernandez Bedoya, J.P. Fernández Ramos, A. Ferrando, J. Flix, M.C. Fouz, P. Garcia-Abia, O. Gonzalez Lopez, S. Goy Lopez, J.M. Hernandez, M.I. Josa, G. Merino, J. Puerta Pelayo, I. Redondo, L. Romero, J. Santaolalla, M.S. Soares, C. Willmott

Universidad Autónoma de Madrid, Madrid, Spain

C. Albajar, G. Codispoti, J.F. de Trocóniz

Universidad de Oviedo, Oviedo, Spain

J. Cuevas, J. Fernandez Menendez, S. Folgueras, I. Gonzalez Caballero, L. Lloret Iglesias, J. Piedra Gomez³¹, J.M. Vizán Garcia

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

J.A. Brochero Cifuentes, I.J. Cabrillo, A. Calderon, S.H. Chuang, J. Duarte Campderros, M. Felcini³², M. Fernandez, G. Gomez, J. Gonzalez Sanchez, C. Jorda, P. Lobelle Pardo, A. Lopez Virto, J. Marco, R. Marco, C. Martinez Rivero, F. Matorras, F.J. Munoz Sanchez, T. Rodrigo, A.Y. Rodríguez-Marrero, A. Ruiz-Jimeno, L. Scodellaro, M. Sobron Sanudo, I. Vila, R. Vilar Cortabitarte

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo, E. Auffray, G. Auzinger, P. Baillon, A.H. Ball, D. Barney, C. Bernet⁵, G. Bianchi, P. Bloch, A. Bocci, A. Bonato, H. Breuker, T. Camporesi, G. Cerminara, T. Christiansen, J.A. Coarasa Perez, D. D'Enterria, A. De Roeck, S. Di Guida, M. Dobson, N. Dupont-Sagorin, A. Elliott-Peisert, B. Frisch, W. Funk, G. Georgiou, M. Giffels, D. Gigi, K. Gill, D. Giordano, M. Giunta, F. Glege, R. Gomez-Reino Garrido, P. Govoni, S. Gowdy, R. Guida, M. Hansen, P. Harris, C. Hartl, J. Harvey, B. Hegner, A. Hinzmann, V. Innocente, P. Janot, K. Kaadze, E. Karavakis, K. Kousouris, P. Lecoq, P. Lenzi, C. Lourenço, T. Mäki, M. Malberti, L. Malgeri, M. Mannelli, L. Masetti, F. Meijers, S. Mersi, E. Meschi, R. Moser, M.U. Mozer, M. Mulders, E. Nesvold, M. Nguyen, T. Orimoto, L. Orsini, E. Palencia Cortezon, E. Perez, A. Petrilli, A. Pfeiffer, M. Pierini, M. Pimiä, D. Piparo, G. Polese, L. Quertenmont, A. Racz, W. Reece, J. Rodrigues Antunes, G. Rolandi³³, T. Rommerskirchen, C. Rovelli³⁴, M. Rovere, H. Sakulin, F. Santanastasio, C. Schäfer, C. Schwick, I. Segoni, S. Sekmen, A. Sharma, P. Siegrist, P. Silva, M. Simon, P. Sphicas³⁵, D. Spiga, M. Spiropulu⁴, M. Stoye, A. Tsiros, G.I. Veres¹⁷, J.R. Vlimant, H.K. Wöhri, S.D. Worm³⁶, W.D. Zeuner

Paul Scherrer Institut, Villigen, Switzerland

W. Bertl, K. Deiters, W. Erdmann, K. Gabathuler, R. Horisberger, Q. Ingram, H.C. Kaestli, S. König, D. Kotlinski, U. Langenegger, F. Meier, D. Renker, T. Rohe, J. Sibille³⁷

Institute for Particle Physics, ETH Zurich, Zurich, Switzerland

L. Bäni, P. Bortignon, M.A. Buchmann, B. Casal, N. Chanon, Z. Chen, A. Deisher, G. Dissertori, M. Dittmar, M. Dünser, J. Eugster, K. Freudenreich, C. Grab, P. Lecomte, W. Luster, M. A.C. Marini, P. Martinez Ruiz del Arbol, N. Mohr, F. Moortgat, C. Nägeli³⁸, P. Nef, F. Nessi-Tedaldi, L. Pape, F. Pauss, M. Peruzzi, F.J. Ronga, M. Rossini, L. Sala, A.K. Sanchez, A. Starodumov³⁹, B. Stieger, M. Takahashi, L. Tauscher[†], A. Thea, K. Theofilatos, D. Treille, C. Urscheler, R. Wallny, H.A. Weber, L. Wehrli

Universität Zürich, Zurich, Switzerland

E. Aguilo, C. AMSLER, V. Chiochia, S. De Visscher, C. Favaro, M. Ivova Rikova, B. Millan Mejias, P. Otiougova, P. Robmann, H. Snoek, S. Tupputi, M. Verzetti

National Central University, Chung-Li, Taiwan

Y.H. Chang, K.H. Chen, A. Go, C.M. Kuo, S.W. Li, W. Lin, Z.K. Liu, Y.J. Lu, D. Mekterovic, A.P. Singh, R. Volpe, S.S. Yu

National Taiwan University (NTU), Taipei, Taiwan

P. Bartalini, P. Chang, Y.H. Chang, Y.W. Chang, Y. Chao, K.F. Chen, C. Dietz, U. Grundler, W.-S. Hou, Y. Hsiung, K.Y. Kao, Y.J. Lei, R.-S. Lu, D. Majumder, E. Petrakou, X. Shi, J.G. Shiu, Y.M. Tzeng, M. Wang

Cukurova University, Adana, Turkey

A. Adiguzel, M.N. Bakirci⁴⁰, S. Cerci⁴¹, C. Dozen, I. Dumanoglu, E. Eskut, S. Girgis, G. Gokbulut, I. Hos, E.E. Kangal, G. Karapinar, A. Kayis Topaksu, G. Onengut, K. Ozdemir, S. Ozturk⁴², A. Polatoz, K. Sogut⁴³, D. Sunar Cerci⁴¹, B. Tali⁴¹, H. Topakli⁴⁰, L.N. Vergili, M. Vergili

Middle East Technical University, Physics Department, Ankara, Turkey

I.V. Akin, T. Aliev, B. Bilin, S. Bilmis, M. Deniz, H. Gamsizkan, A.M. Guler, K. Ocalan, A. Ozpineci, M. Serin, R. Sever, U.E. Surat, M. Yalvac, E. Yildirim, M. Zeyrek

Bogazici University, Istanbul, Turkey

M. Deliomeroğlu, E. Gülmez, B. Isildak, M. Kaya⁴⁴, O. Kaya⁴⁴, S. Ozkorucuklu⁴⁵, N. Sonmez⁴⁶

Istanbul Technical University, Istanbul, Turkey

K. Cankocak

National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine

L. Levchuk

University of Bristol, Bristol, United Kingdom

F. Bostock, J.J. Brooke, E. Clement, D. Cussans, H. Flacher, R. Frazier, J. Goldstein, M. Grimes, G.P. Heath, H.F. Heath, L. Kreczko, S. Metson, D.M. Newbold³⁶, K. Nirunpong, A. Poll, S. Senkin, V.J. Smith, T. Williams

Rutherford Appleton Laboratory, Didcot, United Kingdom

L. Basso⁴⁷, K.W. Bell, A. Belyaev⁴⁷, C. Brew, R.M. Brown, D.J.A. Cockerill, J.A. Coughlan, K. Harder, S. Harper, J. Jackson, B.W. Kennedy, E. Olaiya, D. Petyt, B.C. Radburn-Smith, C.H. Shepherd-Themistocleous, I.R. Tomalin, W.J. Womersley

Imperial College, London, United Kingdom

R. Bainbridge, G. Ball, R. Beuselinck, O. Buchmuller, D. Colling, N. Cripps, M. Cutajar, P. Dauncey, G. Davies, M. Della Negra, W. Ferguson, J. Fulcher, D. Futyan, A. Gilbert, A. Guneratne Bryer, G. Hall, Z. Hatherell, J. Hays, G. Iles, M. Jarvis, G. Karapostoli, L. Lyons, A.-M. Magnan, J. Marrouche, B. Mathias, R. Nandi, J. Nash, A. Nikitenko³⁹, A. Papageorgiou, J. Pela¹, M. Pesaresi, K. Petridis, M. Pioppi⁴⁸, D.M. Raymond, S. Rogerson, N. Rompotis, A. Rose, M.J. Ryan, C. Seez, P. Sharp[†], A. Sparrow, A. Tapper, M. Vazquez Acosta, T. Virdee, S. Wakefield, N. Wardle, T. Whyntie

Brunel University, Uxbridge, United Kingdom

M. Barrett, M. Chadwick, J.E. Cole, P.R. Hobson, A. Khan, P. Kyberd, D. Leggat, D. Leslie, W. Martin, I.D. Reid, P. Symonds, L. Teodorescu, M. Turner

Baylor University, Waco, USA

K. Hatakeyama, H. Liu, T. Scarborough

The University of Alabama, Tuscaloosa, USA

C. Henderson, P. Rumerio

Boston University, Boston, USA

A. Avetisyan, T. Bose, C. Fantasia, A. Heister, J. St. John, P. Lawson, D. Lazic, J. Rohlf, D. Sperka, L. Sulak

Brown University, Providence, USA

J. Alimena, S. Bhattacharya, D. Cutts, A. Ferapontov, U. Heintz, S. Jabeen, G. Kukartsev, G. Landsberg, M. Luk, M. Narain, D. Nguyen, M. Segala, T. Sinthuprasith, T. Speer, K.V. Tsang

University of California, Davis, Davis, USA

R. Breedon, G. Breto, M. Calderon De La Barca Sanchez, S. Chauhan, M. Chertok, J. Conway, R. Conway, P.T. Cox, J. Dolen, R. Erbacher, M. Gardner, R. Houtz, W. Ko, A. Kopecky, R. Lander, O. Mall, T. Miceli, R. Nelson, D. Pellett, B. Rutherford, M. Searle, J. Smith, M. Squires, M. Tripathi, R. Vasquez Sierra

University of California, Los Angeles, Los Angeles, USA

V. Andreev, D. Cline, R. Cousins, J. Duris, S. Erhan, P. Everaerts, C. Farrell, J. Hauser, M. Ignatenko, C. Plager, G. Rakness, P. Schlein[†], J. Tucker, V. Valuev, M. Weber

University of California, Riverside, Riverside, USA

J. Babb, R. Clare, M.E. Dinardo, J. Ellison, J.W. Gary, F. Giordano, G. Hanson, G.Y. Jeng⁴⁹, H. Liu, O.R. Long, A. Luthra, H. Nguyen, S. Paramesvaran, J. Sturdy, S. Sumowidagdo, R. Wilken, S. Wimpenny

University of California, San Diego, La Jolla, USA

W. Andrews, J.G. Branson, G.B. Cerati, S. Cittolin, D. Evans, F. Golf, A. Holzner, R. Kelley, M. Lebourgeois, J. Letts, I. Macneill, B. Mangano, J. Muelmenstaedt, S. Padhi, C. Palmer, G. Petrucciani, M. Pieri, R. Ranieri, M. Sani, V. Sharma, S. Simon, E. Sudano, M. Tadel, Y. Tu, A. Vartak, S. Wasserbaech⁵⁰, F. Würthwein, A. Yagil, J. Yoo

University of California, Santa Barbara, Santa Barbara, USA

D. Barge, R. Bellan, C. Campagnari, M. D'Alfonso, T. Danielson, K. Flowers, P. Geffert, J. Incandela, C. Justus, P. Kalavase, S.A. Koay, D. Kovalskyi¹, V. Krutelyov, S. Lowette, N. Mccoll, V. Pavlunin, F. Rebassoo, J. Ribnik, J. Richman, R. Rossin, D. Stuart, W. To, C. West

California Institute of Technology, Pasadena, USA

A. Apresyan, A. Bornheim, Y. Chen, E. Di Marco, J. Duarte, M. Gataullin, Y. Ma, A. Mott, H.B. Newman, C. Rogan, V. Timciuc, P. Traczyk, J. Veverka, R. Wilkinson, Y. Yang, R.Y. Zhu

Carnegie Mellon University, Pittsburgh, USA

B. Akgun, R. Carroll, T. Ferguson, Y. Iiyama, D.W. Jang, Y.F. Liu, M. Paulini, H. Vogel, I. Vorobiev

University of Colorado at Boulder, Boulder, USA

J.P. Cumalat, B.R. Drell, C.J. Edelmaier, W.T. Ford, A. Gaz, B. Heyburn, E. Luiggi Lopez, J.G. Smith, K. Stenson, K.A. Ulmer, S.R. Wagner

Cornell University, Ithaca, USA

L. Agostino, J. Alexander, A. Chatterjee, N. Eggert, L.K. Gibbons, B. Heltsley, W. Hopkins, A. Khukhunaishvili, B. Kreis, N. Mirman, G. Nicolas Kaufman, J.R. Patterson, A. Ryd, E. Salvati, W. Sun, W.D. Teo, J. Thom, J. Thompson, J. Vaughan, Y. Weng, L. Winstrom, P. Wittich

Fairfield University, Fairfield, USA

D. Winn

Fermi National Accelerator Laboratory, Batavia, USA

S. Abdullin, M. Albrow, J. Anderson, L.A.T. Bauerdick, A. Beretvas, J. Berryhill, P.C. Bhat, I. Bloch, K. Burkett, J.N. Butler, V. Chetluru, H.W.K. Cheung, F. Chlebana, V.D. Elvira, I. Fisk, J. Freeman, Y. Gao, D. Green, O. Gutsche, A. Hahn, J. Hanlon, R.M. Harris, J. Hirschauer, B. Hooberman, S. Jindariani, M. Johnson, U. Joshi, B. Kilminster, B. Klima, S. Kunori, S. Kwan, C. Leonidopoulos, D. Lincoln, R. Lipton, L. Lueking, J. Lykken, K. Maeshima, J.M. Marraffino, S. Maruyama, D. Mason, P. McBride, K. Mishra, S. Mrenna, Y. Musienko⁵¹, C. Newman-Holmes, V. O'Dell, O. Prokofyev, E. Sexton-Kennedy, S. Sharma, W.J. Spalding, L. Spiegel, P. Tan, L. Taylor, S. Tkaczyk, N.V. Tran, L. Uplegger, E.W. Vaandering, R. Vidal, J. Whitmore, W. Wu, F. Yang, F. Yumiceva, J.C. Yun

University of Florida, Gainesville, USA

D. Acosta, P. Avery, D. Bourilkov, M. Chen, S. Das, M. De Gruttola, G.P. Di Giovanni, D. Dobur, A. Drozdetskiy, R.D. Field, M. Fisher, Y. Fu, I.K. Furic, J. Gartner, J. Hugon, B. Kim, J. Konigsberg, A. Korytov, A. Kropivnitskaya, T. Kypreos, J.F. Low, K. Matchev, P. Milenovic⁵², G. Mitselmakher, L. Muniz, R. Remington, A. Rinkevicius, P. Sellers, N. Skhirtladze, M. Snowball, J. Yelton, M. Zakaria

Florida International University, Miami, USA

V. Gaultney, L.M. Lebolo, S. Linn, P. Markowitz, G. Martinez, J.L. Rodriguez

Florida State University, Tallahassee, USA

T. Adams, A. Askew, J. Bochenek, J. Chen, B. Diamond, S.V. Gleyzer, J. Haas, S. Hagopian, V. Hagopian, M. Jenkins, K.F. Johnson, H. Prosper, V. Veeraraghavan, M. Weinberg

Florida Institute of Technology, Melbourne, USA

M.M. Baarmand, B. Dorney, M. Hohlmann, H. Kalakhety, I. Vodopyanov

University of Illinois at Chicago (UIC), Chicago, USA

M.R. Adams, I.M. Anghel, L. Apanasevich, Y. Bai, V.E. Bazterra, R.R. Betts, J. Callner, R. Cavanaugh, C. Dragoiu, O. Evdokimov, E.J. Garcia-Solis, L. Gauthier, C.E. Gerber, D.J. Hofman, S. Khalatyan, F. Lacroix, M. Malek, C. O'Brien, C. Silkworth, D. Strom, N. Varelas

The University of Iowa, Iowa City, USA

U. Akgun, E.A. Albayrak, B. Bilki⁵³, K. Chung, W. Clarida, F. Duru, S. Griffiths, C.K. Lae, J.-P. Merlo, H. Mermerkaya⁵⁴, A. Mestvirishvili, A. Moeller, J. Nachtman, C.R. Newsom, E. Norbeck, J. Olson, Y. Onel, F. Ozok, S. Sen, E. Tiras, J. Wetzel, T. Yetkin, K. Yi

Johns Hopkins University, Baltimore, USA

B.A. Barnett, B. Blumenfeld, S. Bolognesi, D. Fehling, G. Giurgiu, A.V. Gritsan, Z.J. Guo, G. Hu, P. Maksimovic, S. Rappoccio, M. Swartz, A. Whitbeck

The University of Kansas, Lawrence, USA

P. Baringer, A. Bean, G. Benelli, O. Grachov, R.P. Kenny Iii, M. Murray, D. Noonan, V. Radicci, S. Sanders, R. Stringer, G. Tinti, J.S. Wood, V. Zhukova

Kansas State University, Manhattan, USA

A.F. Barfuss, T. Bolton, I. Chakaberia, A. Ivanov, S. Khalil, M. Makouski, Y. Maravin, S. Shrestha, I. Svintradze

Lawrence Livermore National Laboratory, Livermore, USA

J. Gronberg, D. Lange, D. Wright

University of Maryland, College Park, USA

A. Baden, M. Boutemour, B. Calvert, S.C. Eno, J.A. Gomez, N.J. Hadley, R.G. Kellogg, M. Kirn, T. Kolberg, Y. Lu, M. Marionneau, A.C. Mignerey, A. Peterman, K. Rossato, A. Skuja, J. Temple, M.B. Tonjes, S.C. Tonwar, E. Twedt

Massachusetts Institute of Technology, Cambridge, USA

G. Bauer, J. Bendavid, W. Busza, E. Butz, I.A. Cali, M. Chan, V. Dutta, G. Gomez Ceballos, M. Goncharov, K.A. Hahn, Y. Kim, M. Klute, Y.-J. Lee, W. Li, P.D. Luckey, T. Ma, S. Nahn, C. Paus, D. Ralph, C. Roland, G. Roland, M. Rudolph, G.S.F. Stephans, F. Stöckli, K. Sumorok, K. Sung, D. Velicanu, E.A. Wenger, R. Wolf, B. Wyslouch, S. Xie, M. Yang, Y. Yilmaz, A.S. Yoon, M. Zanetti

University of Minnesota, Minneapolis, USA

S.I. Cooper, P. Cushman, B. Dahmes, A. De Benedetti, G. Franzoni, A. Gude, J. Haupt, S.C. Kao, K. Klapoetke, Y. Kubota, J. Mans, N. Pastika, R. Rusack, M. Sasseville, A. Singovsky, N. Tambe, J. Turkewitz

University of Mississippi, University, USA

L.M. Cremaldi, R. Kroeger, L. Perera, R. Rahmat, D.A. Sanders

University of Nebraska-Lincoln, Lincoln, USA

E. Avdeeva, K. Bloom, S. Bose, J. Butt, D.R. Claes, A. Dominguez, M. Eads, P. Jindal, J. Keller, I. Kravchenko, J. Lazo-Flores, H. Malbouisson, S. Malik, G.R. Snow

State University of New York at Buffalo, Buffalo, USA

U. Baur, A. Godshalk, I. Iashvili, S. Jain, A. Kharchilava, A. Kumar, S.P. Shipkowski, K. Smith

Northeastern University, Boston, USA

G. Alverson, E. Barberis, D. Baumgartel, M. Chasco, J. Haley, D. Trocino, D. Wood, J. Zhang

Northwestern University, Evanston, USA

A. Anastassov, A. Kubik, N. Mucia, N. Odell, R.A. Ofierzynski, B. Pollack, A. Pozdnyakov, M. Schmitt, S. Stoynev, M. Velasco, S. Won

University of Notre Dame, Notre Dame, USA

L. Antonelli, D. Berry, A. Brinkerhoff, M. Hildreth, C. Jessop, D.J. Karmgard, J. Kolb, K. Lannon, W. Luo, S. Lynch, N. Marinelli, D.M. Morse, T. Pearson, R. Ruchti, J. Slaunwhite, N. Valls, J. Warchol, M. Wayne, M. Wolf, J. Ziegler

The Ohio State University, Columbus, USA

B. Bylsma, L.S. Durkin, C. Hill, R. Hughes, P. Killewald, K. Kotov, T.Y. Ling, D. Puigh, M. Rodenburg, C. Vuosalo, G. Williams, B.L. Winer

Princeton University, Princeton, USA

N. Adam, E. Berry, P. Elmer, D. Gerbaudo, V. Halyo, P. Hebda, J. Hegeman, A. Hunt, E. Laird, D. Lopes Pegna, P. Lujan, D. Marlow, T. Medvedeva, M. Mooney, J. Olsen, P. Piroué, X. Quan, A. Raval, H. Saka, D. Stickland, C. Tully, J.S. Werner, A. Zuranski

University of Puerto Rico, Mayaguez, USA

J.G. Acosta, E. Brownson, X.T. Huang, A. Lopez, H. Mendez, S. Oliveros, J.E. Ramirez Vargas, A. Zatserklyaniy

Purdue University, West Lafayette, USA

E. Alagoz, V.E. Barnes, D. Benedetti, G. Bolla, D. Bortoletto, M. De Mattia, A. Everett, Z. Hu, M. Jones, O. Koybasi, M. Kress, A.T. Laasanen, N. Leonardo, V. Maroussov, P. Merkel,

D.H. Miller, N. Neumeister, I. Shipsey, D. Silvers, A. Svyatkovskiy, M. Vidal Marono, H.D. Yoo, J. Zablocki, Y. Zheng

Purdue University Calumet, Hammond, USA

S. Guragain, N. Parashar

Rice University, Houston, USA

A. Adair, C. Boulahouache, V. Cuplov, K.M. Ecklund, F.J.M. Geurts, B.P. Padley, R. Redjimi, J. Roberts, J. Zabel

University of Rochester, Rochester, USA

B. Betchart, A. Bodek, Y.S. Chung, R. Covarelli, P. de Barbaro, R. Demina, Y. Eshaq, A. Garcia-Bellido, P. Goldenzweig, Y. Gotra, J. Han, A. Harel, S. Korjenevski, D.C. Miner, D. Vishnevskiy, M. Zielinski

The Rockefeller University, New York, USA

A. Bhatti, R. Ciesielski, L. Demortier, K. Goulios, G. Lungu, S. Malik, C. Mesropian

Rutgers, the State University of New Jersey, Piscataway, USA

S. Arora, A. Barker, J.P. Chou, C. Contreras-Campana, E. Contreras-Campana, D. Duggan, D. Ferencek, Y. Gershtein, R. Gray, E. Halkiadakis, D. Hidas, D. Hits, C. Kilic⁵⁵, A. Lath, S. Panwalkar, M. Park, R. Patel, V. Rekovic, A. Richards, J. Robles, K. Rose, S. Salur, S. Schnetzer, C. Seitz, S. Somalwar, R. Stone, S. Thomas

University of Tennessee, Knoxville, USA

G. Cerizza, M. Hollingsworth, S. Spanier, Z.C. Yang, A. York

Texas A&M University, College Station, USA

R. Eusebi, W. Flanagan, J. Gilmore, T. Kamon⁵⁶, V. Khotilovich, R. Montalvo, I. Osipenkov, Y. Pakhotin, A. Perloff, J. Roe, A. Safonov, T. Sakuma, S. Sengupta, I. Suarez, A. Tatarinov, D. Toback

Texas Tech University, Lubbock, USA

N. Akchurin, J. Damgov, P.R. Duderod, C. Jeong, K. Kovitanggoon, S.W. Lee, T. Libeiro, Y. Roh, I. Volobouev

Vanderbilt University, Nashville, USA

E. Appelt, D. Engh, C. Florez, S. Greene, A. Gurrola, W. Johns, P. Kurt, C. Maguire, A. Melo, P. Sheldon, B. Snook, S. Tuo, J. Velkovska

University of Virginia, Charlottesville, USA

M.W. Arenton, M. Balazs, S. Boutle, B. Cox, B. Francis, J. Goodell, R. Hirosky, A. Ledovskoy, C. Lin, C. Neu, J. Wood, R. Yohay

Wayne State University, Detroit, USA

S. Gollapinni, R. Harr, P.E. Karchin, C. Kottachchi Kankanamge Don, P. Lamichhane, A. Sakharov

University of Wisconsin, Madison, USA

M. Anderson, M. Bachtis, D. Belknap, L. Borrello, D. Carlsmith, M. Cepeda, S. Dasu, L. Gray, K.S. Grogg, M. Grothe, R. Hall-Wilton, M. Herndon, A. Hervé, P. Klabbers, J. Klukas, A. Lanaro, C. Lazaridis, J. Leonard, R. Loveless, A. Mohapatra, I. Ojalvo, G.A. Pierro, I. Ross, A. Savin, W.H. Smith, J. Swanson

†: Deceased

1: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

- 2: Also at National Institute of Chemical Physics and Biophysics, Tallinn, Estonia
- 3: Also at Universidade Federal do ABC, Santo Andre, Brazil
- 4: Also at California Institute of Technology, Pasadena, USA
- 5: Also at Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France
- 6: Also at Suez Canal University, Suez, Egypt
- 7: Also at Cairo University, Cairo, Egypt
- 8: Also at British University, Cairo, Egypt
- 9: Also at Fayoum University, El-Fayoum, Egypt
- 10: Now at Ain Shams University, Cairo, Egypt
- 11: Also at Soltan Institute for Nuclear Studies, Warsaw, Poland
- 12: Also at Université de Haute-Alsace, Mulhouse, France
- 13: Now at Joint Institute for Nuclear Research, Dubna, Russia
- 14: Also at Moscow State University, Moscow, Russia
- 15: Also at Brandenburg University of Technology, Cottbus, Germany
- 16: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- 17: Also at Eötvös Loránd University, Budapest, Hungary
- 18: Also at Tata Institute of Fundamental Research - HECR, Mumbai, India
- 19: Now at King Abdulaziz University, Jeddah, Saudi Arabia
- 20: Also at University of Visva-Bharati, Santiniketan, India
- 21: Also at Sharif University of Technology, Tehran, Iran
- 22: Also at Isfahan University of Technology, Isfahan, Iran
- 23: Also at Shiraz University, Shiraz, Iran
- 24: Also at Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Teheran, Iran
- 25: Also at Facoltà Ingegneria Università di Roma, Roma, Italy
- 26: Also at Università della Basilicata, Potenza, Italy
- 27: Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- 28: Also at Università degli studi di Siena, Siena, Italy
- 29: Also at University of Bucharest, Faculty of Physics, Bucuresti-Magurele, Romania
- 30: Also at Faculty of Physics of University of Belgrade, Belgrade, Serbia
- 31: Also at University of Florida, Gainesville, USA
- 32: Also at University of California, Los Angeles, Los Angeles, USA
- 33: Also at Scuola Normale e Sezione dell' INFN, Pisa, Italy
- 34: Also at INFN Sezione di Roma; Università di Roma "La Sapienza", Roma, Italy
- 35: Also at University of Athens, Athens, Greece
- 36: Also at Rutherford Appleton Laboratory, Didcot, United Kingdom
- 37: Also at The University of Kansas, Lawrence, USA
- 38: Also at Paul Scherrer Institut, Villigen, Switzerland
- 39: Also at Institute for Theoretical and Experimental Physics, Moscow, Russia
- 40: Also at Gaziosmanpasa University, Tokat, Turkey
- 41: Also at Adiyaman University, Adiyaman, Turkey
- 42: Also at The University of Iowa, Iowa City, USA
- 43: Also at Mersin University, Mersin, Turkey
- 44: Also at Kafkas University, Kars, Turkey
- 45: Also at Suleyman Demirel University, Isparta, Turkey
- 46: Also at Ege University, Izmir, Turkey
- 47: Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- 48: Also at INFN Sezione di Perugia; Università di Perugia, Perugia, Italy

49: Also at University of Sydney, Sydney, Australia

50: Also at Utah Valley University, Orem, USA

51: Also at Institute for Nuclear Research, Moscow, Russia

52: Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia

53: Also at Argonne National Laboratory, Argonne, USA

54: Also at Erzincan University, Erzincan, Turkey

55: Now at University of Texas at Austin, Austin, USA

56: Also at Kyungpook National University, Daegu, Korea