

Russell et al

Egocentric perspective in episodic memory

Self-perspective in episodic memory after parietal damage and in healthy ageing

Highlights:

- Damage to right parietal cortex leads to a specific deficit in episodic memory
- This deficit is in recognition of the perspective from which a memory was encoded
- MVPA in neuroimaging shows parietal involvement in recalling from own perspective
- Supramarginal and angular gyri regions crucial for these aspects of episodic memory

1
2
3 **Self-perspective in episodic memory after parietal damage and in healthy ageing**
4
5

6
7 **Charlotte Russell^{1*}, Sarah Davies², Korina Li³, Anna-Sofia Musil¹,**
8

9
10 **Pareesh A. Malhotra^{3**} & Adrian L. Williams^{2**}**
11
12

13
14
15 1. Department of Psychology, Institute of Psychiatry, Psychology and Neuroscience,
16
17 King's College London, SE1 1UL
18

19
20 2. Psychology Division, Department of Life Sciences, Brunel University London, UB8 3PH
21

22
23 3. Division of Brain Sciences Department of Medicine, Imperial College London, W6 8RP
24
25

26
27
28 * corresponding author
29

30
31 ** denotes equal contribution
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

1
2
3
4 Although there is strong support from functional imaging studies for lateral parietal lobe
5 involvement in episodic memory, patients with damage to these regions do not appear to
6 suffer from severe deficits in this cognitive domain. As such there has been no definitive
7 explanation of this area's precise involvement. Here, we hypothesised that parietal regions
8 play a crucial role in episodic memory - specifically in recollecting details from an egocentric
9 perspective. In order to test this hypothesis systematically, we designed a novel
10 experimental task utilising a head-mounted camera to record images from the participant's
11 perspective, enabling us to evaluate the integrity of memory from the individual's own point
12 of view. In the first study we examined patients with parietal damage and in a second
13 study, using fMRI, we examined young and older healthy participants. Right-hemisphere
14 patients with parietal damage were able to recall information accurately when recollecting
15 what items had been present and where these items had been. However, patients were
16 significantly impaired when attempting to judge from which perspective they had viewed
17 the scenes. Critically, the patient group showed no evidence of impairment on standard
18 tests of episodic and working memory. Examination of healthy participants in the second
19 study utilised multi-voxel pattern analysis on neural activity during the recognition phase of
20 a similar task. This revealed sensitivity to be highest around the angular gyrus of the lateral
21 parietal cortex for our critical comparison - that is, when viewing stimuli that were the same
22 as their egocentric view during encoding versus the identical scene but presented from an
23 alternative angle. Our results provide important evidence that parietal cortex is directly
24 involved in egocentric spatial perspective aspects of episodic memory and demonstrate for
25 the first time a specific deficit in episodic memory in patients with right parietal damage.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 **Key words:** Episodic memory; parietal cortex, neuropsychology, ageing
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

1
2
3 Patients with damage confined to lateral parietal cortex are not thought to suffer from
4
5 episodic memory problems. Right hemisphere lesions affecting this region often result in
6
7 the striking attentional deficits that form the core symptoms of spatial neglect (e.g., Mort *et*
8
9 *al.*, 2003; Corbetta and Shulman, 2011), with left hemisphere damage leading to other
10
11 deficits including those considered part of Gerstmann's syndrome or aphasias (e.g.,
12
13 Fridriksson *et al.*, 2010; Rusconi *et al.*, 2010). There has been a surge of interest in the
14
15 apparent contradiction between activity consistently seen in lateral parietal cortex during
16
17 functional neuroimaging studies of episodic memory (Cabeza *et al.*, 2008; Schooler *et al.*, 2011
18
19 and see Rugg & King, 2017; Sestieri *et al.*, 2017 for recent reviews) and the assumption,
20
21 followed by accruing evidence, of preserved episodic recall skills in patients with damage to
22
23 this region (Ally *et al.*, 2008; Simons *et al.*, 2008). This contrasts with the clear relationship
24
25 between neural activity in medial temporal regions elicited during episodic recall and the
26
27 debilitating long-term memory loss associated with damage to those regions (Scoville and
28
29 Milner, 1957; Wagner *et al.*, 1998; Squire *et al.*, 2004; Rugg and Vilberg, 2013).
30
31
32
33
34
35
36
37
38
39
40

41 Despite the fact that lateral parietal cortex lesions do not reduce recall accuracy in standard
42
43 episodic memory tasks, careful investigation by a number of groups has delineated subtle
44
45 impairments. For example, even with preserved overall recall accuracy, patients with
46
47 bilateral parietal lesions show an atypical relationship between self-rated confidence in
48
49 their performance and their actual performance. In some paradigms they rate themselves
50
51 less confident than controls despite accurate performance, whereas in other paradigms
52
53 they rate themselves as confident despite failures to recognise previously presented items
54
55 (e.g., Berryhill *et al.*, 2009; Olson and Berryhill, 2009; Simons *et al.*, 2010; Hower *et al.*,
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2014). Perhaps relating to this altered confidence, patients make fewer false alarms than healthy controls and are less susceptible to falsely recalling semantically related lures (Drowos *et al.*, 2010). In autobiographical tasks where patients are asked to freely recall an event from their own life, patients with parietal damage can recall personal memories but description of these events is independently rated as vague and imprecise, suggestive of impairments in vividly recalling the event (Berryhill *et al.*, 2007; Davidson *et al.*, 2008; Berryhill, 2012). Consistent with this suggestion is evidence that individuals with parietal damage are less likely to rate the responses given in a recall task as explicit 'Remember' versus the implicit 'Know' (e.g., Drowos *et al.*, 2010), as are reports from patients that even when accurately recalling an autobiographical event they do not have the subjective feeling of having experienced the memory themselves (see Davidson *et al.*, 2008).

Complementary to this neuropsychological work a number of groups have examined disruption to relevant parts of posterior parietal cortex in healthy participants, focusing on the left angular gyrus. Repetitive TMS (Sestieri *et al.*, 2013) and continuous theta burst stimulation (Yazar *et al.* 2014) has suggested that disrupting this region affects the subjective experience of remembering without loss of overall accuracy of recall. Further, using functional imaging Bonnici *et al.* (2016) demonstrated that classification accuracy in multi-voxel pattern analysis increased within angular gyrus as participants reported greater vividness of recall.

The nature of these neuropsychological impairments and the results of experimental disruption make it pertinent to consider well-documented functions of parietal cortex. Computational and neuroimaging evidence suggests a key role for parietal cortex in

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

egocentric spatial representation of visual input (see Burgess *et al.*, 2001; Burgess, 2008; Lambrey *et al.*, 2012). In order to create an accurate image of the world around us- from our own first-person perspective - we rely on parietal cortex. Evidence of impaired route learning when only egocentric (as opposed to allocentric) cues are available in parietal patients further supports this (Weniger *et al.*, 2009). As does evidence that deficits in visuo-spatial awareness commonly seen after right hemisphere parietal damage predominately affect egocentric rather than allocentric space (Rorden *et al.*, 2012). Consideration of the features of episodic memory highlights the potentially critical role for egocentric spatial recall in this function. If we are asked to remember the event of having dinner last night, we would not feel that we were accurately recalling our memory if we did not construct a visuospatial mental image of the scene as we perceived it at the time. Accurate recall of egocentric perspective appears critical to episodic recall. Several studies have shown that as memory declines in ageing there is increased recall from a third person 'observer' perspective and that this is associated with a lack of vivid recalling of autobiographical events (Piolino *et al.*, 2006; Piolino *et al.*, 2009). Ciaramelli and colleagues directly examined topographical memory for route planning tasks in parietal patients and their data revealed more severe impairments on the tasks loading on egocentric representations as compared to allocentric representations (Ciaramelli *et al.*, 2010). In a recent study Ciaramelli and colleagues (2017) in an assessment on patients with parietal damage, confirmed that although memory for word stimuli were accurately recalled, patients (compared to healthy controls) were less likely to select 'Remember' as opposed to 'Know', suggestive of a changed subjective experience of the recalled memory after this damage.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Related to this, St Jacques and colleagues have examined, in healthy individuals shifting the perspective from which they encoded an event at retrieval (St Jacques et al, 2018; St Jacques, Szpunar & Schacter, 2017; Marcotti & St Jacques, 2018). These studies have revealed that posterior parietal cortex appears intrinsically involved in this shift and that the requirement to shift visual perspective at recall reduces the overall accuracy of the memory and – crucially- this effect leads to a reduction in reported subjective vividness of the memory.

In the studies presented here we aim to further probe the functions of parietal cortex – in particular the angular gyrus and nearby regions –in relation to recalling novel scenes from a self – *egocentric*- perspective. Within our design we are able to directly compare memory for egocentric perspective with memory for allocentric spatial relationships. This allows us to examine differences in recall between these conditions in patients with damage to these parietal regions and, in a second study, in older versus younger participants. Our intention to additionally examine the effect of ageing on memory for egocentric perspective was motivated by the seeming harmony between descriptions of memory change in parietal patients with those in the ageing literature – albeit more dramatic in the patient groups. For example, in older people episodic recall is also frequently reported as lacking detail and specificity (e.g, Levine et al, 2002; Addis et al, 2011; Schacter et al, 2013). Further, the evidence for an increase in memories reported from an ‘observer’s viewpoint rather than ‘field’ perspective harmonises with the processes we are directly probing with our paradigm (see, Piolino et al, 2006; 2009). Additionally, the second study was a direct attempt to address the apparent contradiction of parietal involvement in episodic memory in functional neuroimaging despite there being only subtle changes to these processes in patients with

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

damage here. Using fMRI based multi-voxel pattern analysis (MVPA) we will decode which areas of parietal cortex differentiate between self-perspective of a previously presented scene compared to the same scene from another perspective. This analysis enables us to examine whether specific patterns of activity can predict whether the participant is currently viewing their own view of the scene or an alternative perspective. Note that this analysis differs from univariate analysis in which we might predict parietal activity during recollection of a scene but without this activity necessarily discriminating whether they are viewing their own view or that of another. Using MVPA analysis with healthy participants we aim to confirm that parietal cortex is involved in the aspects of episodic memory that we hypothesise to be affected by acquired damage to these regions.

In summary, in our first study we were interested in two things. First, whether right hemisphere damage to parietal cortex can result in deficits in episodic memory. Second, whether these deficits are specifically related to memory for egocentric self-perspective. In Experiment 2 we adapt our neuropsychological paradigm to examine changes in this aspect of episodic memory with age and to directly examine whether areas of interest within parietal cortex are involved in differentiating our own self-perspective of an episodic memory.

2. Experiment 1

2. 1 Materials and Methods

2.1.1 Participants

Six patients (3 females) with chronic lesions affecting parietal cortex were recruited. All had been patients on the stroke unit at Imperial College Healthcare NHS Trust. All previously suffered from spatial neglect, which had recovered at the time of testing – confirmed at the start of the session with the Behavioural Inattention Test Star Cancellation task (Wilson *et al.*, 1987). They were aged 66 to 79 (mean = 71.2 years) and had a right hemisphere stroke more than 6 months previously. They were compared to a group of eight healthy age-matched participants (6 females), aged 64 to 83 (mean = 71.6 years). Inclusion criteria for patients were: a lesion affecting right parietal cortex; no current visuo-spatial neglect or extinction; no reports of memory problems; no other neurological impairment; no diagnosis of psychiatric impairment. Individuals were excluded if these criteria were not fulfilled. Healthy participants confirmed they had no current diagnosis of neurological or psychiatric illness and no memory problems. All participants also completed the mini-mental state exam (MMSE), no individual scored below 27/30 (patient group mean = 29.4; healthy group mean = 29.9), anyone scoring below 27 was excluded. All participants gave written consent according to the Declaration of Helsinki, with the study having been approved by the local research ethics committee (Figure 1).




Figure 1

Individual Lesions and Lesion Overlap

Four of the six patients' lesions were mapped directly onto patients' native MRI scans using MRIcron software (www.mccauslandcenter.sc.edu/mricro/mricron). The anatomical scan and lesions were subsequently mapped onto stereotaxic space using Clinical Toolbox (www.mccauslandcenter.sc.edu/CRNL/clinical-toolbox) for spatial normalisation, implemented via SPM8 software (www.fil.ion.ucl.ac.uk/spm/software/spm8). The remaining two lesions were plotted using onto a T1 weighted template in MRIcron software (www.mccauslandcenter.sc.edu/crnl/mricron/) by direct comparison with the native scan. All scans were mapped by trained neurologists (KL and PM).

A. Lesions mapped for each of 6 patients individually

B. Lesion overlap. Lesions are represented in purple, with regions damaged in increasing numbers of patients shown in brighter shades. All patients had strokes affecting the middle cerebral artery territory, with varying degrees of frontal and temporal involvement. The

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

single region of maximal overlap (all 6 patients affected) was in the right supramarginal gyrus, centred around MNI coordinates (46, -33, 26) extending laterally to (59, -29, 26) and superiorly to (46,-33,27).

2.1.2 Standardised Memory tasks

These tasks were used to assess episodic and spatial working memory:

Rivermead Behavioural Memory Test – Line drawing delayed recognition (RBMT-3). Line drawings of common objects were shown for three seconds each. A delayed recognition task followed after 30 minutes; *Corsi Block Tapping Task*. Participants observed a sequence of taps and then repeated this in the identical ('Forward span') or reverse order ('Backward span'). The task starts with a short sequence increasing in length to a maximum of nine or until the participant is incorrect in both trials of a particular length.

2.1.3 Experimental task

Participants were presented with 14 novel 3D scenes to remember. Each scene consisted of two items positioned in separate squares of a 2 x 2 grid pattern, placed on the table in front of the participant (see Figure 2). For half of the scenes participants sat to the left of the grid, for the other half to the right, seating position was randomly allocated across the trials with an equal distribution of both angles – the order in which left and right were used was counterbalanced across participants. Participants were asked to move seats for this manipulation. If they were in a wheelchair, their wheelchair was moved for them. Viewing position, items used and the order of presentation were counterbalanced across participants. Pilot testing with a similar group of patients had confirmed suitability of this number of scenes.




Figure 2

Experimental task stimuli and conditions

Pictures shown in column A are indicative of the original scenes shown to participants. Each scene consisted of two items taken from a total of 14 individual categories. Examples of possible categories included kitchen items, toy vehicles, models of musical instruments, and fruit. There were 8 potential exemplars from each category. No category was used more than once. In each encoding trial, two items from separate categories were placed on the grid in front of participants. Note that during encoding photographs were not used but the actual 3-D image, test images were purported to be from the head camera participants wore during encoding. Seen in this figure are examples from the animal and sports equipment categories. In the recognition task, items such as those shown in column B were used alongside the veridical images of what they had seen as shown in column A. During the 'Item shift' condition shown in the top panel of Figure 2, the lure image was taken from the same perspective but one of the items had moved by one square. In the critical 'Perspective Shift' condition in the lure image, the items occupied the same squares as in original image but the photograph was taken from the opposite angle.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

During encoding, the two items for each scene were placed onto the grid by the experimenter and left for 1 minute. Participants were told that their memory for the scenes and what they looked like to them would be probed later. Each scene was explicitly given a category name, e.g., 'Stationery'. Scenes were presented in two blocks of seven. After each block, participants were cued with the category name of each scene and then asked to recall the items, their positions, and the viewing perspective they experienced that scene from. For example, they could be asked, 'when we showed you the stationery scene can you tell us what items we showed to you, where they were on the grid and where you viewed this scene from?'

All participants were wearing a head camera during encoding. They were shown this camera and it was explained that a still would be taken from each scene, which they would need to identify in the memory test. In reality we pre-prepared stills of each scene – facilitating the later memory test and making the images as clear as possible on the screen. In our pilot studies the head camera images were too unclear to faithfully represent the scene as shown, e.g., similar items like a leopard and tiger were frequently confused. Importantly, no participant in either group questioned that the images were taken from their head camera or mentioned being suspicious during debrief when we explained what had been done. The head camera was used to emphasise that it was how the scenes appeared to the participants that was crucial and not – for example – what objects were in the scene.

Recognition tests took place approximately 2 hours after encoding. Two images were presented on the screen simultaneously. Participants were required to select the image taken from their head camera and that therefore represented the scene that they had seen,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

as they remembered it. One of the images was identical to a pseudo-image from the participants' head camera while viewing one of the scenes, i.e., the same scene (same objects in same grid squares) taken from the same viewing angle as presented to that particular participant. The identity of the other simultaneously presented scenes were split into two experimental categories. Half were 'Item Shift' lures; that is, the same exemplars from the category were used with the same viewing angle but one item had moved squares within the grid. This condition was designed to assess the maintenance of allocentric spatial information from the scenes as one object had changed in its spatial relationship to the other object within the grid. Alternatively, the lure could be a 'Perspective Shift'; that is, the same category exemplars were presented on the same grid squares but the picture was taken from the opposite viewing angle. Note that inherent in this design is that the perceptual demands are similar for the two conditions ('Perspective shift' and 'Item shift') and that simultaneous presentation of the previously seen image and the 'lure' scenes minimised any requirement for mental rotation.

Participants verbally selected which image they believed they had seen (the correct image was randomly presented on the left or right side of the screen). There were 14 trials, one for each unique scene. Participants were given as long as they needed and the images remained on the screen while they decided. They were then asked how confident they were of their decision, rating their confidence on a scale of 1-4 (4 being most confident).

2.2 Results and Discussion

Analysis was carried out using non-parametric statistics as Shapiro-Wilk tests revealed that some data were not normally distributed (see Table 1 and Figure 3).

2.2.1 Standard Memory Tasks

Patients and controls were statistically equivalent ($U = 30, p = .49$) in the line-drawing delayed recognition test. Spatial working memory performance (Corsi Block task) was also equivalent across the groups both for the 'Forward' ($U = 21, p = .76$) and the 'Backward span' conditions ($U = 36, p = .14$).

	Patients	Healthy Controls
<i>Line drawing – memory task</i>		
<i>(proportion)</i>	.97 (.03)	.99 (.02)
<i>Corsi blocks forward span</i>	6.83 (1.72)	6.50 (1.51)
<i>Corsi blocks backwards span</i>	6.17 (1.94)	7.88 (1.64)
Experimental Task		
<i>Item Shift</i>		
<i>Accuracy (proportion)</i>	.83 (.17)	.97 (.07)
<i>Confidence (/4)</i>	3.60 (.27)	3.73 (.46)
<i>Perspective Shift</i>		
<i>Accuracy (proportion)</i>	.53 (.20)	.86 (.15)
<i>Confidence(/4)</i>	3.39 (.50)	3.46 (.61)
Recap Condition in encoding		
session (proportion)		
<i>Item identity</i>	.89 (.08)	
<i>Item position</i>	.88 (.14)	N/A
<i>Viewing angle</i>	.87 (.10)	

Table 1:

Data from all tasks in Experiment 1, standard deviations in brackets.

Additional information on percentage correct recall in the recap questions asked during the encoding

session:

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

'Item Identity' indicates responses to the question: *'Which items did you see in the (e.g.) Sports equipment trial?'*

For *'Item position'* the question was: *'Where were the items shown in (e.g.) sports equipment trial?'*

For the *'Viewing Angle'* question they were asked: *"Where were you sitting when you saw the (e.g.) sports equipment trial?"*

2.2.2 Experimental Task

First, patients' performance in the encoding session was assessed. Analysis was carried out on their performance against ceiling (i.e., examining whether scores were significantly different from 100%, see Table 1) as, although the control group answered the same questions at the same time and experienced the encoding sessions in exactly the same way, data were not systematically collected. Patients scored significantly lower than ceiling when recalling the actual items ($U = 44, p = .01$) and repeating positions on the grid ($U = 40, p = .04$). Given the failure of data collection from the older group, this is a conservative estimation of patient performance. It is important to note that any mistakes were corrected in this part of the encoding session. For the element of greatest interest (perspective from which they viewed the scene) their scores did not significantly differ from 100% ($U = 30, p = .07$).

Analysis was then carried out on scores in the two conditions of the main task. Between-subjects analysis compared the two groups' scores in the Item Shift condition to those obtained in the Perspective Shift condition. These data revealed that patients and controls were equivalently accurate in the Item Shift condition ($U = 37, p = .11$). However, in the

condition assessing memory for personal perspective (Perspective Shift) the patient group performance was significantly worse than the healthy controls ($U = 43.5, p = .01$).

Data were then compared within-subject group. Whereas the control group were equivalently accurate in the Item Shift and Perspective Shift conditions ($W = 10, p = .06$), the Patient group were significantly more accurate in the Item Shift condition ($W = 15, p = .04$).




Figure 3

- A. Overall number correct for each group in the line drawing recognition task (standard error mean bars)
- B. Spatial span in Corsi blocks task. Both the forwards and backwards conditions are shown for each group (standard error mean bars).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

C. *Performance in the Perspective Shift task. Dots plot scores of individual participants. Red diamond indicates mean. Red lines are standard error bars and the pink diamond indicates median.*

D. *Performance in the Item Shift task. Red diamond indicates mean. Red lines are standard error bars and the pink diamond indicates median.*

Five out of the six patients demonstrated the expected deficit –performing more accurately in the Item Shift task compared to the Perspective shift task. When subtracting the patients' scores in Perspective Shift from those achieved in Item Shift the decrement in performance from the Item Shift to the Perspective Shift task was sizable, ranging from 14% to 57% ($M = 36.80$; median = 29). The only patient who did not show this effect was accurate on only 57% of trials for both tasks - their performance can be seen as the lowest dot on the Item Shift task for this group in this condition. This abnormally weak performance in Item Shift might indicate a wider memory problem or reflect occipital damage, which was not present in the other patients.

Finally, we assessed whether the patients' confidence in their responses differed from that of controls. Self-rated scores of confidence were equivalent across groups for both the Perspective Shift and Item Shift conditions ($U = 29$, $p = .57$; $U = 33.5$, $p = .23$ respectively).

The patient group's confidence in their performance also did not differ for the two experimental conditions ($W = 10$, $p = .50$). However, the control group were significantly more confident in their answers to the Item Shift condition compared to the Perspective Shift condition ($W = 20$, $p = .05$).

Experiment 1 reveals impairments in self-perspective aspects of episodic memory in patients with damage to right parietal cortex. These individuals were not impaired on a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

standard episodic memory task, were at ceiling in a task assessing encoding of scenes, were unimpaired in spatial working memory tasks and – crucially - were not impaired in an experimental memory condition in which allocentric spatial relationships were manipulated (i.e. Item Shift trials). However, they were impaired when compared to neurologically healthy volunteers when judging which image - purporting to be from the head camera they wore during encoding- was the one they had experienced *only* when they were required to differentiate their own versus another image of the same scene taken from another angle. This is important, as the use of these egocentric visuo-spatial mechanisms in episodic memory has not been previously studied in research interrogating the role of these regions in long-term memory. Here we directly assessed this aspect of episodic memory and demonstrate that it is impaired following parietal damage. These novel findings are potentially important for these patients. This group has been previously overlooked in analysis of episodic memory impairments, and delineating long-lasting cognitive deficits after right hemisphere stroke is important as evidence suggests that these have been underestimated (Bonato, 2012).

The two conditions within the novel paradigm directly compare the ability to select a participant's own scene in the presence of a lure that varies in either perspective information (Perspective Shift) or in allocentric spatial information (Item Shift). It was in the Perspective Shift condition *only* that the patient group revealed a deficit. This suggests that parietal damage does not result in an overall impairment in episodic memory or indeed visuo-spatial aspects of episodic memory *per se* (as tested by Item Shifts), which harmonises with previous research. In addition, these patients performed the standard test of spatial working memory – the Corsi block task- at a level not significantly different to controls. As a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

result, the data cannot be explained by impairments in spatial working memory (Malhotra *et al.*, 2005). Further, overall task difficulty is unlikely to explain the results as the Corsi block task is challenging, particularly the backwards condition, and here patients' performance was statistically equivalent to healthy control participants.

We were concerned with delineating whether patients with right-hemisphere damage might suffer from specific and subtle episodic deficits rather than outlining precise lesion sites; however it is interesting to examine the lesion anatomy. The region of damage overlapping in all patients centres on the right supramarginal gyrus (Brodmann's area 40). This is relevant as evidence suggests these ventral parietal regions are involved in explicit 'Remember' as opposed to more implicit 'Know' responses in standard episodic tasks (e.g., Wheeler and Buckner, 2004). There is a large body of evidence implicating ventral lateral parietal cortex in episodic retrieval - specifically with the *quality* of the memory, as involvement of these regions enables retrieval of richer more vivid information (see Vilberg & Rugg, 2008 and Rugg & King, 2017). Although this relationship is often shown in the left hemisphere, frequently the episodic tasks used to investigate neural response use word stimuli as the to-be-remembered items (e.g., Wheeler and Buckner, 2004; Gonzalez *et al.*, 2015) which might be one factor leading to greater recruitment of left-hemisphere regions. The data here suggest that it is likely that the act of explicitly visually recalling an event (leading to accurate 'Remember' responses) relies in part on the ventral parietal region damaged in our patients. Here, we suggest that in right hemisphere the 'Remember' act involves accurate recall of visuospatial relations from an egocentric viewpoint.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Data examining the self-rated confidence of our participants complements findings suggesting that accurate insight of episodic recall accuracy is altered after parietal damage (e.g., Berryhill *et al.*, 2009; Simons *et al.*, 2010). Patients were equivalently confident in their responses to both perspective and item shift conditions despite their poor performance in the first condition.

The task developed in Experiment 1 reveals a specific deficit in right hemisphere stroke patients in episodic recall from self-perspective. The area of maximal overlap is consistent with evidence for the lateral posterior parietal cortex and vividness of episodic recall. An interesting avenue to develop this paradigm is with healthily ageing adults. Older adults' episodic recall is frequently reported as lacking specificity and detail, as has been reported in parietal patients (e.g., see Levine *et al* 2002; Addis *et al*, 2011; Schacter *et al*, 2013). There is also an increase in memories being recalled from an 'observer' rather than 'field' perspective (Piolino *et al* 2006; 2009), which correlates with reports of a weaker subjective feeling of experiencing the memory and a lack of rich detail (e.g., Piolino *et al*, 2006). The task used here is suited to probing which areas of parietal cortex are involved in discriminating between one's own versus another perspective in episodic memory. By introducing a critical design adaptation (simply showing only one image to the participants at a time in the recognition task) with MVPA we can use it to decode parts of the parietal cortex that are involved in this discrimination.

3. Experiment 2

3. 1 Materials and Methods

3. 1. 1 Participants

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Twenty-eight individuals took part: 16 healthy younger participants (11 females) between the age of 19 and 24 years ($M = 20.87$, $SD = 1.73$) and 12 healthy older participants (8 females) aged from 62 to 81 years ($M = 69.25$, $SD = 6.18$). Younger participants were recruited from the local student population and older participants were recruited from an over 50s exercise class and from the local University of the Third Age (U3A). Data are presented for 23 of these individuals. Two people opted out of going in the MRI scanner on day 2, a further two experienced technical problems on the second day and one person's performance was below chance in all memory tests and conditions. The remaining 25 were 13 younger participants (19-24 year olds, $M = 21$) and 10 older adults (62-81 year olds, $M = 69.30$). Each participant was reimbursed £15 in total for their attendance costs. All participants were asked to confirm that they had no current diagnosis of neurological or psychiatric illness. They all gave written consent according to the Declaration of Helsinki, with the study having been approved by the local research ethics committee.

3.1.2 Standardised Memory tasks

As in the previous study, the following tasks were used to assess episodic and spatial working memory: *Rivermead Behavioural Memory Test – Line drawing delayed recognition* (RBMT-3); *Corsi Block Tapping Task – forwards and backwards*. This experiment ran over two days, permitting further examination. Extra assessments were, *Buschke's Selective Reminding Test* (Buschke, 1973) and the *Rey-Osterrieth Complex Figure (ROCF) test*. The Buschke task consisted of a series of 12 unrelated words presented over 12 trials, or until the subject was able to recall the entire list on three consecutive trials. A delayed recall test was given without warning 30 minutes after completion. In the *ROCF* participants were asked to copy a complex visuo-spatial design as accurately as possible. Once complete, the

1
2
3 design was removed and participants were asked to immediately draw it again from
4
5
6
7 memory. After 20-30 minutes they had a delayed-recall task for the figure.

8 **3.1.2 Experimental Task**

9
10 The task was modified in several ways. First, in order to make the memory test more
11
12 challenging for healthy participants, double the number of scenes were shown at encoding.
13
14 The same 14 categories were used with the same 8 exemplars but two different scenes from
15
16 each category were presented (both contained different exemplars). For example, in the
17
18 stationery category one scene might contain a stapler and a highlighter pen and the other a
19
20 roll of tape and a ruler. Scenes from the same category were not presented in the same
21
22 encoding block. Encoding was split into 4 blocks of 7 scenes. As before, at the end of each
23
24 block participants were cued with the category name and then asked to recall the items,
25
26 their positions, and the viewing angle they experienced that scene from. To increase the
27
28 difficulty further the recognition test in the scanner for the novel scenes took place the day
29
30 after encoding, rather than on the same day.
31
32
33
34
35
36
37
38
39
40

41 Again all participants wore a head camera during encoding. They were shown this camera
42
43 and it was explained to them that a still would be taken of each scene to be used in a
44
45 memory test in the scanner on the following day. As in Experiment 1, we had pre-prepared
46
47 stills of each scene. On the first day, encoding was followed by administration of
48
49 standardised tests. At the end of the session participants were presented again with the to-
50
51 be-encoded scenes, but for only 15 seconds. This was added, as piloting revealed this
52
53
54
55
56
57 facilitated maintenance of the scenes for the next day in the older group.
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In the recognition test in the scanner on day 2 the task was adapted to enable us to interrogate the imaging data. We sought to decode neural activity in response to images from within specific conditions and it was therefore necessary to have only one visible at a time. Participants saw 28 individual scenes, two times during the scanning session. Ten of these scenes were identical to the ones shown in the lab the day before (i.e., the scene that they saw from the angle in which they had viewed it, *identical* condition), 10 were of the same scene but taken from the opposite viewing perspective (*perspective shift* condition), 4 were taken from the same viewing angle with the same items but they were placed in different position (*item shift* condition), and 4 were from the same viewing angle, the same category and same squares used but contained different exemplars (*object change* condition).

Images were displayed for five seconds. After the stimulus disappeared a question was presented on screen: “Did you see this scene in the lab yesterday?” It was emphasised that the viewing angle of these images was not relevant for the first question. They were required to respond whether those items in those grid positions were seen during encoding. If participants responded ‘yes’ then they were asked: “Was the scene taken from your viewpoint (i.e. it is the image from your head-camera)?” The task took approximately eight minutes, after which an anatomical scan of the participants’ brain was taken for 4 and half minutes. The participant then repeated the recall experiment in order to increase discrimination power.

3.1.3 Imaging & data analysis

1 Data were acquired at the Combined Universities Brain Imaging Centre (CUBIC) with a 3T
2 MRI scanner (Magnetom Trio, Siemens, Erlangen, Germany) using a standard Siemens eight-
3 channel array headcoil. Functional images were acquired with a gradient-echo, echoplanar
4 (EPI) sequence (TR 3000 ms, TE 31 ms, voxel size 3 x 3 x 3 mm) comprising 41 axial slices (64
5 x 64 matrix) covering the entire brain, and acquired continuously during each experimental
6 run (231 timepoints). A high-resolution (1 x 1 x 1 mm) anatomical scan of the whole brain
7 was also acquired.
8
9

10 All data were pre-processed using SPM8 (www.fil.ion.ucl.ac.uk/spm/) and included slice-
11 time correction, motion correction, and spatial smoothing using a Gaussian kernel with
12 FWHM of 6mm. A general linear model approach was used to model each individual correct
13 trial as a separate regressor, and convolved with the canonical hemodynamic response
14 function. Modelled trials represented the 5s presentation of the visual scene plus the time
15 leading to their response. The 'object change' and item shift' trials were also included as
16 regressors of no interest, as were the realignment parameters as calculated during motion
17 correction. Participant-specific parameter estimates were calculated for each regressor and
18 transformed into t-values resulting in a series of t-maps representing trial-specific activation
19 for use in the subsequent MVPA analysis.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 A searchlight approach was adopted to identify those regions of the brain sensitive to
50 differences between 'identical' and 'perspective shift' scenes. MVPA was performed with
51 custom written Matlab scripts utilizing a linear support vector machine (LIBSVM;
52 www.csie.ntu.edu.tw/~cjlin/libsvm/). A spherical ROI (radius = 3 voxels) was defined that
53 moved sequentially through every voxel in the functional volume space providing 123 t-
54
55
56
57
58
59
60
61
62
63
64
65

1 values from each ROI for classification purposes. Thus the feature matrix for pattern
2 classification comprised 40x123 elements (20 exemplars of the 'identical' response and 20
3 exemplars of the 'perspective shift' response [from 2 recording sessions], 123 voxels ROI),
4 with each feature normalised to unit length. The information content of each searchlight
5 ROI was assessed using a k-fold cross validation approach (k=5). Partitioning of the feature
6 matrix into training and test sets was balanced such that there were always two examples of
7 each condition from both sessions. Classifier accuracy was calculated by averaging over the
8 5 folds. In order to generalise the results and assess overall classification performance, the
9 final three-dimensional maps of decoder accuracies for each participant were normalised to
10 MNI space and smoothed with a 6mm Gaussian filter.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28 Group analysis, testing whether decoding performance was above chance between
29 subjects, was conducted using a voxelwise non-parametric permutation test (FSL
30 Randomise; Winkler et al, 2014) and corrected for multiple comparisons using threshold-
31 free cluster enhancement. In order to investigate the functional role of the parietal lobe as
32 well as the lesioned areas as identified in Experiment 1, we restricted this group analysis to
33 two regions of interest: The lesion region as determined from the patient study (Figure 1),
34 and a broader bi-lateral parietal lobe ROI defined using WFU Pickatlas (Ver. 3.0.4; Maldjian
35 et al, 2003).
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 **3. 2 Results**

53 **3.2.1 Standardised Tests (Table 2)**

54 *Rivermead Behavioural Memory Test:*
55
56
57
58
59
60
61
62
63
64
65

1
2
3 Recognition by older participants did not significantly differ from younger participants after
4 a 30-minute delay, $U = 49$, $z = -1.29$, $p = .31$.

5 *Corsi Block Tests:*

6
7 Scores on the forward tapping task did not significantly differ between younger and older
8 participants, $t(21) = -1.34$, $p = .19$. This was also true in the backwards condition, $t(21) = -.97$,
9
10
11
12
13 $p = .34$.

14
15 *Bushcke Selective Reminding Task:*

16
17 Younger participants were able to recall significantly more words after a 30-minute delay
18
19 than the older participants, $U = 23$, $z = -2.80$, $p = .01$.

20
21
22 *Rey Complex Figure:*

23
24
25 Figures were scored according to the Boston Qualitative Scoring System (Stern et al, 1994).
26
27
28 There were no significant differences between the groups $t(17) = -.20$, $p = .84$.

	Older Group	Younger Group
<i>Line drawing – memory task (proportion)</i>	.96 (.56)	.99 (.39)
<i>Corsi blocks forward span</i>	6.20 (1.32)	6.92 (1.26)
<i>Corsi blocks backwards span</i>	5.80 (1.55)	6.31 (.95)
<i>Bushcke Selective Reminding Task (30 min delay - /12)</i>	9.3 (2.98)	11.69 (.48)
<i>Rey-Osterrith Complex Figure Task (delayed retention %tile)</i>	47.25 (15.63)	48.91 (18.82)
Experimental Task		
Question 1	d' 2.96 (.80)	d' 3.86 (.43)
'Did you see this scene in the lab yesterday?'	Hit rate .94 (.06) False alarm .16 (.16)	Hit rate .97 (.02) False alarm .04 (.06)
Question 2		
'Is this the image from your own view (your head camera)?'	d' 1.10 (.94) Hit rate .71 (.15) False alarm .34 (.17)	d' 2.51 (.94) Hit rate .90 (.08) False alarm .17 (.13)

Table 2

Data from behavioural measures in Experiment 2 (standard deviations in brackets).




Figure 4

Dots plot scores of individual participants: Red diamonds indicate mean; red lines are standard error bars; pink diamonds indicate median.

3.2.2 Experimental task

First, responses to Question 1 ('Did you see this scene yesterday?') were analysed. There were 4 conditions (Identical; Perspective shift; Object Change; Item shift). The correct answer to this question for Identical and Perspective shift should be 'yes' and 'no' for the other two conditions. The paradigm lends itself well to analysis with a sensitivity measure such as d' prime. Questions such as 'Is this the image from your head camera?', are potentially affected by response bias – somebody might be more prone to respond 'yes' and achieve a similar percentage correct to another participant who is more selective. Therefore, we analysed target sensitivity (d') according to signal detection theory, taking account of proportion for hit rates and false alarms in each question (Table 2). For question 1 'Did you see this scene in the lab yesterday?' saying 'yes' to the 'Identical' and 'Perspective-Shift' trials was correct –the hit rate - whereas saying 'yes' to object change

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

and item shift constituted false alarms. Independent samples t-test on this sensitivity analysis revealed the older group to be significantly worse than that younger group ($t(21) = -3.50, p = .002$).

Responses to the second question (Is this the image from your head camera?) were then analysed. As participants were only asked this question if they answered that they had seen the scene yesterday, data here were analysed only for the relevant conditions –‘Identical’ and ‘Perspective shift’ stimuli. Here responding ‘yes’ to ‘Identical’ trials constituted the hit rate whereas responding ‘yes’ to ‘Perspective shift’ trials was considered a false alarm. Here, the older group were again significantly worse than the younger group ($t(21) = -3.59, p = .002$). Although the older participants were not significantly worse than the younger group in the Rivermead recognition task, Rey Complex Figure task and the forwards and backwards Corsi blocks, they were less sensitive than the younger group to the correct answers for both questions in our experimental paradigm. They also recalled significantly fewer words in the Bushcke task than younger participants.

3.2.3 Imaging Data

MVPA searchlight analysis was carried out to identify areas of the brain sensitive to changes in viewing angle (i.e. brain areas whose responses differ such that they differentiate between identical and perspective shift scenes). We restricted the group level analysis to two ROIs: the lesion region and bilateral parietal lobes. The statistical maps derived from this group analysis of both the young and old individual classifier accuracy maps (testing for classifier accuracy > 50% and thresholded at $p < 0.05$, FWE corrected for multiple comparisons) are shown in Figure 5.




Figure 5

Group classifier maps indicating areas of significant decoder performance in differentiating the neural response associated with correctly identified identical/perspective-shift scenes (classifier accuracy > 50%; threshold $p < 0.05$, FWE corrected for multiple comparisons). These are all superimposed on a template brain in the MNI coordinate space. Blue circles highlight statistically significant cluster peaks. (A) Young group; (B) & (C) Older group; (D) Blue overlay represents the lesion map ROI, orange/yellow overlay represents the group classifier map established in the older group parietal ROI (as shown in B & C), white overlay represents the group classifier map from the lesion ROI in the older group.

Independent t -tests comparing each cluster across groups revealed that there were no significant differences between old and young for the left hemisphere circled cluster in 5A (Cluster 1 Young; $t(21) = -.88$, $p = .39$) and the left hemisphere circled anterior cluster in 5B (Cluster 1 Old; $t(21) = .62$, $p = .55$). Whereas the classifier accuracy in the other three areas of significance in the old group

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

significantly differed between the two groups: posterior circled region in 5B (Cluster 2; $t(21) = -2.70$, $p = .01$); circled region in 5C (Cluster 3; $t(21) = -2.24$, $p = .04$); and the precuneus region (Cluster 4; $t(21) = 2.37$, $p = .03$).

See Supplementary methods for whole brain analysis.

The analysis within the larger bilateral parietal ROI reveals a number of distinct clusters in both groups. In the young participants, 'Cluster 1 Young' emerges with a peak within the inferior parietal lobe of the left hemisphere (Fig 5A, mean classifier accuracy at statistical peak 66%; $t=3.98$; MNI -33, -37, 58). The older group in contrast shows much larger networks of the parietal region sensitive to differences between identical and perspective shift scenes. In the left hemisphere, two lateral parietal clusters can be seen (Fig 5B) - one more anterior, close to the postcentral gyrus ('Cluster 1 Old', accuracy 61%; $t=4.89$; MNI -45, -28, 58) and corresponding to that observed in the younger group; the second in the inferior parietal lobe and encompassing the angular gyrus ('Cluster 2 Old', accuracy 53%; $t=4.86$; MNI -42, -70, 46). A similar cluster to this is observed in the right hemisphere again in the region of inferior parietal and angular gyrus (Fig 5C, 'Cluster 3 Old', accuracy 55%; $t=4.47$; MNI 45, -64, 46). Finally, a midline cluster spanning both hemispheres is observed in the region of the precuneus (Fig 5C 'Cluster 4 Old', accuracy 53%; $t=4.18$; MNI -3, -67, 37).

When the group analysis is restricted to the ROI defined by the lesion map, a smaller cluster centred around the same right hemisphere angular gyrus region as identified above is evident in the older group. This is shown in Fig 5D as the small white cluster overlayed on the parietal lobe map as described above as well as the lesion ROI (shown in blue).

Unsurprisingly, no clusters were observed in the younger group within this ROI.

1
2
3 Results revealed by MVPA show that specific patterns of activity in areas of parietal cortex
4
5 were indicative of whether participants were viewing either a scene they encoded the
6
7 previous day *from the view in which it was experienced* or the same scene from a *different*
8
9 *perspective*. This reinforces the proposal, supported by the results of Experiment 1, that the
10
11 role for parietal cortex within episodic memory is directly related to recalling from one's
12
13 own eyes –egocentric – encoding perspective. In the young group it would appear that left
14
15 parietal cortex is vital– for example the area of supramarginal gyrus indicted in our
16
17 classification analysis - whereas in the older group we revealed evidence of a wider
18
19 network, which encompasses the angular gyrus in both hemispheres. The involvement of
20
21 angular gyri in our task harmonises with evidence from previous neuropsychological, neural
22
23 disruption and functional imaging studies that these regions are involved with the subjective
24
25 experience of recalling an episodic memory (eg.,Davidson et al 2008; Sestieri et al, 2013;
26
27 Marcotti and St Jacques, 2018). We suggest that recall of our memory from an egocentric
28
29 perspective – as we experienced it at the time – is crucial to this sense of subjective
30
31 experience.
32
33

34
35
36 Regarding the results from the younger group there is a question regarding why
37
38 classification accuracy in decoding differences between the key conditions did not activate
39
40 the larger network of areas identified in the older group. First, we draw attention to
41
42 functional imaging studies of memory with standard univariate analysis comparing young
43
44 and older groups. There are many examples of greater neural recruitment in older adults,
45
46 potentially as compensation for less efficiency (e.g., Angel et al, 2014, Stern et al, 2012;
47
48 Zarahn et al, 2007). Morcom & Friston (2012), used an analysis similar to that adopted here,
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

to decode neural activity patterns associated with successful recall in younger and older adults. Their analysis revealed wider involvement of bilateral networks in the older as compared to the younger group. In contrast, Wang et al (2016), using MVPA, revealed no differences between young and old adults but this was in the context of no group differences in the behavioural assessment of memory – in contrast to our study. We would suggest that a key reason for our results is that the young group found the task rather easy as, in order to achieve reasonable accuracy in the older group across a complicated set of stimuli and a long time gap, there were only 28 scenes and these scenes were shown twice. It is possible that the younger group did not need to use the same strategy as the older group – a strategy that is key to our design i.e. recreating a mental image of the scene when encoded. A hint of this is that at debrief some younger participants reported recalling which chair they sat in for the different scenes as a mnemonic - older participants never reported this strategy. We believe that in follow-up work, if this strategy is counteracted and the task more challenging for younger participants, they would show the same regions are being used for this task.

4. General Discussion

In the two studies presented here we have introduced a novel paradigm that demonstrates a key role for ventral parietal regions in egocentric perspective information within episodic memory. This is important as accurately recalling experienced events from one's own perspective is linked with increased accuracy and stronger sense of subjective experience (e.g., Marcotti and St Jacques, 2018). Our data support research implying a role for vividness of recall in activation of ventral regions of parietal cortex, and suggest, an alternative or additional account to those currently posited-that these regions are involved in multimodal

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

representations and processing of context per se in episodic memory (e.g., Bonnici et al 2016 and Ramanan, Piquet & Irish, 2017). It appears likely from this evidence (where modality was the same across conditions) that the visual-spatial representation from one's own encoding view is perhaps a critical context processed by these areas of parietal cortex.

In Experiment 1, our paradigm revealed a deficit in egocentric visuospatial memory in patients with right parietal damage. This is the first demonstration of a specific episodic memory deficit in this patient group using a new task. The next step is to carry out a larger study clarifying whether this deficit in accurate recall from an egocentric perspective might explain previous findings in patients with parietal damage who lacked vivid detail in recalling and had a reduced feeling of recalling a personally experienced event when remembering (e.g., Drowos et al, 2010). The results described here help to explain the role of parietal cortex in episodic memory and suggest that patients with parietal damage may have long-lasting deficits in this domain. Another important avenue for further exploration, would involve assessment of patients with damage to analogous left-hemisphere regions. This would be valuable as evidence suggests that the right hemisphere may be specialised for egocentric processing (e.g. Iachini, Ruggiero, Conson & Trojano, 2009).

It is difficult to draw strong conclusions about the relative roles of the left and right hemispheres from the existing literature concerning episodic memory following parietal damage. Some studies have only tested very small groups of patients with bilateral damage (n=2: e.g., Berryhill et al, 2009; 2007; Drowos et al, 2010; Hower et al, 2014). Or their critical impairment has only been found in a bilateral group and not in either group of unilateral patients (e.g., Simons et al, 2010). Davidson et al (2008) assessed a group of four left

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

hemisphere patients and one right hemisphere patient, precluding separate analysis of the two groups. Recently Ciaramelli et al (2017) tested six patients in their main analysis from an original group of five with left and two with right hemisphere damage but did not state which hemisphere was damaged in the patient who did not complete the study and analysis was not separated according to lesion side. To our knowledge only one study has separately analysed patients with left (n = 3) versus right (n = 4) parietal damage (Ciaramelli et al, 2010), revealing those with left hemisphere damage to be more impaired on two topographical tasks loading on egocentric as opposed to allocentric memory. However, pertinent to our current paradigm, there were similar impairments in both groups when patients were directly asked if they recalled their route from a first ('field') or third ('observer') person perspective (only two left hemisphere patients and two right hemisphere patients used a first person perspective) and both groups were abnormal when asked if they 're-experienced' the route during recall. Only one individual reported 're-experiencing' the memory and they had left hemisphere damage. This was a small group but it is relevant that no right hemisphere patient reported re-experiencing the memory. Right hemisphere regions might be particularly important for re-experiencing an event, for which it is crucial to reconstruct the visual spatial image of the event as we experienced it, from our own eyes' perspective.

Our study is an important development as, to our knowledge, this is the largest patient study examining episodic memory in patients with parietal damage in which they are matched for lesion hemisphere. Given the disparity in stroke outcome after left versus right hemisphere damage in attention as demonstrated by the far greater number of patients with spatial neglect after right hemisphere damage (e.g., Stone et al, 1993), it is important

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

to analyse patients with damage to each hemisphere separately. Further, evidence suggests that any spatial neglect after left hemisphere stroke impairs allocentric spatial processing more than egocentric (see Kleinman et al, 2007).

In the second experiment healthy older participants were impaired in judging whether a presented scene was from the same perspective they viewed it from - in contrast to performance in most other tests of episodic and working memory. Functional imaging using MVPA in older participants during our task revealed that ventral parietal regions, including angular gyrus, were sensitive to differences between stimuli from a self or other perspective. In the older group, ROI analysis of bilateral parietal cortex revealed the angular gyrus to be involved in correctly discriminating whether a scene at retrieval was the same as one's own perspective. Therefore, both left and right hemisphere are involved in making the judgment critical to our task. Regions around the angular gyri bilaterally were significantly accurate in distinguishing between self versus another's visual perspective in previously encoded scenes. It is clear that the left hemisphere is critically involved in episodic memory and the previous focus on this area is not simply a result of the nature of the stimuli or language requirements in the task – in our task verbal encoding of the scenes would be very unlikely to encapsulate the visuo-spatial differences between one's own versus another's angle of view. This raises the question of why were patients had an intact left hemisphere impaired in our task? Taken together, our results might suggest that although the left hemisphere is important, the right hemisphere regions are necessary to reliably make the judgement required here. As discussed above, examination of neuropsychological evidence for the role of the different hemispheres of parietal cortex in episodic memory is not fully informative at present. Though we would draw attention to evidence from other studies

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

that these regions in the right hemisphere might be concerned with recalling personally experienced episodic information regarding the self both visuo-spatially (e.g. St Jacques et al, 2016) and auditorily (e.g., Levine et al 2004). This is consistent with the separate analysis of right and left hemisphere patients in Ciaramelli et al (2010) – that is, the right hemisphere patients did not ‘re-experience’ the memory and were poor with first-person recall. As well as examining left-hemisphere patients, study of the connectivity between these regions in a task similar to ours might enable further clarification. Bellana et al (2016) carried out analysis of bilateral angular gyrus connectivity with the default mode network (DMN). They found that left and right angular gyri were strongly connected to this DMN at retrieval, with right angular gyrus being more strongly connected to medial temporal lobe (MTL) regions and the left angular gyrus with other units of the DMN. This provides converging evidence that the hemispheres have differing roles during episodic recall.

33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In conclusion, our studies reveal egocentric episodic deficits in patients with right parietal damage and delineate the role for ventral lateral parietal regions in egocentric representation in a functional imaging study using classification analysis. The results presented here provide important evidence that parietal cortex activity is directly associated with egocentric spatial perspective aspects of episodic memory and demonstrate for the first time a specific deficit in this cognitive process in patients with right hemisphere parietal damage.

5. Acknowledgements

We thank all our participants for their incredibly valuable help. This work was supported by a grant from the Bial Foundation, Portugal (155/10) and the National Institute for Health Research (NIHR) Imperial Biomedical Research Centre.

6. References

- Addis DR, Roberts RP, Schacter DL. Age-related neural changes in autobiographical remembering and imagining. *Neuropsychologia* 2011; 49 (13):3656-69
- Ally BA, Simons JS, McKeever JD, Peers PV, Budson AE. Parietal contributions to recollection: electrophysiological evidence from aging and patients with parietal lesions. *Neuropsychologia* 2008; 46(7): 1800-12.
- Angel L, Bastin C, Genon S, Balteau E, Phillips C, Luxen A, Maquet P, Salmon E, Collette F. Differential effects of aging on the neural correlates of recollection and familiarity. 2013. *Cortex*; 49. 10.1016.
- Bellana B, Liu Z, Anderson JAE, Moscovitch M, Grady CL. Laterality effects in functional connectivity of the angular gyrus during rest and episodic retrieval. *Neuropsychologia*. 2016 Jan 8;80:24-34
- Berryhill ME. Insights from neuropsychology: pinpointing the role of the posterior parietal cortex in episodic and working memory. *Front Integr Neurosci* 2012; 6: 31.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Berryhill ME, Drowos DB, Olson IR. Bilateral parietal cortex damage does not impair associative memory for paired stimuli. *Cogn Neuropsychol* 2009; 26(7): 606-19.

Berryhill ME, Phuong L, Picasso L, Cabeza R, Olson IR. Parietal lobe and episodic memory: bilateral damage causes impaired free recall of autobiographical memory. *J Neurosci* 2007; 27(52): 14415-23.

Bonato M. Neglect and extinction depend greatly on task demands: a review. *Front Hum Neurosci* 2012; 6: 195.

Bonnici HM, Kumaran D, Chadwick MJ, Weiskopf N, Hassabis D, Maguire EA Multi-voxel pattern analysis in human hippocampal subfields *Hippocampus*. 2012 22:1143–1153

Bonnici HM, Richter FR, Yazar Y, Simons JS. Multimodal Feature Integration in the Angular Gyrus during Episodic and Semantic Retrieval. *J Neurosci*. 2016 May 18;36(20):5462-71

Burgess N. Spatial cognition and the brain. *Ann N Y Acad Sci* 2008; 1124: 77-97.

Burgess N, Becker S, King JA, O'Keefe J. Memory for events and their spatial context: models and experiments. *Philos Trans R Soc Lond B Biol Sci* 2001; 356(1413): 1493-503.

Cabeza R, Ciaramelli E, Olson IR, Moscovitch M. The parietal cortex and episodic memory: an attentional account. *Nat Rev Neurosci* 2008; 9(8): 613-25.

1
2
3 Chadwick MJ, Hassabis D, Maguire EA Decoding overlapping memories in the medial
4
5
6
7
8
9 temporal lobes using high-resolution fMRI. *Learn. Mem.* 2011 18: 742-746

10 Chadwick MJ, Hassabis D, Weiskopf N, Maguire EA. Decoding individual episodic memory
11
12 traces in the human hippocampus. *Current Biology.* 2010; 20(6): 544-7.
13
14
15
16

17 Ciaramelli E, Faggi G, Scarpazza C, Mattioli F, Spaniol J, Ghetti S, Moscovitch M. Subjective
18
19
20
21
22
23
24
25
26
27 recollection independent from multifeatural context retrieval following damage to the
28
29
30
31
32
33
34
35
36
37 posterior parietal cortex. *Cortex.* 2017 Jun;91:114-125

38 Ciaramelli E, Grady C, Levine B, Ween J, Moscovitch M. Top-down and bottom-up attention
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65 to memory are dissociated in posterior parietal cortex: neuroimaging and
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

38 Corbetta M, Shulman GL. Spatial neglect and attention networks. *Annu Rev Neurosci* 2011;
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65 34: 569-99.

46 Davidson PS, Anaki D, Ciaramelli E, Cohn M, Kim AS, Murphy KJ, *et al.* Does lateral parietal
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

56 Drowos DB, Berryhill M, Andre JM, Olson IR. True memory, false memory, and subjective
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
46

1
2
3 Fridriksson J, Kjartansson O, Morgan PS, Hjaltason H, Magnusdottir S, Bonilha L, *et al.*

4
5 Impaired speech repetition and left parietal lobe damage. *J Neurosci* 2010; 30(33): 11057-
6
7
8 61.

9
10
11
12 Gonzalez A, Hutchinson JB, Uncapher MR, Chen J, LaRocque KF, Foster BL, *et al.*

13
14
15 Electrocortigraphy reveals the temporal dynamics of posterior parietal cortical activity
16
17 during recognition memory decisions. *Proc Natl Acad Sci U S A* 2015; 112(35): 11066-71.

18
19
20
21
22
23 Hower KH, Wixted J, Berryhill ME, Olson IR. Impaired perception of mnemonic oldness, but
24
25 not mnemonic newness, after parietal lobe damage. *Neuropsychologia* 2014; 56: 409-17.

26
27
28
29
30
31 Iachini T, Ruggiero G, Conson M, Trojano L. Lateralization of egocentric and allocentric
32
33 spatial processing after parietal brain lesions. *Brain Cogn.* 2009 Apr;69(3):514-20.

34
35
36
37
38
39 Kleinman JT, Newhart M, Davis C, Heidler-Gary J, Gottesman RF, Hillis AE. Right hemispatial
40
41 neglect: frequency and characterization following acute left hemisphere stroke. *Brain Cogn.*
42
43 2007;64(1):50-9.

44
45
46
47
48
49 Lambrey S, Doeller C, Berthoz A, Burgess N. Imagining being somewhere else: neural basis of
50
51 changing perspective in space. *Cereb Cortex* 2012; 22(1): 166-74.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Levine B, Svoboda E, Hay JF, Winocur G, Moscovitch M. Aging and autobiographical memory: dissociating episodic from semantic retrieval. *Psychol Aging*. 2002 Dec;17(4):677-89.

Levine B, Turner GR, Tisserand D, Hevenor SJ, Graham SJ, McIntosh AR. The functional neuroanatomy of episodic and semantic autobiographical remembering: a prospective functional MRI study. *Journal of Cognitive Neuroscience*. 2004; 16(9): 1633-46

Malhotra P, Jager HR, Parton A, Greenwood R, Playford ED, Brown MM, *et al*. Spatial working memory capacity in unilateral neglect. *Brain* 2005; 128(Pt 2): 424-35.

Maldjian, JA, Laurienti, PJ, Burdette, JB, Kraft RA. An Automated Method for Neuroanatomic and Cytoarchitectonic Atlas-based Interrogation of fMRI Data Sets. *NeuroImage* 2003. 19:1233-1239.

Marcotti P, St Jacques PL. Shifting visual perspective during memory retrieval reduces the accuracy of subsequent memories. *Memory*. 2018 Mar;26(3):330-341.

Morcom AM, Friston KJ. Decoding episodic memory in ageing: a Bayesian analysis of activity patterns predicting memory. *Neuroimage*. 2012;59(2):1772-82.

Mort DJ, Malhotra P, Mannan SK, Rorden C, Pambakian A, Kennard C, *et al*. The anatomy of visual neglect. *Brain* 2003; 126(Pt 9): 1986-97.

1
2
3 Olson IR, Berryhill M. Some surprising findings on the involvement of the parietal lobe in
4 human memory. *Neurobiol Learn Mem* 2009; 91(2): 155-65.

5
6
7
8 Piolino P, Desgranges B, Clarys D, Guillery-Girard B, Taconnat L, Isingrini M, *et al.*
9
10 Autobiographical memory, auto-noetic consciousness, and self-perspective in aging. *Psychol*
11
12
13 *Aging* 2006; 21(3): 510-25.

14
15
16
17
18 Piolino P, Desgranges B, Eustache F. Episodic autobiographical memories over the course of
19
20
21 time: cognitive, neuropsychological and neuroimaging findings. *Neuropsychologia* 2009;
22
23
24 47(11): 2314-29.

25
26
27
28 Ramanan S, Piguet O, Irish M. Rethinking the Role of the Angular Gyrus in Remembering the
29
30
31 Past and Imagining the Future: The Contextual Integration Model. *Neuroscientist*. 2017 Oct
32
33
34 1:1073858417735514

35
36
37
38 Rorden C, Hjalton H, Fillmore P, Fridriksson J, Kjartansson O, Magnúsdóttir S, *et al.*
39
40
41 Allocentric neglect strongly associated with egocentric neglect. *Neuropsychologia* 2012;
42
43
44 50(6): 1151-7.

45
46
47
48 Rugg MD, King DR. Ventral lateral parietal cortex and episodic memory retrieval. *Cortex*.
49
50
51 2017 Jul 25. pii: S0010-9452(17)30237-X. doi: 10.1016/j.cortex.2017.07.012.

52
53
54
55
56 Rugg MD, Vilberg KL. Brain networks underlying episodic memory retrieval. *Curr Opin*
57
58
59 *Neurobiol* 2013; 23(2): 255-60.

1
2
3 Rusconi E, Pinel P, Dehaene S, Kleinschmidt A. The enigma of Gerstmann's syndrome
4
5 revisited: a telling tale of the vicissitudes of neuropsychology. *Brain* 2010; 133(Pt 2): 320-32.
6
7

8
9
10 St Jacques PL, Carpenter AC, Szpunar KK, Schacter DL. Remembering and imagining
11
12 alternative versions of the personal past. *Neuropsychologia*. 2018 Feb;110:170-179.
13
14
15

16
17
18 St Jacques PL, Szpunar KK, Schacter DL. Shifting visual perspective during retrieval shapes
19
20 autobiographical memories. *Neuroimage*. 2017 Mar 1;148:103-114.
21
22
23

24
25
26 Schacter DL, Gaesser B, Addis DR. Remembering the past and imagining the future in the
27
28 elderly. *Gerontology*. 2013;59(2):143-51
29
30
31

32
33
34 Schoo LA, van Zandvoort MJ, Biessels GJ, Kappelle LJ, Postma A, de Haan EH. The posterior
35
36 parietal paradox: Why do functional magnetic resonance imaging and lesion studies on
37
38 episodic memory produce conflicting results? *J Neuropsychol* 2011; 5(Pt 1): 15-38.
39
40
41

42
43
44 Scoville WB, Milner B. Loss of recent memory after bilateral hippocampal lesions. *J Neurol*
45
46 *Neurosurg Psychiatry* 1957; 20(1): 11-21.
47
48
49

50
51
52 Sestieri C, Capotosto P, Tosoni A, Luca Romani G, Corbetta M. Interference with episodic
53
54 memory retrieval following transcranial stimulation of the inferior but not the superior
55
56 parietal lobule. *Neuropsychologia*. 2013 Apr;51(5):900-6.
57
58
59

1
2
3 Sestieri C, Shulman GL, Corbetta M. The contribution of the human posterior parietal cortex
4 to episodic memory. *Nat Rev Neurosci.* 2017 Feb 17;18(3):183-192.

5
6
7
8 Simons JS, Peers PV, Hwang DY, Ally BA, Fletcher PC, Budson AE. Is the parietal lobe
9 necessary for recollection in humans? *Neuropsychologia* 2008; 46(4): 1185-91.

10
11
12
13
14
15 Simons JS, Peers PV, Mazuz YS, Berryhill ME, Olson IR. Dissociation between memory
16 accuracy and memory confidence following bilateral parietal lesions. *Cereb Cortex* 2010;
17 20(2): 479-85.

18
19
20
21
22
23
24
25
26 Spiers HJ, Burgess N, Hartley T, Vargha-Khadem F, O'Keefe J. Bilateral hippocampal
27 pathology impairs topographical and episodic memory but not visual pattern matching.
28
29
30
31 *Hippocampus* 2001; 11(6): 715-25.

32
33
34
35
36 Squire LR, Stark CE, Clark RE. The medial temporal lobe. *Annu Rev Neurosci* 2004; 27: 279-
37
38 306.

39
40
41
42
43
44 Stern R, Singer EA, Duke LM, Singer NG, Morey C, Daughtrey EW, Kaplan E. The Boston
45
46
47
48 Qualitative Scoring System for the Rey-Osterrieth Complex Figure. *Clinical*
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Tu S, Spiers HJ, Hodges JR, Piguet O, Hornberger M. Egocentric versus Allocentric Spatial Memory in Behavioral Variant Frontotemporal Dementia and Alzheimer's Disease. *J Alzheimers Dis.* 2017;59(3):883-892.

Tulving E, Episodic Memory: From Mind to Brain. *Annual Review of Psychology* 2002 53:1, 1-25

Rugg MD, Vilberg KL. Brain networks underlying episodic memory retrieval. *Curr Opin Neurobiol.* 2013 Apr;23(2):255-60

Stone SP, Halligan PW, Greenwood RJ. The incidence of neglect phenomena and related disorders in patients with an acute right or left hemisphere stroke. *Age Ageing.* 1993 Jan;22(1):46-52.

Wagner AD, Schacter DL, Rotte M, Koutstaal W, Maril A, Dale AM, *et al.* Building memories: remembering and forgetting of verbal experiences as predicted by brain activity. *Science* 1998; 281(5380): 1188-91.

Weniger G, Ruhleder M, Wolf S, Lange C, Irle E. Egocentric memory impaired and allocentric memory intact as assessed by virtual reality in subjects with unilateral parietal cortex lesions. *Neuropsychologia* 2009; 47(1): 59-69.

Wheeler ME, Buckner RL. Functional-anatomic correlates of remembering and knowing. *Neuroimage* 2004; 21(4): 1337-49.

1
2
3 Wilson B, Cockburn J, Halligan P. Development of a behavioral test of visuospatial neglect.

4
5 Arch Phys Med Rehabil 1987; 68(2): 98-102.
6
7

8
9
10 Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation inference for the
11
12 general linear model. Neuroimage. 2014 May 15;92:381-97.
13
14
15

16
17
18 Yazar Y, Bergström ZM, Simons JS. Reduced multimodal integration of memory features
19
20 following continuous theta burst stimulation of angular gyrus. Brain Stimul. 2017 May -
21
22 Jun;10(3):624-629.
23
24
25

26
27
28 Zarahn E, Rakitin B, Abela D, Flynn J, Stern Y. Age-related changes in brain activation during
29
30 a delayed item recognition task. Neurobiol Aging. 2007 May;28(5):784-98
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65