

Manuscript Details

Manuscript number	AQTOX_2019_61
Title	Combined effects of environmental xeno-estrogens within multi-component mixtures: comparison of in vitro human- and zebrafish-based estrogenicity bioassays
Article type	Research Paper

Abstract

In vitro bioassays based on estrogen receptor (ER) activation are commonly used to monitor the environmental contamination by xeno-estrogens. However, recent studies showed that fish- and human-based bioassays may have distinct responses to environmental samples, highlighting not only the need to better understand bioassay-specific ER response to environmentally more realistic mixtures of individual chemicals, but also how well these mixture responses can be explained by the default additivity model of concentration addition (CA). For this purpose, we investigated experimentally a 12-compound mixture in two different mixture ratios (M1 and M2) by testing the combination of (1) all 12 compounds, (2) only the ER activators present in this mixture, (3) only the ER inhibitors, and (4) ER activators and inhibitors combined. The mixture included well-known ER ligands such as bisphenol A (BPA) and genistein (GEN), but also non-estrogenic compounds that were considered as representatives of a freshwater background contamination. Studies were conducted on zebrafish (zf) liver cells stably expressing zfER α (ZELH α cells) or zfER β 2 (ZELH β 2 cells) and human ER reporter gene (MELN) cells, with the main aim (1) to assess the robustness of CA, and (2) to evaluate the potentially confounding influence of environmental chemicals on additivity. The testing of individual chemicals revealed a higher prevalence of ER inhibiting chemicals in zebrafish than human cells (e.g. propiconazole, benzo(b)fluoranthene). We also identified chemicals that activated hER but inhibited zfER response (e.g. benzo(a)pyrene, triphenylphosphate). In MELN cells, the estrogenic activity of both 12-compound mixtures M1 and M2 was well predicted by CA. However, in ZELH β 2 cells, the same mixtures induced significantly lower estrogenic responses than expected by CA. In contrast, if only the two ER ligands BPA and GEN were tested as binary mixture, their mixture effects were in good agreement with CA expectations. The stepwise experimental approach of testing subgroups of only ER activators or/and inhibitors indicate that the observed deviation from additivity is due to ZELH-specific inhibiting chemicals. Thus, the very distinct responses of human- and zebrafish cell lines to M1 and M2 can entirely be explained by the presence of ER inhibiting chemicals selectively active in zebrafish cells. Overall, this study provides novel information on the ability of environmental pollutants to interfere positively or negatively with zfER-signalling and shows that the response to a complex mixture of xeno-estrogens can be influenced by the presence of other (non- or anti-estrogenic) chemicals in a bioassay-specific manner.

Keywords	estrogenicity; anti-estrogen; mixture; in vitro reporter gene; human; zebrafish
Taxonomy	Estrogen Receptor, Danio Rerio, Mixture Model, In Vitro Toxicology, Chemical Mixture, Endocrine Activity
Corresponding Author	Selim Ait-Aissa
Corresponding Author's Institution	ineris
Order of Authors	Hélène Serra, Martin Scholze, Rolf Altenburger, Wibke Busch, Helene Budzinski, François BRION, Selim Ait-Aissa
Suggested reviewers	Taisen Iguchi, José M. Navas, Daniel Schlenk

Submission Files Included in this PDF

File Name [File Type]

AquaTox_Serra_COVERING_LETTER-012119.pdf [Cover Letter]

AquaTox_Serra_MS_012119.pdf [Manuscript File]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Research Data Related to this Submission

There are no linked research data sets for this submission. The following reason is given:
Data will be made available on request

Aït-Aïssa Selim, PhD
National Institute of Industrial Environment and Risks (INERIS)
Unit of Ecotoxicology in vitro and in vivo - UMR I SEBIO 02
Parc Alata - BP2, f-60550 Verneuil-en-Halatte, FRANCE
Phone : (33) 3 44 556 511
E-mail: selim.ait-aïssa@ineris.fr

To: Editor of *Aquatic Toxicology*

January 18th, 2019

Subject: Original manuscript submission for publication in *Aquatic Toxicology*.

Dear Editor,

With this letter, we are submitting the manuscript “Combined effects of environmental xeno-estrogens within multi-component mixtures: comparison of in vitro human- and zebrafish-based estrogenicity bioassays” by Hélène Serra, Martin Scholze, Rolf Altenburger, Wibke Busch, Hélène Budzinski, François Brion and Selim Aït-Aïssa, for publication as a research article in *Aquatic Toxicology*.

In this study, we investigated differences between human and zebrafish cell-based assays in assessing estrogenic activity of mixtures of aquatic contaminants. By using a stepwise experimental approach based on the concentration addition model, we newly identify ER-response inhibiting chemicals in zebrafish cells and demonstrate that they negatively influenced the zebrafish cell response to two 12-compound mixtures. Our study confirms previously reported differences between human and zebrafish bioassays in response to environmental pollutants and complex mixtures, and illustrates such differences using model aquatic pollutants. Another major outcome is the assessment of the estrogenic effects of mixtures including both ER activators and inhibitors, which has been rarely reported in such a methodological way, and the demonstration that deviation of additivity is likely to occur when present in environmental mixtures. These findings may have implication in environmental monitoring, i.e. need to consider bioassays that are specific to aquatic vertebrates when assessing estrogenic potency of samples issued from the aquatic environment, but also, more generally, in the assessment of mixture estrogenic effect, which can vary depending on the examined tissue or species.

An assurance is given that the material has not been published or submitted elsewhere.

We hope our paper will reach the standards allowing it to be published in *Aquatic Toxicology* and are looking forward to hearing from you.

Yours sincerely,

S. Aït-Aïssa

1
2
3 **1 Combined effects of environmental xeno-estrogens within multi-component mixtures:**
4
5 **2 comparison of *in vitro* human- and zebrafish-based estrogenicity bioassays**
6
7

8 3 H el ene Serra^{1,2}, Martin Scholze³, Rolf Altenburger⁴, Wibke Busch⁴, H el ene Budzinski², Fran ois
9 4 Brion¹, Selim A it-A issa^{1,*}
10
11

12 5
13
14 ¹Institut National de l'Environnement Industriel et des risques (INERIS), Unit  Ecotoxicologie *in*
15 6 *vitro* et *in vivo*, UMR-I 02 SEBIO, 60550 Verneuil-en-Halatte, France
16 7
17

18 8 ² UMR-CNRS EPOC/LPTC, Universit  de Bordeaux, Talence, France
19

20 9 ³Brunel University London, Uxbridge, United Kingdom
21

22 10 ⁴UFZ– Helmholtz Centre for Environmental Research, Leipzig, Germany
23
24

25 11

26 12 *Corresponding author. Email: selim.ait-aissa@ineris.fr
27
28

29 13
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59

60
61
62 **ABSTRACT**
63
64

15 *In vitro* bioassays based on estrogen receptor (ER) activation are commonly used to monitor the
66
67 environmental contamination by xeno-estrogens. However, recent studies showed that fish- and
68
69 human-based bioassays may have distinct responses to environmental samples, highlighting not
70
71 only the need to better understand bioassay-specific ER response to environmentally more realistic
72
73 mixtures of individual chemicals, but also how well these mixture responses can be explained by
74
75 the default additivity model of concentration addition (CA). For this purpose, we investigated
76
77 experimentally a 12-compound mixture in two different mixture ratios (M1 and M2) by testing the
78
79 combination of (1) all 12 compounds, (2) only the ER activators present in this mixture, (3) only
80
81 the ER inhibitors, and (4) ER activators and inhibitors combined. The mixture included well-known
82
83 ER ligands such as bisphenol A (BPA) and genistein (GEN), but also non-estrogenic compounds
84
85 that were considered as representatives of a freshwater background contamination. Studies were
86
87 conducted on zebrafish (zf) liver cells stably expressing zfER α (ZELH α cells) or zfER β 2 (ZELH β 2
88
89 cells) and human ER reporter gene (MELN) cells, with the main aim (1) to assess the robustness of
90
91 CA, and (2) to evaluate the potentially confounding influence of environmental chemicals on
92
93 additivity. The testing of individual chemicals revealed a higher prevalence of ER inhibiting
94
95 chemicals in zebrafish than human cells (e.g. propiconazole, benzo(b)fluoranthene). We also
96
97 identified chemicals that activated hER but inhibited zfER response (e.g. benzo(a)pyrene,
98
99 triphenylphosphate). In MELN cells, the estrogenic activity of both 12-compound mixtures M1 and
100
101 M2 was well predicted by CA. However, in ZELH β 2 cells, the same mixtures induced significantly
102
103 lower estrogenic responses than expected by CA. In contrast, if only the two ER ligands BPA and
104
105 GEN were tested as binary mixture, their mixture effects were in good agreement with CA
106
107 expectations. The stepwise experimental approach of testing subgroups of only ER activators or/and
108
109 inhibitors indicate that the observed deviation from additivity is due to ZELH-specific inhibiting
110
111 chemicals. Thus, the very distinct responses of human- and zebrafish cell lines to M1 and M2 can
112
113 entirely be explained by the presence of ER inhibiting chemicals selectively active in zebrafish
114
115
116
117
118

119
120
121 40 cells. Overall, this study provides novel information on the ability of environmental pollutants to
122
123 41 interfere positively or negatively with zfER-signalling and shows that the response to a complex
124
125 42 mixture of xeno-estrogens can be influenced by the presence of other (non- or anti-estrogenic)
126
127 43 chemicals in a bioassay-specific manner.

129 44 **KEY WORDS:** estrogenicity, anti-estrogen, mixture, *in vitro* reporter gene, human, zebrafish
130
131

132 45 **HIGHLIGHTS (IF NEEDED):**
133
134

- 135 46 - Human and zebrafish cells showed distinct estrogenic response to 12-component mixtures
136
137 47 containing bisphenol A and genistein
138
139 48 - Several ER inhibiting chemicals were identified only in zebrafish cells
140
141 49 - Using a stepwise experimental approach, we showed that these inhibiting chemicals influenced
142
143 50 negatively the zebrafish cells response to xeno-estrogens mixtures
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177

1. Introduction

The occurrence of numerous endocrine disrupting chemicals (EDC) in aquatic ecosystems has raised concern over their potential adverse effects in aquatic organisms, such as fish (Sumpter, 2005). Many EDCs, such as natural and synthetic hormones, pesticides or industrial chemicals, are xeno-estrogens, *i.e.* they bind the estrogen receptors (ERs) and subsequently alter the transcription of target genes involved in key physiological functions (Sumpter, 2005). *In vitro* bioassays based on ER transactivation have been used to assess the estrogenic activity of chemicals, but also of environmental samples (Könemann et al., 2018; Zacharewski, 1997). In case of environmental monitoring, they are expected to enable an integrative detection of various ER-active contaminants within complex environmental mixtures considering both known and unknown xeno-estrogens. They provide a unique quantitative response which may be summarized as estradiol-equivalent (E2-Eq, Kase et al., 2018).

To date, a large majority of *in vitro* bioassays used in environmental bio-monitoring are based on mammalian or yeast cell systems that stably express a reporter gene which expression is controlled by the human ER subtype α (hER α) (Könemann et al., 2018; Kunz et al., 2015). However, the relevance of using human-based assay to assess hazard and risk for aquatic species is a question of concern in environmental assessment (Hotchkiss et al., 2008). For instance, humans express two ER subtypes, ER α and ER β , but most teleost fish express at least three ER subtypes, ER α , ER β 1 and ER β 2 (Menuet et al., 2002; Tohyama et al., 2015). Fish and human ER have relatively low sequence homologies in their ligand binding domain (Menuet et al., 2002; Tohyama et al., 2015). These structural differences are believed to contribute to the distinct sensitivity to certain xeno-estrogens (Miyagawa et al., 2014), along with other factors linked to the cell specificities, such as cell metabolic capacities (Le Fol et al., 2015), presence/absence of transcriptional cofactors or cross-talks with other signalling pathways (Navas and Segner, 2000; Ohtake et al., 2003).

237
238
239 75 In a recent study, we reported that some surface water samples were active on a zebrafish
240
241 76 liver cell line stably expressing zebrafish ER β 2 (zfER β 2), the ZELH β 2 cells, but not on human
242
243 77 breast cancer MELN cells that endogenously express hER α (Sonavane et al., 2016). Similarly, some
244
245 78 effluent extracts from sewage treatment plants produced very different *in vitro* responses in cells
246
247 79 expressing human or medaka ER α (Ihara et al., 2014). These differences were further confirmed *in*
248
249 80 *vivo* by measuring vitellogenin induction in exposed male medaka (Ihara et al., 2015). In the latter
250
251 81 study, the estrogenic chemicals identified were not sufficient to explain the distinct response of fish
252
253 82 bioassays. However, the authors showed that the anti-estrogenic activity measured in the samples
254
255 83 may contribute to the different responses of medaka and human ER.
256
257

258
259 84 Several studies have addressed the combined effect of ER ligands in reconstituted mixtures,
260
261 85 generally concluding on their additive effects based on concentration addition (CA) predictions
262
263 86 (Kortenkamp, 2007). However, xeno-estrogens occur in the aquatic ecosystem together with other
264
265 87 chemicals that have various and distinct modes of action (e.g. Escher et al., 2014; Neale et al., 2015,
266
267 88 Busch et al., 2016). To date, few studies have investigated additive effects of xeno-estrogens in
268
269 89 more diverse exposure scenarios, such as with non- or weak estrogenic chemicals (Evans et al.,
270
271 90 2012) or with anti-estrogenic chemicals (Yang et al., 2015). Recently, a mixture of 12 selected
272
273 91 environmental chemicals was tested in zebrafish and human-based bioassays as part of a larger
274
275 92 round-robin study. The aim was to investigate whether the estrogenic activity of the ER ligands in
276
277 93 this mixture (e.g. genistein and bisphenol A) was detectable against the background of the other
278
279 94 environmental pollutants (Altenburger et al., 2018). This study concluded that in human MELN
280
281 95 cells the overall estrogenic activity of the mixtures was accurately predicted by an assumed
282
283 96 additivity of the estrogenic chemicals. However, in zebrafish ZELH β 2 cells the measured estrogenic
284
285 97 response of the mixture was lower than expected. The reasons of this discrepancy between human
286
287 98 and zebrafish-based ER-reporter gene assays were unknown, and therefore raised the question about
288
289 99 potential limitations of a presumed CA additivity.
290
291
292
293
294
295

296
297
298 100 In this context, the present study was designed as a follow-up of Altenburger et al. (2018)
299
300 101 to investigate the different responses of zebrafish- and human-based *in vitro* reporter gene assays.
301
302 102 We hypothesized that estrogenic chemicals within environmental mixtures have additive effects
303
304 103 following default model of CA that are well detected by zebrafish and human-based bioassays. In
305
306 104 such way, we investigated (1) the additivity of xeno-estrogens in zebrafish and human-based
307
308 105 bioassays and (2) the influence of non-estrogenic chemicals of the mixtures. As in Altenburger et
309
310 106 al. (2018), we used the same 12-compound mixture in two different mixture ratios (M1 and M2),
311
312 107 which included xeno-estrogens (e.g. bisphenol A and genistein), and non-estrogenic chemicals
313
314 108 representatives of a freshwater contamination background. The general experimental set-up design
315
316 109 is outlined in Figure 1. Firstly, each chemical was tested for both estrogenic and anti-estrogenic
317
318 110 activities in zebrafish-and human-based bioassays. Secondly, combinations of chemicals that
319
320 111 proved to be active at M1 and M2 mixture ratios (either ER activating, ER inhibiting, or both) were
321
322 112 tested and then discussed in relation to the outcomes from the 12-component mixture response. The
323
324 113 concentration addition model was used to evaluate the additivity of active chemicals in each mixture
325
326 114 scenario.

328 329 330 115 **2. Material and methods**

331 332 333 116 ***2.1 Chemical selection, mixtures design and experimental approach***

334
335 117 Twelve environmentally relevant chemicals were selected following (1) a prioritization
336
337 118 exercise based on occurrence, hazard and available environmental quality standard (Busch et al.,
338
339 119 2016), and (2) a screening of prioritized contaminants through multiple bioassays (Neale et al.,
340
341 120 2017a). As a result, two fixed-ratio mixtures of 12 chemicals with dissimilar mode of actions were
342
343 121 designed (Table SI-1) and tested as part of a benchmarking exercise (Altenburger et al., 2018). The
344
345 122 first mixture ratio (M1) was composed in such way that the diverse bioactivities of the individual
346
347 123 chemicals had a chance to be detected experimentally by an array of 19 bioassays. The second
348
349 124 mixture ratio (M2) was chosen to mimic a realistic freshwater contamination scenario. In the current
350
351
352
353
354

355
356
357 125 study, all 12 chemicals were tested individually for their capacity to induce or inhibit ER-mediated
358
359 126 luciferase response in different cellular assays. Based on the information on the activity of
360
361 127 individual chemicals in each bioassay, chemicals predicted to contribute to M1 and M2 responses
362
363 128 based on CA prediction were identified. Subgroup mixtures were then designed containing either
364
365 129 only ER activators or only ER inhibitors, or both ER activators and inhibitors (Figure 1, Table 1).
366
367 130 These mixtures were designed such that their relative concentration ratios agreed to that from the
368
369 131 original M1 and M2 mixtures (i.e. real sub-mixtures), to allow the best possible comparison to the
370
371 132 outcomes from the 12 compound mixtures.
372
373
374

375 133 ***2.2 Chemicals and reagents***

376 134 17 β -estradiol (E2, CAS#50-28-2, purity of >98%), triclosan (TCS, CAS#3380-34-5, purity
377
378 of 97% - 103%), bisphenol A (BPA, CAS#80-05-7, purity of 97%), genistein (GEN, CAS#446-72-
379 135 0, purity of > 98%), propiconazole (CAS#60207-90-1, purity of >98%), diclofenac (CAS#15307-
380 136 0, purity of > 98%), propiconazole (CAS#60207-90-1, purity of >98%), diclofenac (CAS#15307-
381 136 0, purity of > 98%), propiconazole (CAS#60207-90-1, purity of >98%), diclofenac (CAS#15307-
382 136 0, purity of > 98%), propiconazole (CAS#60207-90-1, purity of >98%), diclofenac (CAS#15307-
383 137 79-6), diazinon (CAS#333-41-5, purity of >98%), diuron (CAS#330-54-1, purity >98%), cyprodinil
384
385 138 (CAS#121552-61-2, purity of >98%), triphenylphosphate (TPP, CAS#115-86-6, purity >99%),
386
387 139 benzo(a)pyrene (BaP, CAS#50-32-8, purity >96%), benzo(b)fluoranthene (BbF, CAS#205-99-2,
388
389 140 purity of 98%), chlorophene (CAS#120-32-1, purity of 95%), hydroxy-tamoxifen (OH-TAM,
390
391 141 CAS#68392-35-8, purity of >98%) and dimethylsulfoxide (DMSO) were purchased from Sigma-
392
393 142 Aldrich (France). The cell culture medium and reagents Leibovitz 15 culture medium (L-15), fetal
394
395 143 calf serum (FCS), 4-(2-hydroxy-ethyl)-1-piperazineethanesulfonic acid (HEPES), epidermal
396
397 144 growth factor (EGF), G418, 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazoliumbromide (MTT)
398
399 145 and D-luciferin were purchased from Sigma Aldrich (St-Quentin Fallavier, France); Dulbecco's
400
401 146 Modified Eagle Medium (DMEM), DMEM High Glucose (DMEM HG) powder, F-12 nutrient
402
403 147 mixture (Ham's F12) powder, penicillin and streptomycin were purchased from Gibco (France);
404
405 148 insulin, hygromycin B and sodium bicarbonate were purchased from Dominique Dutscher (France).
406
407

408 149 ***2.3 In vitro bioassays: cell lines, luciferase and cell viability assays***

414
415
416 150 The zebrafish *in vitro* assays have been derived from the zebrafish liver (ZFL) cell line
417
418 151 (Cosnefroy et al., 2012). ZFL were stably transfected, first, with an ERE-driven firefly luciferase
419
420 152 gene, yielding the ZELH cell line, and then either with zfER α subtype, yielding the ZELH α cell
421
422 153 line, or with zfER β subtype yielding the ZELH β 2 cell line (Cosnefroy et al., 2012). Establishment
423
424 154 of these cell models and their response to different classes of well-known xeno-estrogens have been
425
426 155 previously described (Cosnefroy et al., 2012; Sonavane et al., 2016). The human-derived MELN
427
428 156 cell line (Balaguer et al., 1999) was kindly provided by Dr Patrick Balaguer (INSERM Montpellier,
429
430 157 France). It is derived from the breast cancer MCF-7 cells, which endogenously express the hER α ,
431
432 158 but no functional hER β (P. Balaguer, *personal communication*). MELN cells were stably
433
434 159 transfected with an ERE-driven firefly luciferase reporter gene.
435
436
437

438 160 Conditions for routine cell culture have been detailed previously (Balaguer et al., 1999;
439
440 161 Cosnefroy et al., 2012). The cells used were pathogen-free and controlled on a regular basis. For
441
442 162 exposure experiments, ZELH-derived cells were seeded in 96-well white opaque culture plates
443
444 163 (Greiner CellStarTM, Dutscher, France) at 25,000 cells per well in phenol red-free LDF-DCC
445
446 164 medium (containing L-15 50%, DMEM HG 35%, Ham's F12 15%, HEPES 15 mM, 0.15 g/L
447
448 165 sodium bicarbonate, 0.01 mg/mL insulin, 50 ng/mL EGF, 50 U/mL penicillin and streptomycin
449
450 166 antibiotics, 5% v/v stripped serum). MELN were seeded at 80,000 cells per well in phenol red-free
451
452 167 DMEM medium containing 5% v/v stripped serum. Cells were left to adhere for 24h. Then, they
453
454 168 were exposed in triplicates to serial dilutions of test compound for either 72h at 28°C for zebrafish
455
456 169 cells or 16h at 37°C for MELN cells. Each plate included both solvent and positive controls (in two
457
458 170 triplicates each). E2 was used as a positive quality control for ER activation, and hydroxy-tamoxifen
459
460 171 (OH-TAM) for ER inhibition. In addition, a serial dilution of 7 to 8 concentrations of E2 was tested
461
462 172 in each experiment. At the end of exposure, the culture medium was removed and replaced by 50 μ L
463
464 173 per well of medium containing 0.3 mM luciferin. The luminescence signal was measured in living
465
466 174 cells using a microtiter plate luminometer (Synergy H4, BioTek).
467
468
469
470
471
472

473
474
475 175 The cell viability was assessed by using the 3-(4,5-dimethyl-thiazol-2-yl)-2,5diphenyl
476
477 176 tetrazolium bromide (MTT) assay (Mosmann, 1983). After cell exposure, the culture medium was
478
479 177 removed and replaced by 100 μ L of medium containing 0.5 mg/mL MTT. Cells were incubated for
480
481 178 3h. In metabolically active cells, MTT is reduced onto a blue formazan precipitate, which is
482
483 179 dissolved by adding 100 μ L of DMSO after removal of MTT-containing medium. Plates were read
484
485 180 at 570 nm against a 640 nm reference wavelength on a microplate reader (KC-4, BioTek
486
487 181 Instruments, France) and results are expressed as absorbance units relative to control cells.
488
489

490 182 ***2.4 Testing of multi-component mixtures***

491
492 183 The mixture compositions are given in Table SI-1, SI-2 and SI-3. The two 12-component
493
494 184 mixtures were prepared in methanol (as part of a round robin study on bioassays, Altenburger et al.,
495
496 185 2018). Stocks solutions and serial dilutions of single chemicals and 2-, 3-, 4- and 5-component
497
498 186 mixtures were prepared in DMSO. The response of MELN cells to TPP and BPA using either
499
500 187 DMSO or methanol as vehicle were similar (data not shown), thus, no significant effect of the
501
502 188 solvent was to expect. To investigate the anti-estrogenic activity of the chemicals or mixtures, the
503
504 189 cells were exposed in the presence of E2 at a concentration leading to 80% of maximal response,
505
506 190 i.e. 0.1 nM in MELN and ZELH β 2 and 1 nM in ZELH α assays. The ZELH cells, that correspond
507
508 191 to the parent cell line of ZELH α and ZELH β 2 cells but lack functional ER, were used additionally
509
510 192 as a control for non-specific luciferase modulation. As for the other cell lines, cytotoxicity was
511
512 193 measured in parallel in the way previously described. Final solvent concentrations in culture
513
514 194 medium were 0.1% v/v (agonist assay) or 0.15% v/v (in case of co-exposure with E2), which do
515
516 195 not affect luciferase expression or cell viability. Stock solutions of chemicals in DMSO and
517
518 196 methanol were maintained at -20°C for up to three months.
519
520

521 522 197 ***2.5 Data analysis***

523 524 525 198 ***2.5.1 Data treatment and analysis***

532
533
534 199 Luciferase activity (LUC) was normalized to a response range between 0 and 1 on an
535
536 200 experiment-to-experiment basis as follows:

539 201
$$Response = \frac{LUC_{chemical} - LUC_{control}}{LUC_{E2} - LUC_{control}} \quad (1)$$

543 202 where $LUC_{chemical}$ is the luminescent signal induced by the tested chemical, $LUC_{control}$ is the average
544
545 203 luminescent signal of the solvent controls and LUC_{E2} is the average luminescent signal of the E2
546
547 204 positive controls. Concentration-effect data analysis was performed in the same way for individual
548
549 205 compounds and mixtures. In short, a nonlinear regression model best-fit approach was used to
550
551 206 describe pooled data sets in the best possible way (Scholze et al., 2001). If different regression
552
553 207 functions led to similar goodness-of-fits, the logit model (which is a re-parameterised form of the
554
555 208 Hill equation) was given preference. To account for inter-study variations we included experiments
556
557 209 as random factor in the best-fit data analysis (nonlinear mixed effect model). A detailed description
558
559 210 can be found in Altenburger et al. (2018).

562 211 ***2.5.2 Mixture prediction and uncertainty assessment***

564 212 The combined response from individual substances was assumed to follow the concept of
565
566 213 concentration addition (CA). Here we used the standard form of non-interaction, i.e.:

570 214
$$\sum_{i=1}^n \left(\frac{C_i}{EC_{xi}} \right) = 1 \quad (2)$$

573 215 where C_i is the concentration of the i^{th} substance in the mixture expected to produce a mixture
574
575 216 response X, and EC_{xi} the concentration of the i^{th} substance leading to the same response X as
576
577 217 expected for the mixture.

580 218 To account for the statistical uncertainty in the CA prediction, a combination of Monte-Carlo (MC)
581
582 219 simulations and bootstrapping nonlinear regression functions (Tibshirani and Efron, 1993) was
583
584 220 conducted to simulate approximate 95% confidence limits around the predicted mean response of
585
586 221 the mixture. Here the MC step is responsible for linking the data input from the single compounds

591
592
593 222 (i.e. estimates about ECs or individual effects) to the mixture prediction, and the bootstrapping step
594
595 223 is responsible for generating data information relevant for input variables (i.e. uncertainty
596
597 224 distributions around the single substance EC's or effects). We followed a parametric bootstrap with
598
599 225 resamples drawn from the fitted nonlinear mixed effect model. Differences between predicted and
600
601 226 observed mixture effects (concentration) were deemed statistically significant when the 95%
602
603 227 confidence belts of the prediction did not overlap with those of the experimentally observed mixture
604
605 228 effects (Altenburger et al., 2018). The comparative assessment was performed on mixture
606
607 229 concentrations leading to 20% ER activation (EC20) or inhibition (IC20).
608
609

610 230 **3. Results**

611 231 *3.1. Activation and inhibition of ER response by single chemicals*

612
613
614 231
615
616
617 232 The results of ER activation and inhibition by all 12 chemicals and the reference compounds (E2
618
619 233 and OH-TAM) on MELN, ZELH α and ZELH β 2 cells are presented in Table 2, and the
620
621 234 concentration-response data are provided in supplementary information (Figure SI-1 for ER
622
623 235 activation and SI-2 for ER inhibition).
624
625

626 236 As expected, genistein and BPA were active in all cell lines, but at different sensitivity and
627
628 237 efficacy levels. MELN cells responded to BPA with an EC20 of 0.12 μ M and a maximal induction
629
630 238 of 86% of the positive E2 control response, while ZELH α and ZELH β 2 cells showed a lower
631
632 239 sensitivity with an EC20 of 2.1 μ M and 5.0 μ M, respectively, and a maximum luciferase induction
633
634 240 around 30 % (Table 2). In case of genistein, MELN (EC20 of 0.0121 μ M) and ZELH β 2 cells (EC20
635
636 241 of 0.015 μ M) were more responsive than ZELH α cells (EC20 of 1.4 μ M). BaP, TPP and diazinon
637
638 242 weakly induced luciferase activity in MELN cells with an EC20 of 0.57 μ M, 4.1 μ M and 15 μ M,
639
640 243 respectively, whereas no activity was recorded at non-cytotoxic concentrations in zebrafish cells.
641
642 244 No other chemicals showed any estrogenic response up to 30 μ M in any bioassays.
643
644
645
646
647
648
649

650
651
652 245 The inhibition of ER response by the 12 chemicals revealed distinct response between the
653
654 246 bioassays (Table 2). Overall, several chemicals were identified as new ER inhibitors, mainly in
655
656 247 ZELH-zfERs cells. TPP and BaP decreased ER response in ZELH α and ZELH β 2 cells at
657
658 248 concentrations where they did not affect cell viability or the luciferase activity in the ER-negative
659
660 249 ZELH cells. Conversely, benzo(b)fluoranthene and propiconazole decreased E2-induced luciferase
661
662 250 activity up to 90% in ZELH α and ZELH β 2 and in ER-negative ZELH cells. Cyprodinil decreased
663
664 251 E2-induced luciferase activity across all the cell lines with similar sensitivity, suggesting a likely
665
666 252 non-specific effect of this chemical on luciferase activity (Table 2, Figure SI-3).
667
668

669 253 ***3.2. Combined effects of xeno-estrogens in multi-component mixtures***

670
671
672 254 The concentration-response curves estimated for the single chemicals were used to predict
673
674 255 the ER activation and ER inhibition of M1 and M2 mixtures using the CA model. Since CA can
675
676 256 describe only ER activation or ER inhibition, but not their co-occurrence, the additive response of
677
678 257 a mixture containing both ER activators and inhibitors is predicted solely from the ER activators in
679
680 258 case of ER activation or from the ER inhibitors in case of ER inhibition. Therefore, the chemicals
681
682 259 expected to induce ER activation or ER inhibition in M1 and M2 mixtures were identified for each
683
684 260 cell line based on CA prediction. They were then tested as subgroup mixtures containing either ER
685
686 261 activating (M1_A, M2_A), ER inhibiting (M1_I, M2_I), or both ER activating and inhibiting
687
688 262 chemicals (M1_A+I, M2_A+I) (Table 1). The relative concentration ratios were always kept in
689
690 263 accordance to the 12-compound mixtures M1 and M2. All subgroup mixture results are presented
691
692 264 in Figure 2 (mixture composition according to M1) and Figure 3 (mixture composition according
693
694 265 to M2), together with the outcomes for M1 and M2 (Altenburger et al., 2018). Details about the
695
696 266 mixture composition are given in Tables SI-1 (12-component mixtures) and in SI-2 and SI-3
697
698 267 (subgroup mixtures).
699
700

701 702 268 ***3.2.1 Additivity of ER activating or inhibiting chemicals***

709
710
711 269 Regarding subgroup mixtures of ER activating chemicals, there was overall a good
712
713 270 agreement between observed and predicted EC20 across all cell lines and for both mixtures M1 and
714
715 271 M2 compositions. In MELN cells, TPP, BPA and genistein at M1 mixture ratio had additive effects
716
717 272 very well predicted by CA model with a ratio between observed and predicted EC20 of 1.3
718
719 273 (M1_A_{MELN}, Figure 2A, Table 3). In comparison, the measured estrogenic activity of BPA and
720
721 274 genistein in M2_A_{MELN} was below the predicted response, although not statistically significant
722
723 275 (M2_A_{MELN}, Figure 3A, Table 4). BPA and genistein were the only two identified estrogenic
724
725 276 chemicals in ZELH α and ZELH β 2 cells. Their binary mixture induced an estrogenic response in a
726
727 277 good agreement with CA prediction at M1 and M2 concentration ratios in ZELH α (Figure 2E and
728
729 278 3E) and ZELH β 2 cells (Figure 2I and 3I). The ratio of observed against predicted EC20 was of 0.40
730
731 279 and 0.55 in ZELH α cells, and 0.71 and 0.73 in ZELH β 2 cells for M1 and M2, respectively.
732
733

734
735 280 As observed for single chemicals, ER inhibiting chemicals were more prevalent in ZELH α
736
737 281 and ZELH β 2 cells than in MELN cells. In MELN cells, cyprodinil was predicted to inhibit E2
738
739 282 response in M1, but only at high concentrations (M1_I_{MELN}, Figure 2B), and no inhibiting chemical
740
741 283 was identified for M2. In contrast, TPP, chlorophene and propiconazole were identified as ER
742
743 284 inhibiting chemicals of M1 in ZELH α and ZELH β 2 cells. In subgroup mixtures, they induced a
744
745 285 strong ER inhibition in ZELH α (M1_I_{ZELH α} , Figure 2F) and ZELH β 2 cells (M1_I_{ZELH β 2}, Figure 2J),
746
747 286 well predicted by the CA model (EC20 ratio of 0.87 and 0.83, respectively). Similarly, the subgroup
748
749 287 mixtures of ER inhibitors based on M2 mixture ratio induced a strong inhibition, well predicted by
750
751 288 CA model (M2_I_{ZELH α} , figure 3F and M2_I_{ZELH β 2}, Figure 3J, respectively). Overall, the combined
752
753 289 effects of ER activating or ER inhibiting chemicals were in good agreement with CA predictions
754
755 290 for both M1 and M2 mixture ratios and across all cell lines.
756
757

758 291 ***3.2.2 Estrogenic response to the 12-component mixtures: influence of inhibiting***
759
760 292 ***chemicals***
761
762
763
764
765
766
767

768
769
770 293 For each cell line, the combined effects of activator and inhibitor subgroup mixtures
771
772 294 (M1_A+I and M2_A+I) were determined and compared to the results of the 12 component mixtures
773
774 295 M1 and M2 (Figures 2 and 3, right part). The observed and predicted EC20 or IC20 of each mixture
775
776 296 are presented in Tables 3 (M1) and 4 (M2).
777

778
779 297 In MELN cells, the estrogenic activity of M1_A+I_{MELN} (Figure 2C) was well predicted by
780
781 298 CA, and this accuracy was not impacted negatively by the presence of 9 other environmental
782
783 299 substances (M1, Figure 2D). No active ER inhibitors were present at non-cytotoxic concentration
784
785 300 in the mixture M2, and therefore a mixture of activators and inhibitors was not tested. Nevertheless,
786
787 301 the mixture effect of all 12 substances was well explained by the additivity of the only two
788
789 302 estrogenic chemicals identified, BPA and genistein (M2, Figure 3D).
790

791
792 303 In zebrafish ZELH α cells, M1 was not expected to induce any estrogenic response in the
793
794 304 range of tested concentrations, and indeed no estrogenic response was observed neither with the 5-
795
796 305 component mixture (M1_A+I_{ZELH α} , Figure 2G) nor with the 12-component mixture M1 (Figure
797
798 306 2H). Conversely, a strong ER inhibiting response was measured (up to 80% inhibition) for both the
799
800 307 5- and 12-component mixtures, which was well predicted by the CA model (IC20 ratio of 0.74 and
801
802 308 0.95, respectively). Thus, the ER inhibition measured remained unaffected by addition of estrogenic
803
804 309 and inactive chemicals. In case of M2, the estrogenic activity of ER activating and inhibiting
805
806 310 chemicals was correctly predicted by CA model (Figures 3G and 3H). However, the estrogenic
807
808 311 activity measured was lower than that of BPA and genistein binary mixture results (Figure 3E),
809
810 312 suggesting an influence of ER inhibiting compounds.
811

812
813
814 313 In zebrafish ZELH β 2 cells, an estrogenic response was expected according to CA for the
815
816 314 mixture of activators and inhibitors, as supported by the additive outcomes from the binary mixture
817
818 315 of BPA and genistein (M1_A_{ZELH β 2}, Figure 2I). However, M1_A+I_{ZELH β 2} did not induce any
819
820 316 estrogenic response at test concentrations (Figure 2K). Instead, a strong inhibition of ER response
821
822 317 was measured, which was in line with the M1_I_{ZELH β 2} results and CA prediction (Figure 2J). As
823

827
828
829 318 observed for the subgroup mixture of ER activating and inhibiting chemicals (M1_A+I_{ZELHβ2}), M1
830
831 319 mixture did not induce any estrogenic activity but inhibited E2-induced response (Figure 2H).
832
833 320 Hence, these results indicate that inhibiting chemicals in M1 indeed influenced ER response in
834
835 321 ZELHβ2 cells. Compared with M1, the estrogenic activity measured for the subgroup mixture of
836
837 322 ER activators and inhibitors corresponding to M2 mixture ratio was well predicted by CA model
838
839 323 (M2_A+ I_{ZELHβ2}, Figure 3K), although the maximal efficacy observed was well below the one of
840
841 324 the BPA and genistein binary mixture (M2_A_{ZELHβ2}, Figure 3I). When ER activating and inhibiting
842
843 325 chemicals were grouped with inactive chemicals in M2, the estrogenic activity was well predicted
844
845 326 by CA up to 20% (Figure 3L), but the maximal estrogenic response remained lower than expected
846
847 327 based on the M2_A_{ZELHβ2} mixture results (Figure 3I). In comparison, the inhibition of ER response
848
849 328 was well predicted by CA for both M2_A+I_{ZELHβ2} (Figure 3K) and M2 (figure 3L). The results of
850
851 329 the 4-component mixture M2_A+I_{ZELHβ2} on ZELHβ2 cells are very similar to M2 results,
852
853 330 considering both ER activation and inhibition (Figure 3K and 3L).
854
855

856 331 **4. DISCUSSION**

857
858
859
860 332 The current study investigated the distinct responses of zebrafish ZELHα and ZELHβ2 and human
861
862 333 MELN cells ER reporter gene bioassays to 12-component mixtures composed of xeno-estrogens
863
864 334 and other environmental relevant chemicals (Altenburger et al., 2018). By using a stepwise
865
866 335 experimental approach from individual chemicals to subgroup mixture testing, we were able to
867
868 336 explain the distinct response of human and zebrafish bioassays to the same 12-component mixtures.
869
870

871 337 ***4.1. Distinct responses of human and zebrafish cell lines to individual chemicals***

872
873
874 338 BPA and genistein are well-known ER agonist ligands and were indeed active in all ER-
875
876 339 based bioassays, in agreement with previous studies using the same cellular models (Balaguer et
877
878 340 al., 1999; Cosnefroy et al., 2012; Le Fol et al., 2017; Sonavane et al., 2016). Apart from these two
879
880
881
882
883
884
885

886
887
888 341 compounds, the screening of individual chemicals highlighted some marked differences between
889
890 342 cell assays for some of the 10 chemicals.
891
892

893 343 One major outcome relates to the higher prevalence of chemicals inhibiting E2-induced
894
895 344 luciferase activity in ZELH-zfERs cells than in MELN cells (Table 2). Some chemicals had opposite
896
897 345 responses in zebrafish and human cells. For instance, BaP -a known AhR-ligand- and TPP were
898
899 346 estrogenic in MELN cells but decreased E2-induced response in ZELH α and ZELH β 2 cells. The
900
901 347 mechanistic interaction between AhR and ER signalling pathways has been documented in human
902
903 348 (Matthews and Gustafsson, 2006; Ohtake et al., 2003) and in fish (e.g. Navas and Segner, 2000).
904
905 349 The prototypical AhR ligand TCDD was shown to induce a weak estrogenic response in MELN
906
907 350 cells (Balaguer et al., 1999) while it decreased E2 response in all ZELH-zfER cells (Sonavane,
908
909 351 2015). The distinct responses to BaP in ZELH-zfERs and MELN cells might thus be explained, at
910
911 352 least partially, by AhR-ER interactions. In comparison, less information is available on the ability
912
913 353 of TPP to interact with ER signalling. Previous studies have reported a weak agonist effect on hER α
914
915 354 transactivation (Kojima et al., 2013), as observed in the current study in MELN cells, while some
916
917 355 TPP metabolites are reported to have an anti-estrogenic activity on hER β transactivation (Kojima
918
919 356 et al., 2016). However, TPP was unable to induce the ER-regulated brain aromatase expression gene
920
921 357 in transgenic cyp19a1b-GFP zebrafish embryos (Neale et al., 2017a). Considering the anti-
922
923 358 estrogenic activity of TPP evidenced in zebrafish liver cells, further research would be warranted
924
925 359 to assess whether TPP (or metabolites) either binds directly zfERs or alters zfER transactivation
926
927 360 through cross-talk(s) with other signaling pathways.
928
929
930

931 361 Other chemicals, such as propiconazole and cyprodinil, decreased E2-induced estrogenic
932
933 362 activity in an ER non-specific manner, i.e. they decreased firefly luciferase also in the parent cell
934
935 363 line ZELH that does not express functional zfER (Table 2, Figure SI-5). Such inhibition may reflect
936
937 364 either a direct effect on luciferase enzyme or an indirect effect on baseline transcriptional machinery
938
939 365 in the promoter region of the reporter gene, irrespectively of ER activity. Despite a weak estrogenic
940
941
942
943
944

945
946
947 366 activity on hER α reported *in vitro* (Medjakovic et al., 2014; Schlotz et al., 2017), cyprodinil
948
949 367 decreased firefly luciferase activity in all cells, irrespectively of E2 addition. The structural
950
951 368 similarities of cyprodinil with known firefly luciferase inhibitor (Auld and Inglese, 2004) and its
952
953 369 capacity to interfere with ATP production (Coleman et al., 2012) suggest a possible effect on the
954
955 370 reporter gene system. In case of propiconazole, a weak hER α agonist activity was reported in the
956
957 371 high μ M range in MVLN cells (Kjeldsen et al., 2013) and anti-proliferative effects measured in
958
959 372 MCF-7 cells (Kjaerstad et al., 2010). In fish, interference of propiconazole with estrogen signalling
960
961 373 pathway has been reported *in vivo* (Skolness et al., 2013) but no information on ER agonist or
962
963 374 antagonist activity is available. Thus, additional assays would be warranted to assess the specific
964
965 375 activity of propiconazole and cyprodinil on ER-signalling pathway in zebrafish.
966
967

968 969 376 **4.2. Deciphering cell-specific response to xeno-estrogen mixtures** 970

971
972 377 BPA and genistein were the main drivers for ER agonistic response in M1 and M2. When
973
974 378 combined as binary mixture, they induced in all zebrafish and human-based bioassays responses
975
976 379 that were in good agreement with CA predictions. This additivity is consistent with several previous
977
978 380 studies which reported additive effects of selected estrogens on different biological models such as
979
980 381 mammalian cells (Ghisari and Bonefeld-Jorgensen, 2009; Heneweer et al., 2005) or *in vitro* fish
981
982 382 cells (Le Page et al., 2006; Petersen and Tollefsen, 2011) and *in vivo* in fish (Brian et al., 2005;
983
984 383 Brion et al., 2012). Furthermore, our results demonstrate for the first time the suitability of the
985
986 384 ZELH-zfER cell line to investigate mixture effects of ER agonists at the receptor level in a zebrafish
987
988 385 cell context.
989
990

991 386 The screening for anti-estrogenic activity showed that some inhibiting chemicals active on
992
993 387 ZELH-zfER cells were present at effective concentrations in M1 and M2, e.g. TPP and
994
995 388 propiconazole. Although the underlying mechanism of ER inhibition remains unclear, the subgroup
996
997 389 mixtures of inhibiting chemicals had additive effects in ZELH α and ZELH β 2 cells, in all co-
998
999 390 exposure scenario, i.e. with inactive and/or estrogenic chemicals. In case of M1, a decreased
1000

1004
1005
1006 391 luciferase activity was also observed in ZELH cells, well predicted by the additive effects of TPP
1007
1008 392 and propiconazole (Figure SI-4). These results indicate that the inhibition observed in ZELH-zfERs
1009
1010 393 cells for M1 may involve non-ER specific luciferase inhibition.
1011
1012

1013 394 Interestingly, we observed in ZELH β 2 cells that the addition of the inhibiting chemicals to
1014
1015 395 the binary mixture of BPA and genistein resulted in a decrease in the expected estrogenic response
1016
1017 396 to a similar level as observed in the 12-component mixtures M1 and M2. In case of M1, the presence
1018
1019 397 of inhibiting chemicals silenced entirely the estrogenic activity expected, whereas in M2, only the
1020
1021 398 efficacy of the response was decreased. To a lesser extent, a similar trend was observed for M2 in
1022
1023 399 ZELH α cells. The experimental approach consisting of testing ER activating and inhibiting
1024
1025 400 chemicals separately and then together allowed us to evidence the role of inhibiting chemicals in
1026
1027 401 the deviation from expected additivity of genistein and BPA in ZELH β 2 cells. The experimental
1028
1029 402 results from the stepwise testing approach demonstrate that the response to the 12-chemical
1030
1031 403 mixtures in each bioassay can entirely be explained by the individual responses of the 12 chemicals.
1032
1033

1034 1035 404 *4.3. Differences between zebrafish and human-based bioassay responses*

1036
1037

1038 405 Our results highlight marked differences between human and zebrafish cells responses. Each
1039
1040 406 cell line displays cell-specific features, such as co-activator recruitment or metabolic capacities. For
1041
1042 407 instance, ZELH cells originate from zebrafish liver cells and have retained some metabolic
1043
1044 408 capacities qualitatively similar to zebrafish hepatocytes but distinct from MELN cells (Le Fol et al.,
1045
1046 409 2015), which may have played a role in the specific response to inhibiting chemicals in our study.
1047
1048 410 Indeed, metabolism has been previously suggested to negatively influence the response to xeno-
1049
1050 411 estrogen mixtures in rainbow trout hepatocytes (Petersen and Tollefsen, 2011) and in the E-
1051
1052 412 SCREEN assay (Evans et al., 2012). The characterization of internal concentrations of chemicals
1053
1054 413 in MELN and ZELH-zfER cells would be needed to estimate the influence of metabolism on the
1055
1056 414 xeno-estrogen response.
1057
1058
1059
1060
1061
1062

1063
1064
1065 415 To further investigate the relevance of the estrogenic mixture response in fish, both M1 and
1066
1067 416 M2 were tested on transgenic zebrafish embryos expressing GFP under control of *cyp19a1b*
1068
1069 417 promoter in radial glial cells in the EASZY assay (Brion et al., 2012). Indeed, in previous studies,
1070
1071 418 we showed that ZELH-zfER response profile to individual chemicals or environmental samples was
1072
1073 419 better correlated than the MELN assay with *in vivo* estrogenic activity measured in the EASZY
1074
1075 420 assay (Neale et al., 2017b; Sonavane et al., 2016). As a result, no estrogenic activity was measured
1076
1077 421 for both M1 and M2 mixtures because of a high embryo mortality, especially for M1 (Altenburger
1078
1079 422 et al., 2018). Thus, we could not confirm *in vitro* combined effects in zebrafish *in vivo*.
1080

1081 1082 1083 423 **4.4. Implication for quantifying the estrogenic activity of samples** 1084

1085
1086 424 A consistent body of literature exist regarding the assessment of additivity of xeno-estrogens
1087
1088 425 according to CA. However, very few studies investigated the robustness and validity of CA model
1089
1090 426 in more complex and realistic mixture scenarios. In the current study, the main factors
1091
1092 427 differentiating zebrafish and human ER response to M1 and M2 was the presence of inhibiting
1093
1094 428 chemicals that had higher influence on zfER activation in zebrafish cells. This agrees well with the
1095
1096 429 findings of Ihara et al. (2014) that evidenced that anti-estrogenic activity in wastewater treatment
1097
1098 430 plant extracts was a key factor to explain the different estrogenic activity measured in human and
1099
1100 431 medaka ER α transactivation *in vitro*.
1101

1102
1103 432 The 12-component mixtures were designed to mimic a simplified scenario of environmental
1104
1105 433 surface water contamination. To assess whether the mixture context would have influenced the
1106
1107 434 quantification of estrogenic activity mediated by xeno-estrogens, the mixture results were used to
1108
1109 435 quantify estradiol-equivalents (E2-Eq) in each bioassay (Table SI-4). Overall, M2 was predicted to
1110
1111 436 be more estrogenic (mean E2-Eq > 10 μ M) than M1 (mean E2-Eq < 1 μ M). In MELN cells, the
1112
1113 437 estrogenicity of M1 and M2 was almost not affected by the mixture context: the ratio of observed
1114
1115 438 to predicted E2-Eq was close to 1 for both mixtures. In contrast, ZELH α and ZELH β 2 responses to
1116
1117 439 xeno-estrogens in this specific mixture scenario were more susceptible to co-occurrence of
1118

1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180

440 inhibiting chemicals: the estrogenic activity was underestimated in M1 and M2, whenever
441 quantified. In case of ZELH β 2 cells, similar IC20 were derived for both M1 and M2, however, the
442 inhibiting chemicals abolished the estrogenic response in case of M1, while they only partially
443 decreased the maximal efficacy level in case of M2, without altering significantly the EC20
444 measured. These results suggest the presence of a balance between estrogenic and ER inhibiting
445 chemicals which can influence the detection, and thus the quantification, of xeno-estrogens in
446 ZELH β 2 cells.

447 **5. CONCLUSION**

448 In summary, this study demonstrates that BPA and genistein had additive effects *in vitro* in
449 zebrafish bioassays, comforting their use to assess combined effects of xeno-estrogens. In addition,
450 we show that the distinct responses of zebrafish and human-based bioassays to a 12-component
451 mixture were due to newly identified ER inhibiting chemicals selectively active in ZELH α and
452 ZELH β 2 cells (e.g. TPP, propiconazole) and altering zfER response to xeno-estrogens. In the
453 context of water bio-monitoring, this study illustrates the need for a mindful consideration of the
454 bioassay specificities (e.g. fish vs human ER, cell context) to ensure a proper interpretation of
455 results, as environmental chemicals may interfere with ER response, positively or negatively, in a
456 cell-specific manner.

457 **ACKNOWLEDGMENT**

458 This work was supported by the EU Seventh Framework Programme as a part of SOLUTIONS
459 project (FP7-ENV-2013-two-stage) under grant agreement number 603437, and by the French
460 Ministry of Ecology (P190-Ecotoxicologie, P181-DRC50). We wish to thank Emmanuelle Maillot-
461 Maréchal for her precious technical help with the cell cultures.

462 **DECLARATIONS OF INTEREST**

463 The authors declare that no conflict of interest regarding the publication of this paper.

1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239

464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499

AUTHOR CONTRIBUTIONS:

H.S., M.S., R.A., W.B., H.B., F.B and S.A conceived and designed the experiments; H.S. has performed the experiments; H.S. and M.S. analysed the data; H.S., M.S. and S.A. have written the manuscript; all authors have read and approved the final manuscript.

REFERENCES

Altenburger, R., Scholze, M., Busch, W., Escher, B.I., Jakobs, G., Krauss, M., Krüger, J., Neale, P.A., Ait-Aissa, S., Almeida, A.C., Seiler, T.-B., Brion, F., Hilscherová, K., Hollert, H., Novák, J., Schlichting, R., Serra, H., Shao, Y., Tindall, A., Tolefsen, K.-E., Umbuzeiro, G., Williams, T.D., Kortenkamp, A., 2018. Mixture effects in samples of multiple contaminants – An inter-laboratory study with manifold bioassays. *Environ. Int.* 114, 95–106. <https://doi.org/10.1016/j.envint.2018.02.013>

Auld, D.S., Inglese, J., 2004. Interferences with Luciferase Reporter Enzymes, in: Sittampalam, G.S., Coussens, N.P., Nelson, H., Arkin, M., Auld, D., Austin, C., Bejcek, B., Glicksman, M., Inglese, J., Iversen, P.W., Li, Z., McGee, J., McManus, O., Minor, L., Napper, A., Peltier, J.M., Riss, T., Trask, O.J., Weidner, J. (Eds.), *Assay Guidance Manual*. Eli Lilly & Company and the National Center for Advancing Translational Sciences, Bethesda (MD).

Balaguer, P., Francois, F., Comunale, F., Fenet, H., Boussioux, A.M., Pons, M., Nicolas, J.C., Casellas, C., 1999. Reporter cell lines to study the estrogenic effects of xenoestrogens. *Sci. Total Environ.* 233, 47–56. [https://doi.org/10.1016/S0048-9697\(99\)00178-3](https://doi.org/10.1016/S0048-9697(99)00178-3)

Brian, J.V., Harris, C.A., Scholze, M., Backhaus, T., Booy, P., Lamoree, M., Pojana, G., Jonkers, N., Runnalls, T., Bonfà, A., Marcomini, A., Sumpter, J.P., 2005. Accurate Prediction of the Response of Freshwater Fish to a Mixture of Estrogenic Chemicals. *Environ. Health Perspect.* 113, 721–728. <https://doi.org/10.1289/ehp.7598>

Brion, F., Le Page, Y., Piccini, B., Cardoso, O., Tong, S.-K., Chung, B., Kah, O., 2012. Screening Estrogenic Activities of Chemicals or Mixtures In Vivo Using Transgenic (cyp19a1b-GFP) Zebrafish Embryos. *Plos One* 7, e36069. <https://doi.org/10.1371/journal.pone.0036069>

Busch, W., Schmidt, S., Kuehne, R., Schulze, T., Krauss, M., Altenburger, R., 2016. Micropollutants in European rivers: A mode of action survey to support the development of effect-based tools for water monitoring. *Environ. Toxicol. Chem.* 35, 1887–1899. <https://doi.org/10.1002/etc.3460>

Coleman, M.D., O’Neil, J.D., Woehrling, E.K., Ndunge, O.B.A., Hill, E.J., Menache, A., Reiss, C.J., 2012. A Preliminary Investigation into the Impact of a Pesticide Combination on Human Neuronal and Glial Cell Lines In Vitro. *PLoS ONE* 7. <https://doi.org/10.1371/journal.pone.0042768>

1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298

- 500 Cosnefroy, A., Brion, F., Maillot-Maréchal, E., Porcher, J.-M., Pakdel, F., Balaguer, P., Ait-Aïssa, S.,
501 2012. Selective activation of zebrafish estrogen receptor subtypes by chemicals by using
502 stable reporter gene assay developed in a zebrafish liver cell line. *Toxicol. Sci. Off. J. Soc.*
503 *Toxicol.* 125, 439–449. <https://doi.org/10.1093/toxsci/kfr297>
- 504 Escher, B.I., Allinson, M., Altenburger, R., Bain, P.A., Balaguer, P., Busch, W., Crago, J., Denslow,
505 N.D., Dopp, E., Hilscherova, K., Humpage, A.R., Kumar, A., Grimaldi, M., Jayasinghe, B.S.,
506 Jarosova, B., Jia, A., Makarov, S., Maruya, K.A., Medvedev, A., Mehinto, A.C., Mendez, J.E.,
507 Poulsen, A., Prochazka, E., Richard, J., Schifferli, A., Schlenk, D., Scholz, S., Shiraish, F.,
508 Snyder, S., Su, G., Tang, J.Y.M., van der Burg, B., van der Linden, S.C., Werner, I.,
509 Westerheide, S.D., Wong, C.K.C., Yang, M., Yeung, B.H.Y., Zhang, X., Leusch, F.D.L., 2014.
510 Benchmarking Organic Micropollutants in Wastewater, Recycled Water and Drinking
511 Water with In Vitro Bioassays. *Environ. Sci. Technol.* 48, 1940–1956.
512 <https://doi.org/10.1021/es403899t>
- 513 Evans, R.M., Scholze, M., Kortenkamp, A., 2012. Additive Mixture Effects of Estrogenic Chemicals
514 in Human Cell-Based Assays Can Be Influenced by Inclusion of Chemicals with Differing
515 Effect Profiles. *PLOS ONE* 7, e43606. <https://doi.org/10.1371/journal.pone.0043606>
- 516 Ghisari, M., Bonefeld-Jorgensen, E.C., 2009. Effects of plasticizers and their mixtures on estrogen
517 receptor and thyroid hormone functions. *Toxicol. Lett.* 189, 67–77.
518 <https://doi.org/10.1016/j.toxlet.2009.05.004>
- 519 Heneweer, M., Muusse, M., van den Berg, M., Sanderson, J.T., 2005. Additive estrogenic effects of
520 mixtures of frequently used UV filters on pS2-gene transcription in MCF-7 cells. *Toxicol.*
521 *Appl. Pharmacol.* 208, 170–177. <https://doi.org/10.1016/j.taap.2005.02.006>
- 522 Hotchkiss, A.K., Rider, C.V., Blystone, C.R., Wilson, V.S., Hartig, P.C., Ankley, G.T., Foster, P.M.,
523 Gray, C.L., Gray, L.E., 2008. Fifteen Years after “Wingspread”—Environmental Endocrine
524 Disruptors and Human and Wildlife Health: Where We are Today and Where We Need to
525 Go. *Toxicol. Sci.* 105, 235–259. <https://doi.org/10.1093/toxsci/kfn030>
- 526 Ihara, M., Ihara, M.O., Kumar, V., Narumiya, M., Hanamoto, S., Nakada, N., Yamashita, N.,
527 Miyagawa, S., Iguchi, T., Tanaka, H., 2014. Co-occurrence of Estrogenic and Antiestrogenic
528 Activities in Wastewater: Quantitative Evaluation of Balance by in Vitro ER α Reporter Gene
529 Assay and Chemical Analysis. *Environ. Sci. Technol.* 48, 6366–6373.
530 <https://doi.org/10.1021/es5014938>
- 531 Ihara, M., Kitamura, T., Kumar, V., Park, C.-B., Ihara, M.O., Lee, S.-J., Yamashita, N., Miyagawa, S.,
532 Iguchi, T., Okamoto, S., Suzuki, Y., Tanaka, H., 2015. Evaluation of Estrogenic Activity of
533 Wastewater: Comparison Among In Vitro ER α Reporter Gene Assay, In Vivo Vitellogenin
534 Induction, and Chemical Analysis. *Environ. Sci. Technol.* 49, 6319–6326.
535 <https://doi.org/10.1021/acs.est.5b01027>
- 536 Kase, R., Javurkova, B., Simon, E., Swart, K., Buchinger, S., Könemann, S., Escher, B.I., Carere, M.,
537 Dulio, V., Ait-Aïssa, S., Hollert, H., Valsecchi, S., Polesello, S., Behnisch, P., di Paolo, C.,
538 Olbrich, D., Sychrova, E., Gundlach, M., Schlichting, R., Leborgne, L., Clara, M.,
539 Scheffknecht, C., Marneffe, Y., Chalon, C., Tusil, P., Soldan, P., von Danwitz, B., Schwaiger,
540 J., Palao, A.M., Bersani, F., Perceval, O., Kienle, C., Vermeirssen, E., Hilscherova, K.,

1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357

- 541 Reifferscheid, G., Werner, I., 2018. Screening and risk management solutions for steroidal
542 estrogens in surface and wastewater. *TrAC Trends Anal. Chem.* 102, 343–358.
543 <https://doi.org/10.1016/j.trac.2018.02.013>
- 544 Kjaerstad, M.B., Taxvig, C., Andersen, H.R., Nellemann, C., 2010. Mixture effects of endocrine
545 disrupting compounds in vitro. *Int. J. Androl.* 33, 425–433. <https://doi.org/10.1111/j.1365-2605.2009.01034.x>
- 547 Kjeldsen, L.S., Ghisari, M., Bonfeld-Jørgensen, E.C., 2013. Currently used pesticides and their
548 mixtures affect the function of sex hormone receptors and aromatase enzyme activity.
549 *Toxicol. Appl. Pharmacol.* 272, 453–464. <https://doi.org/10.1016/j.taap.2013.06.028>
- 550 Kojima, H., Takeuchi, S., Itoh, T., Iida, M., Kobayashi, S., Yoshida, T., 2013. In vitro endocrine
551 disruption potential of organophosphate flame retardants via human nuclear receptors.
552 *Toxicology* 314, 76–83. <https://doi.org/10.1016/j.tox.2013.09.004>
- 553 Kojima, H., Takeuchi, S., Van den Eede, N., Coyaci, A., 2016. Effects of primary metabolites of
554 organophosphate flame retardants on transcriptional activity via human nuclear
555 receptors. *Toxicol. Lett.* 245, 31–39. <https://doi.org/10.1016/j.toxlet.2016.01.004>
- 556 Könemann, S., Kase, R., Simon, E., Swart, K., Buchinger, S., Schlüsener, M., Hollert, H., Escher, B.I.,
557 Werner, I., Aït-Aïssa, S., Vermeirssen, E., Dulio, V., Valsecchi, S., Polesello, S., Behnisch, P.,
558 Javurkova, B., Perceval, O., Di Paolo, C., Olbrich, D., Sychrova, E., Schlichting, R., Leborgne,
559 L., Clara, M., Scheffknecht, C., Marneffe, Y., Chalon, C., Tušil, P., Soldà, P., von Danwitz,
560 B., Schwaiger, J., San Martín Becares, M.I., Bersani, F., Hilscherová, K., Reifferscheid, G.,
561 Ternes, T., Carere, M., 2018. Effect-based and chemical analytical methods to monitor
562 estrogens under the European Water Framework Directive. *TrAC Trends Anal. Chem.* 102,
563 225–235. <https://doi.org/10.1016/j.trac.2018.02.008>
- 564 Kortenkamp, A., 2007. Ten Years of Mixing Cocktails: A Review of Combination Effects of
565 Endocrine-Disrupting Chemicals. *Environ. Health Perspect.* 115, 98–105.
566 <https://doi.org/10.1289/ehp.9357>
- 567 Kunz, P.Y., Kienle, C., Carere, M., Homazava, N., Kase, R., 2015. In vitro bioassays to screen for
568 endocrine active pharmaceuticals in surface and waste waters. *J. Pharm. Biomed. Anal.*
569 106, 107–115. <https://doi.org/10.1016/j.jpba.2014.11.018>
- 570 Le Fol, V., Aït-Aïssa, S., Cabaton, N., Dolo, L., Grimaldi, M., Balaguer, P., Perdu, E., Debrauwer, L.,
571 Brion, F., Zalko, D., 2015. Cell-specific biotransformation of benzophenone-2 and
572 bisphenol-s in zebrafish and human in vitro models used for toxicity and estrogenicity
573 screening. *Environ. Sci. Technol.* 49, 3860–3868. <https://doi.org/10.1021/es505302c>
- 574 Le Fol, V., Aït-Aïssa, S., Sonavane, M., Porcher, J.-M., Balaguer, P., Cravedi, J.-P., Zalko, D., Brion,
575 F., 2017. In vitro and in vivo estrogenic activity of BPA, BPF and BPS in zebrafish-specific
576 assays. *Ecotoxicol. Environ. Saf.* 142, 150–156.
577 <https://doi.org/10.1016/j.ecoenv.2017.04.009>
- 578 Le Page, Y., Scholze, M., Kah, O., Pakdel, F., 2006. Assessment of Xenoestrogens Using Three
579 Distinct Estrogen Receptors and the Zebrafish Brain Aromatase Gene in a Highly

1358
1359
1360 580 Responsive Glial Cell System. *Environ. Health Perspect.* 114, 752–758.
1361 581 <https://doi.org/10.1289/ehp.8141>
1362
1363 582 Matthews, J., Gustafsson, J.-Å., 2006. Estrogen receptor and aryl hydrocarbon receptor signaling
1364 583 pathways. *Nucl. Recept. Signal.* 4. <https://doi.org/10.1621/nrs.04016>
1365
1366 584 Medjakovic, S., Zoehling, A., Gerster, P., Ivanova, M.M., Teng, Y., Klinge, C.M., Schildberger, B.,
1368 585 Gartner, M., Jungbauer, A., 2014. Effect of Nonpersistent Pesticides on Estrogen Receptor,
1369 586 Androgen Receptor, and Aryl Hydrocarbon Receptor. *Environ. Toxicol.* 29, 1201–1216.
1370 587 <https://doi.org/10.1002/tox.21852>
1371
1372 588 Menuet, A., Pellegrini, E., Anglade, I., Blaise, O., Laudet, V., Kah, O., Pakdel, F., 2002. Molecular
1373 589 characterization of three estrogen receptor forms in zebrafish: Binding characteristics,
1374 590 transactivation properties, and tissue distributions. *Biol. Reprod.* 66, 1881–1892.
1375 591 <https://doi.org/10.1095/biolreprod66.6.1881>
1376
1377 592 Miyagawa, S., Lange, A., Hirakawa, I., Tohyama, S., Ogino, Y., Mizutani, T., Kagami, Y., Kusano, T.,
1378 593 Ihara, M., Tanaka, H., Tatarazako, N., Ohta, Y., Katsu, Y., Tyler, C.R., Iguchi, T., 2014.
1380 594 Differing species responsiveness of estrogenic contaminants in fish is conferred by the
1381 595 ligand binding domain of the estrogen receptor. *Environ. Sci. Technol.* 48, 5254–5263.
1382 596 <https://doi.org/10.1021/es5002659>
1383
1384 597 Mosmann, T., 1983. Rapid Colorimetric Assay for Cellular Growth and Survival - Application. *J.*
1385 598 *Immunol. Methods* 65, 55–63. [https://doi.org/10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4)
1386
1387 599 Navas, J.M., Segner, H., 2000. Modulation of trout 7-ethoxyresorufin-O-deethylase (EROD) activity
1388 600 by estradiol and octylphenol. *Mar. Environ. Res.* 50, 157–162.
1389
1390 601 Neale, P.A., Ait-Aïssa, S., Brack, W., Creusot, N., Denison, M.S., Deutschmann, B., Hilscherova, K.,
1391 602 Hollert, H., Krauss, M., Novak, J., Schulze, T., Seiler, T.-B., Serra, H., Shao, Y., Escher, B.I.,
1392 603 2015. Linking in Vitro Effects and Detected Organic Micropollutants in Surface Water Using
1393 604 Mixture-Toxicity Modeling. *Environ. Sci. Technol.* 49, 14614–14624.
1394 605 <https://doi.org/10.1021/acs.est.5b04083>
1395
1396 606 Neale, P.A., Altenburger, R., Ait-Aïssa, S., Brion, F., Busch, W., de Aragão Umbuzeiro, G., Denison,
1398 607 M.S., Du Pasquier, D., Hilscherová, K., Hollert, H., Morales, D.A., Novák, J., Schlichting, R.,
1399 608 Seiler, T.-B., Serra, H., Shao, Y., Tindall, A.J., Tollefsen, K.E., Williams, T.D., Escher, B.I.,
1400 609 2017a. Development of a bioanalytical test battery for water quality monitoring:
1401 610 Fingerprinting identified micropollutants and their contribution to effects in surface water.
1402 611 *Water Res.* 123, 734–750. <https://doi.org/10.1016/j.watres.2017.07.016>
1403
1404 612 Neale, P.A., Munz, N.A., Ait-Aïssa, S., Altenburger, R., Brion, F., Busch, W., Escher, B.I., Hilscherová,
1405 613 K., Kienle, C., Novák, J., Seiler, T.-B., Shao, Y., Stamm, C., Hollender, J., 2017b. Integrating
1406 614 chemical analysis and bioanalysis to evaluate the contribution of wastewater effluent on
1407 615 the micropollutant burden in small streams. *Sci. Total Environ.* 576, 785–795.
1408 616 <https://doi.org/10.1016/j.scitotenv.2016.10.141>
1409
1410 617 Ohtake, F., Takeyama, K., Matsumoto, T., Kitagawa, H., Yamamoto, Y., Nohara, K., Tohyama, C.,
1411 618 Krust, A., Mimura, J., Chambon, P., Yanagisawa, J., Fujii-Kuriyama, Y., Kato, S., 2003.

1417
1418
1419 619 Modulation of oestrogen receptor signalling by association with the activated dioxin
1420 620 receptor. *Nature* 423, 545–550. <https://doi.org/10.1038/nature01606>
1421
1422 621 Petersen, K., Tollefsen, K.E., 2011. Assessing combined toxicity of estrogen receptor agonists in a
1423 622 primary culture of rainbow trout (*Oncorhynchus mykiss*) hepatocytes. *Aquat. Toxicol.*
1424 623 *Amst. Neth.* 101, 186–195. <https://doi.org/10.1016/j.aquatox.2010.09.018>
1425
1426
1427 624 Schlotz, N., Kim, G.-J., Jäger, S., Günther, S., Lamy, E., 2017. In vitro observations and in silico
1428 625 predictions of xenoestrogen mixture effects in T47D-based receptor transactivation and
1429 626 proliferation assays. *Toxicol. In Vitro* 45, 146–157.
1430 627 <https://doi.org/10.1016/j.tiv.2017.08.017>
1431
1432 628 Scholze, M., Boedeker, W., Faust, M., Backhaus, T., Altenburger, R., Grimme, L.H., 2001. A general
1433 629 best-fit method for concentration-response curves and the estimation of low-effect
1434 630 concentrations. *Environ. Toxicol. Chem.* 20, 448–457.
1435
1436
1437 631 Skolness, S.Y., Blanksma, C.A., Cavallin, J.E., Churchill, J.J., Durhan, E.J., Jensen, K.M., Johnson, R.D.,
1438 632 Kahl, M.D., Makynen, E.A., Villeneuve, D.L., Ankley, G.T., 2013. Propiconazole inhibits
1439 633 steroidogenesis and reproduction in the fathead minnow (*Pimephales promelas*). *Toxicol.*
1440 634 *Sci. Off. J. Soc. Toxicol.* 132, 284–297. <https://doi.org/10.1093/toxsci/kft010>
1441
1442 635 Sonavane, M., 2015. Intérêt d'une approche combinants bioessais in vitro et in vivo chez le poisson
1443 636 zèbre pour l'identification de perturbateurs endocriniens dans l'environnement
1444 637 aquatique. PhD dissertation, AgroParisTech, Paris.
1445
1446 638 Sonavane, M., Creusot, N., Maillot-Maréchal, E., Péry, A., Brion, F., Aït-Aïssa, S., 2016. Zebrafish-
1447 639 based reporter gene assays reveal different estrogenic activities in river waters compared
1448 640 to a conventional human-derived assay. *Sci. Total Environ.* 550, 934–939.
1449 641 <https://doi.org/10.1016/j.scitotenv.2016.01.187>
1450
1451
1452 642 Sumpster, J.P., 2005. Endocrine disrupters in the aquatic environment: An overview. *Acta*
1453 643 *Hydrochim. Hydrobiol.* 33, 9–16. <https://doi.org/10.1002/aheh.200400555>
1454
1455 644 Tibshirani, R.J., Efron, B., 1993. An introduction to the bootstrap. *Monogr. Stat. Appl. Probab.* 57,
1456 645 1–436.
1457
1458 646 Tohyama, S., Miyagawa, S., Lange, A., Ogino, Y., Mizutani, T., Tatarazako, N., Katsu, Y., Ihara, M.,
1459 647 Tanaka, H., Ishibashi, H., Kobayashi, T., Tyler, C.R., Iguchi, T., 2015. Understanding the
1460 648 molecular basis for differences in responses of fish estrogen receptor subtypes to
1461 649 environmental estrogens. *Environ. Sci. Technol.* 49, 7439–7447.
1462 650 <https://doi.org/10.1021/acs.est.5b00704>
1463
1464
1465 651 Yang, R., Li, N., Rao, K., Ma, M., Wang, Z., 2015. Combined action of estrogen receptor agonists
1466 652 and antagonists in two-hybrid recombinant yeast in vitro. *Ecotoxicol. Environ. Saf.* 111,
1467 653 228–235. <https://doi.org/10.1016/j.ecoenv.2014.09.025>
1468
1469 654 Zacharewski, 1997. In Vitro Bioassays for Assessing Estrogenic Substances. *Environ. Sci. Technol.*
1470 655 31, 613–623. <https://doi.org/10.1021/es960530o>
1471
1472
1473
1474
1475

1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534

656 **TABLES AND FIGURES**

657

658 **Table 1:** Overview of mixtures and their abbreviations tested on four different cell lines. More
 659 details about the composition of the mixtures are provided in the Supplementary Information
 660 (Tables SI 1-3). ⁽¹⁾ published in Altenburger et al; ⁽²⁾ corresponds to cyprodinil which was the only
 661 ER inhibitor.

662

mixture	ER activation		ER inhibition	
	M1	M2	M1	M2
MELN				
activators	M1_A _{MELN}	M2_A _{MELN}	-	-
inhibitors	-	-	M1_I _{MELN} ⁽²⁾	-
activators + inhibitors	M1_A+I _{MELN}	-	M1_A+I _{MELN}	-
activators + inhibitors + inactives	M1 ⁽¹⁾	M2 ⁽¹⁾	M1	-
ZELHα				
activators	M1_A _{ZELHα}	M2_A _{ZELHα}	-	-
inhibitors	-	-	M1_I _{ZELHα}	M2_I _{ZELHα}
activators + inhibitors	M1_A+I _{ZELHα}	M2_A+I _{ZELHα}	M1_A+I _{ZELHα}	M2_A+I _{ZELHα}
activators + inhibitors + inactives	M1 ⁽¹⁾	M2 ⁽¹⁾	M1	M2
ZELHβ2				
activators	M1_A _{ZELHβ2}	M2_A _{ZELHβ2}	-	-
inhibitors	-	-	M1_I _{ZELHβ2}	M2_I _{ZELHβ2}
activators + inhibitors	M1_A+I _{ZELHβ2}	M2_A+I _{ZELHβ2}	M1_A+I _{ZELHβ2}	M2_A+I _{ZELHβ2}
activators + inhibitors + inactives	M1 ⁽¹⁾	M2 ⁽¹⁾	M1	M2
ZELH				
inhibitors	-	-	M1_I _{ZELH}	M2_I _{ZELH}
inhibitors + inactives	-	-	M1	M2

663

Table 2: ER activation (EC20) and inhibition (IC20) of 12 test substances in MELN, ZELH α , ZELH β 2 and ZELH cells. Results are expressed in EC20 (activation) or IC20 (inhibition) are expressed in M concentration. E2 and OH-TAM were the positive control substances for ER activation and inhibition, respectively. Data originate from at least 2 independent experiments done in triplicates. Chemicals were tested in the 0.01 – 30 \times 10⁻⁶ M range, except for genistein (from 10⁻⁹ M). All concentration-response data are presented in SI-1 and SI-2.

	ER activation (EC20)			ER inhibition (IC20)			
	MELN mean (95% CI)	ZELH α mean (95% CI)	ZELH β 2 mean (95% CI)	MELN mean (95% CI)	ZELH α mean (95% CI)	ZELH β 2 mean (95% CI)	ZELH mean (95% CI)
E2	3.4 \times 10 ⁻¹² (2.6 \times 10 ⁻¹² - 4.3 \times 10 ⁻¹²)	1.3 \times 10 ⁻¹⁰ (1.1 \times 10 ⁻¹⁰ - 1.6 \times 10 ⁻¹⁰)	6.0 \times 10 ⁻¹² (4.74 \times 10 ⁻¹² - 7.7 \times 10 ⁻¹²)	-	-	-	-
OH-TAM	-	-	-	5.2 \times 10 ⁻⁹ (4.5 \times 10 ⁻⁹ - 6.0 \times 10 ⁻⁹)	1.8 \times 10 ⁻⁹ (9.4 \times 10 ⁻¹⁰ - 3.4 \times 10 ⁻⁹)	1.9 \times 10 ⁻⁹ (1.4 \times 10 ⁻⁹ - 2.8 \times 10 ⁻⁹)	> 3 \times 10 ⁻⁵
Bisphenol A	1.2 \times 10 ⁻⁷ (8.2 \times 10 ⁻⁸ - 1.7 \times 10 ⁻⁷)	2.1 \times 10 ⁻⁶ (1.3 \times 10 ⁻⁶ - 3.6 \times 10 ⁻⁶)	5.0 \times 10 ⁻⁶ (2.4 \times 10 ⁻⁶ - 6.1 \times 10 ⁻⁶)	> 3 \times 10 ⁻⁵	2.02 \times 10 ⁻⁵ (1.1 \times 10 ⁻⁵ - 3.6 \times 10 ⁻⁵)	8.8 \times 10 ⁻⁶ (8.7 \times 10 ⁻⁷ - 1.3 \times 10 ⁻⁵)	> 3 \times 10 ⁻⁵
Genistein	1.21 \times 10 ⁻⁸ (6.0 \times 10 ⁻⁹ - 2.9 \times 10 ⁻⁸)	1.4 \times 10 ⁻⁶ (9.5 \times 10 ⁻⁷ - 1.9 \times 10 ⁻⁶)	1.5 \times 10 ⁻⁸ (6.9 \times 10 ⁻⁹ - 3.1 \times 10 ⁻⁸)	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵
Diazinon	1.5 \times 10 ⁻⁵ (1.2 \times 10 ⁻⁵ - 1.9 \times 10 ⁻⁵)	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵
Triphenylphosphate	4.1 \times 10 ⁻⁶ (2.9 \times 10 ⁻⁶ - 5.7 \times 10 ⁻⁶)	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	8.0 \times 10 ⁻⁶ (3.2 \times 10 ⁻⁷ - 1.3 \times 10 ⁻⁵)	1.7 \times 10 ⁻⁶ (8.3 \times 10 ⁻⁷ - 3.5 \times 10 ⁻⁶)	1.1 \times 10 ⁻⁵ (3.0 \times 10 ⁻⁷ - 1.3 \times 10 ⁻⁵)
Benzo(a)pyrene	5.7 \times 10 ⁻⁷ (4.6 \times 10 ⁻⁷ - 7.2 \times 10 ⁻⁷)	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	4.2 \times 10 ⁻⁶ (2.5 \times 10 ⁻⁶ - 7.3 \times 10 ⁻⁶)	1.4 \times 10 ⁻⁶ (7.7 \times 10 ⁻⁷ - 2.4 \times 10 ⁻⁶)	> 3 \times 10 ⁻⁵
Benzo(b)fluorantene	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	1.95 \times 10 ⁻⁶ (1.1 \times 10 ⁻⁶ - 3.4 \times 10 ⁻⁶)	1.5 \times 10 ⁻⁶ (5.4 \times 10 ⁻⁷ - 4.1 \times 10 ⁻⁶)	1.8 \times 10 ⁻⁶ (7.2 \times 10 ⁻⁷ - 4.4 \times 10 ⁻⁶)
Chlorophene	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	1.0 \times 10 ⁻⁵ (2.6 \times 10 ⁻⁶ - 1.7 \times 10 ⁻⁵)	6.2 \times 10 ⁻⁶ (3.4 \times 10 ⁻⁶ - 9.8 \times 10 ⁻⁶)	> 1 \times 10 ⁻⁵
Propiconazole	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	8.1 \times 10 ⁻⁶ (3.1 \times 10 ⁻⁶ - 1.9 \times 10 ⁻⁵)	4.4 \times 10 ⁻⁶ (2.6 \times 10 ⁻⁶ - 7.7 \times 10 ⁻⁶)	2.4 \times 10 ⁻⁶ (3.7 \times 10 ⁻⁷ - 1.4 \times 10 ⁻⁵)
Cyprodinil	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	4.9 \times 10 ⁻⁶ (3.0 \times 10 ⁻⁶ - 8.1 \times 10 ⁻⁶)	2.0 \times 10 ⁻⁶ (1.2 \times 10 ⁻⁶ - 3.4 \times 10 ⁻⁶)	4.2 \times 10 ⁻⁶ (1.4 \times 10 ⁻⁶ - 1.3 \times 10 ⁻⁵)	4.1 \times 10 ⁻⁶ (2.6 \times 10 ⁻⁶ - 1.6 \times 10 ⁻⁵)
Triclosan	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵
Diuron	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵
Diclofenac	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵	> 3 \times 10 ⁻⁵

1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616

Table 3: Observed and predicted ER activation and inhibition for mixture M1 and its subgroups in MELN, ZELH α and ZELH β 2 cells. All concentrations are in M. (-) not tested as none of the individual compounds showed activity below its cytotoxic concentration range. (n.a.): the calculation is not applicable. Star indicates statistical significance (p<0.05). ^(a) re-calculated from Altenburger et al., 2018; ^(b) corresponds to cyprodinil which was the only ER inhibitor; ^(c) above cytotoxic concentration range.

Cell line	Mixture (name)	ER activation (EC20)			ER inhibition (IC20)		
		Observed Mean (95% CI)	Predicted Mean (95% CI)	Ratio obs/pred	Observed Mean (95% CI)	Predicted Mean (95% CI)	Ratio obs/pred
MELN	M1_A _{MELN}	1.2×10^{-6} (9.3×10^{-7} - 1.6×10^{-6})	8.9×10^{-7} (5.9×10^{-7} - 1.3×10^{-6})	1.3	-	-	-
	M1_I _{MELN}	-	-	-	4.9×10^{-6} ^(b) (3.0×10^{-6} - 8.2×10^{-6})	4.9×10^{-6} ^(b) (3.0×10^{-6} - 8.2×10^{-6})	1
	M1_A+I _{MELN}	2.1×10^{-6} (1.5×10^{-6} - 2.9×10^{-6})	2.6×10^{-6} (1.7×10^{-6} - 3.8×10^{-6})	0.81	$> 2 \times 10^{-5}$ ⁽⁴⁾	8.3×10^{-5} (5.0×10^{-5} - 1.4×10^{-4})	n.a.
	M1	6.1×10^{-6} ^(a) (3.9×10^{-6} - 9.2×10^{-6})	6.7×10^{-6} ^(a) (4.4×10^{-6} - 9.5×10^{-6})	0.91	3.4×10^{-5} (1.1×10^{-5} - 1.0×10^{-4})	5.9×10^{-4} ^(c) (3.6×10^{-4} - 9.8×10^{-4})	0.058*
ZELH α	M1_A _{ZELHα}	8.2×10^{-7} (6.5×10^{-7} - 1.6×10^{-6})	2.0×10^{-6} (1.0×10^{-6} - 3.0×10^{-6})	0.41	-	-	-
	M1_I _{ZELHα}	-	-	-	2.7×10^{-6} (1.9×10^{-6} - 3.6×10^{-6})	3.1×10^{-6} (1.2×10^{-6} - 1.2×10^{-5})	0.87
	M1_A+I _{ZELHα}	$> 4 \times 10^{-5}$ ^(c)	2.1×10^{-4} (1.3×10^{-4} - 3.2×10^{-4})	n.a.	4.2×10^{-6} (1.9×10^{-6} - 9.5×10^{-6})	5.7×10^{-6} (2.4×10^{-6} - 2.3×10^{-5})	0.74
	M1	$> 10^{-5}$ ^(c)	3.0×10^{-4} (1.8×10^{-4} - 4.6×10^{-4})	n.a.	4.2×10^{-6} (2.0×10^{-6} - 8.7×10^{-6})	4.4×10^{-6} (1.7×10^{-6} - 1.7×10^{-5})	0.95
ZELH β 2	M1_A _{ZELHβ2}	8.6×10^{-8} (3.7×10^{-8} - 1.8×10^{-7})	1.2×10^{-7} (5.5×10^{-8} - 2.4×10^{-7})	0.71	-	-	-
	M1_I _{ZELHβ2}	-	-	-	2.9×10^{-6} (2.0×10^{-6} - 4.0×10^{-6})	3.5×10^{-6} (2.1×10^{-6} - 5.1×10^{-6})	0.83
	M1_A+I _{ZELHβ2}	$> 2 \times 10^{-5}$ ^(c)	1.3×10^{-5} (5.8×10^{-6} - 2.5×10^{-5})	n.a.	4.4×10^{-6} (3.0×10^{-6} - 6.3×10^{-6})	6.4×10^{-6} (4.0×10^{-6} - 9.0×10^{-6})	0.69
	M1	$> 3 \times 10^{-5}$ ^(c)	1.8×10^{-5} (8.1×10^{-6} - 3.5×10^{-5})	n.a.	3.7×10^{-6} (2.1×10^{-6} - 6.3×10^{-6})	5.0×10^{-6} (3.0×10^{-6} - 7.1×10^{-6})	0.74

Table 4: Observed and predicted ER activation and inhibition for mixture M2 and its subgroups. All concentrations are in M. (-) not tested as none of the individual compounds showed activity below its cytotoxic concentration range. (n.a.): the calculation is not applicable. Star indicates statistical significance ($p < 0.05$). ^(a) re-calculated from Altenburger et al., 2018; ^(b) maximal induction measured below 20%.

Cell line	Mixture (name)	ER activation (EC20)			ER inhibition (IC20)		
		Observed Mean (95% CI)	Predicted Mean (95% CI)	Ratio obs/pred	Observed Mean (95% CI)	Predicted Mean (95% CI)	Ratio obs/pred
MELN	M2_A _{MELN}	1.6×10^{-7} (8.2×10^{-8} - 2.9×10^{-7})	6.4×10^{-8} (4.0×10^{-8} - 9.5×10^{-8})	2.5	-	-	-
	M2	1.5×10^{-7} ^(a) (6.8×10^{-8} - 2.8×10^{-7})	2.08×10^{-7} ^(a) (1.3×10^{-7} - 3.3×10^{-7})	0.72	-	-	-
ZELH α	M2_A _{ZELHα}	1.1×10^{-6} (7.4×10^{-7} - 1.7×10^{-6})	2.0×10^{-6} (1.2×10^{-6} - 3.1×10^{-6})	0.55	-	-	-
	M2_I _{ZELHα}	-	-	-	6.7×10^{-6} (2.9×10^{-6} - 1.3×10^{-5})	6.1×10^{-6} (2.2×10^{-6} - 1.1×10^{-5})	1.1
	M2_A+I _{ZELHα}	1.5×10^{-6} (7.8×10^{-7} - 2.8×10^{-6})	4.9×10^{-6} (3.0×10^{-6} - 7.5×10^{-6})	0.31*	7.6×10^{-6} (5.3×10^{-6} - 1.0×10^{-5})	1.0×10^{-5} (3.7×10^{-6} - 1.7×10^{-5})	0.76
	M2	$> 1.5 \times 10^{-7}$ ^(b)	6.6×10^{-6} (4.0×10^{-6} - 1.0×10^{-5})	n.a.	8.3×10^{-6} (6.0×10^{-6} - 1.1×10^{-5})	1.4×10^{-5} (5.3×10^{-6} - 2.4×10^{-5})	0.59
ZELH β 2	M2_A _{ZELHβ2}	1.1×10^{-7} (3.3×10^{-8} - 3.2×10^{-7})	1.5×10^{-7} (7.0×10^{-8} - 3.0×10^{-7})	0.73	-	-	-
	M2_I _{ZELHβ2}	-	-	-	7.5×10^{-6} (5.3×10^{-6} - 1.0×10^{-5})	6.6×10^{-6} (1.7×10^{-6} - 8.2×10^{-6})	1.1
	M2_A+I _{ZELHβ2}	1.2×10^{-6} (2.9×10^{-7} - 4.5×10^{-6})	3.7×10^{-7} (1.7×10^{-7} - 7.3×10^{-7})	3.2	7.7×10^{-6} (2.1×10^{-6} - 1.8×10^{-5})	6.8×10^{-6} (1.8×10^{-6} - 8.6×10^{-6})	1.1
	M2	1.8×10^{-6} (3.2×10^{-7} - 6.6×10^{-6})	5.0×10^{-7} (2.3×10^{-7} - 9.8×10^{-7})	3.6	4.1×10^{-6} (3.2×10^{-6} - 5.1×10^{-6})	9.2×10^{-6} (2.4×10^{-6} - 1.2×10^{-5})	0.44

1658
1659
1660 **Figure 1:** Experimental approach selected to study the combined effects of ER activating and
1661 inhibiting chemicals within the 12-component mixtures.
1662
1663
1664
1665




Figure 2: Predicted and measured effects of multi-component mixtures based on M1 concentration ratios. Data represent the mean (+/- SD) of a minimum of 3 independent experiments done in triplicates and pooled together. The green line represents CA prediction for ER activation and the orange line ER inhibition, and their respective dotted line represent the 95% CI belt. Cytotoxic concentrations (measured by MTT) were removed.





Figure 3: Predicted and measured effects of multi-component mixtures based on M2 concentration ratios. Data represent the mean (+/- SD) of a minimum of 3 independent experiments done in triplicates and pooled together. The green line represents CA prediction for ER activation and the orange line ER inhibition, inhibition, and their respective dotted line represent the 95% CI belt. Cytotoxic concentrations (measured by MTT) were removed.



1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893

SUPPLEMENTARY INFORMATION

Combined effects of environmental xeno-estrogens within multi-component mixtures: comparison of *in vitro* human- and zebrafish-based estrogenicity bioassays

Hélène Serra^{1,2}, Martin Scholze³, Rolf Altenburger⁴, Wibke Busch⁴, Hélène Budzinski², François Brion¹, Selim Aït-Aïssa^{1,*}

¹Institut National de l'Environnement Industriel et des risques (INERIS), Unité Ecotoxicologie *in vitro* et *in vivo*, UMR-I 02 SEBIO, 60550 Verneuil-en-Halatte, France

² UMR-CNRS EPOC/LPTC, Université de Bordeaux, Talence, France

³Brunel University London, Uxbridge, United Kingdom

⁴UFZ– Helmholtz Centre for Environmental Research, Leipzig, Germany

*Corresponding author. Email: selim.ait-aissa@ineris.fr

Content:

Table SI 1: Composition of the 12 compound mixtures M1 and M2 and the highest substance concentration tested *in vitro*.

Table SI 2: Composition of mixtures of ER activator (M1_A), ER inhibitors (M1_I) or combined ER activators and inhibitors (M1_A+I) tested in MELN, ZELH α , ZELH β 2 and ZELH cells.

Table SI 3: Composition of mixtures of ER activator (M2_A), ER inhibitors (M2_I) or combined ER activators and inhibitors (M2_A+I) tested in MELN, ZELH α , ZELH β 2 and ZELH cells.

Table SI 4: Estrogenic activity of the M1 and M2 expressed in estradiol equivalent.

Figure SI 1: Response of the 12 chemicals on ER activation in MELN, ZELH α and ZELH β 2 cells.

Figure SI 2: Response of the 12 chemicals on E2-induced ER inhibition in MELN, ZELH α , ZELH β 2 and ZELH cells.

Figure SI 3: Cyprodinil response in MELN, ZELH α , ZELH β 2 and ZELH cells.

Figure SI 4: Predicted and observed effects of inhibiting chemicals on ZELH cells.

Table SI 1: Composition of the 12 compound mixtures M1 and M2 and the highest substance concentration tested *in vitro*.

	M1		M2	
	Concentration (M)	proportion ¹⁾	Concentration (M)	proportion ¹⁾
Benzo(a)pyrene	6,00E-08	0,05%	9,47E-09	0,06%
Benzo(b)fluorantene	1,00E-07	0,08%	9,51E-09	0,06%
Bisphenol A	7,00E-07	0,58%	4,17E-06	27,70%
Chlorophene	9,00E-06	7,50%	6,40E-06	42,51%
Cyprodinil	1,00E-06	0,83%	1,87E-07	1,24%
Diazinon	6,00E-09	0,00%	1,96E-08	0,13%
Diclofenac	3,00E-05	24,99%	2,90E-06	19,26%
Diuron	6,00E-07	0,50%	2,08E-07	1,38%
Genistein	1,00E-07	0,08%	4,47E-07	2,97%
Propiconazole	6,00E-05	49,97%	8,48E-08	0,56%
Triphenylphosphate	1,50E-05	12,49%	2,32E-07	1,54%
Triclosan	3,50E-06	2,92%	3,89E-07	2,58%
Mixture	1.2E-4	100%	1.51E-5	100%

¹⁾ mixture composition according to Altenburger et al., (2018)

Table SI 2: Composition of mixtures of ER activator (M1_A), ER inhibitors (M1_I) or combined ER activators and inhibitors (M1_A+I) tested in MELN, ZELH α , ZELH β 2 and ZELH cells. The mixture composition is based on their relative proportion in the 12-compound mixture M1 (Table SI 1).

Type of mixture	MELN			ZELH α / ZELH β 2			ZELH
	Activators	Inhibitors	Inhibitors + activators	Activators	Inhibitors	Inhibitors + activators	Inhibitors
Mixture name	M1_A _{MELN}	M1_I _{MELN}	M1_A+I _{MELN}	M1_A _{ZELHα} , M1_A _{ZELHβ2}	M1_I _{ZELHα} , M1_I _{ZELHβ2}	M1_I+A _{ZELHα} , M1_I+A _{ZELHβ2}	M1_I _{ZELH}
Genistein	1%	-	1%	13%	-	0.2%	-
Bisphenol A	4%	-	4%	87%	-	0.8%	-
Triphenylphosphate	95%	-	89%	-	18%	17.7%	20%
Cyprodinil	-	100%	6%	-	-	-	1%
Diclofenac	-	-	-	-	-	-	-
Chlorophene	-	-	-	-	11%	10.6%	-
Propiconazole	-	-	-	-	71%	70.8%	79%
Total	100%	100%	100%	100%	100%	100%	100%

Table SI 3: Composition of mixtures of ER activator (M2_A), ER inhibitors (M2_I) or combined ER activators and inhibitors (M2_A+I) tested in MELN, ZELH α , ZELH β 2 and ZELH cells. The mixture composition is based on their relative proportion in the 12-compound mixture M2 (Table SI 1).

	MELN		ZELH α		ZELH β 2			ZELH
Type	Activators	Activators	Inhibitors	Inhibitors + activators	Activators	Inhibitors	Inhibitors + activators	Inhibitors
Name	M2_A _{MELN}	M2_A _{ZELHα}	M2_I _{ZELHα}	M2_I+A _{ZELHα}	M2_A _{ZELHβ2}	M2_I _{ZELHβ2}	M2_I+A _{ZELHβ2}	M2_I _{ZELH}
Genistein	10%	10%		4,0%	10%		4,0%	
Bisphenol A	90%	90%		37,2%	90%	38,6%	37,2%	
Triphenylphosphate			3,5%	2,0%		2,1%	2,1%	77%
Chlorophene			96,4%	56,6%		59,3%	57,1%	
Propiconazole								23%
Benzo(a)pyrene			0,14%	0,08%				
Total	100%	100%	100%	100%	100%	100%	100%	100%

2071
 2072
 2073
 2074
 2075
 2076
 2077
 2078
 2079
 2080
 2081
 2082
 2083
 2084
 2085
 2086
 2087
 2088
 2089
 2090
 2091
 2092
 2093
 2094
 2095
 2096
 2097
 2098
 2099
 2100
 2101
 2102
 2103
 2104
 2105
 2106
 2107
 2108
 2109
 2110
 2111
 2112
 2113
 2114
 2115
 2116
 2117
 2118
 2119
 2120
 2121
 2122
 2123
 2124
 2125
 2126
 2127
 2128
 2129


Table SI 4: Estrogenic activity of the M1 and M2 expressed in estradiol equivalent.

Estradiol-equivalents (E2-Eq, in μM) were calculated for the 12-component mixtures on the bases of their predicted and observed EC20s in relation to the EC20 of E2 (derived from all pooled control data). The E2-Eq(observed) is the ratio between the EC20(E2) and the regression-estimated EC20(mixture), and E2-Eq(predicted) is the ratio between the EC20(E2) and the CA predicted EC20(mixture). n.a.: not applicable (not estrogenic activity measured).










	M1 E2-Equivalent (μM)			M2 E2-Equivalent (μM)		
	Observed Mean	Predicted Mean	Ratio Observed/Predicted	Observed Mean	Predicted Mean	Ratio Observed/Predicted
MELN	0.56	0.51	1.1	22.7	16.3	1.39
ZELH α	n.a.	0.43	n.a.	n.a.	19.7	n.a.
ZELH β 2	n.a.	0.33	n.a.	3.33	12	0.278

2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170










Figure SI 1: Response of the 12 chemicals on ER activation in MELN, ZELH α and ZELH β 2 cells. Data represent each replicate and their mean (red dash) of at least 2 independent experiments done in triplicates. Chemicals were tested in the 10 nM - 30 μ M range, except for genistein (from 1 nM). 17 β -estradiol (E2) was used as positive control. The horizontal dotted line at 20% figures the threshold of effect.












2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211






2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240
2241
2242
2243
2244
2245
2246
2247
2248
2249
2250
2251
2252



2253
2254
2255
2256
2257
2258
2259
2260
2261
2262
2263
2264
2265
2266
2267
2268
2269
2270
2271
2272
2273
2274
2275
2276
2277
2278
2279
2280
2281
2282
2283
2284
2285
2286
2287
2288
2289
2290
2291
2292
2293




2294
2295
2296
2297
2298
2299
2300
2301
2302
2303
2304
2305
2306
2307
2308
2309
2310
2311
2312
2313
2314
2315
2316
2317
2318
2319
2320
2321
2322
2323
2324
2325
2326
2327
2328
2329
2330
2331
2332
2333
2334




2335
2336
2337
2338
2339
2340
2341
2342
2343
2344
2345
2346
2347
2348
2349
2350
2351
2352
2353
2354
2355
2356
2357
2358
2359
2360
2361
2362
2363
2364
2365
2366
2367
2368
2369
2370
2371
2372
2373
2374
2375


Figure SI 2: Response of the 12 chemicals on E2-induced ER inhibition in MELN, ZELH α , ZELH β 2 and ZELH cells. Data represent each replicate and the mean (red dash) of at least 2 independent experiments done in triplicates. Chemicals were tested in the 10 nM - 30 μ M range. MELN and ZELH β 2 cells were co-exposed with 0.1 nM E2, and ZELH α and ZELH cells with 1 nM E2. Cell viability (MTT) was measured for at least one experiment and is represented in green full circles (mean +/- SD). The horizontal dotted line at 80% figures the threshold of effect. Hydroxy-tamoxifen (OH-TAM) was used as positive control.




2376
2377
2378
2379
2380
2381
2382
2383
2384
2385
2386
2387
2388
2389
2390
2391
2392
2393
2394
2395
2396
2397
2398
2399
2400
2401
2402
2403
2404
2405
2406
2407
2408
2409
2410
2411
2412
2413
2414
2415
2416




2417
2418
2419
2420
2421
2422
2423
2424
2425
2426
2427
2428
2429
2430
2431
2432
2433
2434
2435
2436
2437
2438
2439
2440
2441
2442
2443
2444
2445
2446
2447
2448
2449
2450
2451
2452
2453
2454
2455
2456
2457



2458
2459
2460
2461
2462
2463
2464
2465
2466
2467
2468
2469
2470
2471
2472
2473
2474
2475
2476
2477
2478
2479
2480
2481
2482
2483
2484
2485
2486
2487
2488
2489
2490
2491
2492
2493
2494
2495
2496
2497
2498



2499
 2500
 2501 **Figure SI 3: Cyprodinil response in MELN, ZELH α , ZELH β 2 and ZELH cells.** The
 2502 response was measured with cyprodinil alone (ER, luciferase induction relative to DMSO
 2503 control) or in presence of E2 (antiER, luciferase induction relative to E2 positive control). Data
 2504 represent the mean (\pm SD) of a minimum of 2 independent experiments done in triplicates and
 2505
 2506
 2507
 2508
 2509
 2510 pooled together.



2558
2559
2560
2561
2562
2563
2564
2565
2566
2567
2568
2569
2570
2571
2572
2573
2574
2575
2576
2577
2578
2579
2580
2581
2582
2583
2584
2585
2586
2587
2588
2589
2590
2591
2592
2593
2594
2595
2596
2597
2598
2599
2600
2601
2602
2603
2604
2605
2606
2607
2608
2609
2610
2611
2612
2613
2614
2615
2616

Figure SI 4: Predicted and observed effects of inhibiting chemicals on ZELH cells. Results of subgroup mixtures M1_I_{ZELH} (A), M2_I_{ZELH} (B), and 12-component mixtures M1 (B) and M2 (D). Mixture effects were predicted according to CA model (orange line, 95% CI belt). Luciferase (LUC) activity was measured in absence (black circles) or in presence of E2 (co-exposure with E2 at 1 nM, grey open circles). The data (mean +/- SD) originate from at least 2 independent experiments done in triplicates and pooled together. Cytotoxic concentrations (measured by MTT) were removed.

