

1 **(a) Article Title:** Interpreting measures of fundamental movement skills and their
2 relationship with health-related physical activity and self-concept

3 **(b) Authors names:** Stuart Jarvis¹, Morgan Williams¹, Paul Rainer¹, Eleri Sian Jones², John
4 Saunders³, Richard Mullen¹

5

6 **Affiliations:**

7 ¹Faculty of Life Sciences and Education, University of South Wales, Pontypridd, CF37-1DL,
8 United Kingdom

9 ²School of Sport Health and Exercise Science, Bangor University, Bangor, LL57-2PZ,
10 United Kingdom

11 ³School of Exercise Science (QLD), Australian Catholic University, Brisbane, Australia

12

13 **(c) Running Head:** FMS AND HEALTH RELATED PHYSICAL ACTIVITY

14

15 **(d) Future Correspondence:** Stuart Jarvis, Faculty of Life Sciences and Education,
16 University of South Wales, Pontypridd, CF37-1DL, United Kingdom.

17 Email: stuart.jarvis@southwales.ac.uk

18

19 **Abstract**

20 The aims of this study were to determine proficiency levels of fundamental movement skills
21 (FMS) using cluster analysis in a cohort of UK primary school children; and to further
22 examine the relationships between FMS proficiency and other key aspects of health-related
23 physical activity behaviour. Participants were 553 primary children aged between 9 and 12,
24 294 boys and 259 girls, who were assessed across eight different FMS. Physical activity
25 behaviours included markers of physical fitness, recall of physical activity behaviour and
26 physical self-concept. Hierarchical cluster analysis was used to classify groups based on FMS
27 proficiencies and discriminant analysis to predict FMS proficiency based upon the physical
28 activity variables. This interpretation of FMS performance revealed distinct groups of FMS
29 proficiency in both genders with several gender specific components of physical activity
30 shown to discriminate children with differing levels of FMS proficiency ($p < .05$, $r > .40$).

31 *Keywords:* Fundamental movement skills, physical activity, self-concept, children.

32

33

34

35

36

37

38

39

40

41

42

43 **Introduction**

44 Despite compelling evidence that both the physical fitness and health status of children
45 and adolescents are substantially enhanced by regular physical activity, it is still unclear why
46 some youth are more physically active than others (Stodden & Holfelder, 2013). In response,
47 the concept of physical literacy has emerged in contemporary sport development policy and
48 practice (Lloyd, Colley, & Tremblay, 2010). However, despite the efforts of Whitehead
49 (2010) and others the lack of clarity in the current models used to operationalise the
50 theoretical concept (Giblin, Collins, & Button, 2014) has led to physical literacy in many
51 programmes being operationalised as the development of physical competency and often just
52 as fundamental movement skills (FMS; Keegan, Keegan, Daley, Ordway, & Edwards, 2013).
53 Although physical competency is recognised as an important dimension of physical literacy
54 (Whitehead, 2010) the exact balance of movement capacities (i.e., fundamental, combined
55 and complex) required to attain physical competency has yet to be clearly expressed (Giblin
56 et al., 2014). Despite this lack of conceptual clarity, FMS are viewed as the building blocks
57 for more complex motor skills and patterns and represent the underlying performance
58 competencies required for adequate participation in many forms of physical activity (Cliff,
59 Okely, Smith, & McKeen, 2009). FMS are common motor activities comprised of a series of
60 observable movement patterns, consisting of locomotor skills (e.g., run, hop and jump),
61 manipulative skills (e.g., catch, throw and kick), and stability skills (e.g., static and dynamic
62 balance; Gallahue & Donnelly, 2003). Acquiring proficiency in FMS during childhood has
63 been suggested as a vital component of children's physical, cognitive and social
64 development, (Malina, 2009).

65 Over the past decade, an overall decline in both children's motor skill performance and
66 physical activity has been reported (Hardy, Barnett, Espinel, & Okely, 2013). The underlying
67 explanations for this decline are unclear (Tompsett, Burkett, & McKean, 2014), and the

68 causes are clearly multidimensional in nature. One potential obstacle to an increased
69 understanding of this decline may be linked to our interpretation of FMS proficiency, an
70 accurate evaluation of which is critical for assessing and shaping pedagogical decisions for
71 enhancing physical literacy in children. Researchers have attempted to address this issue
72 through the use of standardized means for calculating individual item scores. Thus, several
73 studies have calculated a total score for each individual FMS skill, based on a criterion of
74 *mastery* if all components of the skill are demonstrated, *near mastery* if one component is
75 absent and *poor* if two or more components are not evident from a set number of trials (e.g.,
76 Van Beurden, Zask, Barnett, & Dietrich, 2002). A number of FMS scoring systems focus
77 either on distinct categories of motor competencies such as locomotor skills, object control
78 skills or use a combination of categories to aggregate FMS scores. For example, catching and
79 throwing, are summarized as an object control score, and presented as a single result (e.g.,
80 Cohen, Morgan, Plotnikoff, Barnett, & Lubans 2015). However, grouping skills into these
81 distinct categories may mask some individual skill performance with the result that
82 inadequacies in specific movement skills that require greater focus can go unnoticed by
83 practitioners.

84 As a result, Giblin et al. (2014) suggested that more research was required to refine the
85 procedures used in assessing and classifying FMS to enable more accurate interpretation of
86 the results obtained and greater effectiveness in their use in promoting skill proficiency. More
87 recently, Barnett, Miller, Laukkanen, and Morgan (2016) emphasised the need for FMS
88 assessment to accurately identify specific FMS deficits in individuals and contribute to the
89 provision of a learning environment that is developmentally appropriate, This may, for
90 example, necessitate that an individual FMS be learnt and practiced initially in a closed
91 environment (e.g., without the influence of other skills or such pressures as competition and
92 outcome scores), before being integrated with other FMS within a more advanced learning

93 environment (e.g., sport specific contexts). Given such suggestions, this study used cluster
94 analysis, as a means to categorize individuals that displayed similar characteristics, when
95 taking into account the full range of skills measured. This analysis enables a necessary
96 discrimination to be made between individuals who may have registered a similar aggregate
97 score, but one achieved across a very different range of skills.

98 In addition to investigating effective means for assessing overall FMS proficiency, this
99 research also focused on the relationship between FMS proficiency and other aspects of
100 children's physical activity behaviour that form the building blocks of physical literacy.
101 Stodden and colleagues' (2008) spiral model of engagement-disengagement in physical
102 activity points towards a dynamic and reciprocal relationship between FMS competence and
103 physical activity behaviours in mid childhood (ages 8 to 10) years) and onwards towards
104 adolescence. They advocate that in this developmental model it is important to substantiate
105 which variables of health-related physical activity (i.e., physical fitness, physical self-
106 concept, physical activity, and weight status) have the potential to impact FMS performance
107 as any future intervention to promote and sustain health outcomes should have a clear
108 strategy to address all of these elements.

109 In other literature, it has been suggested that a significant inverse association exists
110 between FMS proficiency and both weight status (Cliff, Okely, & Magarey, 2011) and
111 cardio-respiratory fitness (Hardy et al., 2013). It has also been suggested that muscular
112 strength is critical for successful FMS development and performance (Behringer, Vom
113 Heede, Matthews, & Mester, 2011). Physical self-concept (i.e., an individual's perception of
114 his/her own physical competence) has been shown to be an important correlate of FMS
115 proficiency in children (Robinson, 2011). Further, Barnett, Van Beurden, Morgan, Brooks,
116 and Beard (2008b) suggested that children's physical activity behaviour may also be partially
117 attributed to their actual FMS competence.

118 Considering these issues and the potential importance of FMS as a means to both
119 understanding dimensions of children's physical literacy and explaining their lifelong
120 involvement with health-related physical activity, the purpose of the present study is to:
121 examine a more discriminating classification of FMS performance, and apply it to an
122 exploration of the relationships between FMS proficiency and other key aspects of physical
123 activity behaviour in a cohort of 9-12-year-old UK school children. It is hypothesised that
124 children with more proficient FMS profiles will demonstrate more favourable measures of the
125 associated physical activity variables.

126 Method

127 Participants and Settings

128 Following the granting of ethical approval, 591 children, aged between 9 and 12, from 18
129 schools in the South-East Wales region of the UK, attended the test centre; 553 complete data
130 sets were recorded comprising 294 males (M age = 10.9 years, SD = 0.62), and 259 females
131 (M age = 10.7 years, SD = 0.64). Parental consent and child assent were obtained for each
132 participant. All data were collected during normal school hours.

133 Instruments and Measures

134 **Fundamental movement skills.** FMS proficiencies were assessed using selected process
135 orient checklists taken from the Australian resource 'Get Skilled: Get Active' (NSW
136 Department of Education and Training, 2000). The resource includes checklists of skills from
137 all categories of FMS (locomotor, manipulative and stability; Gallahue & Donnelly, 2003)
138 and is valid for use with both children and adolescents. The checklist, contains eight
139 individual FMS, including four locomotor skills (run, vertical jump, side gallop, leap) three
140 manipulative skills (catch, overhand throw, kick) and one stability skill (static balance). The
141 reliability and validity of the skills and their components have been previously established
142 (Okely & Booth, 2000). Get Skilled: Get Active was preferred to other measures of FMS

143 (e.g., the TGMD-2; Ulrich, 2000) as it includes a stability component of FMS assessment and
144 is culturally acceptable for use with children in this population (Foweather, 2010).

145 **Anthropometry and physical fitness.** Anthropometric and physical fitness assessments
146 were conducted with the High Priority battery from the ALPHA (Assessing Levels of
147 Physical Activity and Fitness) Health-Related Fitness Test Battery for Children and
148 Adolescents Test Manual (Ruiz et al., 2011). The battery includes assessments of body
149 composition (weight, height, BMI), cardio-respiratory fitness (20m multi-stage test) and
150 musculoskeletal fitness (handgrip strength, standing long jump). In addition, the study
151 included a separate motor fitness measure the 20-metre sprint, which has previously been
152 reported to be a valid and reliable measure of speed in children (Morrow, Jackson, Disch, &
153 Mood, 2005).

154 **Physical activity.** The Physical Activity Questionnaire for Children (PAQ-C; Crocker,
155 Bailey, Faulkner, Kowalski, & McGrath, 1997) was used as an indicator of the children's
156 'typical' level of physical activity behaviour (cf. Welk & Eklund, 2005). The instrument uses
157 nine multiple choice questions to assess a child's physical activity over the previous seven
158 days. The PAQ-C has been shown to have adequate test-retest reliability (range: $r = 0.75 -$
159 0.82) and validity (range: $r = 0.45 - 0.53$; Crocker et al., 1997). The choice of the PAQ-C for
160 use with this population was based on a review of physical activity self-report measures by
161 Biddle, Gorley, Pearson, & Bull (2011), who supported its validity, reliability, and
162 practicality for use with children and adolescents. The instrument has also been
163 recommended by the ALPHA Health-Related Fitness Test Battery for Children and
164 Adolescents (Ruiz et al., 2011) for use with European samples of young people.

165 **Physical self-concept.** The Children and Youth Physical Self Perception Profile was used
166 to examine participants' perceptions of Global Self-Worth (GSW), Physical Self-Worth and
167 its sub-domains of Sports Competence (SC), Physical Conditioning (PC), Body

168 Attractiveness (BA) and Physical Strength (PS). Each scale is assessed by six items scored on
169 a four-point scale with the average score used to represent the value for the scale. Previous
170 work by Welk, Corbin, Dowell, and Harris (1997) and Welk and Eklund (2005) have
171 demonstrated adequate reliability and a good fit for the CY-PSPP measurement model. In
172 addition, Welk and Eklund also established that the instrument can be used in research with
173 children as young as nine years of age. As it has not been used with a population of children
174 from South-East Wales, we conducted a confirmatory factor analysis (CFA) of the CY-PSPP
175 to establish its utility for the present sample.

176 **Procedure**

177 Data were collected by an experienced FMS practitioner and a team of trained research
178 assistants. The FMS assessments were video recorded (Sony video camera, Sony, UK) and
179 analysed using performance analysis software (Studio Code, NSW, Australia) in accordance
180 with the 'Get Skilled: Get Active' guidelines. A process oriented checklist was used to
181 determine the total number of components performed correctly for each skill attempt and
182 analysed by the study author. If there was any uncertainty about whether a feature was
183 consistently present or not, it was checked as absent. For reliability of the FMS assessment
184 inter- and intra-rater reliability analysis was performed on a randomly selected sample of
185 completed FMS sets by the author and a second experienced FMS practitioner and
186 determined using linear weighted Kappa (Fleiss, Levin, & Paik, 2003). Physical fitness
187 assessments and data collection followed the procedures described in the High Priority
188 ALPHA Test Battery (Ruiz et al., 2011). The 20 metre sprint efforts followed the procedures
189 outlined by Oliver and Meyers (2009) and were recorded with Smart Speed dual beam
190 electronic timing gates (Fusion Sport, Queensland, Australia). The CY-PSPP and the PAQ-C
191 survey instruments were administered in a classroom at the test centre, to small groups (no
192 greater than 6 participants), of same gender. The purpose of both the survey instruments was

193 explained to the children and it was stressed that there were no right or wrong answers. Each
194 item in both the surveys was read to the children with research assistants circulating
195 throughout the room to provide extra assistance. Prior to administration of the CY-PSPP,
196 example items were provided and demonstrated to the participants based on Whitehead's
197 (1995) recommendations for its use with young children.

198 **Statistical Analysis**

199 **Confirmatory factor analysis.** The factorial validity of the CY-PSPP was examined
200 using CFA with the Mplus statistical programme (Muthen & Muthen, 2010). The
201 demographic variables used were gender; male ($n = 294$), female ($n = 259$) and group ($n =$
202 553). The CFA models were fitted for each group separately to test for configurable
203 invariance. Global model fit indices were examined at each stage of the CFA, along with
204 detailed assessment of the completely standardized factor loadings, the standardized
205 residuals, and the modification indices. All CFAs were conducted using the robust maximum
206 likelihood estimation procedure with a Satorra–correction ($S-B\chi^2$) and fit indices corrected
207 for robust estimation based on the recommendation of Hu and Bentler (1999) amongst others
208 who suggest that multiple criteria be used to evaluate the fit of the overall model to the data.
209 These fit indices, in addition to the normed chi-square test (χ^2), included the chi-square to
210 degrees of freedom ratio (χ^2/df), the Comparative Fit Index (CFI; Bentler, 1990), the Tucker-
211 Lewis Index (TLI), the Root Mean Square Error of Approximation (RMSEA; Steiger, 1990)
212 and the Standardized root mean square residual (SRMR; Bollen, 1989). Hu and Bentler's
213 (1999) recommendations for good fit were adopted, with a χ^2/df ratio of 3:1 or less indicating
214 good fit, and cut off values of 0.95 for CFI, 0.06 for RMSEA and 0.08 for SRMR. To
215 examine whether the CY-PSPP displayed equivalence of measures across genders, a
216 measurement invariance approach was employed via multi-group CFA. Measurement
217 invariance assessed invariance of construct, factor loading, item intercepts and error

218 variances in a hierarchical ordering with increased constraints from one model to the next. As
219 a result, a model is only tested if the previous model in the hierarchical ordering has been
220 shown to be equivalent across groups. In addition to the fit indices described above, the
221 Bayesian Information Criterion (BIC) and Akaike Information Criterion (AIC) indices were
222 used to indicate model fit.

223 **FMS group classification and proficiency.** Intra- and inter-rater reliability for all FMS
224 measures displayed a level of agreement that was good or above (Kw range = 0.68 to 0.93)
225 and (Kw range = 0.61 to 0.81) respectively, based on Altman's (1991) thresholds to describe
226 reliability. Data were split by gender and preliminary analyses confirmed these two groups
227 differed ($p < .05$). Ward's two-way hierarchical cluster analysis (JMP version 10.02; SAS
228 Institute, Marlow, UK) was used to classify groups based on the FMS item scores. The
229 number of clusters was determined at the point where the scree plot of the distance values
230 plateaued. To verify the classification analysis, differences between the clusters on total FMS
231 score were assessed using t-test for females (2 groups) and ANOVA for males (3 groups)
232 with Tukey's post hoc. To describe the features that best described the clusters, a decision
233 tree induction (DTI) method was used (Morgan, Williams, & Barnes, 2013). The DTI was
234 split then pruned to retain the r^2 minimising the likelihood of over fitting. Finally, the validity
235 of the model was assessed via inspection of the ROC curve, area under the curve and the
236 corresponding confusion matrix.

237 **Discriminant analysis.** Following FMS group classification, discriminant analysis was
238 used to examine which of the FMS groups scored more highly on the other physical activity
239 variables. Initial screening of dependent variables revealed non-normal distributions and
240 outlying cases were modified by assigning the outlying case(s) a raw score that was one unit
241 larger (or smaller) than the next most extreme score. The analysis was then reassessed to
242 confirm the assumptions corresponding to linearity, normality, multicollinearity and

243 heterogeneity of variance-covariance matrices. For the discriminant analysis, loadings > 0.4
244 were considered significant, based on Stevens' (1992) conservative recommendation. A
245 classification matrix was constructed to assess the predictive accuracy of the discriminant
246 functions using a proportional chance criterion of > 56% (Hair, Anderson, Tatham, & Black,
247 1998). Classification accuracy was examined using Press's Q statistic, compared to the χ^2
248 critical value of > 6.63.

249 **Results**

250 **Confirmatory Factor Analysis**

251 The results of analysis conducted to evaluate CY-PSPP measurement model fit are
252 presented in Table 1. A χ^2/df ratio of 3:1 or less is successfully demonstrated in each model.
253 The CFI indexes exceeded the 0.90 criterion, RMSEA values were below .06, and SRMR
254 were below .10, all indicating an adequate overall fit of the model. All questionnaire items
255 loaded onto their designated factors with non-zero loadings. Median loadings for the full
256 group, boys subsample and girls subsample were 0.76 (range = 0.59 – 0.92), 0.75 (range =
257 0.61 – 0.92) and 0.75 (range = 0.55 – 0.95), respectively. These findings suggest an adequate
258 fit for the CY-PSPP measurement model to these data and reasonable psychometric
259 properties. Inter correlations amongst sub domains signified zero cross loadings on all other
260 factors. In general, the correlations among the sub domains (SC, PC, BA, and PS) were
261 moderate to strong across the full group ($r = 0.57 - 0.93$), boys sub group ($r = 0.56 - 0.96$),
262 and girls sub group ($r = 0.51 - 0.93$). As expected, the sub domains demonstrated stronger
263 associations with the PSW than with GSW in all groups. The correlations between GSW and
264 PSW were higher than the correlations between GSW and the other CY-PSPP sub domains
265 for all groups.

266 Measurement invariance across boys and girls sub groups to evaluate the CY-PSPP factor
267 structure for gender sensitivity is shown in Table 1. The fit indices in Table 1 confirm an

268 excellent fit of the independent factor structure; Model 1 provides excellent multiple fit
269 indices to the data (χ^2/df , CFI index, RMSEA, SRMR, AIC/BIC value) indicating that the
270 factorial structure of the construct is equal across groups. As configural invariance was
271 supported, coefficients were then constrained to be equal to test for metric invariance. Model
272 2 has good fit indices; therefore, constraining the factor loading to be the same across the
273 groups. The scalar invariance model (Model 3) provided a good fit to the data as did the error
274 variance invariance model (Model 4). The overall goodness of fit indices and the tests of
275 differences in fit between adjacent models therefore support measurement invariance. Taken
276 together, the data provide supportive evidence for the validity of the CY-PSPP with this
277 population.

278 Insert Table 1 here

279 **FMS Classification**

280 **Boys.** Three groups emerged from the analysis; Low (total FMS = 18 ± 4), Intermediate
281 (total FMS = 25 ± 4), and High (total FMS = 31 ± 3) Proficiency. When total FMS scores for
282 these groups were compared, all the group means differed significantly, Low versus High =
283 13 (95% CI = 11-14); Low versus Intermediate = 7 (95% CI = 5-8), and Intermediate versus
284 High = 6 (95% CI = 5-7). Figure 1 shows the frequency distribution of FMS performance of
285 the cluster groups on each FMS. The final DTI model (Figure 2) had a total of seven splits.
286 From the column contributions, the FMS with the largest difference between the cluster
287 groups was vertical jump ($G^2 = 78.03$) followed by the overhand throw ($G^2 = 64.26$), then
288 leap ($G^2 = 31.19$). Side gallop ($G^2 = 23.06$), static balance ($G^2 = 18.58$) and the catch ($G^2 =$
289 18.49) also featured, but to a lesser extent. The FMS of run and kick made no contribution
290 between the groups. The high proficiency cluster demonstrated strongest performances for
291 the splits on vertical jump; overhand throw, static balance, catch and side gallop. The low
292 proficiency group were poor in the vertical jump and poorest in the splits of side gallop and

293 the leap. The intermediate proficiency group demonstrated lower performance than the high
294 proficiency group but better performance than the low group across all splits except for the
295 catch. In summary, whether the child scored high or not on vertical jump (first split),
296 subsequent skills identified the high proficiency cluster as being the most competent of the
297 groups across the identified splits.

298 **Girls.** Two groups (Low and High Proficiency) were identified. The Low and High
299 Proficiency group had total FMS scores of $21, \pm 4$ and $28, \pm 3$, which were significantly
300 different, mean difference = 6, 95% CI = 5-7. Figure 1 shows the frequency distribution of
301 scores of the two clusters on each FMS. Comparisons between the groups showed the high
302 proficiency group were the most proficient across all FMS. The final girls' DTI model
303 (Figure 2) had five splits ($r^2 = 0.48$) that differentiated between the two clusters. Static
304 balance ($G^2 = 84.36$) was the FMS variable with the largest contribution to the model. The
305 catch ($G^2 = 44.51$), vertical jump ($G^2 = 27.34$) and leap ($G^2 = 10.84$) followed but their
306 impact was much smaller. Run, side gallop, kick and overhand throw made no contribution
307 and did not feature in the final model. Girls who scored higher on the static balance and the
308 vertical jump demonstrated higher probability of being in the high cluster group. Girls who
309 scored lower on the static balance but higher on the catch, static balance and the leap splits
310 also demonstrated higher probability of being in the high cluster group. In contrast, the low
311 cluster group demonstrated poorer skill proficiency across all splits. In summary, whether
312 good performance was observed in static balance (first split), subsequent skills identified the
313 high proficiency group as being the most proficient.

314

Insert Figures 1 and 2 here

315 **Descriptive Statistics**

316 Descriptive statistics for male and female FMS groups for all the independent variables are
317 reported in Table 2. In boys, the high proficiency group demonstrated better performance

318 measures of physical fitness, physical activity recall and physical self-perception than both
319 the intermediate and low proficiency groups. The low group demonstrated the lowest
320 performance scores across all these measures. In girls, the high proficiency group
321 demonstrated higher scores on measures of physical fitness, physical activity recall and
322 physical self-perception than the low group.

323 Insert Table 2 here

324 **Discriminant Analysis**

325 **Boys.** Analysis revealed two discriminant functions. The first function explained 86.7% of
326 the variance, canonical $R^2 = 0.26$, whereas the second function explained only 13.3%,
327 canonical $R^2 = 0.05$. In combination, these discriminant functions significantly differentiated
328 the cluster groups, $\Lambda = 0.70$, $\chi^2(24) = 102.73$, $p < .001$; although removing the first function
329 indicated that the second function did not significantly differentiate the groups, $\Lambda = 0.95$, χ^2
330 $(11) = 15.27$, $p = 0.17$. Closer analysis of the discriminant loadings in Table 3, reveals that
331 Sprint, MSFT, SLJ and CY-PSPP Condition sub scale exceeded the criterion on the first
332 function (> 0.40). The discriminant function plot showed that the first function discriminated
333 the high group from the intermediate group and the low group. With 67.3% of the original
334 grouped cases correctly classified, the intermediate group were 87.2% correctly classified, the
335 high group were 34.2% and the low group were 29%, Press's $Q = 17.69$ (> 6.63), $p < 0.05$.
336 The classification ratio exceeds the proportional chance criterion of 56 % demonstrating
337 predictive accuracy of the discriminant function (Hair et al., 1998).

338 **Girls.** A single discriminant function that explained all the variance was identified,
339 canonical $R^2 = 0.14$. The function significantly differentiated the groups, $\Lambda = 0.86$, $\chi^2(12) =$
340 36.65 , $p < .001$. Closer analysis of the discriminant loadings in Table 3 revealed that Sprint,
341 SLJ, HG, PAQ-C, and MSFT, were significant predictors of group membership ($> .40$).
342 Classification results showed that 69.5 % of original grouped cases were correctly classified

343 (low group = 47.1%, high group = 84.1%, Press's $Q = 39.39 (> 6.63)$, $p < .05$. The
344 classification ratio exceeds the proportional chance criterion of 56 % demonstrating
345 predictive accuracy of the discriminant function (Hair, et al., 1998).

346 Insert Table 3 here

347 Discussion

348 The novel approach of using cluster analysis to examine FMS proficiency successfully
349 identified groups with different proficiency levels. In addition, discriminant analysis revealed
350 that FMS proficiency level could be predicted by a combination of several physical activity
351 related variables for both males and females. Specifically, these were cardio respiratory
352 fitness and lower body musculoskeletal strength in both boys and girls and upper body
353 musculoskeletal strength in girls. Physical activity recall was a significant predictor for girls,
354 whereas for boys, the physical condition subscale of the CY-PSPP was prominent.

355 For both boys and girls, FMS proficiency levels were low (based on similar reporting of
356 FMS proficiency in children) and not dissimilar to levels demonstrated in other UK based
357 studies with similar aged children (e.g., Foweather, 2010). This is concerning given the
358 importance placed on FMS in enhancing physical literacy and promoting health (Tompsett et
359 al., 2014). It is generally believed that most children should master the less complex FMS
360 (i.e., sprint run, vertical jump, catch, side gallop and over-arm throw) by age nine and more
361 complex FMS (i.e., leap and kick) by age ten (Hardy, King, Espinel, Cosgrove, & Bauman,
362 2010). Attainment of movement proficiency at this level is purported to form a foundation for
363 physical literacy, the absence of which might lead to activity avoidance and the associated
364 implications for health (Stodden et al., 2008). As highlighted earlier, it is the *interpretation* of
365 the FMS scores that may be important in revealing insights into children's FMS proficiency.
366 The classification method adopted in this study was effective in distinguishing group
367 membership and provides practitioners with more precise details of FMS proficiency that

368 avoids misclassification which in turn may help those children most in need of additional
369 support.

370 In addition to identifying FMS group membership and a more refined focus on FMS
371 ability with boys and girls it is also mindful to recognise FMS differentials that exist across
372 genders. In this study, it was shown that girls displayed poorer proficiency in specific
373 manipulative skills (i.e., overarm throw and kick) compared to boy's groups. These findings
374 support previous research in gender differentials across FMS (e.g., Hardy et al., 2013)
375 amongst others who suggest boys tend to possess higher proficiency in manipulative skills
376 than girls although this divide is not as clear within locomotor skills.

377 A subsidiary aim of this study was to directly test the factorial validity of the CY-PSPP.
378 For this population, CFA clearly supported the hierarchical structure of the CY-PSPP and
379 yielded a clean factor structure, supporting claims by Welk and Eklund (2005) that young
380 children can judge themselves differently according to the physical domain of their lives
381 being addressed.

382 The second major aim of the present study was to examine the relationship between FMS
383 proficiency and the potential impact of several key health related measures of physical
384 activity involvement (Stodden et al., 2008) at what has been suggested to be a critical
385 developmental age (Malina, 2009). In this study, discriminant analysis revealed that for both
386 boys and girls, measures of physical fitness were significant predictors of FMS proficiency.
387 More specifically in both genders, these measures included cardio respiratory fitness, the
388 sprint run and musculoskeletal fitness (i.e., upper body strength in girls and lower body
389 strength in boys and girls). A positive relationship between FMS ability and cardio
390 respiratory fitness levels has previously been demonstrated (Barnett, Van Beurden, Morgan,
391 Brooks, & Beard, 2008a; Okely, Booth, & Patterson, 2001). In addition, Hardy, Reinten-
392 Reynolds, Espinel, Zask, and Okely (2012) confirmed a clear and consistent association

393 between low competency in FMS and inadequate cardio-respiratory fitness in children.
394 Although this relationship appears robust, the directionality of this relationship is unclear. For
395 example, Cohen et al. (2015) have suggested that improvements in overall FMS competency
396 may act as a causal mechanism for physical activity behaviour change and subsequent
397 improvements in cardio respiratory fitness. Despite this uncertainty, promoting both FMS and
398 cardio respiratory fitness would seem to be beneficial for children.

399 Regarding musculoskeletal fitness Stodden, True, Langendorfer, and Gao (2013) have
400 suggested that a certain level of force production and force attenuation is needed to
401 proficiently perform many ballistic FMS (e.g., throwing, kicking, striking, jumping, running,
402 and leaping). Behringer et al. (2011) have identified that children showed greater training
403 induced gains in the skills of jumping, running, and throwing after a programme of strength
404 training. At present, levels of muscular fitness appear to be declining in UK children (Cohen
405 et al., 2011), which might negatively impact FMS proficiency as witnessed in this study.
406 Further, the development of strength is closely related to sprint performance, another
407 significant predictor in this study. This is consistent with the finding that the development of
408 sprint speed has been shown to be a distinguishing characteristic of successful participation in
409 physical activities in both children and adults (Hammami, Makhlouf, Chtara, Padulo, &
410 Chaouachi, 2015).

411 It is important to note here that BMI was not related to FMS performance in boys and
412 girls, which is consistent with the studies of Castelli and Valley (2007) and Hume et al.
413 (2008). However, these findings contrast with several studies that reported that elevated BMI
414 has a negative effect on FMS performance (Cliff et al., 2009; Okely, Booth, & Chey, 2004;
415 Southall, Okely, & Steele, 2004). Most apparent in these studies is the seemingly negative
416 relationship between BMI and locomotor FMS (e.g., run, hop, side gallop). Locomotor skills
417 may be more related to BMI than object control skills as these skills require more ‘whole

418 body' movement and transfer of body weight, and so are more difficult to perform given
419 overweight and obese childrens' increased overall mass (Okely et al., 2004). Okely and
420 colleagues (2004) suggested that the relationship between skill competence and being
421 overweight may be reciprocal. Therefore, although BMI might be an important measure in
422 terms of health and physical activity its actual relationship with FMS remains unclear and
423 further investigation is clearly needed.

424 In this study, it was shown that for girls, but not boys, involvement in physical activity
425 significantly discriminated between the FMS groups. Okely et al. (2001) and Raudsepp and
426 Pall (2006) have also found that the relationship between FMS and time in organised physical
427 activity outside of the school environment was stronger for girls than boys. A distinction
428 between organised (i.e., involving adult interventions such as in club sport and other
429 instructional activity) and non-organised activity did not form part of the present study.
430 Future research would benefit from differentiating between these types of activity.

431 The physical condition (PC) subscale of the CY-PSPP differentiated between the boys'
432 FMS groups. Physical condition represents the individual's perceptions regarding the level of
433 their physical condition, physical fitness, stamina, their ability to maintain exercise and how
434 confident they feel in the exercise and fitness setting. Spiller (2009) suggests that through
435 participation many boys learn that the optimal functionality and performance of their bodies
436 (i.e., physical condition) is more important than other facets such as appearance and
437 participation in physical activity, typically providing a better 'fit' for the development of
438 males' identity and FMS skill acquisition. In addition, Fowweather (2010) suggests that with
439 advancing age children are more able to make informed judgements about their level of
440 physical condition and so it is likely that the relationship between physical activity and motor
441 competence will strengthen in those with advanced levels of physical condition. No other
442 CY-PSPP subscales significantly predicted FMS proficiency in boys or girls.

443 The present study holds several limitations. The PAQ-C only assesses general levels of
444 physical activity for individuals in the school system. It does not provide an estimate of
445 frequency, time and intensity nor does it differentiate between organised and non-organised
446 activity. In addition, subjectivity, social desirability bias, and variable recall ability especially
447 in young people are considered limitations of the physical activity self-report instrument used
448 in this study. To increase the strength and accuracy of reported physical activity behaviour
449 Chinapaw, Mokkink, Van Poppel, Van Mechelen, and Terwee (2010) suggested that a
450 combination of self-report and accelerometry be adopted. Children's motivation during field
451 tests of physical fitness depends upon several factors such as motivation, understanding and
452 perceived success (Fairclough et al., 2016). For these reasons, the physical fitness test results
453 in this study should be interpreted with caution. The failure to confirm an association with
454 some of the associated physical activity involvement variables may be due to the more
455 conservative 0.40 cut off value used in the discriminant analysis of this study. While other
456 research has used 0.30 as the cut-off point, the authors believe based on Stevens' (1992)
457 suggestion that 0.40 is justified as it identifies only the key variables that contribute the most
458 to the discriminating function.

459 In summary, the novel interpretation of FMS performance in this study has provided
460 researchers with an alternative method of portraying FMS competence. Having the provision
461 to identify and specifically target the weakest FMS might better inform practitioners trying to
462 improve movement skills. The present study also identified gender-specific components of
463 physical activity that discriminate children with differing levels of FMS proficiency.

References

- 464
465 Altman, D. G. (1991). *Practical statistics for medical research*. Boca Raton, FL: Chapman &
466 Hall.
- 467 Barnett, L. M., Van Beurden, E., Morgan, P. J., Brooks, L. O., & Beard, J. R. (2008a). Does
468 childhood motor skill proficiency predict adolescent fitness? *Medicine and Science in*
469 *Sports and Exercise*, 40(12), 2137–2144. doi:10.1249/MSS.0b013e31818160d3.
- 470 Barnett, L. M., Van Beurden, E., Morgan, P. J., Brooks, L. O., & Beard, J. R. (2008b).
471 Perceived sports competence mediates the relationship between childhood motor skill
472 proficiency and adolescent physical activity and fitness: a longitudinal assessment.
473 *International Journal of Behavioural Nutrition and Physical activity*, 5, 40. doi:10.1186
474 /1479-5868-5-40
- 475 Barnett, L. M., Miller, A., Laukkanen, A., & Morgan, P. J. (2016). Fundamental movement
476 skills: An important focus. *Journal of Teaching in Physical Education*, 3(3), 219-225.
- 477 Behringer, M., Vom Heede, A., Matthews, M., & Mester, J. (2011). Effects of strength
478 training on motor performance skills in children and adolescents: a meta-analysis.
479 *Paediatric Exercise Science*; 23(2), 186-206.
- 480 Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological Bulletin*,
481 107, 238–246. doi:10.1037/0033-2909.107.2.238
- 482 Biddle, S., Gourley, T., Pearson, N., & Bull, F. C. (2011). An assessment of self-reported
483 physical activity instruments in young people for population surveillance: project ALPHA,
484 *International Journal of Behavioural Nutrition and Physical Activity*, 8, 1.
- 485 Bollen, K. A. (1989). A new incremental fit index for general structural equation models.
486 *Sociological Methods and Research*, 17(3), 303-316. doi: 10.1177/0049124189017003004
- 487 Castelli, D. M., & Valley, J. A. (2007). The relationship of physical fitness and motor
488 competence to physical activity. *Journal of Teaching in Physical Education*, 26(4),

- 489 358–374.
- 490 Chinapaw, M. M., Mokkink, L. B., Van Poppel, M. N., Van Mechelen, W., & Terwee, C. B.
491 (2010). Physical activity questionnaires for youth: A systematic review of measurement
492 properties. *Sports Medicine*, *40*, 539-563. doi:10.2165/11530770
- 493 Cliff, D. P., Okely, A. D., & Magarey, A. M. (2011). Movement skill mastery in a clinical
494 sample of overweight and obese children. *International Journal of Paediatric Obesity*,
495 *6*(5-6), 473-475. doi:10.3109/17477166.2011.575154
- 496 Cliff, D. P., Okely, A. D., Smith, L. M., & McKeen, K. (2009). Relationships between
497 fundamental movement skills and objectively measured physical activity in pre-school
498 children. *Paediatric Exercise Science*, *21*(4), 436-449.
- 499 Cohen, D., Voss, C., Taylor, M., Delextrat, A., Ogunleye, A., & Sandercock, G. (2011). Ten-
500 year secular changes in muscular fitness in English children. *Acta Paediatrica*, *100*(10),
501 175-177. doi:10.1111/j.1651-2227.2011.02318.x
- 502 Cohen, K. E., Morgan, P. J., Plotnikoff, R. C., Barnett, L. M., & Lubans, D. R. (2015).
503 Improvements in fundamental movement skill competency mediate the effect of the
504 SCORES intervention on physical activity and cardiorespiratory fitness in children.
505 *Journal of Sports Sciences*. doi:101080/02640414.2015.1017734
- 506 Crocker, P. R. E., Bailey, D. A., Faulkner, R. A., Kowalski, K., & McGrath, R. (1997).
507 Measuring general levels of physical activity: Preliminary evidence for the Physical
508 Activity Questionnaire for Older Children. *Medicine & Science in Sports & Exercise*,
509 *29*(10), 1344-1349.
- 510 Fairclough, S. J., McGrane, B., Sanders, G., Taylor, S., Owen, M., & Curry, W. (2016). A
511 non-equivalent group pilot trial of a school based physical activity and fitness intervention
512 for 10-11-year-old English children; born to move. *BMC Public Health*, *16*(1), 861.
513 doi:10.1186/s12889-016-3550-7.

- 514 Fleiss, J. L., Levin, B., & Paik, M. C. (2003) *Statistical methods for ratters and proportions*
515 (3rd ed.). Hoboken, NJ: John Wiley & Sons.
- 516 Foweather, L. (2010). The effect of interventions on fundamental movement skills, physical
517 activity and psychological well-being among children. Thesis (PhD). Liverpool John
518 Moores University
- 519 Gallahue, D. L., & Donnelly, F. C. (2003). *Developmentally physical education for all*
520 *children* (4th ed.). Champaign, IL: Human Kinetics
- 521 Giblin, S., Collins, D. & Button, C. (2014). Physical literacy: Importance, assessment and
522 future directions. *Sports Medicine*, 44(9), 1177-1184. doi:10.1007/s40279-014-0205-7
- 523 Hair, J. F., Anderson, R. E., Tatham, R. L., & Black, W. C. (1998). *Multivariate data*
524 *analysis* (5th ed.). Upper Saddle River, NJ: Prentice Hall.
- 525 Hammami, R., Makhlouf, I., Chtara, M., Padulo, J. & Chaouachi, A. (2015). The contribution
526 of vertical explosive strength to sprint performance in children. *Sports Sci Health*, 11(1),
527 37-42. doi: 10.1007/s11332-014-0200-2.
- 528 Hardy, L. L., Barnett, L., Espinel, P., & Okely, A.D., (2013). Thirteen-year trends in child
529 and adolescent fundamental movement skills: 1997-2010. *Medicine and Science in Sports*
530 *and Exercise*, 45(10), 1965-70. doi:10.1249/MSS.0b013e318295a9fc
- 531 Hardy, L., King, L., Espinel, P., Cosgrove, C., & Bauman, A., (2010). *NSW schools' physical*
532 *activity and nutrition survey*: Full report. Sydney: NSW Ministry of Health.
- 533 Hardy L. L., Reinten-Reynolds, T., Espinel, P., Zask, A., & Okely, A. D. (2012). Prevalence
534 and correlates of low fundamental movement skill competency in children. *Pediatrics*,
535 130(2), e390-e398. doi:10.1542/peds.2012-0345
- 536 Hu, L., & Bentler, P. M. (1999). Cut off criteria for fit indexes in covariance structure
537 analysis: Conventional criteria versus new alternatives. *Structural Equation Modelling*,
538 6(1), 1-55. doi:10.1080/10705519909540118

- 539 Hume, C., Okely, A., Bagley, S., Telford, A., Booth, M., Crawford, D., & Salmon, J.
540 (2008). Does weight status influence associations between children's fundamental
541 movement skills and physical activity. *Research Quarterly for Exercise and Sport*, 79; (2),
542 158–165. doi: 10.1080/02701367.2008.10599479
- 543 Keegan, R. J., Keegan, S. L., Daley, S., Ordway, C., & Edwards, A. (2013). Getting Australia
544 moving: Establishing a physically literate and active nation (Game Plan). Canberra,
545 Australia: University of Canberra.
- 546 Lloyd, M., Colley, R. C., & Tremblay, M. S. (2010). Advancing the debate on 'fitness testing
547 for children: Perhaps we're riding the wrong animal. *Paediatric Exercise Science*, 22(2),
548 176-182.
- 549 Malina, R. M. (2009). Children and adolescents in the sport culture: The overwhelming
550 majority to the select few. *Journal of Exercise Science Fitness*, 7(2), 1-10.
- 551 Morgan, S., Williams, M. D., & Barnes, C. (2013). Applying decision tree induction for
552 identification of important attributes in one-versus-one player interactions: A hockey
553 exemplar. *Journal of Sport Sciences*, 31(10), 1031-1037.
- 554 Morrow, J. R., Jackson, A. W., Disch, J. G., & Mood, D. P. (2005). Measurement and
555 evaluation in human performance (3rd ed.). Champaign, IL: Human Kinetics.
- 556 Muthen, L. K., & Muthen, B. O. (2010). Mplus user's guide (6th ed.). Los Angeles, CA:
557 Muthen & Muthen.
- 558 New South Wales Department of Education and Training (2000). *Get Skilled: Get Active*.
559 Sydney, Australia: New South Wales Department of Education and Training.
- 560 Okely, A. D., Booth, M., & Chey, T. (2004). Relationship between body composition and
561 fundamental movement skills among children and adolescents. *Research Quarterly for*
562 *Exercise and Sport*, 75(3), 238-247. doi:10.1080/02701367.2004.10609157
- 563 Okely, A. D., Booth, M., & Patterson, J. W. (2001). Relationship of physical activity to

- 564 fundamental movement skills among adolescents. *Medicine and Science in Sports and*
565 *Exercise*; 33(11), 1899-1904. doi:10.1097/00005768-200111000-00015
- 566 Okely, A. D., & Booth, M. L. (2000). The development and validation of an instrument to
567 assess children's fundamental movement skill ability. In 2000 *Pre-Olympic Congress*
568 *Book of Abstracts*, 245.
- 569 Oliver, J. L., & Meyers, R. W. (2009). Reliability and generality of measures of acceleration,
570 planned agility, and reactive agility. *International Journal of Sports Physiology and*
571 *Performance*, 4(3), 345-354.
- 572 Raudsepp, L., & PaII, P. (2006). The relationship between fundamental motor skills and
573 outside school physical activity of elementary school children. *Pediatric Exercise Science*,
574 18(4), 426-435.
- 575 Robinson, L. E. (2011). The relationship between perceived physical competence and
576 fundamental motor skills in preschool children. *Child Care Health Dev*, 37(4), 589-596.
- 577 Ruiz, J. R., Castro-Pinero, J., Espana-Romero, V., Artero, E., Ortega, F. B., & Cuenca, M. M.
578 (2011). Field-based fitness assessment in young people: the ALPHA health-related fitness
579 test battery for children and adolescents. *British Journal of Sports Medicine*, 45(6), 518-
580 524. doi:10.1136/bjism.2010.075341
- 581 Southall, J., Okely, A., & Steele, J. R. (2004). Actual and perceived physical competence in
582 overweight and non-overweight children. *Pediatric Exercise Science*, 16, 15-24.
- 583 Spiller, V. (2009). Young people's perceptions and experiences of their physical bodies.
584 Thesis (PhD). Victoria University.
- 585 Steiger, J. (1990). Structural model evaluation and modification: an interval estimation
586 approach. *Multivariate Behaviour Research*, 25(2), 173-180.
- 587 Stevens, J. P. (1992). *Applied multivariate statistics for the social sciences* (4thed.). Mahwah,
588 NJ: Lawrence Erlbaum Associates.

- 589 Stodden, D. F., Goodway, J. D., Langendorfer, S. J., Robertson, M. A., Rudisill, M. E., Garcia,
590 C., & Garcia, L. E. (2008). A developmental perspective on the role of motor skill
591 competence in physical activity: An emergent relationship. *QUEST*, *60*(2), 290-306.
- 592 Stodden, F. D., & Holfelder, B. (2013). No child left behind: The role of motor skill
593 development. *Zeitschrift fur Sportpsychologie*, *20*(1), 10-17. doi:10.1026/1612-5010
594 /a000088.
- 595 Stodden, F. D., True, L. K., Langendorfer, S. J., & Gao, Z. (2013). Associations among
596 selected motor skills and health related fitness: Indirect evidence for Seefeldt's proficiency
597 barrier in young adults. *Research Quarterly for Exercise and Sport*, *84*(3), 397-403.
- 598 Tompsett, C., Burkett, B., & McKean, M. R. (2014). Development of physical literacy and
599 movement competency: A literature review. *Journal of Fitness Research*, *3*(2), 53-74.
- 600 Ulrich, D. A. (2000). *Test of gross motor development. Examiner's Manual* (2nd Ed.). Austin,
601 Texas: Pro-ED. Inc.
- 602 Van Beurden, E., Zask, A., Barnett, L. M., & Dietrich, U. C. (2002). Fundamental movement
603 skills – how do primary school children perform? The 'Move it Grove it' program in rural
604 Australia. *Journal of Science and Medicine in Sport*, *5*(3), 244-252.
- 605 Welk, G. J., Corbin, C. B., Dowell, M. N., & Harris, H. (1997). The validity and reliability of
606 two different versions of the children and youth physical self-perception profile.
607 *Measurement in Physical Education and Exercise Science*, *1*, 163-177.
- 608 Welk, G. J., & Eklund, B. (2005). Validation of the children and youth physical self-
609 perceptions profile for young children. *Psychology of Sport and Exercise*, *6*(1), 51-65.
- 610 Whitehead, J. R. (1995). A study of children's physical self-perceptions using an adapted
611 physical self-perception profile questionnaire. *Paediatric Exercise Science*, *7*, 132-151.
- 612 Whitehead, M. (2010). *Physical literacy: Throughout the life course*. Abingdon, UK:
613 Routledge

Table 1. Measurement model (A) fit of CFA for the full group, male and female sub groups, and Measurement invariance (B) of the CY-PSPP factor structure.

Measurement Model (A)	<i>n</i>	SB- χ^2	χ^2	<i>df</i>	<i>P</i> <	CFI	TLI	RMSEA (90% CI)	SRMR
Full group	553	1362.507	2.35	579	0.001	0.950	0.898	0.048 (0.043-0.052)	0.038
Boys subgroup	294	920.885	1.59	579	0.001	0.906	0.898	0.047 (0.042-0.052)	0.059
Girls subgroup	259	1128.288	1.94	579	0.001	0.934	0.928	0.055 (0.050-0.061)	0.044

Invariance Model (B)	SB- χ^2	χ^2	<i>df</i>	<i>P</i> <	CFI	TLI	RMSEA	SRMR	BIC	AIC
Model 1	1297.741	-	579	0.001	0.900	0.892	0.046	0.051	46801.951	46264.242
Model 2	2084.538	797.74	1188	0.001	0.882	0.875	0.051	0.066	47198.541	46254.273
Model 3	2130.261	752.42	1218	0.001	0.880	0.876	0.051	0.067	47051.415	46238.295
Model 4	2256.413	717.38	1274	0.001	0.801	0.867	0.050	0.065	4694.312	46198.654

Note. CY-PSPP=Children and Youth Physical Self-Perception Profile; **SB- χ^2** : Satorra-Bentler scaled goodness of fit chi-square statistic; *df*: degrees of freedom for chi-square statistic; CFI: comparative fit index; TLI: Tucker-Lewis Index; RMSEA: Root mean squared error of approximation; 90% CI: 90% confidence interval of the point estimate; SRMR: Standardized root mean square residual; BIC: Bayesian Information Criterion; AIC: Akaike Information Criterion. Model 1: testing equivalence of measurement model across gender; Model 2: CFA analysis for Boys and Girls with measurement invariance of factor loadings; Model 3: CFA analysis for Boys and Girls of factor loadings and intercepts; Model 4: CFA analysis for Boys and Girls with measurement of factor loadings, intercepts, and residuals.

Table 2. Means and standard deviations of physical characteristics and performance measures for boys and girls FMS group

classification

Variables	Descriptive group data (mean \pm SD)						
	Boys				Girls		
	Total Group (<i>n</i> = 294)	Low Group (<i>n</i> = 31)	Inter. Group (<i>n</i> = 187)	High Group (<i>n</i> = 76)	Total Group (<i>n</i> = 259)	Low Group (<i>n</i> = 102)	High Group (<i>n</i> = 157)
BMI	18.5 \pm 2.9	19.5 \pm 4.9	18.4 \pm 2.7	18.2 \pm 2.3	19.1 \pm 3.1	19.07 \pm 3.43	19.03 \pm 2.81
SLJ (cm)	143 \pm 22	129 \pm 20.7	141 \pm 20.4	153 \pm 19.9	131 \pm 18	125 \pm 17.17	135 \pm 18.13
DHG (Kg)	18.5 \pm 3.4	17.7 \pm 3.1	18.1 \pm 3.4	19.8 \pm 3.3	17.1 \pm 3.3	16.17 \pm 3.57	17.74 \pm 3.01
MSFT (m)	821 \pm 400	506 \pm 339	773 \pm 360	1066 \pm 389	612 \pm 304	539 \pm 263	659 \pm 320
SPRINT (sec)	4.14 \pm 0.33	4.50 \pm 0.41	4.15 \pm 0.28	3.96 \pm 0.29	4.31 \pm 0.34	4.44 \pm 0.37	4.24 \pm 0.30
PAQ-C	3.44 \pm 0.65	3.06 \pm 0.71	3.46 \pm 0.64	3.53 \pm 0.58	3.22 \pm 0.65	3.06 \pm .065	3.33 \pm 0.63
CY-PSPP	18.91 \pm 3.03	17.32 \pm 3.38	18.90 \pm 2.94	19.60 \pm 2.88	18.0 \pm 3.11	17.49 \pm 3.00	18.29 \pm 3.14
CY-SC	3.16 \pm 0.65	2.85 \pm 0.78	3.14 \pm 0.64	3.31 \pm 0.54	2.97 \pm 0.65	2.85 \pm 0.63	3.04 \pm 0.65
CY-PC	3.14 \pm 0.63	2.76 \pm 0.70	3.11 \pm 0.60	3.36 \pm 0.70	2.98 \pm 0.65	2.86 \pm 0.64	3.06 \pm 0.65
CY-BA	2.95 \pm 0.75	2.72 \pm 0.89	2.97 \pm 0.76	2.99 \pm 0.74	2.79 \pm 0.75	2.73 \pm 0.74	2.82 \pm 0.75
CY-PS	2.91 \pm 0.68	2.71 \pm 0.71	2.89 \pm 0.68	3.04 \pm 0.65	2.75 \pm 0.65	2.68 \pm 0.61	2.80 \pm 0.67
CY-PSW	3.27 \pm 0.57	2.98 \pm 0.60	3.29 \pm 0.56	3.37 \pm 0.54	3.10 \pm 0.62	3.02 \pm 0.66	3.15 \pm 0.59
CY-GSW	3.50 \pm 0.50	3.31 \pm 0.64	3.50 \pm 0.48	3.53 \pm 0.49	3.39 \pm 0.55	3.34 \pm 0.55	3.42 \pm 0.55

Note. BMI = Body mass index; SLJ = Standing long jump; DHG = Dominant handgrip; MSFT = Multistage fitness test; PAQ-C = Physical activity questionnaire children; CY-PSPP = Children and youth physical self-perception profile; CY-PSPP- SC = Sport competence subscale; CY-PSPP –PC = Physical condition subscale; CY-PSPP –BA = Body attractiveness subscale; CY-PSPP –PS = Physical strength subscale; CY-PSPP –PSW = Physical self-worth subscale; CY-PSPP –GSW = Global self-worth subscale





Figure 1. Frequency distribution of boys and girls FMS skill components present via group classification on each FMS.

Boys



Girls




Figure 2. Final decision trees including the 7 splits for boys FMS groups (Level 1 = Low group; Level 2 = Intermediate group; Level 3 = High group) and the 5 splits for girls FMS groups (Level 1 = Low group; Level 2 = High group).