



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Ever Graze-Prime Lamb Production on Perennial Based Grasslands in Southwest Australia

Paul Sanford

*Department of Agriculture and Food Western Australia, Australia*

J. Ryan

*Department of Agriculture and Food Western Australia, Australia*

P. Jakobsson

*Department of Agriculture and Food Western Australia, Australia*

E. Dobbe

*Department of Agriculture and Food Western Australia, Australia*

E. McCready

*Department of Agriculture and Food Western Australia, Australia*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/9-2/7>

**The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.**

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Ever Graze-Prime lamb production on perennial based grasslands in southwest Australia

P.Sanford ,J.Ryan ,P.Jakobsson ,E.Dobbe , E.McCready

Western Australian Department of Agriculture and Food and Cooperative Research Centre for Future Farming Industries ,  
444 Albany Highway Albany Australia 6330 . E-mail : psanford@agric.wa.gov.au

**Key words :** perennials ,prime lamb ,farming systems ,simulation

**Introduction** Traditional livestock production in south west Australia is based on annual plants that have been shown to result in land degradation through salinization ,soil erosion ,waterlogging and soil acidification .Novel livestock systems including deep-rooted summer active perennial plants can significantly reduce land degradation . However the adoption of these new systems is dependent on improved and more profitable livestock production .This paper reports on the findings of a study investigating prime lamb production on perennial forages .

**Materials and methods** Computer simulations suggested that a Merino prime lamb production system based on summer-active perennials (future farming system) could substantially increase farm profit while reducing the risk of salinity (Sanford & Young 2005) .To test this system in 2005 a 60ha site was chosen in the Albany Hinterland catchment and sown to tall fescue (*Festuca arundinacea*) cv .Quantum Max P (16 ha) ,lucerne (*Medicago sativa*) cv .Sardi 10 (8 ha) ,kikuyu (*Pennisetum clandestinum*) cv .Whittet (18 ha) ,mixture of setaria (*Setaria sphacelata*) cv .Splenda and green panic (*Megathyrus maximus*) (3 ha) and chicory (*Cichorium intybus*) cv .Puna (15 ha) in spring .The soil at the site was sand over gravel/clay . Merino ewes were joined to Poll Dorset rams in March 2006 ,lambling commenced in early August ,lambs were weaned in early November and removed from the field site .Rotational grazing was employed except during lambing and periods when feed was low in which case the kikuyu pasture was set stocked .The ewes were in a feedlot from 7<sup>th</sup> April to 30<sup>th</sup> May and 24<sup>th</sup> November to 31<sup>st</sup> December Supplement consisted of 88% pellets (68% DMD ,14% CP) and 12% lupins (87% DMD ,34% CP) .

**Results and discussion** Annual rainfall for the site in 2006 was 290mm well below the long term average of 500mm .As a consequence of the drought conditions 559kg/ha of supplement was fed ,464kg/ha more than predicted for the future farming system (Table 1) .The reproductive efficiency achieved in the field was higher than the all of the simulated farming systems at 7.8 lambs weaned per ha (Table 1) .However low pasture growth meant lambs were turned off at weaning and therefore achieved only 195 kg lamb per ha compared to the future farming system goal of 260kg lamb per ha (Table 1) .Given normal rainfall and pasture growth the demonstration could potentially produce 354kg of lamb per ha (based on a liveweight of 51kg) .

**Table 1** Livestock production data in 2006 compared to a simulated traditional farming system ,current best practice and future farming system .

	Livestock production from field site 2006	Simulated traditional farming system(30% crop , 70% annual pasture)	Simulated current best practice (30% crop , 23% annual pasture . 47% perennial pasture)	Simulated future farming system(30% crop ,70% perennial pasture)
Stocking rate (dse/ha)	9.9	8.5	10.0	12.0
Supplementary feed (kg/ha)	559	279	83	95
Lambing %	153	89	92	104
Weaning %	119	89	92	104
No of lambs weaned per ha	7.8	4.6	5.5	7.3
Weaning liveweight (kg)	25	25	25	25
Total lamb produced (kg/ha)	195	161	193	260

**Conclusions** Drought conditions substantially reduced the livestock benefits of perennials with low pasture growth resulting in a large increase in supplementary feed .How this system performs in an average or better season ,will determine if it is more profitable in the longer term than current farming systems .

### Reference

Sanford P. & J.Young (2005) .Are new farming systems based on perennial pastures in south west Australia more profitable ?  
*Proceedings of the XX International Grassland Congress ,Dublin ,Ireland , 847 .*