

## University of Kentucky **UKnowledge**

International Grassland Congress Proceedings

21st International Grassland Congress / 8th International Rangeland Congress

Cultivating the Surface Soil to Renovate a Grama Rhodes (Chloris gayana K. cv. Callide) Pasture in Northwest Argentine (NOA). 2. **Effect of Nitrogen Fertilization** 

R. N. Berti INTA, Argentina

Follow this and additional works at: https://uknowledge.uky.edu/igc



Part of the Plant Sciences Commons, and the Soil Science Commons

This document is available at https://uknowledge.uky.edu/igc/21/9-2/26

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

## Cultivating the surface soil to renovate a Grama Rhodes (Chloris gayana K .cv .Callide) pasture in northwest Argentine (NOA) 2 Effect of nitrogen fertilization

Berti R. N. INTA EEA SALTA Argentine Republic .rberti@correo .inta .gov .ar

Key words: Grama Rhodes ,nitrogen fertilization ,forage production ,weeds ,Northwest Argentine region .

Introduction Grama Rhodes is a pasture employed in several regions of NOA . Its production is strongly to associate to availability of nitrogen in soils (Berti 2006) . Methods to improve the supply of nitrogen to grass on NOA are being sought .

Material and Methods At Cerrillos Salta (24°54′S;65°29′W;1250 m.), Argentina on a grama Rhodes pasture seeded in 1994 a trial was carried out in November 2000 ,which was related in another paper (Berti,2008). Total rainfall for 2000/01;01/02;02/03;03/04 and 04/05 ,covering the experimental period (May-November) were 844.7;566.3;548.2;562.4 and 356 mm. The historical annual average for the site was 643 mm. A split-split plot design ,with three replications were employed. Two levels of nitrogen fertilization ,(0). Control ,no fertilizer and (1),100 kg. N. ha<sup>-1</sup> were used. Nitrogen was applied annually ,with broadcasting urea after cultivation in the end of November ,before the first rain. Seven harvest of plant material were done between Nov 2000 and Mar 2005. The plant samples were dried ,weighed ,grounded and Nitrogen (N) and Dry Matter (DM) content was determined (A.O. A.C.,1980). In the last period samples were taken from 0 to 20 cm layers of soil ,with a 2.5 cm coring tube at fifteen random position in each plot at the end of the growing season. These samples were dried and N-NO3 determined (Bremner et al ,1966).

Results and Discussion The fertilization increased aerial biomass of Grama ,weeds ,and nitrogen content of forage (Table 1) . At the end of trial the content in soil of N-Nitrate (N-NO $^3$ ) in treatment fertilized was significantly higher . The mean response obtained (20 kg DM by unit of N) in grama is similar to the one previously reported (Berti 2006) . During the periods 03/04 and mainly in 04/05 ,lack of rains occurred after cultivation resulting in increased of weeds in treatments with N .

Table 1 Effect of fertilization treatments on the total yields and N content of grama weeds and N-nitrate in soil.

Fertilization	Grama D .M . ,kg ha <sup>-1</sup>	Grama N . ,kg ha <sup>-1</sup>	Weeds D .M . ,kg $ha^{-1}$	Soil N-NO3 ,ppm
0 ,No fertilizer	12411a	122a	174 <b>a</b>	12 5a
1 ,100 kg N year ha <sup>-1</sup>	22441b	261b	789Ь	17 .4b
MSE z	2633 .5	36.2	548	3.9

a ,b : Values on the same column with different letters are different ,P<0 .05 .Duncan Test .

Conclusion Nitrogen fertilization significantly increased aerial biomass of Grama and weeds .

## References

A.O. A.C., 1980. Ass. Off. Agr. Chemistry. Official Methods of Analysis. Washington, D.C...

Berti ,R .N .2006 .Effect of soil scarification and nitrogen fertilization of Chloris gayana Cv .Callide ,on forage accumulation . RA PA ,Vol .26 ,Supl : 121-122 .

Berti ,R .N 2008a .Cultivating the surface soil to renovate a Grama Rhodes (Chloris gayana K .cv .Callide) pasture in Northwest Argentine (NOA) .1 .Effect of implements (In this Congress) .

Brenmer J. M. and Keeney ,D. R. ,1966 .Exchangeable ammonium nitrate and nitrite by extraction-distillation methods .Soil Sci Am .Proc . 30 577-582 .

z : Standard error of the mean