

**The Nelson Mandela African Institution of Science and Technology**

**<http://dspace.nm-aist.ac.tz>**

---

Materials, Energy, Water and Environmental Sciences

Research Articles [MEWES]

---

2018-04-05

# Crustacean derived calcium phosphate systems: Application in defluoridation of drinking water in East African rift valley

Wagutu, Agatha W.

Elsevier B.V.

---

<https://doi.org/10.1016/j.jhazmat.2017.12.049>

*Downloaded from Nelson Mandela-AIST's institutional repository*

# Crustacean derived calcium phosphate systems: Application in defluoridation of drinking water in East African rift valley

Agatha W. Wagutu, Revocatus Machunda, Yusufu Abeid Chande Jande

To download full text click that link

DOI: <https://doi.org/10.1016/j.jhazmat.2017.12.049>

## Abstract

Calcium phosphate adsorbents, derived from prawns and crabs shell biomass wastes have been developed using wet chemistry and low temperature treatment. The adsorbents were characterized by X-ray diffractometry and Fourier transform infrared spectroscopy. Batch adsorption test were carried out to investigate their effectiveness in adsorption of fluoride from ground and surface waters. Adsorption capacities were compared with bone char and synthetic hydroxyapatite (CCHA). Results indicate that prawns derived adsorbent (PHA) formed hexagonal structure with phases identifiable with hydroxyapatite while crabs based adsorbent (CHA) formed predominantly monoclinic structure with crystalline phase characteristic of brushite. Vibrational analysis and kinetic studies predicted defluoridation occurred mainly by ion exchange and ion adsorption mechanisms. Defluoridation capacity of the adsorbents was found to be superior compared to bone char and CCHA. CHA was the most effective with efficiencies above 92% and highest capacity of 13.6 mg/g in field water with fluoride concentration of 5–70 mg/L. PHA had highest capacity of 8.5 mg/g which was still better than 2.6 mg/g recorded by CCHA and bone char. Adsorption was best described by pseudo 2nd order kinetics. The findings indicate that crustacean derived calcium phosphate systems have better potential for defluoridation than traditional bone char and synthetic systems.

## Keywords

Adsorption; Crabs; Fluoride; Prawns; Kinetics