

The Nelson Mandela African Institution of Science and Technology

<http://dspace.nm-aist.ac.tz>

---

Life sciences and Bio-engineering

Research Articles [LISBE]

---

2018-04-03

# Effects of clipping and irrigation on carbon storage in grasses: implications for CO<sub>2</sub> emission mitigation in rangelands

Tuffa, Samuel

Taylor & Francis Online

---

<https://doi.org/10.1080/17583004.2018.1451963>

*Downloaded from Nelson Mandela-AIST's institutional repository*

# Effects of clipping and irrigation on carbon storage in grasses: implications for CO<sub>2</sub> emission mitigation in rangelands

Samuel Tuffa, Dana Hoag & Anna C. Treydte

To download full text click that link

DOI: <https://doi.org/10.1080/17583004.2018.1451963>

## Abstract

Understanding how individual grasses respond to herbivory and rainfall has been hampered by the difficulty of quantifying above- and belowground carbon (C) storage in grasses. Particularly by restoring degraded rangelands through reseeding, their C storage potential can be greatly enhanced. The responses of reseeded grasses to the effects of herbivory and precipitation were assessed to evaluate the potential of individual grasses for C storage as a technique for climate change mitigation. Clipping experiments were conducted on mature grass tufts of two native grass species, *Chloris gayana* and *Cenchrus ciliaris*, in the semi-arid Borana rangelands, Ethiopia. Further, above- and belowground C storage of young grasses of the same species in pot and field plot trials was experimentally quantified under simulated grazing and variable rainfall. The results showed that aboveground C was significantly 4 times lower in the clipped compared to unclipped mature grasses. In contrast, 3 times higher C was found in young reseeded grasses that were clipped compared to unclipped ones. Clipping and irrigation in combination significantly influenced belowground C in young grasses, with reduced irrigation overriding clipping effects. The paper concludes that moderate grazing should be encouraged to enhance CO<sub>2</sub> uptake, consequently contributing to climate change mitigation in rangelands.

**KEYWORDS:** Borana, herbaceous layer restoration, herbivory, livestock management, rainfall variability