

Animal welfare, etológia és tartástechnológia



Animal welfare, ethology and housing systems

Volume 16

Issue 2

Gödöllő
2020

DIGESTIBILITY AND PERFORMANCE OF BROILER CHICKENS FED DIETS WITH HUMIC SUBSTANCES

Marcin Andrej, Nad' Pavel, Bujňák Lukáš, Skalická Magdaléna

University of Veterinary Medicine and Pharmacy in Košice,
Department of Nutrition and Animal Breeding
041 81 Košice (SK), Komenského 73
andrej.marcin@uvlf.sk

Received – Érkezett: 15. 11. 2019.
Accepted – Elfogadva: 19.10.2020.

Abstract

The objective of the study was to investigate the effects of dietary intake of humic substances (HS) on the ileal digestibility of dry matter (DM) and crude protein (CP). One-day-old broiler chickens (Cobb 500, n=120, average weight 50 g) were divided at random into 4 equal groups (A, B, C / negative control). Chickens were fed with mixtures with the content of CP - BR1 230.20, BR2 222.20, BR3 209.40 g.kg⁻¹ DM for 37 days. The applied additives HN/HNM (Humac Ltd., Slovak Republic) contained humic acids (HA) min. 650/570 and fulvic acids (FA) min. 50/50 g.kg⁻¹. The humic substances were added into diets of experimental groups according to the content of HA/FA in the feed additives A 4.55/0.35, B 3.99/0.35, C 2.85/0.25 g.kg⁻¹. The body weights (BW) and feed consumption were assessed once a week. The average daily weight gains (ADWG), average daily feed intake (ADFI) and the feed conversion ratio (FCR) were calculated. The ileal chymus, sampled from 48 birds after slaughter, was used for the quantification of DM, CP, ash and ash insoluble in HCl on day 37. The production parameters of chickens were from day 1 – 37 as follows. The average finishing body weights were A 2506.67 ± 160.47, B 2490.25 ± 166.39, C 2377.75 ± 133.19 and control 2319.42 ± 92.55 g. The average weight gains achieved the values A 66.39 ± 16.85, B 65.95 ± 17.77, C 62.91 ± 18.06 and control 61.34 ± 17.31 g.day⁻¹. The average daily feed intakes were A 108.97 ± 50.11, B 116.91 ± 55.35, C 108.85 ± 51.44 and control 106.83 ± 46.97 g.day⁻¹ from day 1 – 37. The addition of humic substances caused the increase of ileal digestibility in B group compared to control: DM 38.32 ± 0.18 / 30.94 ± 2.41% (p < 0.05) and CP 40.39 ± 0.05 / 34.29 ± 0.26% (p < 0.05). The dietary intake of feed additives containing HS seems to have a positive effect on finishing BW and the ileal digestibility of DM and CP in the case of the minimal inclusion level of HA/FA 3.99/0.35 g.kg⁻¹.

Keywords: gut of poultry, humates, dry matter, crude protein, ileum

Introduction

The feed additives with the antimicrobial characteristics were used for the improvement of production parameters of livestock and poultry (*Dibner and Buttin, 2002*). The antibiotic growth promoters were banned in the EU with the Regulation 1831/2003 (*Castanon, 2007*) because of the negative residual effects of animal products used in human nutrition on health and the increased bacterial resistance.

The alternative growth promoters to antimicrobials which are used nowadays are organic acids (*Ur Rehman et al., 2016*), plant extracts (*Abd El-Hack and Alagawany, 2015; Dhama et al., 2015*), probiotics and prebiotics (*Alagawany et al., 2018*) and enzymes (*Griggs and Jacob, 2005*). As far as the essential oils are concerned, these are important as growth promoters of poultry (*Levkut et al., 2010, 2011*).

The additional feed additives with a positive effect on the production parameters of animals are could be the humic substances (HS) too. The enhancement of growth ability of animal depends on the sources of humic acids (HA) and fulvic acids (FA) present in HS. According to already published results, the improvements were observed in the case of production of broiler chickens in various parameters such as body weight, feed conversion, the retention of ashes, nitrogen, and energy as well as reduced crypt depth and increased length of the villi of the jejunal mucosa (*Gomez-Rosales and Angeles, 2015; Arif, M. et al., 2019*).

According to the International Humic Substances Society (2007), HS are major components of the natural organic matter (NOM) in soil, water and in geological organic deposits such as lake sediments, peats, brown coals and shales. These are responsible for the characteristic brown colour of decaying plant debris and contribute to the brown or black colour in surface soils. As for formation, HS are complex and heterogeneous mixtures of polydispersed materials created by biochemical and chemical reactions during the decay and transformation of plant and microbial remains in the process called humification. The important components of this process are plant lignin, its transformation products, polysaccharides, melanin, cutin, proteins, lipids, nucleic acids and fine char particles. HS in soils and sediments are divided into following fractions: HA, FA and humin. The strong base (NaOH or KOH) are used for the extraction of HA and FA from the soil and other solid phase sources. Because of the insolubility of HA at low pH, they are precipitated by adding strong acid.

According to the results of *Marcin et al. (2020)*, the addition of HS combined with urea had a positive effect on the amylolytic and cellulolytic activities in the rumen fluid of sheep. Higher activities of digestive enzymes (amylase, lipase, protease) were observed after feeding of broiler chickens with FA (*Mao, Y., 2019*). It is in agreement with the length of the mucosal villi of the jejunum and the increased gut length due to dietary inclusions of HS into broiler feed (*Vašková et al., 2018; Taklimi et al., 2012*).

The scientific hypothesis for the performed experiment was based on the positive effects of HS on the enzymatic activities in the gastrointestinal apparatus which have potential to improve digestion of proteins, starch and cellulose in the gut of poultry.

Therefore, the study aimed to investigate the effects of dietary intake of a preparation containing HS on the basic production parameters and the ileal digestibility of the dry matter (DM) and the crude protein (CP) of broiler chickens.

Materials and methods

Chickens and diets

One hundred and twenty one-day-old broiler chickens of hybrid COBB 500 (Mach Drubež, a.s., Litomyšl, Czech Republic) (average weight 50 g), were delivered from a commercial hatchery. Birds were divided into 4 groups of 30 animals (A, B, C / negative control). They were housed in four-floor pens located in one experimental hall of the University of Veterinary Medicine and Pharmacy in Košice (Slovak Republic) with constant access to feed and water. The pens were identical concerning the same direction and the same area (0.12 m² per broiler chicken).

All groups were fed with mash diets (Agrocass plus, Ltd., Slovak Republic) with CP content: BR1 – 230.20, BR2 – 222.20, BR3 – 209.40 g.kg⁻¹ DM) for 37 days. The methionine was used as the first limiting amino acid. The diets were prepared and formulated without antibiotics and growth promoters. The anticoccidial agents were added into the starter and grower feed mixtures. The calculations of diets were performed according to the nutrient requirements and nutrient value of feeds for poultry (Kočič et al., 1994).

HS (Humac Ltd., Slovak Republic) were added into feeds of three experimental groups of chickens - Humac natur (HN) for A and Humac natur monogastric (HNM) for B and C group. These additives were absent in the feeds for the control group. The characteristics of the applied preparations HN/HNM were the following: the size of particles up to 100 µm, max. moisture 15%, the content of humic acids (HA) min. 650/570, fulvic acids (FA) min. 50/50 g.kg⁻¹, macroelements Ca 42.28/51.1, Mg 5.11/4.86, Fe 19.05/18.09 g.kg⁻¹ and microelements Cu 15/14.25, Zn 37/35.15, Mn 142/135, Co 1.24/1.18, Se 1.67/1.59 as well as Mo 2.7/2.57, V 42.1/40 mg.kg⁻¹ DM. The inclusion level of the additives in the diets of experimental groups was measured according to the content of HA/FA in the feed additives: A 4.55/0.35, B 3.99/0.35, C 2.85/0.25 g.kg⁻¹. The body weights (BW) of chickens and the feed consumption were evaluated once a week. The average daily weight gains (ADWG), the average daily feed intake (ADFI) and feed conversion ratio (FCR) were calculated.

Feed analysis

The samples of diets were analysed (Table 1) according to the official methods of the Association of Official Analytical Chemists (Cunniff, 1995). There were performed the analyses of DM, CP, crude fat, starch and ash. The fibre was determined with the method by Van Soest et al. (1991). The mineral composition of the feed was analysed by atomic absorption spectrophotometry (AAS) (Van Loon, 1980). The quantitative determination of phosphorus was performed spectrophotometrically (Carvalho et al. 1998). The insoluble portion of ash in HCl was determined in the feed mixture as the residue of ash, after dissolving ash in diluted hydrochloric acid by weighing (Daněk et al. 2005). The metabolisable energy value of diets was calculated with the formula from the Commission Regulation (EC) No 152/2009 (European commission, 2009) according to the method of calculation and expression of energy value.

Check of digestibility

The ileal digestibility was checked after the slaughter of 48 birds on day 37. The samples of ileal chymus from each slaughtered chicken were applied into sterile glass containers. Subsequently, each 30 g sample was pooled from the ileal chymus of two chickens. Therefore the resulting total number of analysed samples was 24. The quantifications of DM, CP, ash and ash insoluble in HCl were performed by the above-mentioned methods.

The ileal digestibility was calculated according to the method of *Kadim and Moughan* (1997). The content of the insoluble portion of ash was used as the indicator. The digestibility was determined by calculating the analysed content of nutrients in feed and ileal chymus according to the formula:

$$\text{Nutrient digestibility (\%)} = \frac{\text{nutrient content in diet/indicator in diet} - \text{nutrient output in ileum/indicator in ileum}}{\text{nutrient content in diet/indicator in diet}} * 100$$

Table 1: Composition of the experimental diets

Analysed nutrients (g.kg ⁻¹)	Diets		
	BR1	BR2	BR3
Dry mater	100.00	100.00	100.00
Crude protein	230,20	222.20	209.40
Crude fat	31.30	83.80	67.70
ND fibre	112.60	122.10	128.80
AD fibre	54.60	62.50	68.10
Ash	57.30	60.60	52.10
Ash insoluble in HCl	2.10	1.80	2.40
Starch	485.60	446.80	448.60
Calcium	4.93	6.00	7.41
Phosphorus	5.73	7.93	5.13
Sodium	2.98	1.93	1.60
Magnesium	2.86	3.06	3.11
Potassium	9.03	8.83	8.49
Copper	0.0275	0.0578	0.0594
Zinc	0.0229	0.0294	0.1336
Manganese	0.0789	0.1472	0.1437
Metabolizable energy (MJ.kg ⁻¹)*	13.27	14.30	13.58

*Calculation based on Commission Regulation (EC) 152/2009

Ingredients in diets	
BR1 / BR2 / BR3	maize, wheat, soybean meal GMO, vegetable oil, limestone, amino acids and their salts, monocalcium phosphate, mineral-vitamin premix
BR1	the dried derivative of pig blood
BR2 / BR3	sunflower extracted meal, salt

Statistical analysis

The data are expressed as means ± standard deviation (SD) of single values (IBM SPSS Statistics, Version 24). Means of the results from the treatments were compared by one-way analysis of variance. Treatment means were statistically compared by Tukey-Kramer multiple comparison test. Significance was declared at $P < 0.05$.

Results and discussion

The considerable interest has been in the use of HS as a feed additive to enhance performance and nutrient digestibility of poultry (Domínguez-Negrete et al., 2019).

The present results indicate that the dietary intake of HS is effective in the enhancement of the performance of broilers. The values of BW and ADWG are shown in Table 2. ADWG was improved by the higher amount of the added HS 0.7% into feed, or the inclusion level of HA/FA in the feed additives A 4.55/0.35 and B 3.99/0.35 by 7.61% in A and by 6.99% in B groups compared to control at day 1 – 37. In the case of individual BW, there were observed the only one significant decrease on day 7 when this parameter was lower in A group by 9.7 g ($p < 0.05$) in comparison to the negative control. The finishing BW was higher in A or B groups compared to control by 187.25 and 170.83 g, respectively.

Table 2: Impact of feeding of humic substances on the growth parameters of broiler chickens

Group	BW (g)							ADWG (g.day ⁻¹)
	d 1	d 7	d 14	d 21	d 28	d 35	d 37	d 1 – d 37
A	50.0 ^a ± 3.75	182.07 ^b ± 4.66	488.97 ^a ± 65.44	963.07 ^a ± 120.89	1571.48 ^a ± 280.39	2311.92 ^a ± 367.72	2506.67 ^b ± 160.47	66.39 ^a ± 16.85
B	50.0 ^a ± 2.65	188.82 ^{ab} ± 5.85	512.75 ^a ± 67.64	986.37 ^a ± 116.20	1636.63 ^a ± 210.66	2326.56 ^a ± 292.20	2490.25 ^b ± 166.39	65.95 ^a ± 17.77
C	50.0 ^a ± 3.35	192.27 ^{ab} ± 7.18	521.5 ^a ± 45.87	989.33 ^a ± 102.5	1585.17 ^a ± 286.5	2281.93 ^a ± 326.51	2377.75 ^{ab} ± 133.19	62.91 ^a ± 18.06
Control	50.0 ^a ± 4.15	191.73 ^a ± 8.41	514.86 ^a ± 54.30	985.57 ^a ± 100.03	1633.0 ^a ± 193.26	2291.79 ^a ± 320.88	2319.42 ^a ± 92.55	61.34 ^a ± 17.31

Means with different superscript letters differ significantly ^{a, b} $P < 0.05$ (mean ± SD), BW – body weight, ADWG – average daily weight gain, d – day

The similar improvement of weight gains was observed by Arif et al. (2016). They determined the increase of this parameter of broilers fed HA at 2.25 g.kg⁻¹ diet as a result of the improved feed conversion. Their dosage of HA was approximately one half compared to our inclusion levels. On the other hand, the statistical insignificance of ADWG could be partially supported by the findings of

Karaoğlu et al. (2004). They observed that supplementation of HS and HA in the diet of broiler chickens did not influence the body weight gains.

ADFI and FCR are presented in *Table 3*. There were not observed significant differences in FCR during the experiment. The improvement of ADFI was evaluated in the experimental groups: A and C on day 7 (by 0.6% and 4.02%), B on day 14 and 21 (by 4.39% and 5.39%), A and B on day 28 (by 4.26% and 7.59%), A, B and C on day 35 (by 6.64%, 17.13% and 12.22%) as well as in B and C on day 37 (by 5.38% and 2.10%).

The positive results of feed utilization are in coincidence with the partial improvement of nutrient digestibility and the protection of commensal gut microflora caused by HS (*Windisch et al.*, 2008). However, gut health is a very important factor in the utilization and conversion of nutrients of growing chickens (*Islam et al.*, 2008).

Table 3: Impact of feeding of humic substances on the feed consumption and conversion of broiler chickens

Group	ADFI (g.day ⁻¹ .chicken ⁻¹)						FCR (kg.kg ⁻¹)
	d 7	d 14	d 21	d 28	d 35	d 37	d 1 – d 37
A	28.11	60.39	105.82	147.09	167.20	145.18	1.65 ^a ± 0.29
B	27.29	66.54	112.17	152.38	188.36	154.71	1.64 ^a ± 0.19
C	29.11	60.42	100.95	135.24	177.83	149.52	1.53 ^a ± 0.26
Control	27.94	63.62	106.12	140.82	156.10	146.38	1.51 ^a ± 0.17

Means with different superscript letters differ significantly ^{a, b} P < 0.05 (mean ± SD), ADFI – average daily feed intake, FCR – feed conversion ratio, d – day

The addition of HS had a positive effect on the increase of ileal digestibility of DM or CP at the inclusion level of HA/FA into diets at minimum 3.99/0.35 g.kg⁻¹ feed (*Table 4*) on day 37. As for DM digestibility, there was observed the higher value in the B group compared to control by 7.38%. The significant difference was observed between groups B and C by 14.66% (P < 0.05).

Similarly, in the case of the CP digestibility, the higher value was determined in the B group compared to control by 6.1% and the significant difference was observed between groups B and C (P < 0.05) by 7.60% too. The digestibility of nutrients present in the chymus depends on the enzymatic activities in the small intestine which could be positively influenced by HS.

This statement can be confirmed by the observation of *Terry et al.* (2018) when HS had a positive effect on the apparent total tract digestibility. HS are can improve the digestibility of energy in the ileum which is dose-dependent but on the contrary, the ileal digestibility of N was not affected (*Gomez-Rosales and Angeles*, 2015).

Table 4: Ileal digestibility of DM and CP analysed in ileal chymus after intake of HS sampled from 48 broiler chickens on 37 day (samples were pooled to n = 6, mean ± SD)

Group	DM (%)	CP (%)
A	33.27 ^{ab} ± 5.26	37.99 ^{ab} ± 1.42
B	38.32 ^a ± 0.18	40.39 ^a ± 0.05
C	23.66 ^b ± 0.56	32.79 ^b ± 0.66
Control	30.94 ^{ab} ± 2.41	34.29 ^b ± 0.26

Means with different superscript letters differed significantly: ^{a,b} P < 0.05, DM – dry matter, CP – crude protein, HS – humic substances

Conclusion

The dietary intake of HS seems to have a positive effect on the ileal digestibility at the inclusion level HA/FA at minimum 3.99/0.35 g.kg⁻¹. There were observed the increased digestibility of DM by 7.38% (p < 0.05) and CP by 6.1% (p < 0.05) compared to negative control. The finishing individual BW of experimental groups were positively influenced in A by 187.25 (p < 0.05) and in B groups by 170.83 g (p < 0.05).

Acknowledgement

This study was supported by the grant from the Scientific Grant Agency of the Ministry of Education of the Slovak Republic for the project VEGA-1/0402/20 - Effect of additives in the nutrition of monogastric animals on production health, production parameters, products quality and environment.

References

- Abd El-Hack, M.E., Alagawany, M.* (2015): Performance, egg quality, blood profile, immune function, and antioxidant enzyme activities in laying hens fed diets with thyme powder. *Journal of Animal and Feed Sciences*, 24. 127-133.
- Alagawany, M., Abd El-Hack, M.E., Farag, M.R., Sachan, S., Karthik, K., Khama, K.* (2018): The use of probiotics as eco-friendly alternatives for antibiotics in poultry nutrition. *Environmental Science and Pollution Research International*, 25. 10611-10618.
- Arif, M., Rehman, A., Saeed, M., Abd El-Hack, M.E., Arain, M.A., Haseebarsad, M., Zakria, H.M., Abbasi, I.H.* (2016): Impacts of dietary humic acid supplementation on growth performance, some blood metabolites and carcass traits of broiler chicks. *Indian Journal of Animal Sciences*, 86. 1073-1078.
- Arif, M., Alagawany, M., Abd El-Hack, M.E., Saeed, M., Arain, M.A., Elnesr, S.S.* (2019): Humic Acid as a Feed Additive in Poultry Diets: A Review. *Iranian Journal of Veterinary Research*, 20. 167–172.
- Carvalho, L. H. M., De Koe, T., Tavares, P. B.* (1998): An improved molybdenum blue method for simultaneous determination of inorganic phosphate and arsenate. *Ecotoxicology and*

- Environmental Restoration, 1. 13–19.
- Castanon, J.I.R. (2007): History of the use of antibiotic as growth promoters in European poultry feeds. *Poultry Science*, 86. 2466–2471.
- Cunniff, P. (1995): *Official Methods of Analysis of Association of Official Analytical Chemists*. 16th edn., AOAC International, Arlington, Va, USA.
- Daněk, P., Paseka, A., Smola, J., Ondráček, J., Bečková, R., Rozkot, M., (2005): Influence of lecithin emulsifier on the utilisation of nutrients and growth of piglets after weaning. *Czech Journal of Animal Science*, 50. 459–465.
- Dhama, K., Latheef, S.K., Mani, S., Abdul Samad, H., Karthik, K., Tiwari, R., Khan, R.U., Alagawany, M., Mayada, R.F., Alam, G.M., Laudadio, V., Tufarelli, V. (2015); Multiple beneficial applications and modes of action of herbs in poultry health and production. A review. *International Journal of Pharmacology*, 11. 152-176.
- Dibner, J.J., Buttin, P. (2002): Use of organic acids as a model to study the impact of gut microflora on nutrition and metabolism. *Journal of Applied Poultry Research*, 11. 453-463.
- Domínguez-Negrete, A., Gómez-Rosales, S., a de Lourdes Angeles, M., Humberto López-Hernández, L., Cesaria Reis-de Souza, T., López-García, Y., Zavala-Franco, A., Téllez-Isaias, G. (2019): Effect of the addition of humic substances as growth promoter in broiler chickens under two Feeding Regimens. *Animals*, 9. 1101
- European Commission (2009): Commission Regulation (EC) No 152/2009 of 27 January 2009 laying down the methods of sampling and analysis for the official control of feed. *Official Journal of European Union*, 54. 1–130.
- Griggs, J.P., Jacob, J.P. (2005): Alternatives to antibiotics for organic poultry production. *Journal of Applied Poultry Research*, 14. 750-756.
- Gomez-Rosales, S., Angeles, M.D. (2015): Addition of a worm leachate as source of humic substances in the drinking water of broiler chickens. *Asian-Australasian Journal of Animal Sciences*, 28. 215-222.
- International Humic Substances Society (IHSS) (2007): What are humic substances ? Newsletter, IHSS, Number 38. 2.
http://humic-substances.org/wp-content/uploads/NL_38_2007.pdf
- Islam, K., Schuhmacher, A., Aupperle, H., Gropp, J. (2008): Fumaric acid in broiler nutrition: a dose titration study and safety aspects. *International Journal of Poultry Science*, 7. 903-907.
- Kadim, I.T., Moughan, P.J. (1997): Development of an ileal amino acid digestibility assay for the growing chicken – effects of time after feeding and site of sampling. *British Poultry Science*, 38, 89-95. 1997.
- Karaoğlu, M., Macit, M., Esenbuğa, N., Durdağ, H., Bilgin, Ö.C. (2004): Effect of dietary humate on performance, slaughter, carcass and meat quality parameters of broilers. *International Journal of Poultry Science*, 6. 406-410.
- Carvalho, L. H. M., De Koe, T., Tavares, P. B. (1998): An improved molybdenum blue method for simultaneous determination of inorganic phosphate and arsenate. *Ecotoxicology and Environmental Restoration*, 1. 13–19.
- Kočí, Š., Kočiová, Z., Zelenka, J., Zeman, L. (1994): *The nutrient requirements and nutrient value of feeds for poultry*. Ed.1, Nitra: Research institute of animal production, 46 pp. ISBN 80-967057-3-3.
- Levkut, M., Marcin, A., Lenhardt, L., Porvaz, P., Revajová, V., Šoltysová, B., Blanár, J., Ševčíková, Z., Pistl, J. (2010): Effect of Sage Extract on Alkaline Phosphatase, Enterocyte Proliferative Activity and Growth Performance in Chickens. *Acta Veterinaria Brno*, 79. 177 – 183.

- Levkut, M., Marcin, A., Revajová, V., Lenhardt, L., Danielovič, I., Hecl, J., Blanár, J., Levkutová, M., Pistl, J. (2011): Influence of oregano extract on the intestine, some plasma parameters and growth performance in Dickens. *Acta Veterinaria (Beograd)*, 61. 215-225.
- Mao, Y. (2019): Modulation of the Growth Performance, Meat Composition, Oxidative Status, and Immunity of Broilers by Dietary Fulvic Acids. *Poultry Science*, 10. 4509–4513.
- Marcin, A., Bujňák, L., Mihok, T., Nad', P. (2020): Effects of humic substances with urea on protozoal population and fermentation in the rumen of sheep. *Bulgarian Journal of Veterinary Medicine*, 23. 60-69.
- Taklimi, S.M.S.M., Ghahri, H., Isakan, M.A. (2012): Influence of different levels of humic acid and esterified glucomannan on growth performance and intestinal morphology of broiler chickens. *Agricultural Sciences*, 3. 663-668.
- Terry, S.A., de Oliveira Ribeiro, G., Gruninger, R.J., Hunerberg, M., Ping, S., Chaves, A.V., Bulet, J., Beauchemin, K.A., McAllister, T.A. (2018): Effect of humic substances on rumen fermentation, nutrient digestibility, methane emissions and rumen microbiota in beef heifers. *Journal of Animal Sciences*, 96. 3863-3877.
- Ur Rehman, Z., Haq, A.U., Akram, N., Abd El-Hack, M.E., Saeed, M., Ur Rehman, S., Meng, C., Alagawany, M., Sayab, M., Dhama, K., Ding, C. (2016). Growth performance, intestinal histomorphology, blood hematology and serum metabolites of broilers chickens fed diet supplemented with graded levels of acetic acid. *International Journal of Pharmacology*, 12. 874-883.
- Van Loon, J. C., (1980): *Analytical Atomic Absorption Spectroscopy, Selected methods*. Academic press, New York, 337 pp.
- Van Soest, P. J., Robertson, J. B., Lewis, B. A., (1991): Methods for dietary fiber, neutral detergent fibre, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, 74. 3583–3597.
- Vaskova, J., Patlevič, P., Žatko, D., Marcinčák, S., Vaško, L., Krempaská, K., Nagy, J. (2018): Effects of Humic Acids on Poultry Under Stress Conditions. *Slovenian Veterinary Research*, 55. 245–253.
- Windisch, W., Schedle, K., Plitzner, C., Kroismayr, A. (2008): Use of phytogenic products as feed additives for swine and poultry. *Journal of Animal Science*, 86. E140-E148.