



Silhol, R., Vickerman, P., Boily, M-C., & al., E. (2021). Quantifying the evolving contribution of HIV interventions and key populations to the HIV epidemic in Yaoundé, Cameroon. *Journal of Acquired Immune Deficiency Syndromes*, 86(4), 396-405.  
<https://doi.org/10.1097/QAI.0000000000002580>

Peer reviewed version

Link to published version (if available):  
[10.1097/QAI.0000000000002580](https://doi.org/10.1097/QAI.0000000000002580)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Lippincott, Williams & Wilkins at [DX.DOI.ORG/10.1097/QAI.0000000000002580](https://dx.doi.org/10.1097/QAI.0000000000002580). Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

# Quantifying the evolving contribution of HIV interventions and key populations to the HIV epidemic in Yaoundé, Cameroon

Romain Silhol, PhD<sup>1,2</sup>, Stefan Baral, MD<sup>3</sup>, Anna L Bowring, PhD<sup>3,4</sup>, Christinah Mukandavire, PhD<sup>2</sup>, Iliassou Mfochive Njindam, MD<sup>3</sup>, Amrita Rao, MSc<sup>3</sup>, Sheree Schwartz, PhD<sup>3</sup>, Ubald Tamoufe, MSc<sup>3,5</sup>, Serge C Billong, MD<sup>6,7</sup>, Oudou Njoya, MD<sup>8</sup>, Anne-Cecile Zoung-Kanyi Bissek, MD<sup>8,9</sup>, Jesus Maria Garcia Calleja, MD<sup>10</sup>, Peter Vikerman, DPhil<sup>11</sup>, Sharmistha Mishra, MD, PhD<sup>12,13\*</sup>, Marie-Claude Boily, PhD<sup>1,2\*</sup>

<sup>1</sup>Department of Infectious Disease Epidemiology, Imperial College London, HPTN Modelling Centre, London, United Kingdom

<sup>2</sup>MRC Centre for Global Infectious Disease Analysis, London, United Kingdom

<sup>3</sup>Johns Hopkins School of Public Health, Department of Epidemiology, Baltimore, MD, USA

<sup>4</sup>Burnet Institute, Melbourne, Australia

<sup>5</sup>Metabiota, Yaoundé, Cameroon

<sup>6</sup>Department of Public Health, Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, Yaoundé, Cameroon

<sup>7</sup>National AIDS Control Committee (NACC/CNLS), Yaoundé, Cameroon

<sup>8</sup>Department of Internal Medicine and Specialties, Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, Yaoundé, Cameroon

<sup>9</sup>Ministry of Health, Division of Operations Research, Yaoundé, Cameroon

<sup>10</sup>World Health Organization, Geneva, Switzerland

<sup>11</sup>Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

<sup>12</sup>St. Michael's Hospital, Li Ka Shing Knowledge Institute, University of Toronto, Toronto, Canada

<sup>13</sup>Department of Medicine, Division of Infectious Disease, University of Toronto, Toronto, Canada

\* These authors have contributed equally to the work.

**Correspondence to:** Romain Silhol, PhD, MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, Imperial College London, St Mary's Hospital, 2 Norfolk Place, W2 1PG London, United Kingdom (email: [r.silhol@imperial.ac.uk](mailto:r.silhol@imperial.ac.uk) , phone: +44207594 3290, fax: +442074023927)

Presented in part as a poster at the Conference on Retroviruses and Opportunistic Infections (CROI): Boston, MA, USA. March 8-11, 2020.

**Conflicts of Interest and Source of Funding:**

The authors have no conflicts of interest to disclose.

This work was partly supported by the HPTN Modelling Centre, which is funded by the U.S. National Institutes of Health (NIH UM1 AI068617) through HPTN. RS and MCB acknowledge funding from MRC Centre for Global Infectious Disease Analysis (MRC GIDA, MR/R015600/1). This award is jointly funded by the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID Concordat agreement and is also part of the EDCTP2 programme supported by the European Union. The program also received support from Linkages across the Continuum of HIV Services for Key Populations Affected by HIV project (LINKAGES, Cooperative Agreement AID-OAA-A-14-00045). PV also acknowledges support from the NIHR Health Protection Research Unit in Behavioural Science and Evaluation at University of Bristol. AB is supported by an Australian National Health and Medical Research Council Early Career Fellowship.

**Running head:** Dynamics of HIV Transmission in Yaoundé

## **Abstract**

**Background:** Key populations (KP) including men who have sex with men (MSM), female sex workers (FSW), and their clients are disproportionately affected by HIV in Sub-Saharan Africa. We estimated the evolving impact of past interventions and contribution of unmet HIV prevention/treatment needs of KP and lower-risk groups to HIV transmission.

**Setting:** Yaoundé, Cameroon.

**Methods:** We parametrised and fitted a deterministic of HIV transmission model to Yaoundé-specific demographic, behavioural, HIV and intervention coverage data in a Bayesian framework. We estimated the fraction of incident HIV infections averted by condoms and antiretroviral therapy (ART) and the fraction of all infections over 10-year periods directly and indirectly attributable to unprotected sex within and between each risk group.

**Results:** Condom use and ART together may have averted 43% (95% uncertainty interval: 31-54) of incident infections over 1980-2018 and 72% (66-79) over 2009-2018. Most onward transmissions over 2009-2018 stemmed from unprotected sex between lower-risk individuals (47% (32-61)), clients (37% (23-51)), and MSM (35% (20-54)) with all their partners. The contribution of commercial sex decreased from 25% (8-49) over 1989-1998 to 8% (3-22) over 2009-2018, due to higher intervention coverage among FSW.

**Conclusion:** Condom use and recent ART scale-up mitigated the HIV epidemic in Yaoundé and changed the contribution of different partnerships to onward transmission over time. Findings highlight the importance of prioritizing HIV prevention and treatment for MSM and clients of FSW whose unmet needs now contribute most to onward transmission, while maintaining services which successfully reduced transmissions in the context of commercial sex.

**Keywords:**

Mathematical model; HIV/AIDS, HIV incidence; Population attributable fraction; Condom use; Key populations

## Introduction

Cameroon has one of the largest, albeit declining, HIV epidemics across West and Central Africa.<sup>1</sup> By 2018, HIV prevalence was 3.6% amongst 15-49 year-old adults, and there were an estimated 540,000 people living with HIV (PLHIV).<sup>1</sup> Modeled estimates of HIV incidence suggest the epidemic peaked in the late 1990s.<sup>1</sup>

In Cameroon, as elsewhere, key populations (KP) such as female sex workers (FSW), their clients, and men who have with men (MSM) experience a high burden of HIV.<sup>2</sup> Despite same-sex practices and sex work being criminalized, research and programs addressing HIV prevention among sex workers have been in place since the emergence of HIV in Cameroon.<sup>3,4</sup> HIV prevalence among FSW and MSM in Cameroon were 21% and 25% in 2016, respectively,<sup>2</sup> with higher (45.1%<sup>2</sup>) HIV prevalence estimates among MSM in Yaoundé (Cameroon's largest city).<sup>5</sup>

FSW have been an important focus for early HIV research and prevention efforts in Yaoundé<sup>6-8</sup> through the World Health Organisation/Global Programme on AIDS which was introduced in Cameroon in 1987.<sup>9</sup> However, FSW received less attention after 2000 as the prevalence increased among pregnant women<sup>9</sup>. For example, only 5% and 2% of all funds allocated to HIV prevention in 2013 were estimated to be spent on programmes on FSW/clients and MSM, respectively.<sup>10</sup>

Clients of FSW have received little attention despite early local studies suggesting an important role due to them acting as a bridge in sexual networks.<sup>6</sup> Little was known about clients' sexual practices and HIV prevalence until a 2017 survey in Yaoundé, which found that HIV prevalence among clients of FSW (3%) was around twice as high as among all males.<sup>11</sup>

Variability in scale-up, reach, and access to condom-based prevention and antiretroviral treatment (ART) has led to differential coverage by risk group. Interventions have increased over time and mainly focused on condom use and ART since male circumcision is ubiquitous in Cameroon.<sup>12-14</sup> Self-reported condom-use at last sex among KP increased to high levels since the 1990s (in 2016, 75% of MSM, and 90% of FSW with their clients<sup>2</sup>) but has remained relatively low among lower-risk populations (~25%).<sup>13,15</sup> In contrast, access to ART became available in 2000,<sup>16,17</sup> with ART coverage among PLHIV reaching 52% in

2018.<sup>1,18</sup> Levels of viral suppression are highest in FSW and lowest in the male population.<sup>2,11,19</sup> Universal HIV testing and treatment was initiated in 2020.<sup>20</sup>

In other West and Central African countries, programs focused on the needs of FSW and their clients have been credited with reducing their HIV epidemics.<sup>21-23</sup> To date, there has been no assessment of the population-level impacts of the delivery of HIV services on the epidemic in Cameroon, and of the evolving contribution of KP to onward transmission, which is rarely estimated. Thus, we conducted a data-driven mathematical modeling study to 1) assess the potential population-level impact of condom-use and ART in risk groups on the HIV epidemic and, 2) assess the contribution of these groups or their partnerships, due to the remaining unmet HIV prevention and treatment needs, to onward HIV transmission over time, in Yaoundé, Cameroon.

## Methods

### Setting

Yaoundé has experienced one of the largest HIV epidemics in Cameroon. The estimated HIV prevalence among all 15-49 years old in the city reached a peak of 8.3% (95% confidence interval 6.7-10.1) in 2004<sup>24</sup> and declined to 4.4% (3.2-5.6) in 2017<sup>19</sup>. In 2016, HIV prevalence among FSW and MSM were 23.9% (20.4-27.8) and 45.1% (39.3-51.0),<sup>2</sup> respectively.

### Mathematical model description

We adapted a published deterministic compartmental model of HIV transmission,<sup>23</sup> and calibrated it to reproduce Yaoundé-specific demographic (15-49 years), behavioural, HIV epidemiological and intervention data over 1980-2018. We modelled a growing population using local census estimates.<sup>25</sup>

Individuals enter the modelled population at age 15 and leave due to ageing (age 50), or via background- and HIV-related mortality. There are six risk groups: lower-risk females and males, FSW, clients of FSW (“clients”), younger (15-24 years) and older (25-49 years) MSM (Figure S1a). The proportion of MSM, FSW and clients is stable over time and fitted to empirical estimates, [with FSW being defined as woman reporting commercial sex as their primary source of income](#).<sup>2,12,13,24-28</sup> In the model, HIV transmission occur within main, casual, and commercial partnerships between risk groups. MSM can have main and casual

partnerships with any female, whereas only FSW and clients engage in commercial sex (Table S1, Figure S1a). [Lower-risk females and males were defined as females and males not engaging in commercial sex nor sex between males \(Table S1, Figure S1a\).](#)

Following infection, PLHIV progress through acute infection to untreated chronic infection (during which they experience HIV-related mortality) (Figure S1b). PLHIV initiate ART, with a fraction of treated PLHIV being virally suppressed and experiencing reduced HIV-related mortality and suppressed infectivity. Individuals may stop and re-start ART.

HIV transmission occurs through insertive/receptive vaginal/anal sex acts. The per-capita rate of acquiring HIV depends on the per-act HIV transmission probability by type of sex act, the annual number/types of new partners, number of sex acts involving and not involving condoms by partnership type (Table S2), sexual mixing between groups by partnership type, the HIV prevalence and probability of viral suppression among partners, and in the case of male partners, circumcision coverage. The per-act efficacy of condoms in reducing HIV acquisition/transmission was 71-98%,<sup>29-31</sup> while the per-partner efficacy of circumcision in reducing HIV acquisition was 38-68%.<sup>14</sup> Pre-Exposure Prophylaxis (PrEP) was not available in Cameroon in 2018 thus was not modelled.

### **Model parameterisation and fitting**

We parametrised and fitted the model within a Bayesian framework using importance resampling method<sup>32,33</sup> to account for uncertainties in parameters and estimates of demographic, epidemiological and intervention fitting outcomes (listed in Tables S1-3, details in supplement).

### **Data sources**

Several data sources informed these parameters and outcomes (Tables S1-3).

Data to inform the demography, sexual behaviour and HIV prevalence for the overall and lower-risk populations were sourced from the Census and United Nations Population Division, several Demographic Health Surveys (DHS) (1997, 2004, 2011, 2018)<sup>13,24,28,34,35</sup> and the 4-city study.<sup>15,27,36</sup> The sexual behaviours and levels of HIV intervention of KPs were based on local surveys spanning over 1990-2018.<sup>2,4,11,26,27,37-44</sup>



The size of each KP was determined based on existing studies in Yaoundé (mainly conducted after 2012),<sup>2,12,13,24-28</sup> which suggested that between 0.5-3.4% of 15-49 years old females were currently FSWs, and 0.5-2.3% and 2-20% (reflecting high heterogeneity in empirical estimates) of 15-49 years old males were MSM and clients of FSW, respectively (Figure S2).

The model reflected reported evolutions in condom-use over time among the different risk groups, and partnership types, the higher condom use among KPs – especially by FSW during commercial sex – compared to the rest of the population, and similar levels of condom use reported by MSM with male and female partners<sup>2</sup> (Figure 1, Table S2). National annual estimates of ART coverage over 2012-2017 were obtained from the Joint United Nations Programme on HIV and AIDS (UNAIDS)<sup>1</sup> which suggested an increase from 24% to 50% over the period. Around 80% of PLHIV on ART were virally suppressed in 2017.<sup>19</sup> The model reflected very high coverage (94-99%) of male circumcision.<sup>12,13,24</sup>

Biological parameters and intervention efficacy parameters related to condom-use, male circumcision, ART, and risk-group- specific level of viral suppression were sourced from published literature (Table S1).<sup>31,45-50</sup>

The model was fitted to 40 empirical fitting outcomes (Table S3) including: total population size (2015), HIV prevalence among all (15-49 years) females and males in Yaoundé (three time points<sup>12,13,24</sup>), HIV prevalence among FSW (five time points<sup>2,6,27,42,51</sup>), and recent HIV prevalence estimates among clients (one time point<sup>11</sup>) and MSM (two time points<sup>2,52,53</sup>). We screened-out simulations where ART coverage among PLHIV was inconsistent with sex-specific UNAIDS national estimates (Figure S3), and fitted the model to recent risk-group-specific estimates of viral suppression levels among PLHIV in Yaoundé<sup>2,11,19</sup> (Figure S4). A set of 1000 posterior fits was resampled (based on their individual log-likelihood<sup>32,33</sup>) from a pool of simulations consistent with each fitting outcomes. Additional available HIV prevalence data not used for model fitting were used for comparison.

### **Impact of condom use and ART**

The fraction of incident HIV infections directly or indirectly averted by past interventions (condoms, ART, and both) over a period $[t_0, t]$  ( $AF_{t_0-t}$ , equation 1), was derived by comparing the model cumulative number of incident HIV infections over $[t_0, t]$  in the baseline scenario with interventions ( $CI_{t_0-t}(int)$ ) with that from a counterfactual scenario without the

intervention (condoms and/or ART) in any of the risk group, or in a specific group over the same period, ( $CI_{t_0-t}(no\ int)$ ).

$$\text{Equation 1: } AF_{t_0-t} = \frac{CI_{t_0-t}(no\ int) - CI_{t_0-t}(int)}{CI_{t_0-t}(no\ int)}$$

The counterfactual scenarios assumed that the intervention had no efficacy in reducing HIV acquisition (condoms) and transmission (condoms and ART) and HIV-related mortality (ART) in relevant risk groups. The cumulative number of incident HIV infections and fraction of incident HIV infections averted were first calculated at the overall population level, then within each risk group, in order to identify which group most benefited from interventions.

We estimated four indicators to describe who acquired and transmitted infections the most (indicators 1,3) and the efficiency in acquisition and transmission (indicators 2,4).

### **Sources of HIV acquisition (indicator 1,2)**

We first estimated the distribution or fraction of all incident HIV infections over 2009-2018 occurring in each risk group (indicator 1), as well as the annual HIV incidence rate (expressed as the number of incident infections per 100 susceptible person-years), in each group (indicator 2).

### **Sources of onward HIV transmission (indicators 3,4)**

We derived the population attributable fraction ( $PAF_{t_0-t}$ , equation 2) as the fraction of all HIV infections directly or indirectly transmitted by specific partnership types between two specific groups over  $[t_0, t]$  (indicator 3). The  $PAF_{t_0-t}$  can also be interpreted as the sources of onward transmission.

$$\text{Equation 2: } PAF_{t_0-t} = \frac{CI_{t_0-t}(risk) - CI_{t_0-t}(no\ risk)}{CI_{t_0-t}(risk)}$$

Here,  $CI_{t_0-t}(risk)$  and  $CI_{t_0-t}(no\ risk)$  are the numbers of incident HIV infections in the presence and absence of transmission during sex between the relevant groups, respectively. The sum of the PAFs over mutually exclusive groups exceeds 100%<sup>54</sup> as it accounts for secondary transmissions which may overlap for different groups.

We derived the per-capita *HIV transmission rate* $_{t_0-t}$  (per 100 infected person-years, equation 3) from a specific risk group by dividing the total number of infections over $[t_0, t]$  due to the specific group by the cumulative number of person-years lived in the infected group over the same period ( $\int_{t_0}^t \# \text{infected (risk)}$ )<sup>55</sup> (indicator 4).

$$\text{Equation 3: } HIV \text{ transmission rate}_{t_0-t} = 100 \times \frac{CI_{t_0-t}(\text{risk}) - CI_{t_0-t}(\text{no risk})}{\int_{t_0}^t \# \text{infected (risk)}}$$

To identify where these direct and indirect transmissions occurred, HIV transmission rates were also calculated as the number of infections due to the specific risk group over $[t_0, t]$  occurring in another specific group.

$AF_{t_0-t}$ ,  $PAF_{t_0-t}$  and *HIV transmission rate* $_{t_0-t}$  were estimated over successive decades over 1989-2028. We report median estimates as well as 95% uncertainty interval (UI, 2.5<sup>th</sup> and 97.5<sup>th</sup> percentiles of the distribution) across the 1000 posterior fits. Uncertainties in  $PAF_{t_0-t}$  estimates are described in the supplementary.

## Results

### Model fitting and epidemic dynamics

Figures 1-2, S2-5 show that the baseline scenario reproduced available demographic data and time trends in HIV prevalence, and intervention coverage data across risk groups.

### Impact of condom use and ART

Over 1980-2018, increases in condom-use and ART may have averted 43.4% (31.4-53.6) of incident HIV infections in the overall population, and 72.2% (65.6-78.5) over 2009-2018 (Figure 1g).

Condom-use alone potentially averted more than a third, 38.7% (26.7-49.9) of incident HIV infections over 1980-2018, and 58.1% (51.5-68.1) over 2009-2018, due to increasing levels of condom use (Figure 2a-c). Condom-use between clients and all their partners was estimated to have averted the largest fraction of overall infections, followed by condom use

between MSM and all partners, and FSW and all partners, and between lower-risk groups with medians  $AF_{2009-2018}$  of 35.5%, 29.3%, 29.2%, 18.7%, respectively (Table 1).

In contrast, ART alone potentially averted 5.8% (3.9-9.1) of infections over 1980-2018, increasing to 32.1% (27.2-36.1) over 2009-2018 (Figure 2e-h, S3-4). ART use among MSM, FSW, clients, and lower-risk groups, respectively, averted 9.4%, 6.6%, 5.6%, and 17.4% of all HIV infections over 2009-2018.

### **Who benefited from HIV interventions**

Over 2009-2018, infections averted from the combination of HIV interventions ( $AF_{2009-2018}$ ) exceeded 71.2% in all groups except MSM. Impact was largest among clients ( $AF_{2009-2018}=81.6\%$  (72.1-88.0)), mainly due to higher levels of interventions among FSW compared to other groups. It was lowest among younger ( $AF_{2009-2018}=54.3\%$  (48.8-62.2)) and older ( $AF_{2009-2018}=42.3\%$  (36.6-49.3)) MSM, despite similar ART coverage and higher condom use as in the lower-risk groups because of higher per-act HIV risk (Figure S6).

### **Sources of HIV acquisition (indicators 1-2)**

Most HIV infections might have been acquired by lower-risk groups despite lower incidence rates (Figure S7). HIV incidence rates over 2009-2018 were almost 3-fold higher among KP than lower-risk groups; and among all females compared to all males. The highest HIV incidence rates were estimated among MSM (Figure S8, Table S4). The model suggested larger incidence declines among KP compared to lower-risk groups. The proportions of PLHIV by risk group over time is shown in Table S5.

### **Contribution of risk groups to onward HIV transmission (indicator 3)**

Over 2009-2018, the largest contributor to onward transmission was sex between lower-risk groups ( $PAF_{2009-2018} = 46.6\%$ ), sex between clients (36.5%), MSM (35.1%) and FSW (13.6%) with all partners (Figure 3, Table 2).

The estimated contribution of KPs was higher at the early stage of the epidemic and declining over time thereafter. The largest estimated decline was for the PAF from sex between FSW and all their partners (from  $PAF_{1989-1998}=33.1\%$  (13.0-57.6)) to  $PAF_{2009-2018}=13.6\%$  (5.9-29.2)) largely because of past increases in condom-use during commercial sex. Conversely, the contribution of sex between lower-risk individuals slightly increased over time (from  $PAF_{1989-1998}=37.5\%$  (23.3-55.6) to  $PAF_{2009-2018}=46.6\%$  (32.0-61.4) over 2009-2018). The high

PAF of MSM with all partners plateaued from the early 2000's (PAF<sub>2009-2018</sub>=35.1% (20.1-54.0)).

#### **Contribution of different partnership types within risk groups to onward transmission**

Over 2009-2018, the PAF of sex between MSM and their female partners (27.5% (15.9-44.5)) was slightly higher than the PAF of sex between MSM and their male partners (23.2% (12.7-36.3)) although the difference was larger earlier in the epidemic (Table 2), with less than <2% of transmissions due to infections from female to MSM.

For FSW, commercial sex initially contributed more to HIV transmission than sex between FSW and their non-commercial sex partners, although this difference attenuated for the last two decades (PAF<sub>2009-2018</sub>=8.3% and 9.0%, respectively), given the large increase in condom use during commercial sex. The large PAF<sub>2009-2018</sub> of sex between clients and all their partners over the past and next decade mainly stems from non-commercial sex rather than commercial sex, with more transmissions occurring from clients to their non-commercial partners (PAF<sub>2009-2018</sub>=26.7%) than from non-commercial partners to clients (PAF<sub>2009-2018</sub>=6.3%) across decades. Estimates of the PAF<sub>2009-2018</sub> of KP were most sensitive to uncertainties in their relative population size (Supplementary and Figures S9-S10).

#### **Efficiency of transmission (indicator 4)**

Despite the PAF of KPs declining over time, sex by KP living with HIV has consistently resulted in more direct and indirect transmissions per year living with the infection (>2-fold) than lower-risk individuals (Table S6). Nearly half of transmissions contributed by FSW over 2009-2018 were secondary infections acquired by lower-risk females (Table S7).

#### **Discussion**

Our study shows that condom use among KP and more recently ART scale-up, achieved a large impact on the HIV epidemic in Yaoundé. Past and current KP programmes have successfully reduced onward transmission in the context of commercial sex. As a result, most onward transmission is now due to unmet prevention/treatment needs of MSM, clients of FSW, and lower-risk groups.

We estimated that to date most of the prevention benefits were due to condom use, particularly among KPs partly because it was introduced and adopted earlier than ART. The magnitude of historical increases in condom use during commercial sex is consistent with estimates from Benin<sup>21</sup>, and Côte d'Ivoire.<sup>22</sup> ART population-level impact recently increased, mainly due to increased coverage in lower-risk population, and this estimated impact is similar to findings from other Western/Central African countries.<sup>22,23</sup> However, our findings suggest that further scale-up would be needed to achieve the UNAIDS target of reducing HIV incidence to below 1 infection per 1000 susceptible-year<sup>56</sup> before 2030 (Figure S5).

Improvements are expected because Cameroon has made ART universally available since 2020,<sup>20</sup> despite possible disruptions in HIV prevention due to COVID-19.

FSW and their clients benefitted the most from past interventions, and despite relatively high condom use, MSM benefited the least. These findings reflect the need for different levels of intervention coverage to address differential vulnerabilities across groups. The reason why the contribution of unmet needs of MSM remain high is a higher per-act transmission probability of anal intercourse compared to vaginal intercourse – which means higher coverage of condom use or other [public health \(e.g. PrEP\) and human rights interventions](#) ~~interventions (e.g. PrEP)~~ are needed to avert transmission. Our findings on the PAF of sex between men resemble that of Dakar, Senegal (42% over 1995-2005), and are higher than those reported in other settings (around 5-10%)<sup>22,23,57,58</sup> where HIV prevalence among MSM is lower. [Our estimate of the HIV incidence among MSM over the 2009-2018 \(8.6 \(7.0-9.9\) infections per 100 susceptible person-year\) is consistent with estimates from empirical studies conducted in Sub-Saharan Africa over the same time period,<sup>59,60</sup> but our predicted HIV incidence markedly decreases to 2.2 infections over 2019-2028 due to reported increases in coverage of interventions among MSM in Yaoundé compared to other settings. Additional empirical studies, using a high-standard sampling design would improve our understanding of these time dynamics and better inform the overall contribution of MSM to the HIV epidemic in Yaoundé.](#)

We found that clients of FSW may play an important role in current onward transmission. The reasons for these are: a) clients comprise a large fraction of the male population; b) complex sexual networks that link commercial sex and casual sex wherein clients have a higher number of casual sex partners than non-client males and thus an added layer of vulnerability beyond just their role in commercial sex networks; c) existing and

disproportionate prevention gaps among clients. Few studies have previously reported on the role of clients in HIV epidemics in Sub-Saharan Africa. In Côte d'Ivoire,<sup>22</sup> partnerships between clients and non-FSW over 2005-2015 contributed to 44% of overall transmissions<sup>22</sup> (34% over 2009-2018 in our study). Taken together, our findings suggest an important benefit of implementing HIV prevention and treatment services to reach clients of FSW, which has unfortunately been found extremely challenging and, to our knowledge, never been deployed at a large-scale.<sup>61-63</sup> Although these efforts would align well with the call to reach men in HIV prevention efforts, and given the challenges in recruiting and sustaining intervention in clients of FSWs, further modelling impact and cost-effectiveness of implementing different prevention interventions, including new promising tools (e.g. HIV self-testing, oral or long-acting injectable PrEP) in different risk populations, may be required. Lower-risk males were assumed to be able to form non-commercial sexual relationships with FSW (at a twice lower rate than clients), and thus the 'lower-risk males' included non-client partners of FSW.

We examined various indicators in our analyses of the contribution of specific partnerships and unmet needs of specific risk groups to the overall HIV epidemic. A key finding is that the per-capita contributions of KP remain high which are a better marker of disproportionate risk because they account for population size as well as differential risks of per-act transmissions. We found that it could be most efficient to prioritize partnerships at disproportionate risk of onward transmission – for example between MSM or clients and their female partners with interventions such as PrEP, increased condom and/or lubricant use, and early and effective ART.

Our results highlight the importance of examining past impact to understand current contribution. All published modelling studies estimated a decline in the contribution of commercial sex (or FSW) to HIV epidemics in Western/Central Africa,<sup>21-23,64,65</sup> with extremely high contributions in the early days of the epidemic. Our results (PAF<sub>2009-2018</sub>=8.3% (3.3-21.9) for the PAF of commercial sex) are in line with estimates for the past decade, between 5% and 19%.<sup>21-23</sup> The reason that unmet needs of FSW play a smaller role in the current epidemic is because of the gains achieved thus far. The significance of this finding is that FSW programmes should be continued in order to keep the HIV epidemic under control, especially in Cameroon where FSWs still experience high levels of stigma and violence, which increases their ability to acquire and transmit HIV.<sup>66</sup>

Our study has several strengths. It was based on a comprehensive search of the published and grey literature complemented with new data for KPs, including data on clients of FSW and size of KPs, which was specific to Yaoundé. We also had access to recent estimates of intervention coverage among KP based on [testing serological measures](#) of viral [load](#) suppression, improving our estimates over time. Our model was fitted into a Bayesian framework which accounted for uncertainties in parameters and outcomes. Our analysis relies on advanced epidemic indicators, such as the “per-capita” transmission rates which have been shown to be useful in furthering epidemic appraisal.

Our study also has limitations and highlighted important data gaps such as city-specific data on clients before 2017 in Yaoundé. Indirect information was traditionally obtained via studies among truck drivers on Cameroonian main roads. [However, these are imperfect proxy population for FSW clients in Yaoundé due to the geographical location of the truck stops \(outside the city of Yaoundé\) and the specific socioeconomic status of this population among all FSW clients, an imperfect proxy population,](#)<sup>51,67,68</sup> and our model was in good agreement with national estimates among clients in the DHS. We predicted high HIV prevalence among MSM before the first empirical survey in 2011,<sup>26</sup> before the reduction in their risk following intervention uptake, but no data were available to validate model predictions. As a result, historical estimates of the contribution of MSM to the HIV epidemic in Yaoundé should be interpreted with caution, whereas our recent estimates are less uncertain since they rely on two consistent local surveys.<sup>2,26</sup> We identified the size of KP as the main factor of uncertainty around estimates of the contribution of KPs to the HIV epidemic. However, KPs size estimate were based on the best data available and the substantial variability across studies was taken into account in the analysis and reflected in the UI of the model estimates (from 2% of clients among males<sup>27</sup> to >20%<sup>28</sup>). [Our estimate of the contribution of MSM to overall transmission which are based on the best available data may nevertheless be under or over-estimated due to data limitation. First, the data used to derive our wide prior range \(0.5% to 2.3%\) of MSM population size estimates may still be an underestimate stemming from measurement bias in a context where sex between men is criminalized. Our results suggested a near 2-fold variation of PAF estimates if MSM pop size varied 2-fold. Second, although our model tried to capture uncertainty by using a Bayesian fitting method, reporting of behavioural data by any population is always subject to desirability bias, whereas assessing the levels and coverage of HIV interventions can also be influenced by selection and reporting biases. These issues were addressed by accounting for the full uncertainty in empirical estimates of levels of condom](#)



use and viral suppression in all populations, and by assuming an over reporting of condom-use consistency (up to 25% relative) for all sex acts, which should have sensibly reduced the effects of these different potential biases. Vertical HIV transmission was not considered in the model based on decreasing rates of vertical transmission over time in Cameroon, alongside low probability of survival into adolescence among infants living with HIV due to limited HIV treatment coverage over the period of time considered,<sup>69</sup> and is unlikely to have influenced PAF estimates. Our model did not represented the population aged 50 years or older, due to large data gaps regarding this age group, but UNAIDS has estimated that only around 5% of new HIV infections in Cameroon occur in this population, thus we assumed that this simplification did not significantly alter our estimates.<sup>1</sup>

In conclusion, our study shows that despite the HIV epidemic in Yaoundé being characterised as “generalised” (with overall HIV prevalence >1%), unmet prevention and treatment needs among KP still contribute to a substantial proportion of transmitted infections. In particular, since women are the ones who acquire most infections, addressing the treatment needs of clients and MSM, who have substantial non-commercial sex or sex with women, would be an efficient and significantly impact the HIV epidemic in Yaoundé, as they contribute the majority of transmissions but represent a minority of the overall population.

## References

1. UNAIDS. AIDSInfo web page. <http://aidsinfo.unaids.org/>. Published 2020. Accessed 14/03/2020.
2. Johns Hopkins School of Public Health C, Metabiota. *2016 Integrated Biological and Behavioral Surveillance (IBBS) survey among female sex workers and men who have sex with men in Cameroon, Final Report*. Yaoundé. 2018. Available from [http://onsp.minsante.cm/sites/default/files/publications/245/IBBS2016\\_preliminaryreport\\_final\\_Mar2017\\_linked.pdf](http://onsp.minsante.cm/sites/default/files/publications/245/IBBS2016_preliminaryreport_final_Mar2017_linked.pdf)
3. Global Network of Sex Work Projects. Sex work in Cameroon. <https://www.nswp.org/news/sex-work-cameroon>. Published 2014. Accessed 23/03/2019.
4. Decker MR, Lyons C, Billong SC, et al. Gender-based violence against female sex workers in Cameroon: prevalence and associations with sexual HIV risk and access to health services and justice. *Sex Transm Infect*. 2016;92(8):599-604.
5. Central Intelligence Agency. The World Factbook: AFRICA :: CAMEROON. [https://www.cia.gov/library/publications/the-world-factbook/geos/print\\_cm.html](https://www.cia.gov/library/publications/the-world-factbook/geos/print_cm.html). Published 2018. Accessed 25/02/2019.
6. Garcia-Calleja JM, Ngoumou P, Boupda A. *HIV Seroprevalence Study among Commercial Sex Workers in Yaounde and Douala*. Yaounde. 1992.
7. Garcia-Calleja J.M, Sam-Abbenyi A., O'Dell V., Bowmans J., Mbouni G., R. S. A review of KABP studies on AIDS and HIV infection in Cameroon. *Bull liais doc - OCEAC*. 1993;53:195-199.
8. Kaptue L, Monny-Lobe M, Zekeng L, Lamptey P, Potts M, Douglas N. HIV-1 infection among prostitutes in Yaoundé (Cameroon). Paper presented at: 3rd International meeting on AIDS in Africa 1988; Tanzania.
9. Mbanya D, Sama M, Tchounwou P. Current status of HIV/AIDS in Cameroon: how effective are control strategies? *Int J Environ Res Public Health*. 2008;5(5):378-383.
10. Comité National de Lutte Contre le Sida. *Estimation des ressources et dépenses 2013 de La lutte contre Le VIH et Sida*. 2014.
11. Tamoufe U, Mfochive I, Bowring A, Fako G, Bennett J. *Review of preliminary data from a study of HIV and risk among clients of female sex workers in Yaounde, Cameroon*. 2018.
12. Auvert B, Buve A, Ferry B, et al. Ecological and individual level analysis of risk factors for HIV infection in four urban populations in sub-Saharan Africa with different levels of HIV infection. *Aids*. 2001;15:S15-S30.
13. République du Cameroun. *Enquête démographique et de santé et à indicateurs multiples (EDS-MICS) 2011*. 2012.
14. Siegfried N, Muller M, Deeks JJ, Volmink J. Male circumcision for prevention of heterosexual acquisition of HIV in men. *Cochrane Database Syst Rev*. 2009(2):CD003362.
15. Lagarde E, Auvert B, Chege J, et al. Condom use and its associations with HIV/sexually transmitted diseases in four urban communities of sub-Saharan Africa. *Aids*. 2001;15:S71-S78.
16. Eboko F, Abé C, Laurent C. *Accès décentralisé au traitement du VIH/sida : évaluation de l'expérience camerounaise*. 2010.
17. Mosoko JJ, Akam W, Weidle PJ, et al. Retention in an antiretroviral therapy programme during an era of decreasing drug cost in Limbe, Cameroon. *Journal of the International Aids Society*. 2011;14.
18. Fonsah JY, Njamnshi AK, Kouanfack C, et al. Adherence to Antiretroviral Therapy (ART) in Yaounde-Cameroon: Association with Opportunistic Infections, Depression, ART Regimen and Side Effects. *Plos One*. 2017;12(1).

19. PEPFAR. *Cameroon Population-Based HIV Impact Assessment, CAMPHIA 2017. Summary Sheet: Preliminary Findings*. 2018.
20. Comité National de Lutte Contre le Sida. Le ministère de la santé publique officialise la gratuite de la prise en charge des patients. <http://www.cnls.cm/actualit%C3%A9s/cameroun-%E2%80%93-vih-sida-le-ministere-de-la-sante-publique-officialise-la-gratuite-de-la-prise>. Published 2019. Accessed 6/09/2019, 2019.
21. Williams JR, Alary M, Lowndes CM, et al. Positive Impact of Increases in Condom Use among Female Sex Workers and Clients in a Medium HIV Prevalence Epidemic: Modelling Results from Project SIDA1/2/3 in Cotonou, Benin. *Plos One*. 2014;9(7).
22. Maheu-Giroux M, Vesga JF, Diabate S, et al. Changing Dynamics of HIV Transmission in Cote d'Ivoire: Modeling Who Acquired and Transmitted Infections and Estimating the Impact of Past HIV Interventions (1976-2015). *J Aids-J Acq Imm Def*. 2017;75(5):517-527.
23. Mukandavire C, Walker J, Schwartz S, et al. Estimating the contribution of key populations towards the spread of HIV in Dakar, Senegal. *J Int AIDS Soc*. 2018;21 Suppl 5:e25126.
24. République du Cameroun. *Enquête Démographique et de Santé, Cameroun 2004*. 2005.
25. The World Bank. *Mapping Female Sex Workers in Cameroon for HIV Program Design and Implementation*. Washington DC. 2016.
26. Ministry of Public Health. *HIV Prevention for Populations at Risk in Cameroon: final report*. Yaoundé. 2014. Available from [https://www.jhsph.edu/research/centers-and-institutes/research-to-prevention/publications/WA\\_FinalReport.pdf](https://www.jhsph.edu/research/centers-and-institutes/research-to-prevention/publications/WA_FinalReport.pdf)
27. Morison L, Weiss HA, Buve A, et al. Commercial sex and the spread of HIV in four cities in sub-Saharan Africa. *AIDS*. 2001;15 Suppl 4:S61-69.
28. République du Cameroun. *Enquête Démographique et de Santé, Cameroun 1998*. 1999.
29. Giannou FK, Tsiara CG, Nikolopoulos GK, et al. Condom effectiveness in reducing heterosexual HIV transmission: a systematic review and meta-analysis of studies on HIV serodiscordant couples. *Expert Rev Pharmacoecon Outcomes Res*. 2016;16(4):489-499.
30. Liu HX, Su YY, Zhu L, Xing JN, Wu J, Wang N. Effectiveness of ART and Condom Use for Prevention of Sexual HIV Transmission in Serodiscordant Couples: A Systematic Review and Meta-Analysis. *Plos One*. 2014;9(11).
31. Weller S, Davis K. Condom effectiveness in reducing heterosexual HIV transmission. *Ugeskrift Laeger*. 2002;164(47):5497-5500.
32. Rubin DB. Using the SIR algorithm to simulate posterior distributions. In: *Bayesian Statistics*. Oxford University Press; 1998.
33. Alkema L, Raftery AE, Brown T. Bayesian melding for estimating uncertainty in national HIV prevalence estimates. *Sex Transm Infect*. 2008;84:111-116.
34. Direction Nationale du Deuxième Recensement Général de la Population et de l'Habitat Y, Cameroun,. *Enquête Démographique et de Santé Cameroun 1991*. 1992.
35. République du Cameroun. *Enquête Démographique et de Santé du Cameroun 2018*. 2019.
36. Ferry B, Carael M, Buve A, et al. Comparison of key parameters of sexual behaviour in four African urban populations with different levels of HIV infection. *AIDS*. 2001;15 Suppl 4:S41-50.
37. Lai Y. *Regional Differences in HIV Epidemic and HIV testing services usage among Cameroonian female sex workers: Comparing Grand North and Grassland Regions*. Baltimore, Maryland. 2015.
38. Ryan KA, Roddy RE, Zekeng L, Weir SS, Tamoufe U. Characteristics associated with prevalent HIV infection among a cohort of sex workers in Cameroon. *Sex Transm Infect*. 1998;74(2):131-135.
39. Kaptue L, Zekeng L, Djoumessi S, Monnylobe M, Nichols D, Debuysscher R. Hiv and Chlamydia Infections among Prostitutes in Yaounde, Cameroon. *Genitourin Med*. 1991;67(2):143-145.

40. Lorente N, Henry E, Fugon L, et al. Proximity to HIV is associated with a high rate of HIV testing among men who have sex with men living in Douala, Cameroon. *Aids Care*. 2012;24(8):1020-1027.
41. Comité National de Lutte contre le VIH/SIDA (CNLS). *Etude intégrée de surveillance comportementale et biologique chez les hommes ayant des rapports sexuels avec des hommes au Cameroun*. Cameroun. 2012.
42. Tamoufe U, Medang R. *Enquete séroépidémiologique et compartmentale sur le VIH/SIDA et à la Syphilis chez les travailleurs du sexe au Cameroun*. 2010.
43. Papworth E, Grosso A, Ketende S, et al. *Examining Risk Factors for HIV and Access to Services among Female Sex Workers (FSW) and Men Who Have Sex with Men (MSM) in Burkina Faso, Togo and Camereroon*. Baltimore, Maryland. 2014.
44. Roddy RE, Zekeng L, Ryan KA, Tamoufe U, Weir SS, Wong EL. A controlled trial of nonoxynol 9 film to reduce male-to-female transmission of sexually transmitted diseases. *N Engl J Med*. 1998;339(8):504-510.
45. Hughes JP, Baeten JM, Lingappa JR, et al. Determinants of per-coital-act HIV-1 infectivity among African HIV-1-serodiscordant couples. *J Infect Dis*. 2012;205(3):358-365.
46. Rodger AJ, Cambiano V, Bruun T. Sexual Activity Without Condoms and Risk of HIV Transmission in Serodifferent Couples When the HIV-Positive Partner Is Using Suppressive Antiretroviral Therapy (vol 316, pg 171, 2016). *Jama-J Am Med Assoc*. 2016;316(19):2048-2048.
47. Hollingsworth TD, Anderson RM, Fraser C. HIV-1 transmission, by stage of infection. *J Infect Dis*. 2008;198(5):687-693.
48. Boily MC, Baggaley RF, Wang L, et al. Heterosexual risk of HIV-1 infection per sexual act: systematic review and meta-analysis of observational studies. *Lancet Infect Dis*. 2009;9(2):118-129.
49. Zwahlen M, Wandel S, Egger M, et al. Duration from seroconversion to eligibility for antiretroviral therapy and from ART eligibility to death in adult HIV-infected patients from low and middle-income countries: collaborative analysis of prospective studies. *Sex Transm Infect*. 2008;84:131-136.
50. Reniers G, Slaymaker E, Nakiyingi-Miiro J, et al. Mortality trends in the era of antiretroviral therapy: evidence from the Network for Analysing Longitudinal Population based HIV/AIDS data on Africa (ALPHA). *Aids*. 2014;28:S533-S542.
51. Mosoko JJ, Macauley IB, Zoungkanyi ACB, Bella A, Koulla-Shiro S. Human Immunodeficiency Virus Infection and Associated Factors among Specific Population Subgroups in Cameroon. *Aids Behav*. 2009;13(2):277-287.
52. Park JN, Papworth E, Kassegne S, et al. HIV prevalence and factors associated with HIV infection among men who have sex with men in Cameroon. *Journal of the International AIDS Society*. 2013;16(4Suppl 3):18752.
53. Holland CE, Papworth E, Billong SC, et al. Access to HIV Services at Non-Governmental and Community-Based Organizations among Men Who Have Sex with Men (MSM) in Cameroon: An Integrated Biological and Behavioral Surveillance Analysis. *Plos One*. 2015;10(4).
54. Rowe AK, Powell KE, Flanders WD. Why population attributable fractions can sum to more than one. *Am J Prev Med*. 2004;26(3):243-249.
55. Silhol R, Boily MC, Dimitrov D, et al. Understanding the HIV epidemic among MSM in Baltimore: a modelling study estimating the impact of past HIV interventions and who acquired and contributed to infections. *J Acquir Immune Defic Syndr*. 2020.
56. UNAIDS. *Making the end of AIDS real: consensus building around what we mean by "epidemic control"*. 2017.
57. Volz EM, Ndemi N, Nowak R, et al. Phylodynamic analysis to inform prevention efforts in mixed HIV epidemics. *Virus Evol*. 2017;3(2).

58. Wirtz AL, Walker DG, Bollinger L, et al. Modelling the impact of HIV prevention and treatment for men who have sex with men on HIV epidemic trajectories in low- and middle-income countries. *Int J Std Aids*. 2013;24(1):18-30.
59. Hessou SPH, Glele-Ahanhanzo Y, Adekpedjou R, et al. HIV incidence and risk contributing factors among men who have sex with men in Benin: A prospective cohort study. *Plos One*. 2020;15(6):e0233624.
60. Nowak RG, Mitchell A, Crowell TA, et al. Individual and Sexual Network Predictors of HIV Incidence Among Men Who Have Sex With Men in Nigeria. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2019;80(4).
61. Lowndes CM, Alary M, Labbe AC, et al. Interventions among male clients of female sex workers in Benin, West Africa: an essential component of targeted HIV preventive interventions. *Sex Transm Infect*. 2007;83(7):577-581.
62. Faugier J, Cranfield S. Reaching male clients of female prostitutes: the challenge for HIV prevention. *Aids Care*. 1995;7 Suppl 1:S21-32.
63. Wulandari LPL, Guy R, Kaldor J. Systematic Review of Interventions to Reduce HIV Risk Among Men Who Purchase Sex in Low- and Middle-Income Countries: Outcomes, Lessons Learned, and Opportunities for Future Interventions. *Aids Behav*. 2020.
64. Low A, Nagot N, Konate I, et al. Potential Impact of Existing Interventions and of Antiretroviral Use in Female Sex Workers on Transmission of HIV in Burkina Faso: A Modeling Study. *J aids-J Acq Imm Def*. 2015;68:S180-S188.
65. Boily MC, Pickles M, Alary M, et al. What Really Is a Concentrated HIV Epidemic and What Does It Mean for West and Central Africa? Insights From Mathematical Modeling. *J aids-J Acq Imm Def*. 2015;68:S74-S82.
66. Lyons CE, Schwartz SR, Murray SM, et al. The role of sex work laws and stigmas in increasing HIV risks among sex workers. *Nat Commun*. 2020;11(1):773.
67. Unknown. HIV and STD prevalence among bus and truck drivers in Cameroon. *AIDS Anal Afr*. 1994;4(5):2.
68. Comité National de Lutte Contre le Sida. *Enquête intégrée de surveillance comportementale et biologique chez les camionneurs de longues distances au Cameroun (2016)*. 2017.
69. UNICEF. *UNICEF Cameroon Country Programme 2018-2020, Strategy Note*. 2017.

## Figure Captions

**Figure 1.** Proportion of sex acts protected by condoms for a) MSM with their male (red) and female (blue) partners, b) FSW and their commercial (red) and non-commercial (blue) partners, c) lower-risk individuals with their main (blue) and casual (red) partners. Condom-use data is described in Table S2; Proportion of PLHIV who are virally suppressed for d) MSM, e) FSW, and f) lower-risk populations. The model is also fitted to UNAIDS estimates of the ART coverage at the national level<sup>1</sup> (Figure S4). Dashed lined represent UNAIDS 90-90 targets (73% virally suppressed among PLHIV). Dots represent data and 95% confidence interval. Line and shaded areas represent median model predictions and 95% uncertainty intervals. In (g) Estimated proportions (AF) of HIV infections in the total population averted by the use of condoms alone, ART alone, and due to condoms and ART together over successive 10-year time-periods. Boxplots represent median (middle line), interquartile range (25 and 75% percentiles and 95% UI (whiskers).

**Figure 2.** Trends in empirical and modelled HIV prevalence among a) all 15-49 years old females, b) all 15-49 years old males, c) all 15-49 years old clients of FSW, d) all 15-49 years old FSW, and all MSM aged e) 15-24 years and f) 25-49 years in Yaoundé, Cameroon. Red squares and interval represent empirical estimates use for model fitting (Table S2), while blue shade represent model 95% uncertainty interval. Grey squares and intervals reflect estimates only used for comparison<sup>7,13,19,24,35</sup>.

**Figure 3.** Model predictions of the contribution of different partnerships to incident HIV infections not covered by interventions over the last three decades in Yaoundé. Boxplots represent median, interquartile range and 95%UI.