



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## MRI cannot replace specialized radiographs prior to unicompartmental knee arthroplasty.

Mortensen, Jacob; Radev, Dimitar; Rasmussen, Lasse; Østgaard, Svend Erik; Kappel, Andreas; Odgaard, Anders

*Published in:*  
Abstracts - Oktober 2020

*Creative Commons License*  
CC BY 4.0

*Publication date:*  
2020

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

### *Citation for published version (APA):*

Mortensen, J., Radev, D., Rasmussen, L., Østgaard, S. E., Kappel, A., & Odgaard, A. (2020). MRI cannot replace specialized radiographs prior to unicompartmental knee arthroplasty. In *Abstracts - Oktober 2020: DOS Bulletin* (pp. 98). [84.] Dansk Ortopædisk Selskab. <https://ortopaedi.dk/abstractbog/2020/98/>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

## MRI cannot replace specialized radiographs prior to unicompartmental knee arthroplasty.

# 84.

*Jacob Mortensen, Dimitar Radev, Lasse Rasmussen, Svend Erik Østgaard, Andreas Kappel, Anders Odgaard*

Orthopedic Department, Hip & Knee Unit, Gentofte Hospital; Radiology Department, Gentofte Hospital;

Orthopedic Department, Hip & Knee Unit, Vejle Sygehus; Orthopedic Department, Hip & Knee Unit, Aalborg University Hospital

**Background:** Choosing the optimal diagnostic approach to knee osteoarthritis could save both the radiation of extra radiographs and costly examinations in the diagnostic process.

**Purpose / Aim of Study:** The purpose of this study was to compare the joint space width of specialized radiography to the cartilage thickness on MRI scans in patients undergoing Unicompartmental and Total Knee Replacements.

**Materials and Methods:** A prospective study, including 60 patients. Specialized radiographs were taken with the Skyline view, the Rosenberg view, and coronal stress radiography. Experienced knee surgeons performed measurements of joint space width (JSW) and minimal joint space width (mJSW) on all radiographs. One experienced radiologist performed measurements of cartilage height on MRI scans. Radiographic measurements of each radiographic technique were used to compare with cartilage height measurements in MRI scans, in each respective knee compartment.

**Findings / Results:** When comparing specialized radiography with MRI, a weak correlation was found in the patellofemoral compartment (Medial facet: JSW/mJSW;  $\rho = 0.39/0,35$  ; CI = 0.07-0.58/0.09-.058 ;  $p < 0.005$ ) (Lateral facet: JSW/mJSW;  $\rho = 0.28/0,32$  ; CI = 0.03- 0.5/0.06-.05 ;  $p < 0.016$ ), a negligible and non- significant correlation was found in the medial compartment, and a moderate to strong correlation in the lateral compartment (Rosenberg view: JSW/mJSW;  $\rho = 0.56/0,62$  ; CI = 0.3-0.8/0.4-.8 ;  $p < 0.000$ ) (Valgus stress: JSW/mJSW;  $\rho = 0.7/0,61$  ; CI = 0.5-0.84/0.4-.77 ;  $p < 0.000$ ).

**Conclusions:** MRI by itself cannot and should not replace these specialized radiographic methods when choosing implant type. MRI should be reserved for more special cases where abnormal radiography or suspicion of atypical clinical findings present themselves. We recommend that a work-up of patients for mUKA include a skyline view with a Rosenberg view projection as a standard, and avoid the extra costs of MRI scan and/or extra radiation of additional special radiographs.