



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Forage-Based Beef Cattle Production in Red Soils Region of Southern China

Y. Y. He

Jiangxi Agricultural University, China

Guoqing Xie

Jiangxi Agricultural University, China

Q. H. Shi

Jiangxi Agricultural University, China

S. L. Wen

Chinese Academy of Agricultural Sciences, China

J. R. Liu

Jiangxi Agricultural University, China

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/9-2/40>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Y. Y. He, Guoqing Xie, Q. H. Shi, S. L. Wen, J. R. Liu, Z. Y. Wu, M. G. Xu, W. J. Zhang, and H. X. Li

Forage-based beef cattle production in red soils region of southern China

He YY¹ Xie GQ^{1*} Shi QH¹ Wen SL² Liu JR¹ Wu ZY³ Xu MG² Zhang WJ⁴ Li HX³

¹ Jiangxi Agricultural University, Nanchang 330045

² Red Soils Experiment Station of CAAS, Hengyang 421001

³ Jiangxi Province Livestock Technology Extension Station, Nanchang 330046

⁴ Taihe Animal Husbandry Bureau, Taihe 343700

* Corresponding author: Xie GQ, E-mail: guoqingxie@163.com

Key words : forage varieties selection, forage and roughage preparation, feeding methods, performances, beef cattle

Materials and methods Materials : (*Pennisetum purpureum* × *P.americanum*) × *P.purpureum* cv .Mott, Dwarf elephant grass, Solander setaria, Chicory, Wynnecassia, Lotononis, rye grass, king grass, rice straw, rice bran, cottonseed meal, ammonia bicarbonate, urea, ammoniated water and beef cattle hybrid. Methods : plot trials, feeding experiments, chemical analysis.

Results 1) Impact of seasonal change on yield and quality of the tested pasture species Among the tested species, (*Pennisetum purpureum* × *P.americanum*) × *P.purpureum* cv .Mott produced the highest leaf yield (14,500 kgDM/ha) and total yield (20,000 kgDM/ha) followed by Dwarf elephant grass and solander setaria. (*Pennisetum purpureum* × *P.americanum*) × *P.purpureum* cv .Mott, Dwarf elephant grass, Solander setaria, Chicory, Wynnecassia and Lotononis can grow from March to November providing available feeds for a period of 6 months from May to October. The protein content ranged from 12.4-19.0% for (*Pennisetum purpureum* × *P.americanum*) × *P.purpureum* cv .Mott, Dwarf elephant grass, Solander setaria and Chicory and from 13.9-19.7% for Lotononis and wynnecassia. The curve of protein content of all the tested pastures in growth season shaped as "V" type (Win et al 2007).

2) Preparation of fresh forage and roughage The effects of ammonia sources, N inclusion, ammoniation duration and chop length on the quality of ammoniated rice straw were studied. Results showed that ammoniated water was the best material to treat rice straw. The nutritive value of ammoniated rice straw increased as the amount of ammoniated water increased and the duration prolonged (He et al 2005). An experiment was conducted to compare the nutritive values between corn silage and king grass silage, with corn silage having the better nutritive values (Xie et al 2004).

3) Study on the techniques of beef cattle year round feeding Beef cattle hybrids fed on ammoniated rice straw supplemented with 7g corn and 3g cottonseed meal (kg LW/d) had the highest ADG (1.44kg) when compared to rice straw, microbial fermented rice straw, corn silage with or without supplements (corn, rice bran, cottonseed meal, urea ect.) treatments (He et al 2004; Xie et al 2005).

Beef cattle hybrids fed on Dwarf elephant grass supplemented with 5g rice bran and 5g cottonseed meal (kgLW/d) had the highest ADG (0.8kg) when compared to King grass, Paspalum atratum and Kudzu treatments with or without supplements (rice bran, cottonseed meal, urea ect.) (Xie et al 2004; Xie et al 2006; He et al 2006).

4) Study on the cycling and utilization of nutrients in ecosystem of forage-cattle-biogas There were no significant differences between the effects of different fertilizers (chemical fertilizer only and chemical fertilizer plus biogas fertilizer) on the yield and nutrient accumulation of rye grass or corn. Inclusion of biogas fertilizer could save synthetic chemical fertilizer by 1/3 and also improve the soil fertility (Liu et al 2003).

Conclusions (*Pennisetum purpureum* × *P.americanum*) × *P.purpureum* cv .Mott, Dwarf elephant grass, Wynnecassia, Lotononis and Kudzu could be recommended to establish in southern China for forage-based beef cattle production; the optimal feeding method for the beef cattle production would be: ammoniated rice straw (treated with ammoniated water (20% NH₃) at an inclusion of 20% or at 5% if with urea, with the addition of rice bran (5g/kgLW/d) and cottonseed meal (1.5g/kgLW/d) in winter; ammoniated rice straw plus rye grass (12gDM/kg LW/d) and rice bran (5g/kg LW/d) in spring; Dwarf elephant grass plus Kudzu vine (5g/kg LW/d), rice bran (5g/kg LW/d) and cottonseed meal (3g/kg LW/d) in summer and autumn.