



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Effect of Different Feeding Methods on Fatty Acids Composition in Muscle of Nanjiang Yellow Goat

Hailing Luo

China Agricultural University, China

Yongfeng Xu

China Agricultural University, China

Yingjun Zhang

China Agricultural University, China

Xiaohong Zhang

China Agricultural University, China

Kai Liu

China Agricultural University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/9-2/48>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Effect of different feeding methods on fatty acids composition in muscle of Nanjiang Yellow Goat

Luo Hailing, Xu Yongfeng, Zhang Yingjun, Zhang Xiaohong, and Liu Kai

College of Animal Science and Technology, China Agricultural University, Beijing 100094, China

Key words : Nanjiang Yellow Goat, kids, feeding method, fatty acids

Introduction Nanjiang Yellow Goat is a new and unique goat bred for mutton through selected-breeding for 40 years in China. It is well known for its high mutton quality. The feeding methods could affect mutton quality, including mutton taste (Arsenos et al., 2002). This study was aimed to investigate the effect of different feeding methods on fatty acids composition in muscle of Nanjiang Yellow Goat.

Materials and methods Nanjiang Yellow Goat Stud is in a semi-tropical climate region at the altitude of 1300-2000 m in Nanjiang county, Sichuan province, China. It is cold and humid in early spring, hot and short in summer and cold and long in winter. The annual averaged temperature, precipitation, relative humidity and evaporation are 16.2°C, 1199 mm, 72% and 1391 mm, respectively.

In this study, 32 two-month old weaning Nanjiang Yellow Goat healthy male kids with similar body weight (BW) were selected and randomly divided into two groups as control (grazing all day) and the treatment (grazing with supplement feeding). Three goat kids from each group were slaughtered to collect longissimus muscle samples at 2, 4 and 6 months for the fatty acid composition measurement. The data were subjected to one-way analysis of variance using GLM procedure in SAS (1989).

Results and discussion The fatty acid composition of longissimus muscle and gluteus of Nanjiang Yellow Goat kids was shown in Table 1. The content of Saturated Fatty Acids (SFA) has decisive influence on nutrition value of the mutton (Callow, 1958). Higher SFA could induce the amount of human blood cholesterol increasing (Grundy et al., 1982). In this study, the contents of SFA increased gradually as the age increases. However, compared with the control, the content of SFA was lower at 4 and 6 months. The results implied that the content of SFA in muscle was reduced by grazing with supplement feeding. The short-chain fatty acids (SCFA) are the metabolism production of the rumen microbe fermenting forage grass. The contents of SCFA between the two groups at the same age were not significantly difference.

Table 1 Fatty acid composition of longissimus muscle and gluteus of Nanjiang Yellow Goat kids.

Muscle	Item	2-month old		4-month old		6-month old	
		Treatment	Control	Treatment	Control	Treatment	Control
Longissimus muscle	SFA	47.74±3.35	47.18±3.79	44.60±2.02	48.99±5.23	46.86±1.93	53.65±7.66
	SCFA (C6-C12)	0.28±0.07 ^b	0.36±0.07 ^b	0.30±0.01 ^b	0.36±0.04 ^b	1.12±0.41 ^{ab}	1.92±1.38 ^a
Gluteus	SFA	49.94±3.26 ^a	43.39±4.15 ^b	54.49±4.3 ^a	55.40±1.10 ^a	56.19±3.07 ^a	58.03±4.3 ^a
	SCFA (C6-C12)	0.32±0.01 ^b	0.33±0.01 ^b	0.36±0.11 ^b	0.45±0.08 ^b	2.37±0.94 ^a	2.24±0.75 ^a

Means with different upper case letters within the same row differ $P < 0.01$. Means with different lower case letters within the same row differ at $P < 0.05$.

Conclusion Grazing with supplement feeding can increase the content of Unsaturated Fatty Acids (UFA) in Nanjiang Yellow Goat kids muscle, decrease the content of short-chain fatty acids.

References

- Arsenos, G., Banos, G., Fortomaris, P., Katsaounis, N., Stamataris, C., Tsaras, L., Zygoiannis, D. 2002. Eating quality of lamb meat: effects of breed, sex, degree of maturity and nutritional management. *Meat Sci.*, 60: 379-387.
- Callow, E. H. 1958. Comparative studies of meat. VI. Factors affecting the iodine number of fat from the fatty and muscular tissue of lambs. *J. Agr. Sci. (Camb.)* 51: 361.
- Grundy, S. M., Bilheimer, D., Blackburn, H., Brown, W. V., Kwiterovich, P. O., Mattson, F., Schonfeld, G., Weidman, W. H. 1982. Rational of the diet-heart statement of the American heart association: *Report of Nutrition committee*. *Circulation* 65: 839-854.