


NPS-MA-93-002

NAVAL POSTGRADUATE SCHOOL

Monterey, California



SOME INVERSE PROBLEMS FOR
JACOBI AND ARROW MATRICES

by

Carlos Borges
Ruggero Frezza
W. B. Gragg

Technical Report For Period
July 1992 - September 1992

Approved for public release; distribution unlimited

FedDocs
D 208.14/2
NPS-MA-93-002

Approved for: Naval Postgraduate School
Monterey, CA 93943

0002
202 13/2
S-K/ fi-93-002

NAVAL POSTGRADUATE SCHOOL
MONTEREY, CA 93943


Rear Admiral R. W. West, Jr.
Superintendent

Harrison Shull
Provost

This report was prepared in conjunction with research conducted for the Naval Postgraduate School and funded by the Naval Postgraduate School.

Reproduction of all or part of this report is authorized.

This report was prepared by:



UNCLASSIFIED
 SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
 OMB No 0704-0188

1a REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
4b DECLASSIFICATION/DOWNGRADING SCHEDULE		5 MONITORING ORGANIZATION REPORT NUMBER(S) NPS-MA-93-002	
1 PERFORMING ORGANIZATION REPORT NUMBER(S) NPS-MA-93-002		5 MONITORING ORGANIZATION REPORT NUMBER(S) NPS-MA-93-002	
5a NAME OF PERFORMING ORGANIZATION Naval Postgraduate School	6b OFFICE SYMBOL (If applicable) MA	7a NAME OF MONITORING ORGANIZATION Naval Postgraduate School	
6c ADDRESS (City, State, and ZIP Code) Monterey, CA 93943		7b ADDRESS (City, State, and ZIP Code) Monterey, CA 93943	
8a NAME OF FUNDING/SPONSORING ORGANIZATION Naval Postgraduate School	8b OFFICE SYMBOL (If applicable) MA	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER O & M,N Direct Funding	
8c ADDRESS (City, State, and ZIP Code) Monterey, CA 93943		10 SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO	PROJECT NO
		TASK NO	WORK UNIT ACCESSION NO
11 TITLE (Include Security Classification) Some Inverse Problems for Jacobi and Arrow Matrices			
12 PERSONAL AUTHOR(S) Carlos Borges, Ruggero Frezza, William B. Gragg			
13a TYPE OF REPORT Technical	13b TIME COVERED FROM 7/92 TO 9/92	14 DATE OF REPORT (Year, Month, Day) 21013	15 PAGE COUNT 9
16 SUPPLEMENTARY NOTATION			
17 COSATI CODES		18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
		Jacobi matrix, Arrow matrix, inverse problems	
19 ABSTRACT (Continue on reverse if necessary and identify by block number) We consider the problem of reconstructing Jacobi matrices and real symmetric arrow matrices from two eigenpairs. Algorithms for solving these inverse problems are presented. We show that there are reasonable conditions under which this reconstruction is always possible. Moreover, it is seen that in certain cases reconstruction can proceed with little or no cancellation. The algorithm is particularly elegant for the tridiagonal matrix associated with a bidiagonal singular value decomposition.			
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a NAME OF RESPONSIBLE INDIVIDUAL CARLOS BORGES		22b TELEPHONE (Include Area Code) (408) 646-2124	22c OFFICE SYMBOL MA/Bc

tridiagonal matrices, and let $\text{UST}_+(n)$ denote that subset of $\text{UST}(n)$ with positive β_i .

We wish to develop an algorithm to reconstruct T from the knowledge of two of its eigenpairs (λ, \mathbf{u}) and (μ, \mathbf{v}) . The eigenvector recurrence for symmetric tridiagonal matrices is

$$\beta_{i-1}u_{i-1} + \alpha_i u_i + \beta_i u_{i+1} = \lambda u_i, \quad (2)$$

where (λ, \mathbf{u}) is any eigenpair of T , u_i is the i th element of \mathbf{u} , and $\beta_0 = \beta_n = 0$. Applying this relation to both eigenpairs gives

$$\begin{aligned} \beta_{i-1}u_{i-1} + \alpha_i u_i + \beta_i u_{i+1} &= \lambda u_i, \\ \beta_{i-1}v_{i-1} + \alpha_i v_i + \beta_i v_{i+1} &= \mu v_i. \end{aligned}$$

Combining these two equations and eliminating α_i gives

$$\beta_{i-1}(v_i u_{i-1} - u_i v_{i-1}) + \beta_i(u_{i+1} v_i - v_{i+1} u_i) = (\lambda - \mu)u_i v_i. \quad (3)$$

Since $\beta_0 = \beta_n = 0$ we get the following initial and terminal conditions

$$\begin{aligned} \beta_1(u_2 v_1 - v_2 u_1) &= (\lambda - \mu)u_1 v_1 \quad (4) \\ \beta_{n-1}(v_n u_{n-1} - u_n v_{n-1}) &= (\lambda - \mu)u_n v_n. \quad (5) \end{aligned}$$

Combining (3) with (4) gives a special case of the Christoffel-Darboux identity,

$$\beta_i(u_{i+1} v_i - v_{i+1} u_i) = (\lambda - \mu) \sum_{k=1}^i u_k v_k \quad (6)$$

for $i = 1, 2, \dots, n-1$. There is also a backward formula,

$$\beta_i(u_{i+1} v_i - v_{i+1} u_i) = -(\lambda - \mu) \sum_{k=i+1}^n u_k v_k, \quad (7)$$

which follows from (3) and (5), or from (6) and the orthogonality of the eigenvectors. In a similar manner, we can show that

$$2\alpha_i u_i v_i = (\lambda + \mu)u_i v_i - \beta_{i-1}(u_i v_{i-1} + v_i u_{i-1}) - \beta_i(u_{i+1} v_i + v_{i+1} u_i). \quad (8)$$

This formula uses all of the available information but it is possible to obtain an equation for the α_i using the β_i and a single eigenpair with the formula

$$\alpha_i v_i = \mu v_i - \beta_{i-1} v_{i-1} - \beta_i v_{i+1}. \quad (9)$$

We can use these equations to reconstruct the original matrix from the two eigenpairs provided that no element of the two eigenvectors is zero

and that $v_i u_{i+1} - u_i v_{i+1} \neq 0$ for any $i = 1, 2, \dots, n-1$. If this is true, then the equations for the α_i simplify to

$$2\alpha_i = (\lambda + \mu) - \beta_{i-1} \left(\frac{v_{i-1}}{v_i} + \frac{u_{i-1}}{u_i} \right) - \beta_i \left(\frac{v_{i+1}}{v_i} + \frac{u_{i+1}}{u_i} \right) \quad (10)$$

or

$$\alpha_i = \lambda - \beta_{i-1} \frac{u_{i-1}}{u_i} - \beta_i \frac{u_{i+1}}{u_i}. \quad (11)$$

Notice that (10) is just the simple average of (11) over both eigenpairs. Using (11), (6), and (7) we can reconstruct the original matrix in $13n - 9$ flops.

In order to determine when these formulas can be applied, we need some additional results. We introduce the following fact from [10].

Fact 1 *Let $T \in \text{UST}_+(n)$ and assume that the eigenvalues are ordered so that $\lambda_1 > \lambda_2 > \dots > \lambda_n$. Then the number of sign changes between consecutive elements of the k th eigenvector of T , denoted s_k , is $k - 1$.*

We refer the reader to [10] for a proof but note that it can be derived from the Sturm sequence property for the characteristic polynomials of the principal submatrices. With this fact in hand we can prove the following theorem.

Theorem 1 *If $T \in \text{UST}_+(n)$ and if (λ, \mathbf{u}) and (μ, \mathbf{v}) are the extremal eigenpairs of T , that is $\lambda = \lambda_1$ and $\mu = \lambda_n$, then $v_i u_{i+1} - u_i v_{i+1} \neq 0$ for any $i = 1, 2, \dots, n-1$.*

Proof. The proof follows trivially by noting that the strict interlacing property for unreduced symmetric tridiagonals (see [12] p. 300) guarantees that none of the numbers $u_i, u_{i+1}, v_i, v_{i+1}$ can be zero. And, since u_i and u_{i+1} must have the same sign and v_i and v_{i+1} must have opposite signs (from fact 1), it follows that both terms in $u_i v_{i+1} - v_i u_{i+1}$ have opposite signs and are nonzero so this difference is really a sum of two strictly positive (negative) numbers and hence is not zero. \square

Hence, if we choose the two extremal eigenpairs of a given element of UST_+ we can always reconstruct the original matrix using the formulas above. Notice that the denominator is computed without cancellation in this case because of the sign pattern. Moreover, if we use the smallest (largest) eigenpair in (9) to get the α_i , then these can be reconstructed from the derived β_i and the data without further cancellation if the matrix is positive (negative) definite. If the matrix is indefinite then there is only one additional cancellation for each of the α_i . If the matrix is singular then choosing the eigenvector associated with the zero eigenvalue prevents further cancellation.

Note that any element of $\text{UST}(n)$ has exactly $2n - 1$ real degrees of freedom and that two eigenpairs contain $2n + 2$ numbers but, in fact, also have $2n - 1$ real degrees of freedom since there are two arbitrary scaling

parameters for the eigenvectors and a single orthogonality condition. The eigenpairs contain precisely the right amount of information.

This algorithm is especially robust when applied to the tridiagonal matrix associated with the bidiagonal SVD. It is well known [7] that the Jordan-Lanczos matrix

$$A = \begin{bmatrix} 0 & B^T \\ B & 0 \end{bmatrix}, \quad (12)$$

where $B \in \mathfrak{R}^{n \times n}$ is an unreduced bidiagonal with positive elements, can be reduced via the *perfect shuffle* to an unreduced tridiagonal T of the form

$$T = \begin{bmatrix} 0 & \beta_1 & & & & \\ \beta_1 & 0 & \beta_2 & & & \\ & \beta_2 & & \ddots & & \\ & & & \ddots & & \\ & & & & & \beta_{n-1} \\ & & & & \beta_{n-1} & 0 \end{bmatrix} \quad (13)$$

The matrix T is $2n \times 2n$ and its eigenvalues occur in plus-minus pairs. It is not difficult to show that if (λ, u) is an eigenpair of T then $(-\lambda, Su)$ is also an eigenpair where S is diagonal with 1 and -1 alternating as the diagonal elements. The reconstruction formula for this matrix simplifies considerably since we need only a single eigenpair. In particular, the β_i are given by

$$\beta_i = \frac{(-1)^i \lambda}{u_{i+1} u_i} \sum_{k=1}^i (-1)^k u_k u_k \quad (14)$$

As a special case of the more general algorithm it is obvious that the denominator $u_{i+1} u_i$ is not zero provided we use the eigenvector associated with the largest eigenvalue. Even more intriguing is that, provided none of the principal submatrices shares an eigenvalue with the full matrix, this denominator will be non-zero for any eigenpair since in this case no element of any eigenvector can be zero. In other words, the reconstruction from any eigenpair is well-posed provided that the given eigenvector has no zero elements. The algorithm requires $5n - 7$ flops working with (14) and the backward equation

$$\beta_i = \frac{(-1)^{i+1} \lambda}{u_{i+1} u_i} \sum_{k=n}^{i+1} (-1)^k u_k u_k. \quad (15)$$

Notice that this matrix has only $n - 1$ real degrees of freedom which is exactly what is given by one eigenpair since the eigenvector contains an arbitrary scaling parameter and must satisfy the special orthogonality condition

$$\sum_{i=1}^n (-1)^i u_i^2 = 0. \quad (16)$$

We point out that this algorithm can be interpreted as the reconstruction of an unreduced bidiagonal B from its largest singular value and both associated singular vectors.

2 Arrow Matrices

We can reconstruct the arrow matrix in a similar manner to that given above. The arrow is of some importance as it occurs in certain divide and conquer schemes for finding the eigenvalues of a tridiagonal matrix [1, 8]. The arrow is also an element of the class of symmetric acyclic matrices (as is the Jacobi matrix) and hence it is possible to find its eigenvalues with "tiny componentwise relative backward error", [5].

The general form of an arrow matrix is

$$A = \begin{bmatrix} \alpha_1 & & & \beta_1 \\ & \alpha_2 & & \beta_2 \\ & & \ddots & \vdots \\ & & & \alpha_{n-1} & \beta_{n-1} \\ \beta_1 & \beta_2 & & \beta_{n-1} & \gamma \end{bmatrix}. \quad (17)$$

If $\beta_i \neq 0$ for $i = 1, 2, \dots, n-1$ and if $\alpha_i \neq \alpha_j$ for any $i \neq j$ then we shall say that $A \in \text{USA}(n)$, where $\text{USA}(n)$ is the set of unreduced symmetric arrow matrices. Proceeding as before, we let (λ, \mathbf{u}) and (μ, \mathbf{v}) be two eigenpairs of A . The eigenvector recurrence is

$$\alpha_i u_i + \beta_i u_n = \lambda u_i, \quad (18)$$

$$\alpha_i v_i + \beta_i v_n = \mu v_i, \quad (19)$$

for $i = 1, 2, \dots, n-1$. Moreover, the eigenvector relation also gives

$$\gamma = \mu - \frac{1}{v_n} \sum_{i=1}^{n-1} \beta_i v_i, \quad (20)$$

for any eigenpair (μ, \mathbf{v}) of A . If we combine (18) and (19) and eliminate α_i , we get

$$\beta_i (v_i u_n - u_i v_n) = (\lambda - \mu) u_i v_i. \quad (21)$$

Similarly, eliminating β_i gives

$$\alpha_i (v_i u_n - u_i v_n) = \mu v_i u_n - \lambda u_i v_n. \quad (22)$$

This gives a very simple, easily vectorizable reconstruction algorithm. The only remaining question is whether the quantities $v_n u_i - u_n v_i$ are all

nonzero. In order to show that this is true under the correct conditions, we need to first establish some facts about the eigenvectors of an unreduced arrow matrix. We begin by noting that

$$A - \lambda I = \begin{bmatrix} D - \lambda I & \mathbf{b} \\ \mathbf{b}^T & \gamma - \lambda \end{bmatrix} \quad (23)$$

where $D = \text{diag}(\alpha_1, \alpha_2, \dots, \alpha_{n-1})$, and $\mathbf{b} = [\beta_1, \beta_2, \dots, \beta_{n-1}]^T$. Following [8] we compute the Gauss factorization

$$\begin{bmatrix} D - \lambda I & \mathbf{b} \\ \mathbf{b}^T & \gamma - \lambda \end{bmatrix} = \begin{bmatrix} I & 0 \\ \mathbf{b}^T(D - \lambda I)^{-1} & 1 \end{bmatrix} \begin{bmatrix} D - \lambda I & \mathbf{b} \\ 0^T & -f(\lambda) \end{bmatrix} \quad (24)$$

where f , the *spectral function*, is given by

$$f(\lambda) = \lambda - \gamma + \sum_{i=1}^{n-1} \frac{\beta_i^2}{\alpha_i - \lambda}. \quad (25)$$

From (24) and (25) the zeros of f are the eigenvalues of A . Furthermore, if A is unreduced, then the eigenvalues of A are strictly interlaced by the α_i . It follows that the eigenvector associated with λ is

$$\mathbf{v}(\lambda) = \begin{bmatrix} (\lambda I - D)^{-1} \mathbf{b} \\ 1 \end{bmatrix}. \quad (26)$$

Note that distinctness of the α_i is critical since it guarantees that $(\lambda I - D)$ is nonsingular. Combining this description of the eigenvectors with the fact that the α_i interlace the eigenvalues, we have the following fact

Fact 2 *Let A be an unreduced arrow matrix with $\beta_i > 0$ for $i = 1, 2, \dots, n-1$. Then the following hold.*

1. *If \mathbf{u} is any eigenvector of A then $u_i \neq 0$ for any $i = 1, 2, \dots, n$.*
2. *If we order the eigenvalues of A so that $\lambda_1 > \lambda_2 > \dots > \lambda_n$ and let \mathbf{u}_k be the eigenvector, from (26), associated with λ_k , then the first $k-1$ elements of \mathbf{u}_k are less than zero, and the last $n-k+1$ elements are greater than zero.*

Proof. The proof of the first fact follows directly from formula (26) and the interlacing property. The second fact follows from formula (26), the interlacing property, and the positivity of the β_i . \square

This simplifies the reconstruction formula since, if we assume that the eigenvectors are normalized so that their last elements are equal to one, the reconstruction formulas can be rewritten as

$$\alpha_i = \lambda - \frac{(\mu - \lambda)v_i}{u_i - v_i}$$

$$\beta_i = \frac{(\mu - \lambda)u_i v_i}{u_i - v_i} \quad (27)$$

$$\gamma = \mu - (\mu - \lambda) \sum_{i=1}^{n-1} \frac{u_i v_i^2}{u_i - v_i}$$

Using these formulas, we can reconstruct the arrow matrix in $10n - 8$ flops. Under the previously mentioned conditions, it is easily shown that none of the denominators in the reconstruction formula are zero and hence we can always reconstruct the matrix from two eigenpairs.

Theorem 2 *If A is an unreduced arrow matrix, and if λ and μ are any two distinct eigenvalues of A with associated eigenvectors \mathbf{u} and \mathbf{v} , normalized to have their last elements equal to one, then $u_i - v_i \neq 0$ for $i = 1, 2, \dots, n - 1$.*

Proof. Assume that $u_i = v_i$. The eigenvector relation implies that

$$\begin{bmatrix} u_i & 1 \\ v_i & 1 \end{bmatrix} \begin{bmatrix} \alpha_i \\ \beta_i \end{bmatrix} = \begin{bmatrix} \lambda u_i \\ \mu v_i \end{bmatrix} \quad (28)$$

which implies that $\lambda = \mu$, but this contradicts the distinctness of the eigenvalues. Hence, it follows that $u_i \neq v_i$. \square

The reconstruction algorithm has another very important property: if the two extremal eigenpairs (λ_1 and λ_n and their associated eigenvectors) are used, then the β_i can be found, up to the scaling factor $\lambda_1 - \lambda_n$, without cancellation. This follows from the fact that if the corresponding eigenvectors are normalized so that their last elements are both one, then all the remaining elements must have opposite signs. This is fortuitous since it means that the differences that appear in the denominator do not involve cancellation. Moreover, if A is indefinite there are no cancellations whatsoever in computing the β_i . Conversely, if A is definite there are no cancellations in computing the α_i . If A is semi-definite (and singular) then there is no cancellation at all, including the computation of γ . The computation of γ involves one cancellation if the matrix is indefinite, and none if it is definite, or semi-definite, provided we choose the correct eigenvector for its computation. In any case, whenever there is cancellation in this algorithm, it is benign.

3 Breakdown of the Jacobi reconstruction

On seeing that the reconstruction algorithm for the arrow is well posed for any two eigenpairs, it is tempting to believe that this might also be the case for the Jacobi matrix algorithm since the same conditions apply – unreduced, no principal submatrix shares an eigenvalue with the full matrix. Unfortunately, it is not true. Consider the matrix

$$\begin{bmatrix} 6 & 2 & 0 & 0 \\ 2 & 4 & 5 & 0 \\ 0 & 5 & 4 & 2 \\ 0 & 0 & 2 & 6 \end{bmatrix} \quad (29)$$

which is an element of UST_+ and satisfies the condition that no principal submatrix shares an eigenvalue with the full matrix. The eigenvalues of this matrix are 10 , $(5 + \sqrt{65})/2$, 5 , $(5 - \sqrt{65})/2$ and the eigenvectors associated with 10 and 5 are $[1 \ 2 \ 2 \ 1]^T$ and $[-2 \ 1 \ 1 \ -2]^T$, respectively. Note that the reconstruction algorithm breaks down for these two eigenpairs and cannot uniquely determine β_2 . Some manipulation of the scalar equations shows that the two eigenpairs in question are eigenpairs of any matrix of the form

$$\begin{bmatrix} 6 & 2 & 0 & 0 \\ 2 & 9 - \gamma & \gamma & 0 \\ 0 & \gamma & 9 - \gamma & 2 \\ 0 & 0 & 2 & 6 \end{bmatrix} \quad (30)$$

Fortunately, we can say a few things about breakdown. First of all, if the algorithm breaks down in the computation of β_i , then it cannot break down in the computation of β_{i-1} or β_{i+1} since this implies that two distinct eigenvalues share the same eigenvector. Second, if there is a breakdown then it is possible to reconstruct a parametrized matrix with the specified eigenpairs by setting $\beta_i = \gamma$ and solving for α_i and α_{i+1} in terms of γ . Setting $\gamma = 0$ will yield a reduced tridiagonal with the specified eigenpairs.

4 Stabilizing divide and conquer algorithms

We note that there are several other important inverse problems for the symmetric arrow matrix. Of interest, is the reconstruction of the symmetric arrow from the eigenvalues and the *shaft* of the arrow (i.e. the elements α_i). In this case it is also possible to reconstruct the arrow in a straightforward manner since we need only determine the β_i and the element γ . Clearly, we can obtain γ from the trace formula, that is

$$\gamma = \sum_{i=1}^n \lambda_i - \sum_{i=1}^{n-1} \alpha_i. \quad (31)$$

The β_i can be computed directly since the $-\beta_i^2$ are the residues of the *partial fraction decomposition*

$$f(\lambda) = \frac{\prod_{i=1}^n (\lambda - \lambda_i)}{\prod_{i=1}^{n-1} (\lambda - \alpha_i)} = \lambda - \gamma + \sum_{i=1}^{n-1} \frac{\beta_i^2}{\alpha_i - \lambda}. \quad (32)$$

Thus we have

$$\beta_j^2 = \lim_{\lambda \rightarrow \alpha_k} (\alpha_k - \lambda) f(\lambda) = - \frac{\prod_{i=1}^n (\alpha_j - \lambda_i)}{\prod_{i \neq j} (\alpha_j - \alpha_i)} \quad (33)$$

This algorithm is used in [3] for the reconstruction of a periodic Jacobi matrix. It can also be applied to stabilize the extension based tridiagonal divide and conquer algorithms [1, 8].

We note that this is very similar with the inverse problem first considered in [2] and then used in [9] to stabilize the modification based Cuppen-Dongarra-Sorensen algorithm [4, 6]. In particular, the zeros of the *spectral function*

$$f(\lambda) = 1 + \sum_{i=1}^n \frac{\beta_i^2}{\alpha_i - \lambda} \quad (34)$$

are the eigenvalues of $D + \mathbf{b}\mathbf{b}^T$. The authors of [9] show that loss of orthogonality in computing the eigenvectors can be avoided by using the computed eigenvalues $\tilde{\lambda}_i$ in the reconstruction formula

$$\tilde{\beta}_j^2 = \frac{\prod_{i=1}^n (\tilde{\lambda}_i - \alpha_j)}{\prod_{i \neq j} (\alpha_i - \alpha_j)} \quad (35)$$

and then computing the eigenvectors of $D + \tilde{\mathbf{b}}\tilde{\mathbf{b}}^T$ from their explicit expressions. The enlightened use of *shifts* of the origin [9] is crucial to both algorithms.

5 Acknowledgements

The first and third authors were supported by direct grant from the Naval Postgraduate School. The third author also acknowledges support from the Interdisciplinary Project Center for Supercomputing at the ETH, Zürich.

References

- [1] P. ARBENZ, *Divide and conquer algorithms for the bandsymmetric eigenvalue problem*, in Parallel Computing '91 (D.J. Evans, G.R. Joubert and H. Liddell, editors), Elsevier Science Publishers B.V., Amsterdam, 1992, pp. 151–158.
- [2] J.L. BARLOW, *Error analysis of update methods for the symmetric eigenvalue problem*, SIAM J. Matrix Anal. Appl., to appear.
- [3] D. BOLEY AND G.H. GOLUB, *A modified method for reconstructing periodic Jacobi matrices*, Math. Comp., 42, 165, (1984), pp. 143–150.
- [4] J.J.M. CUPPEN, *A divide and conquer method for the symmetric tridiagonal eigenproblem*, Numer. Math., 36, (1981), pp. 177–195.

- [5] JAMES W. DEMMEL AND W.B. GRAGG, *On computing accurate singular values of matrices with acyclic graphs*, Linear Algebra Appl., to appear.
- [6] J.J. DONGARRA AND D.C. SORESENSEN, *A fully parallel algorithm for the symmetric eigenvalue problem*, SIAM J. Sci. Statist. Comput., 8, (1987), pp. 139-154.
- [7] G.H. GOLUB AND W. KAHAN, *Calculating the singular values and pseudo-inverse of a matrix*, J. Soc. Indust. Appl. Math. Ser. B Numer. Anal., 2, (1965), pp. 205-224.
- [8] W.B. GRAGG, JOHN R. THORNTON AND DANIEL D. WARNER, *Parallel divide and conquer algorithms for the symmetric tridiagonal eigenproblem and bidiagonal singular value problem*, Modeling and Simulation, Volume 23, Proc. 23rd Annual Pittsburgh Conf., Univ. Pittsburgh School of Engineering, 1992.
- [9] M. GU AND S.C. EISENSTAT, *A stable and efficient algorithm for the rank-one modification of the symmetric eigenproblem*, Department of Computer Science, Yale University, working paper, 1992.
- [10] B. PARLETT, *The symmetric eigenvalue problem*, Prentice Hall Inc., New Jersey, 1980.
- [11] B. PARLETT AND W.-D. WU, *Eigenvector matrices of symmetric tridiagonals*, Numer. Math., 44 (1984), pp. 103-110.
- [12] J.H. WILKINSON, *The Algebraic Eigenvalue Problem*, Clarendon Press, Oxford, England, 1965.

DISTRIBUTION LIST

Director (2)
Defense Tech Information Center
Cameron Station
Alexandria, VA 22314

Research Office (1)
Code 81
Naval Postgraduate School
Monterey, CA 93943

Library (2)
Code 52
Naval Postgraduate School
Monterey, CA 93943

Professor Richard Franke (1)
Department of Mathematics
Naval Postgraduate School
Monterey, CA 93943

Dr. Neil L. Gerr (1)
Mathematical Sciences Division
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000

Dr. Richard Lau (1)
Mathematical Sciences Division
Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000

Harper Whitehouse (Code 743) (1)
NCCOSC RDT&E Division
271 Catalina Blvd.
San Diego, CA 92152-5000

Keith Bromley (Code 7601) (1)
NCCOSC RDT&E Division
271 Catalina Blvd.
San Diego, CA 92152-5000

John Rockway (Code 804) (1)
NCCOSC RDT&E Division
271 Catalina Blvd.
San Diego, CA 92152-5000

Professor Carlos Borges (15)
Department of Mathematics
Naval Postgraduate School
Monterey, CA 93943

DUDLEY KNOX LIBRARY



3 2768 00336645 1