



1975-03

# A Tropical Cyclone Analog Program for the North Indian Ocean

Brand, Samson

---

Monthly Weather Review, Volume 103, pp. 258-261, March 1975.

<http://hdl.handle.net/10945/45749>



Calhoun is a project of the Dudley Knox Library at NPS, furthering the precepts and goals of open government and government transparency. All information contained herein has been approved for release by the NPS Public Affairs Officer.

**Dudley Knox Library / Naval Postgraduate School**  
**411 Dyer Road / 1 University Circle**  
**Monterey, California USA 93943**

<http://www.nps.edu/library>




FIG. 1. October tropical cyclone tracks for the North Indian Ocean (1900-70). Number of tropical cyclones—141.

### 3. Description of the method

The input for the storm to be forecast consists of position and movement as well as the date. The program then searches the history file and selects those storm positions that meet the following criteria:

- a) Within  $\pm 30$  days of current storm
- b) Within  $\pm 3.0^\circ$  latitude/longitude of current storm
- c) Within  $\pm 30^\circ$  of current storm heading (past 12-h direction)
- d) The speed selection (past 12 h) is as follows:
  - 1)  $\pm 3$  kt if current storm speed is  $\leq 7$  kt
  - 2)  $\frac{1}{2}$  to twice current speed if current storm speed is  $> 7$  but  $\leq 13.5$  kt
  - 3)  $\frac{2}{3}$  to  $\frac{3}{2}$  current speed if current storm is  $> 13.5$  kt.

If there are fewer than two acceptable analog candidates, the "window" of acceptance is increased to  $\pm 3.5^\circ$  latitude/longitude. If this criterion fails to gain at least two analogs, no forecast will be produced.

After each analog storm is selected, a 12-h vector is computed based on the speed and heading of the current

storm (persistence) and of the analog storm (climatology). The magnitude of each 12-h vector displacement is defined as

$$\text{DISP} = 12(W_c \cdot C_s + W_a \cdot A_s), \quad (1)$$

where

- $W_c$  = persistence weighting factor
- $C_s$  = mean past 12-h speed of current storm (kt)
- $W_a$  = climatology weighting factor
- $A_s$  = 12-h speed of analog storm (kt).

The direction of the vector is defined as

$$\text{DIR} = W_c \cdot C_h + W_a \cdot A_h, \quad (2)$$

where

- $C_h$  = mean past 12-h heading of current storm
- $A_h$  = heading of analog storm.<sup>1</sup>

This vector is transposed to the location of the current storm position and a new latitude and longitude are determined. The centroid of the new latitude and longitude positions is the forecast position. Up to 72 h

<sup>1</sup> Appropriate corrections are made in the program for 360° circle vector additions and it should be noted that  $W_c + W_a = 1$ .




FIG. 2. Example of 24- and 48-h forecasts, including 50% probability ellipses for INJAH74.

of observational information is stored for each selected analog storm and, since the forecast technique is based on 12-h incremental vectors, the 24-h forecast position is derived from the centroid of the 12-h vector positions,

and so on out to 72 h. The spatial distribution of the positions defines probability ellipses, which are also included as output for INJAH74. Figure 2 shows an example of a 24- and a 48-h forecast, including probability ellipses (50%) for INJAH74.




FIG. 3. Distribution of forecast positions relative to actual verifying position for; (a) 24-h, (b) 48-h, and (c) 72-h forecast test cases. Note: The X's indicate the centers of the distributions.

#### 4. Some additional considerations

In order to determine the general accuracy of the INJAH74 forecast technique, a series of test cases were run for available tropical cyclone data from 1971-73. Thirty forecast situations for 10 Bay of Bengal tropical cyclones were used as the basis for the test.<sup>2</sup>

It became apparent that the technique was sensitive to variations in the weighting scheme (that is, the persistence versus climatology weight for each forecast time). The importance of the climatological factor seemed to be always evident—even, surprisingly, in the early forecast stages.

Not all the test results will be presented or discussed, but Table 1 shows some representative 24-, 48-, and 72-h forecast errors for four distinct weighting schemes. A comparison of total persistence (Column D) with total climatology (Column C)<sup>3</sup> shows the importance of climatology even at 24 hours. Columns A and B show the errors for two tests with an element of persistence for just the first 12 hours. The forecasts thereafter revert toward climatology. Column B with even less

<sup>2</sup> Best track information was used for input and for verification. It should be noted that in an operational mode best track information is not available and this would reduce the accuracy of the forecasts because of possible incorrect analog selections and incorrect motion vectors. The comparative results, in particular with respect to persistence, are nevertheless quite informative and revealing.

<sup>3</sup> Since persistence is inherent in the selection of analogs, this would be a selective climatology. Total climatology as defined here implies no weight for persistence; it is just the centroids of the selected analogs at the specified forecast times.

TABLE 1. Forecast errors for INJAH74 for four separate forecast weighting schemes.

Hours	A						B						C						D					
	12	24	36	48	60	72	12	24	36	48	60	72	12	24	36	48	60	72	12	24	36	48	60	72
Persistence (%)	85	0	0	0	0	0	55	0	0	0	0	0	0	0	0	0	0	0	100	100	100	100	100	100
Climatology (%)	15	100	100	100	100	100	45	100	100	100	100	100	100	100	100	100	100	100	0	0	0	0	0	0
24-Hr Mean Vector Error (nm) [km]	(88.2) [163.4]						(86.7) [160.7]						(84.5) [156.6]						(104.3) [193.3]					
Number of Cases	30						30						30						30					
Standard Deviation (nm) [km]	(54.0) [101.1]						(52.1) [96.6]						(51.5) [95.4]						(60.0) [111.2]					
48-Hr Mean Vector Error (nm) [km]	(213.3) [395.3]						(211.5) [391.7]						(207.8) [385.1]						(288.1) [533.9]					
Number of Cases	28						28						28						28					
Standard Deviation (nm) [km]	(142.2) [263.5]						(139.0) [257.6]						(138.8) [257.2]						(163.8) [303.6]					
72-Hr Mean Vector Error (nm) [km]	(300.2) [556.3]						(299.5) [555.0]						(300.5) [556.9]						(425.6) [788.7]					
Number of Cases	22						22						22						22					
Standard Deviation Error (nm) [km]	(204.0) [378.1]						(204.6) [379.1]						(203.5) [377.1]						(195.3) [361.9]					

weight for persistence in the first 12 hours (55% persistence, 45% climatology) does slightly better, in general, than a heavier weighting toward persistence as in Column A (85% persistence, 15% climatology). Even though total climatology tended to be best in general, for this sample, the weights given in Column B were selected for the final version of INJAH74; for there are times of the year and regions in the North Indian Ocean that will yield very few analogs, and it is at those times that the persistence element becomes important.

To determine if there was any bias in forecast positions relative to the actual verifying positions, a plot was made of the distribution of forecasts for the thirty test cases; this can be seen in Fig. 3. The forecasts for 24 hours seemed to indicate little if any bias. The mean bias for 48 and 72 hours seemed to be more obvious with forecasts tending to be both south and west of the verifying positions. The forecaster might want to subjectively keep in mind the above bias if he is incorporating INJAH74 in his operational forecast.

The forecast method as presented in this paper is a statistical climatological computer forecast technique that can provide objective forecasts for tropical cyclones significantly better than persistence. The importance of climatology is very evident in this tropical cyclone analog technique for this area of the world. Without a significant improvement in the synoptic data base in this region, it is doubtful whether further significant improvements can be achieved from those results presented previously.

INJAH74 is presently running operationally at the U. S. Fleet Weather Central/Joint Typhoon Warning

Center at Guam, the forecast agency responsible for providing tropical cyclone warnings to U. S. Navy and Department of Defense contracted vessels and installations in the western Pacific and Indian Ocean. As the technique is operationally tested, forecast experience will dictate the needs for further modification, development, and tuning of INJAH74.

#### REFERENCES

- Crutcher, H. L., 1971: Atlantic tropical cyclone statistics. NASA CR-61355, National Aeronautics and Space Administration, Huntsville, Ala., 16 pp.
- Gupta, R. N., and R. K. Datta, 1971: Tracking tropical storms in the Bay of Bengal by storm analogue technique using computer. *Vayu Mandal*, 1, 125-127.
- Hodge, W. T., and G. F. McKay, 1970: A computer program to select typhoon analogs and print out their descriptions, including subsequent changes. NWRP Progress Report, National Weather Records Center, Asheville, North Carolina, 41 pp.
- Hope, J. R., and C. J. Neumann, 1970: An operational technique for relating the movement of existing tropical cyclones to past tracks. *Mon. Wea. Rev.*, **98**, 925-933.
- Jarrell, J. D., and W. L. Somervell, Jr., 1970: A computer technique for using typhoon analogs as a forecast aid. Navy Weather Research Facility Tech. Paper No. 6-70.
- , and R. A. Wagoner, 1973: The 1972 typhoon analog program (TYFOON-72). ENVPREDRSCHFAC Tech. Paper No. 1-73. Environmental Prediction Research Facility, Monterey, Calif., 38 pp.
- Neumann, C. J., and J. R. Hope, 1972: Performance analysis of the HURRAN tropical cyclone forecast system. *Mon. Wea. Rev.*, **100**, 245-254.
- Simpson, R. H., 1971: The decision process in hurricane forecasting. Technical Memorandum NWS SR 53, National Weather Service, U. S. Department of Commerce, Miami, Fla., 44 pp.