

RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE

This is the peer reviewed version of an accepted journal article – ‘The effect of tumour necrosis factor- α and insulin on equine digital blood vessel function in vitro.’ The final publication is available at Springer via <https://dx.doi.org/10.1007/s00011-014-0736-2>.

This draft has been made available on the RVC’s Open Access publications repository in accordance with the College’s Open Access policy.

This article may be used for non-commercial purposes in accordance with the publisher’s self-archiving policy, which can be found at <http://www.springer.com/gp/open-access/authors-rights/self-archiving-policy/2124>.

The full details of the published version of the article are as follows:

TITLE: The effect of tumour necrosis factor- α and insulin on equine digital blood vessel function in vitro

AUTHORS: Nicola J. Menzies-Gow, H. Wray, S. R. Bailey, P. A. Harris, J. Elliott

JOURNAL TITLE: Inflammation Research

VOLUME/EDITION: 63/8

PUBLISHER: SpringerLink

PUBLICATION DATE: August 2014

DOI: 10.1007/s00011-014-0736-2

N.J. Menzies-Gow, H. Wray, S.R. Bailey*, P.A. Harris# and J. Elliott

The effect of tumour necrosis factor- α and insulin on equine digital blood vessel function *in vitro*

Royal Veterinary College, London UK. *University of Melbourne, Australia. #Equine Studies Group, WALTHAM Centre for Pet Nutrition, UK.

Corresponding author:

Nicola Menzies-Gow

Email: nmenziesgow@rvc.ac.uk

Telephone: 01707 666297

Fax: 01707 666304

Abstract

Objective and design: Insulin and inflammatory cytokines may be involved in equine laminitis, which might be associated with digital vascular dysfunction. This study determined the effects of TNF- α and insulin on the endothelial-dependent relaxant responses of equine digital blood vessels and on equine digital vein endothelial cell (EDVEC) cGMP production.

Material: Isolated rings of equine digital arteries (EDAs) and veins (EDVs) were obtained and EDVECs were cultured from horses euthanased at an abattoir.

Methods: The effect of incubation with TNF- α (10ng/ml) and/or insulin (1000 μ IU/ml) for 1.5 hours or overnight under hyperoxic and hypoxic conditions on carbachol (endothelium-dependent) induced relaxation was assessed. The time course and concentration dependency of the effect of TNF- α and the effect of insulin (1000 μ IU/ml) on EDVEC cGMP production was determined.

Results: Incubation of EDAs overnight with TNF- α under hypoxic conditions resulted in endothelial-dependent vascular dysfunction. EDVs produced a more variable response. TNF- α increased EDVEC cGMP formation in a time and concentration dependent manner. Insulin had no significant effects.

Conclusions: There is a mismatch between the results obtained from isolated vessel rings and cultured endothelial cells suggesting TNF- α may reduce the biological effect of NO by reducing its bioavailability rather than its formation, leading to endothelial cell dysregulation.

Key Words: Insulin, TNF- α , equine, digital, cGMP, laminitis

Introduction

Laminitis is a common and painful condition of the horse characterised by failure of the attachment of the epidermal cells of the epidermal laminae to the underlying basement membrane of the dermal laminae [1]. Animals at greatest risk of pasture-associated laminitis have a metabolic phenotype including obesity and IR, similar to that seen in human metabolic syndrome (HMS)[2]. Thus the same pathologic mechanisms that underlie the cardiovascular disease associated with HMS, including changes in insulin signalling, inflammatory cytokines and endothelial dysfunction, could contribute to laminitis. Whilst the exact pathogenesis of the disease remains unclear, there is evidence from experimental models of the disease to support roles for inflammation [3-5], vascular and endothelial dysfunction [5-7], insulin resistance [8, 9] and extracellular matrix degradation [10, 11]. However further research is required to fully elucidate the pathways involved and perhaps determine a unifying concept.

In other species, vascular and endothelial dysfunction can be caused by hyperinsulinaemia and/or insulin resistance (IR) [12] and inflammatory mediators [13]. Normally, insulin activates both the phosphatidylinositol (PI3) kinase pathway resulting in stimulation of endothelial nitric oxide synthase (eNOS) and hence vasodilation and the mitogen activated protein (MAP) kinase pathway resulting in enhanced production of endothelin-1 (ET-1) and hence vasoconstriction [14]. IR is characterised by specific impairment of the PI3-kinase-dependent signaling pathway without affecting the MAP-kinase pathway, thus nitric oxide (NO) production is decreased whilst ET-1 production remains unchanged resulting in vasoconstriction [15]. Tumour necrosis factor- α (TNF- α) is a pro-inflammatory cytokine secreted by leukocytes and various other cells, including the non fat cells in adipose tissue of obese individuals, which plays an important role in HMS [13]. This cytokine is implicated in

vascular and endothelial dysfunction, mainly through promotion of redox signalling to inhibit flux through the PI-3 kinase pathway distal to the insulin receptor or via activation of serine kinases [16, 17].

Isolated equine digital blood vessels have been used extensively to study the receptors involved in vasoconstriction and vasodilation [18, 19] and the effects of various drugs [20], mediators [21], cooling [22] and hypoxia [23]. The effect of inflammation on equine digital blood vessels has been mostly investigated previously by exposing the vessels to endotoxin [24-26]. The effect of tumour necrosis factor- α (100pg/ml) on equine digital arterial function has been evaluated once previously [27]. The effects of insulin have also been investigated previously. In one study, equine digital arteries and veins were exposed to extremely high concentrations of insulin (10 μ mol/l or >1,000,000 μ IU/ml) for 30 mins, contracted with phenylephrine and then a second equally high dose of insulin (10 μ mol/l) was added [28]. In separate study, the effect of insulin (1000 μ IU/ml) on the responses of equine small laminar veins to the vasoconstrictors noradrenaline, phenylephrine, ET-1 and 5-hydroxytryptamine (5-HT) was evaluated [29].

The aim of the present study was to determine whether inflammation induced by TNF- α caused impaired vasodilation in equine digital blood vessels, through decreased endothelial nitric oxide (NO) production. This was achieved by investigating the effects of TNF- α on the endothelium-dependent relaxant response of isolated rings of equine digital arteries (EDAs) and veins (EDVs) under hypoxic and hyperoxic conditions and on cGMP (marker of NO production) mediator production by cultured equine digital vein endothelial cells (EDVECs). Furthermore, the effect of insulin was similarly examined to investigate whether high insulin concentrations cause further endothelial dysfunction and exacerbate the effect of TNF- α .

Materials and Methods

Isolated rings of equine digital arteries (EDAs) and veins (EDVs)

The hind limbs of healthy, mixed breed horses, euthanased at an abattoir for purposes other than research, were removed within 10 minutes of death. EDAs were cannulated at the level of the metatarsal-phalangeal joint and flushed with 150 ml cold, oxygenated modified Krebs-Henseleit (Krebs) solution (composition [mM]: CaCl₂ 1.27, MgSO₄ 1.19, NaHCO₃ 25.0, NaCl 118, KH₂PO₄ 1.19, KCl 4.57, glucose 5.55). The legs were then transported to the laboratory. The skin over the lateral aspect of the pastern was removed to reveal the digital artery and vein which were carefully dissected free from surrounding tissues, cleaned of connective tissue and cut into 4-5mm long rings. Vessel rings were stored in pre-oxygenated Krebs at 4°C overnight. Only one EDA and one EDV from each horse were harvested and each was only used in a single study.

Tension Recording in Isolated EDAs and EDVs

Twelve vessel rings could be studied simultaneously in a single study. Vessel rings from the same horse maintained under different incubation conditions were always evaluated during the same experiment. The vessel rings were suspended between two parallel stainless steel wires bathed in 10 ml Krebs solution in an organ bath at 30°C bubbled with either 95% O₂ and 5% CO₂ (designated hyperoxia) or 95% N₂ and 5% CO₂ (designated hypoxia). These gas mixtures provide standard conditions for isolated blood vessel studies [30], as originally described in the work leading to the discovery of the role of nitric oxide in endothelium-dependent relaxation [31]. One of the wires was fixed and the other was connected to an isometric force transducer (HSE force transducer type K30, Linton Instrumentation Ltd, Norfolk, UK). The output of the transducer was fed via an amplifier (HSE type 301; Linton

Instrumentation Ltd) to a data acquisition system (PowerLab; AD Instruments, Oxfordshire, UK). Once mounted, EDA and EDV rings were stretched to 3 g and 2g resting tension, respectively, and allowed to equilibrate for 1 h. Previous studies had shown that this protocol was optimal for the measurement of vasoconstrictor and vasorelaxant responses in this vessel type [18, 32, 33].

The viability of each vessel segment was tested by exchanging the Krebs solution for one in which the sodium chloride had been replaced with potassium chloride to produce a depolarising Krebs solution (DKS; 118 mM KCl). This evaluates the activation of smooth muscle contractile elements to an influx of extracellular Ca^{2+} , without specific receptor activation. Once the tension had increased to reach a plateau, the vessels were washed three times with Krebs solution and allowed to relax to baseline tension. As is standard procedure, vessel rings that failed to increase their tension by >50% of the baseline tension in response to DKS were considered non viable and were discarded [30, 34].

Phenylephrine (PHE, 10^{-6} M; Sigma-Aldrich Company Ltd, Dorset, UK) was then added to the Krebs solution to constrict the vascular smooth muscle. This concentration has previously been shown to produce approximately 50% of maximum tension in equine digital arteries and veins [35, 36]. Once a plateau was reached, carbachol (CCh, 10^{-6} M; Sigma-Aldrich) was added to produce vasodilation. CCh is a cholinergic (muscarinic receptor) agonist that causes endothelial nitric oxide-dependent vasodilation and is thus used to assess the integrity of the endothelium. If the endothelium is damaged during vessel preparation, relaxation to CCh does not occur. The percentage relaxation of PHE-induced tone was calculated, and the vessels were discarded unless a minimum of 50% relaxation to CCh occurred. The vessels were then washed three times with Krebs solution and allowed to relax to baseline tension.

The Effect of TNF- α and Insulin on Relaxant Responses in EDAs and EDVs under Hyperoxic and Hypoxic Conditions

After assessing viability and allowing equilibration back to baseline tension, EDAs (n=6 for each experiment) and EDVs (n=6 for each experiment) were incubated with recombinant equine TNF- α (10ng/ml) and/or insulin (1000 μ IU/ml) for either 1½ hours or overnight (at 37°C) and bubbled with either 95% oxygen and 5% carbon dioxide (designated hyperoxia) or 95% nitrogen and 5% carbon dioxide (designated hypoxia). These gas mixtures provide standard conditions for isolated blood vessel studies [30], as originally described in the work leading to the discovery of the role of nitric oxide in endothelium-dependent relaxation [31]. PHE (10⁻⁶M) was then added to the Krebs to constrict the vessels and, once a plateau of contraction was reached, the carbachol (endothelium-dependent) induced relaxation was again assessed by serially adding increasing concentrations to generate a cumulative concentration response curve.

Equine Digital Vein Endothelial Cell Culture

Equine digital vein endothelial cells (EDVEC) were cultured as previously described [37] from the digits of horses euthanased at an abattoir for purposes other than research. Briefly, as soon as possible post mortem, the digits were flushed to remove the blood with sterile phosphate buffered solution (PBS; 150 ml) by cannulating the medial and lateral digital veins 3 to 4 cm above the coronary band. The medial and lateral digital arteries were then ligated, the digit was infused from the venous side with type II collagenase^a (20ml; 1mg/ml; prewarmed at 37°C) and the limb was incubated in a water bath for 30 min at 37°C. The endothelial cells were then flushed out and collected using sterile PBS before being centrifuged (300 x g for 10min). The supernatant was removed and the cells resuspended in culture medium (Dulbecco's modified Eagles medium^b containing 10% foetal calf serum,

10% newborn calf serum, 100 U/ml penicillin and 100 mg/ml streptomycin) and transferred to a 75cm² flask for incubation at 37°C in 5% CO₂ and 95% air. After 24 hours, erythrocyte contamination was removed with warm sterile PBS and fresh culture medium added. Once 90% confluency was achieved, characterized by the typical cobblestone morphology and positive immunostaining for von Willebrand's factor, the cells were lifted with trypsin and EDTA solution (1 mg/ml and 0.25 mg/ml, respectively), resuspended in culture medium, transferred evenly to 24 well plates and incubated at 37°C for 48 hours to allow the cells to adhere and become confluent.

Effect of TNF- α and Insulin on Mediator Production by EDVEC

Confluent EDVEC monolayers were made quiescent in serum-free medium for 3 hours and pre-incubated with the phosphodiesterase inhibitor, 3-Isobutyl-1-methyl-2,6(1H,3H)-putinedione-Methyl-3-isobutylxanthine^c (IBMX, 1mM) for 30 min at 37°C to inhibit breakdown of cyclic GMP (cGMP). EDVEC were then incubated with DMEM containing 1% bovine serum albumin in the presence or absence of recombinant equine TNF- α (10ng/ml) and/or insulin (1000 μ IU/ml). The culture medium was sampled after 0, 6, 18 and 24 hours incubation. EDVEC were also incubated for 18 hours with varying concentrations of TNF- α alone (0 – 1000 ng/ml).

Protein and Mediator Determination

Cyclic GMP concentrations were measured as an index of endothelial NO production using a commercial enzyme immunoassay system (Biotrak cGMP EIA; Amersham Pharmacia Biotech., Amersham, Buckinghamshire, UK) according to the manufacturer's instructions (Protocol 2) as previously described [38, 39]. Mediator concentrations were expressed per μ g

of protein present in the well, measured following cell lysis using a BCA protein assay kit (Pierce Bio Science Ltd., Tattenhall, Cheshire, UK).

Curve fitting and Statistical Analysis

Statistical analyses were carried out using computer software (Graphpad Prism, version 6.0; GraphPad Inc.). Concentration-response curve data were fitted to a single site sigmoidal response curve model (variable slope) to give values for maximum relaxation (E_{max}) and EC_{50} values. Normality of the distribution of the data was assessed using the Kolmogorov–Smirnov test. Values represent mean \pm sem from $n=6-12$ (see Figure legends), referring to the number of individual animals from which cells or blood vessels were derived. EC_{50} values were presented as the geometric mean with 95% confidence intervals. The effect of TNF- α and insulin on the maximum relaxant responses of EDAs and EDVs under hyperoxic and hypoxic conditions was evaluated using a paired Student's t-test or one way analysis of variance with Dunnett's *post hoc* test. The effect of TNF- α and insulin on EDVEC mediator production was analysed using a linear mixed effect model. Significance was accepted at $p<0.05$.

Results

There was no significant effect of incubation with 10 ng/ml TNF- α and/or 1000 μ IU/ml insulin for 1½ hours, under hyperoxic or hypoxic conditions, on carbachol (endothelium-dependent) induced relaxation of isolated rings of equine digital arteries (EDAs; Figure 1) or equine digital veins (EDVs; Figure 2). The curve fitting parameters calculated from these experiments are presented in Table 1. However, despite the short-term treatment having no appreciable effect, incubation with the same concentration of TNF- α overnight at 37°C under hypoxic conditions significantly ($p<0.05$) decreased the carbachol (endothelium-dependent)

induced relaxation of EDAs (Figure 3). There was no effect of insulin under hyperoxic or hypoxic conditions, even at high concentrations (Figure 4). Finally, incubation with high concentrations of both TNF- α and insulin overnight under hypoxic conditions significantly reduced carbachol (endothelium-dependent) induced relaxation of EDVs; however, again this effect was not observed under hyperoxic conditions (Figure 5).

TNF- α (10ng/ml) stimulated significant increases of 4.0 ± 2.5 and 2.9 ± 3.1 fold ($p < 0.05$) in EDVEC production of cGMP above basal levels after 18 and 24 hours, respectively (Figure 6). This stimulation was TNF- α concentration dependent (Figure 6). Insulin (1000 μ IU/ml) had no significant effect alone or when added in combination with TNF- α (Figure 6).

Discussion

Many studies have hypothesised that reduced lamellar blood flow is a key event in the pathogenesis of laminitis [25, 40-42], with ischaemia and subsequent reperfusion resulting in lamellar damage. However, the results of studies of laminar and digital blood flow using *in vivo* techniques have been conflicting, with increases in blood flow in some studies [43-45] and decreases in others [40, 46-48]. This apparent disagreement may be a consequence of a lack of sensitivity and specificity of the methods used to detect changes in laminar blood flow. *In vitro* studies make it possible to investigate the direct effects of individual factors on the function of specific equine digital blood vessels.

Animals at greatest risk of pasture-associated laminitis have a metabolic phenotype that includes obesity and IR [2]. The prolonged experimental infusion of insulin to induce hyperinsulinaemia resulted in the clinical and histological development of laminitis [49, 50]. Hyperinsulinaemia and IR may predispose to laminitis via vascular function disturbances. Obesity may predispose to laminitis through IR or inflammation as adipocytes are endocrinologically active producing a variety of mediators, some of which antagonise insulin resulting in IR and some of which are pro-inflammatory [51]. TNF- α is one such pro-inflammatory cytokine secreted mainly by the non fat cell within adipose tissue [52], which is associated with endothelial dysfunction [13]. Within a herd of inbred obese ponies, laminitis prone animals had significantly higher plasma TNF- α concentrations compared to normal ponies thus associating laminitis predisposition with increased circulating inflammatory cytokines [53]. Plasma TNF- α concentration is also an independent risk factor for increased IR in horses [54].

TNF- α appears to have variable effects on vascular function in other species. Intracranial injection of TNF- α *in vivo* constricted pial arterioles and reduced cerebral blood flow [55-57]; TNF- α induced pro-contractile effects in coronary arteries [58, 59]; and TNF- α impaired acetylcholine-induced vasodilation of mouse thoracic aorta [60]. In contrast, TNF- α dilated cerebral, mesenteric and cremaster muscle arterioles and relaxed endothelium-denuded aortas [61-66]. In bronchial arteries, TNF- α initially induced dilation, followed by constriction 2 hrs later [67]. Finally, TNF- α (1mM) had no effect on bradykinin-mediated endothelium-dependent relaxant of human omental arteries [68] and TNF- α (10ng/ml) had no effect on the diameter of isolated resistance arteries from the rat cremaster muscle [69]. The effect of TNF- α on equine digital vascular function has only been investigated once previously [27]. Incubation with TNF- α (0.1 or 1ng/ml) for 10 mins significantly decreased the endothelial-dependent relaxation of equine palmar digital arteries to the muscarinic receptor agonist acetylcholine and significantly increased the maximal contraction to noradrenaline [27]. Reported plasma TNF- α concentrations in normal horses range from 1ng/ml [70] to 14ng/ml [71]. In the present study incubation of equine digital arteries for longer (either 90 mins or overnight) with a higher (10ng/ml), but still clinically relevant concentration of TNF- α under hyperoxic conditions had no effect on endothelial-dependent relaxation. However, overnight incubation under hypoxic conditions resulted in endothelial-dependent vascular dysfunction. Equine digital veins showed greater variability in the degree of vasorelaxation but incubation with TNF- α and insulin overnight under hypoxic conditions significantly inhibited vasodilation. The effect of hypoxia on equine digital vascular function has only been investigated once previously; short term (3 hrs) hypoxia enhanced the contractile responses of equine digital arteries [23]. To the authors' knowledge, the effect of TNF- α on vascular function under hypoxic conditions has not been previously evaluated in any species.

Previous studies have produced variable results with respect to the effects of insulin on equine blood vessel function. Infusion of insulin into healthy Standardbreds resulted in mean serum insulin concentrations of approximately 1000 μ IU/ml and the hoof wall surface temperature (HMST) was higher and less variable once hyperinsulinaemia was established, when compared to control horses, suggestive of insulin-mediated increased digital perfusion [50]. In contrast, *in vitro*, following induction of IR using supra physiologic concentrations of insulin (>1,600,000 μ IU/ml), instead of the normal relaxation responses, a further equally high dose of insulin resulted in contraction of isolated rings of equine palmar digital arteries [28] and short-term hyperinsulinaemia (142 μ IU/ml) led to increased vascular resistance within the isolated equine digit in a model of extracorporeal perfusion [72]. Serum insulin concentrations of up to approximately 600 μ IU/ml have been recorded in previously laminitic ponies[73]; thus in the present study an insulin concentration of 600 μ IU/ml was used. This high but still clinically relevant concentration of insulin alone had no effect on endothelium-dependent relaxation.

The combined effect of insulin and TNF- α has been previously evaluated only in humans and rats. Infusion of TNF- α (17ng/min) inhibited endothelium-dependent vasodilation of the brachial artery in humans and the inhibitory effect on vasodilator function was greater with concurrent insulin infusion (0.05mg/kg/min) [74]. Alone neither insulin nor TNF- α had an effect, but TNF- α inhibited the vasodilator but not vasoconstrictor effects of insulin in isolated cremaster muscle resistance arteries, resulting in insulin-mediated vasoconstriction in the presence of TNF- α [69]. In the present study, a high but clinically relevant concentration of insulin did not alter the effects of TNF- α on equine digital vascular function.

The endothelium normally produces vasodilator mediators including nitric oxide (NO), prostacyclin (PGI₂) and endothelium-derived hyperpolarising factor (EDHF) [75] and the potent vasoconstrictor endothelin-1 (ET-1) [76]. Endothelial dysfunction results in an imbalance between the production of vasoconstrictory and vasodilatory mediators predominantly due to reduced endothelium-dependent NO bioavailability [77] and hence favouring vasoconstriction. Endothelial dysfunction may occur as a result of insulin resistance and inflammation [14].

In other species there is a large amount of evidence to suggest that TNF- α is associated with endothelial dysfunction [78-81]. TNF- α (10ng/ml) impaired NO production by bovine aortic endothelial cells in a time and dose-dependent manner due to a reduction in eNOS expression [82-86]. TNF- α (10ng/ml) depressed human aortic endothelial cell NO production via the coordinate down regulation of both eNOS expression and argininosuccinate synthase [87], the enzyme which produces arginine, a key substrate for NO synthase. In the present study cGMP was measured as it is a marker of NO biological activity [88]. TNF- α increased cGMP formation by cultured equine digital endothelial cells in a time and concentration dependent manner.

Insulin increased human umbilical endothelial cell NO production in a dose dependent manner with an ED₅₀ of 500nM [89] and NOS activation in a dose and time dependent manner, an insulin concentration of 600nM was required to activate eNOS and the effect was maximal after 30 mins [90]. Insulin (500nmol/L) also increased eNOS activation in bovine aortic endothelial cells [91]. In the present study, high clinically relevant concentrations of insulin were used. These concentrations of insulin were 5-fold lower than these previous studies and had no effect on endothelial cGMP production over a much longer time period.

A limitation of the current study was that it used large digital vessels. Whilst similar vessels have been used in several other studies attempting to further elucidate the pathogenesis of equine laminitis [18, 41], it must be acknowledged that there are differences in the vascular responses of large conductance or capacitance vessels and small resistance vessels [41]. Thus caution is advised in extrapolating results of the current study to other vessel types within the equine digit.

In conclusion, incubation of equine digital arteries overnight with high clinically relevant concentrations of the inflammatory cytokine TNF- α under hypoxic conditions resulted in endothelial-dependent vascular dysfunction. High clinically relevant concentrations of insulin did not have a similar effect. Equine digital veins showed greater variability in the degree of vasorelaxation but incubation with TNF- α and insulin overnight under hypoxic conditions also significantly inhibited vasodilation. By contrast, clinically relevant concentrations of TNF- α increased cGMP formation by cultured equine digital endothelial cells. Thus, there is a mismatch between results obtained in isolated endothelial cells and intact blood vessels suggesting that TNF- α may reduce the biological effect of NO by reducing its bioavailability rather than its formation, leading to endothelial cell dysregulation. However, the mechanisms involved require further investigation.

Acknowledgements

This study was funded by the Laminitis Trust.

References

1. Pollitt CC. Basement membrane pathology: a feature of acute equine laminitis. *Equine Vet J* 1996; 28:38-46.
2. Treiber KH, Kronfeld DS, Hess TM, Byrd BM, R.K. S, Staniar WB. Evaluation of genetic and metabolic predispositions and nutritional risk factors for pasture-associated laminitis in ponies. *J Am Vet Med Assoc* 2006; 228:1538-1545.
3. Belknap JK, Giguere S, Pettigrew A, Cochran AM, Van Eps AW, Pollitt CC. Lamellar pro-inflammatory cytokine expression patterns in laminitis at the developmental stage and at the onset of lameness: innate vs. adaptive immune response. *Equine veterinary journal* 2007; 39:42-7.
4. Leise BS, Faleiros RR, Watts M, Johnson PJ, Black SJ, Belknap JK. Laminar inflammatory gene expression in the carbohydrate overload model of equine laminitis. *Equine veterinary journal* 2011; 43:54-61.
5. Black SJ, Lunn DP, Yin C, Hwang M, Lenz SD, Belknap JK. Leukocyte emigration in the early stages of laminitis. *Vet Immunol Immunopathol* 2006; 109:161-6.
6. Weiss DJ, Evanson OA, McClenahan D, Fagliari JJ, Dunnwiddie CT, Wells RE. Effect of a competitive inhibitor of platelet aggregation on experimentally induced laminitis in ponies. *Am J Vet Res* 1998; 59:814-7.
7. Hurley DJ, Parks RJ, Reber AJ, Donovan DC, Okinaga T, Vandenplas ML, et al. Dynamic changes in circulating leukocytes during the induction of equine laminitis with black walnut extract. *Vet Immunol Immunopathol* 2006; 110:195-206.
8. Karikoski NP, Horn I, McGowan TW, McGowan CM. The prevalence of endocrinopathic laminitis among horses presented for laminitis at a first-opinion/referral equine hospital. *Domestic Animal Endocrinology* 2011; 41:111-117.
9. Donaldson MT, Jorgensen AJ, Beech J. Evaluation of suspected pituitary pars intermedia dysfunction in horses with laminitis. *J Am Vet Med Assoc* 2004; 224:1123-7.
10. Coyne MJ, Cousin H, Loftus JP, Johnson PJ, Belknap JK, Gradil CM, et al. Cloning and expression of ADAM-related metalloproteases in equine laminitis. *Vet Immunol Immunopathol* 2009; 129:231-41.
11. Wang L, Pawlak E, Johnson PJ, Belknap JK, Alfandari D, Black SJ. Effects of cleavage by a disintegrin and metalloproteinase with thrombospondin motifs-4 on gene expression and protein content of versican and aggrecan in the digital laminae of horses with starch gruel-induced laminitis. *Am J Vet Res* 2012; 73:1047-56.
12. Muniyappa R, Montagnani M, Koh KK, Quon MJ. Cardiovascular actions of insulin. *Endocr Rev* 2007; 28:463-91.
13. Picchi A, Gao X, Belmadani S, Potter BJ, Focardi M, Chilian WM, et al. Tumor necrosis factor-alpha induces endothelial dysfunction in the prediabetic metabolic syndrome. *Circ Res* 2006; 99:69-77.
14. Kim JA, Montagnani M, Koh KK, Quon MJ. Reciprocal relationships between insulin resistance and endothelial dysfunction: molecular and pathophysiological mechanisms. *Circulation* 2006; 113:1888-904.

15. Potenza MA, Marasciulo FL, Chieppa DM, Brigiani GS, Formoso G, Quon MJ, et al. Insulin resistance in spontaneously hypertensive rats is associated with endothelial dysfunction characterized by imbalance between NO and ET-1 production. *Am J Physiol Heart Circ Physiol* 2005; 289:H813-22.
16. Houstis N, Rosen ED, Lander ES. Reactive oxygen species have a causal role in multiple forms of insulin resistance. *Nature* 2006; 440:944-8.
17. Nishikawa T, Kukidome D, Sonoda K, Fujisawa K, Matsuhisa T, Motoshima H, et al. Impact of mitochondrial ROS production in the pathogenesis of insulin resistance. *Diabetes Res Clin Pract* 2007; 77 Suppl 1:S161-4.
18. Bailey SR, Elliott J. Evidence for different 5-HT_{1B/1D} receptors mediating vasoconstriction of equine digital arteries and veins. *Eur J Pharmacol* 1998; 355:175-87.
19. Elliott J, Brady FE. Characterization of vasodilatory adenosine receptors in equine digital veins. *J Vet Pharmacol Ther* 1998; 21:74-81.
20. Stokes AM, Venugopal CS, Hosgood G, Eades SC, Moore RM. Comparison of 2 endothelin-receptor antagonists on in vitro responses of equine palmar digital arterial and venous rings to endothelin-1. *Can J Vet Res* 2006; 70:197-205.
21. Bailey SR, Elliott J. Plasma 5-hydroxytryptamine constricts equine digital blood vessels in vitro: implications for pathogenesis of acute laminitis. *Equine Vet J* 1998; 30:124-30.
22. Zerpa H, Berhane Y, Elliott J, Bailey SR. Cooling augments vasoconstriction mediated by 5-HT₁ and alpha₂-adrenoceptors in the isolated equine digital vein: involvement of Rho kinase. *Eur J Pharmacol* 2007; 569:212-21.
23. Borer KE, Bailey SR, Harris PA, Elliott J. Contractile responses of isolated equine digital arteries under hypoxic or hyperoxic conditions in vitro: role of reactive oxygen species and Rho kinase. *Journal of Veterinary Pharmacology and Therapeutics* 2012:no-no.
24. Bailey SR, Elliott J. The role of prostanoids and nitric oxide in endotoxin-induced hyporesponsiveness of equine digital blood vessels. *Equine Vet J* 1999; 31:212-8.
25. Baxter GM, Laskey RE, Tackett RL, Moore JN, Allen D. In vitro reactivity of digital arteries and veins to vasoconstrictive mediators in healthy horses and in horses with early laminitis. *Am J Vet Res* 1989; 50:508-17.
26. Mallem MY, Gogny M, Gautier F, Bucas V, Desfontis JC. Evaluation of beta₃-adrenoceptor-mediated relaxation in intact and endotoxin-treated equine digital veins. *Am J Vet Res* 2003; 64:708-14.
27. Baxter G. Effects of tumor necrosis factor on in vitro digital arterial responses in horses. *Am J Vet Res* 1994; 55:551-5.
28. Venugopal CS, Eades S, Holmes EP, Beadle RE. Insulin resistance in equine digital vessel rings: an in vitro model to study vascular dysfunction in equine laminitis. *Equine Vet J* 2011; 43:744-9.
29. Keen JA, McGorum BC, Hillier C, Nally JE. Short-term incubation of equine laminar veins with cortisol and insulin alters contractility in vitro: possible implications for the pathogenesis of equine laminitis. *J Vet Pharmacol Ther* 2013; 36:382-8.

30. Borer KE, Bailey SR, Harris PA, Elliott J. Contractile responses of isolated equine digital arteries under hypoxic or hyperoxic conditions in vitro: role of reactive oxygen species and Rho kinase. *J Vet Pharmacol Ther* 2013; 36:267-74.
31. Furchgott RF, Zawadzki JV. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* 1980; 288:373-6.
32. Elliott J, Berhane Y, Bailey SR. Effects of monoamines formed in the cecum of horses on equine digital blood vessels and platelets. *Am J Vet Res* 2003; 64:1124-31.
33. Katz LM, Marr CM, Elliott J. Characterization and comparison of the responses of equine digital arteries and veins to endothelin-1. *Am J Vet Res* 2003; 64:1438-43.
34. Katz LM, Marr CM, Elliott J. Characterisation of the response of equine digital arteries and veins to substance P. *J Vet Pharmacol Ther* 2003; 26:361-8.
35. Elliott J. Alpha-adrenoceptors in equine digital veins: evidence for the presence of both alpha1 and alpha2-receptors mediating vasoconstriction. *J Vet Pharmacol Ther* 1997; 20:308-17.
36. Peroni JF, Moore JN, Noschka E, Lewis TH, Lewis SJ, Robertson TP. Effect of voltage-gated and capacitative calcium entry blockade on agonist-induced constriction of equine laminar blood vessels. *Am J Vet Res* 2007; 68:722-9.
37. Bailey SR, Wheeler-Jones C, Elliott J. Uptake of 5-hydroxytryptamine by equine digital vein endothelial cells: inhibition by amines found in the equine caecum. *Equine Vet J* 2003; 35:164-9.
38. Akarasereenont P, Mitchell JA, Bakhle YS, Thiemermann C, Vane JR. Comparison of the induction of cyclooxygenase and nitric oxide synthase by endotoxin in endothelial cells and macrophages. *Eur J Pharmacol* 1995; 273:121-8.
39. Evans M, Berhane Y, Botham KM, Elliott J, Wheeler-Jones CP. Chylomicron-remnant-like particles modify production of vasoactive mediators by endothelial cells. *Biochem Soc Trans* 2004; 32:110-2.
40. Hood DM, Wagner IP, Brumbaugh GW. Evaluation of hoof wall surface temperature as an index of digital vascular perfusion during the prodromal and acute phases of carbohydrate-induced laminitis in horses. *Am J Vet Res* 2001; 62:1167-72.
41. Peroni JF, Moore JN, Noschka E, Grafton ME, Aceves-Avila M, Lewis SJ, et al. Predisposition for venoconstriction in the equine laminar dermis: implications in equine laminitis. *J Appl Physiol* 2006; 100:759-63.
42. Bailey SR, Marr CM, Elliott J. Current research and theories on the pathogenesis of acute laminitis in the horse. *Vet J* 2004; 167:129-42.
43. Robinson NE, Scott JB, Dabney JM, Jones GA. Digital vascular responses and permeability in equine alimentary laminitis. *Am J Vet Res* 1976; 37:1171-6.
44. Pollitt CC, Davies CT. Equine laminitis: its development coincides with increased sublamellar blood flow. *Equine Vet J* 1998; Supplement 26:125-32.
45. Trout DR, Hornof WJ, Linford RL, O'Brien TR. Scintigraphic evaluation of digital circulation during the developmental and acute phases of equine laminitis. *Equine Vet J* 1990; 22:416-21.

46. Coffman JR, Johnson JH, Guffy MM, Finocchio EJ. Hoof circulation in equine laminitis. *J Am Vet Med Assoc* 1970; 156:76-83.
47. Adair HS, 3rd, Goble DO, Schmidhammer JL, Shires GM. Laminar microvascular flow, measured by means of laser Doppler flowmetry, during prodromal stage of black walnut-induced laminitis in horses. *Am J Vet Res* 2000; 61:862-8.
48. Hood DM, Amoss MS, Hightower D. Equine laminitis I: Radioisotope analysis of the hemodynamics of the foot during the acute disease. *J Equine Med Surg* 1978; 2:439-444.
49. Asplin KE, Sillence MN, Pollitt CC, McGowan CM. Induction of laminitis by prolonged hyperinsulinaemia in clinically normal ponies. *Vet J* 2007; 174:530-5.
50. de Laat MA, McGowan CM, Sillence MN, Pollitt CC. Equine laminitis: induced by 48 h hyperinsulinaemia in Standardbred horses. *Equine Vet J* 2010; 42:129-35.
51. Bujalska IJ, Kumar S, Stewart PM. Does central obesity reflect "Cushing's disease of the omentum"? *Lancet* 1997; 1210-1213.
52. Fain JN. Release of interleukins and other inflammatory cytokines by human adipose tissue is enhanced in obesity and primarily due to the nonfat cells. *Vitam Horm* 2006; 74:443-77.
53. Treiber K, Carter R, Gay L, Williams C, Geor R. Inflammatory and redox status of ponies with a history of pasture-associated laminitis. *Vet Immunol Immunopathol* 2009; 129:216-220.
54. Vick MM, Adams AA, Murphy BA, Sessions DR, Horohov DW, Cook RF, et al. Relationships among inflammatory cytokines, obesity, and insulin sensitivity in the horse. *J Anim Sci* 2007; 85:1144-55.
55. Megyeri P, Abraham CS, Temesvari P, Kovacs J, Vas T, Speer CP. Recombinant human tumor necrosis factor alpha constricts pial arterioles and increases blood-brain barrier permeability in newborn piglets. *Neurosci Lett* 1992; 148:137-40.
56. Sibson NR, Blamire AM, Perry VH, Gauldie J, Styles P, Anthony DC. TNF-alpha reduces cerebral blood volume and disrupts tissue homeostasis via an endothelin- and TNFR2-dependent pathway. *Brain* 2002; 125:2446-59.
57. Tureen J. Effect of recombinant human tumor necrosis factor-alpha on cerebral oxygen uptake, cerebrospinal fluid lactate, and cerebral blood flow in the rabbit: role of nitric oxide. *J Clin Invest* 1995; 95:1086-91.
58. Klemm P, Warner TD, Hohlfeld T, Corder R, Vane JR. Endothelin 1 mediates ex vivo coronary vasoconstriction caused by exogenous and endogenous cytokines. *Proc Natl Acad Sci U S A* 1995; 92:2691-5.
59. Zhang DX, Yi FX, Zou AP, Li PL. Role of ceramide in TNF-alpha-induced impairment of endothelium-dependent vasorelaxation in coronary arteries. *Am J Physiol Heart Circ Physiol* 2002; 283:H1785-94.
60. Huang Y, Yan L, Rong S, Haller H, Kirch T. TNF-alpha induces endothelial dysfunction via PKC-zeta-dependent NADPH oxidase activation. *J Huazhong Univ Sci Technolog Med Sci* 2012; 32:642-7.

61. Baudry N, Vicaud E. Role of nitric oxide in effects of tumor necrosis factor-alpha on microcirculation in rat. *J Appl Physiol* 1993; 75:2392-9.
62. Brian JE, Jr., Faraci FM. Tumor necrosis factor-alpha-induced dilatation of cerebral arterioles. *Stroke* 1998; 29:509-15.
63. Hollenberg SM, Cunnion RE, Parrillo JE. The effect of tumor necrosis factor on vascular smooth muscle. In vitro studies using rat aortic rings. *Chest* 1991; 100:1133-7.
64. Johns DG, Webb RC. TNF-alpha-induced endothelium-independent vasodilation: a role for phospholipase A2-dependent ceramide signaling. *Am J Physiol* 1998; 275:H1592-8.
65. Naziri W, Joshua IG. Role of tumor necrosis factor-alpha in small intestinal microcirculation. *Am Surg* 1998; 64:203-9; discussion 209-10.
66. Shibata M, Parfenova H, Zuckerman SL, Leffler CW. Tumor necrosis factor-alpha induces pial arteriolar dilation in newborn pigs. *Brain Res Bull* 1996; 39:241-7.
67. Wagner EM. TNF-alpha induced bronchial vasoconstriction. *Am J Physiol Heart Circ Physiol* 2000; 279:H946-51.
68. Gillham JC, Myers JE, Baker PN, Taggart MJ. TNF-alpha alters nitric oxide- and endothelium-derived hyperpolarizing factor-mediated vasodilatation in human omental arteries. *Hypertens Pregnancy* 2008; 27:29-38.
69. Eringa EC, Stehouwer CD, Walburg K, Clark AD, van Nieuw Amerongen GP, Westerhof N, et al. Physiological concentrations of insulin induce endothelin-dependent vasoconstriction of skeletal muscle resistance arteries in the presence of tumor necrosis factor-alpha dependence on c-Jun N-terminal kinase. *Arterioscler Thromb Vasc Biol* 2006; 26:274-80.
70. Menzies-Gow NJ, Wray H, Bailey SR, Harris PA, Elliott J. The effect of exercise on plasma concentrations of inflammatory markers in normal and previously laminitic ponies. *Equine Vet J* 2013.
71. Suagee JK, Corl BA, Crisman MV, Pleasant RS, Thatcher CD, Geor RJ. Relationships between Body Condition Score and Plasma Inflammatory Cytokines, Insulin, and Lipids in a Mixed Population of Light-Breed Horses. *Journal of veterinary internal medicine* 2013; 27:157-163.
72. Gauff F, Patan-Zugaj B, Licka TF. Hyperinsulinaemia increases vascular resistance and endothelin-1 expression in the equine digit. *Equine Vet J* 2013; 45:613-8.
73. Bailey SR, Habershon-Butcher J, Ransom K, Elliott J, Menzies-Gow NJ. Hypertension and insulin resistance in a mixed-breed population of ponies predisposed to laminitis: Further characterizing pre-laminitic metabolic syndrome. *Am J Vet Res* 2008; 69:122-129.
74. Rask-Madsen C, Dominguez H, Ihlemann N, Hermann T, Kober L, Torp-Pedersen C. Tumor necrosis factor-alpha inhibits insulin's stimulating effect on glucose uptake and endothelium-dependent vasodilation in humans. *Circulation* 2003; 108:1815-21.
75. Elliott J, Bryant CE, Soydan J. The role of nitric oxide in the responses of equine digital veins to vasodilator and vasoconstrictor agents. *Equine veterinary journal* 1994; 26:378-84.

76. Galley HF, Webster NR. Physiology of the endothelium. *British journal of anaesthesia* 2004; 93:105-13.
77. Potenza MA, Addabbo F, Montagnani M. Vascular actions of insulin with implications for endothelial dysfunction. *Am J Physiol Endocrinol Metab* 2009; 297:E568-77.
78. Pfeiffer A, Janott J, Mohlig M, Ristow M, Rochlitz H, Busch K, et al. Circulating tumor necrosis factor alpha is elevated in male but not in female patients with type II diabetes mellitus. *Horm Metab Res* 1997; 29:111-4.
79. Winkler G, Lakatos P, Salamon F, Nagy Z, Speer G, Kovacs M, et al. Elevated serum TNF-alpha level as a link between endothelial dysfunction and insulin resistance in normotensive obese patients. *Diabet Med* 1999; 16:207-11.
80. Zhang H, Park Y, Zhang C. Coronary and aortic endothelial function affected by feedback between adiponectin and tumor necrosis factor alpha in type 2 diabetic mice. *Arterioscler Thromb Vasc Biol* 2010; 30:2156-63.
81. Zhang H, Park Y, Wu J, Chen X, Lee S, Yang J, et al. Role of TNF-alpha in vascular dysfunction. *Clin Sci (Lond)* 2009; 116:219-30.
82. Alonso J, Sanchez de Miguel L, Monton M, Casado S, Lopez-Farre A. Endothelial cytosolic proteins bind to the 3' untranslated region of endothelial nitric oxide synthase mRNA: regulation by tumor necrosis factor alpha. *Mol Cell Biol* 1997; 17:5719-26.
83. Anderson HD, Rahmutula D, Gardner DG. Tumor necrosis factor-alpha inhibits endothelial nitric-oxide synthase gene promoter activity in bovine aortic endothelial cells. *J Biol Chem* 2004; 279:963-9.
84. Yoshizumi M, Perrella MA, Burnett JC, Jr., Lee ME. Tumor necrosis factor downregulates an endothelial nitric oxide synthase mRNA by shortening its half-life. *Circ Res* 1993; 73:205-9.
85. Kim F, Gallis B, Corson MA. TNF-alpha inhibits flow and insulin signaling leading to NO production in aortic endothelial cells. *Am J Physiol Cell Physiol* 2001; 280:C1057-65.
86. Mohamed F, Monge JC, Gordon A, Cernacek P, Blais D, Stewart DJ. Lack of role for nitric oxide (NO) in the selective destabilization of endothelial NO synthase mRNA by tumor necrosis factor-alpha. *Arterioscler Thromb Vasc Biol* 1995; 15:52-7.
87. Goodwin BL, Pendleton LC, Levy MM, Solomonson LP, Eichler DC. Tumor necrosis factor-alpha reduces argininosuccinate synthase expression and nitric oxide production in aortic endothelial cells. *Am J Physiol Heart Circ Physiol* 2007; 293:H1115-21.
88. Ikeda M, Matsui K, Ishihara Y, Morita I, Murota S, Yuasa T, et al. Cerebellar nitric oxide synthase, cGMP and motor function in two lines of cerebellar mutant mice, Staggerer and Wriggle Mouse Sagami. *Neurosci Lett* 1994; 168:65-8.
89. Zeng G, Quon MJ. Insulin-stimulated production of nitric oxide is inhibited by wortmannin. Direct measurement in vascular endothelial cells. *J Clin Invest* 1996; 98:894-8.

90. Aragonés G, Saavedra P, Heras M, Cabre A, Girona J, Masana L. Fatty acid-binding protein 4 impairs the insulin-dependent nitric oxide pathway in vascular endothelial cells. *Cardiovasc Diabetol* 2012; 11:72.
91. Tanigaki K, Mineo C, Yuhanna IS, Chambliss KL, Quon MJ, Bonvini E, et al. C-reactive protein inhibits insulin activation of endothelial nitric oxide synthase via the immunoreceptor tyrosine-based inhibition motif of FcγRIIB and SHIP-1. *Circ Res* 2009; 104:1275-82.

Table 1. Concentration-response curve parameters for the effect of TNF- α (10ng/ml) and insulin (1000 μ IU/ml) on carbachol-induced relaxation of equine digital arteries and veins, incubated under hyperoxic or hypoxic conditions for 1½ hours. For graphs see Figs 1 and 2. CL—confidence limits. No significant differences were observed between treatment and control groups (oneway ANOVA for arteries and paired *t*-tests for veins).

Treatment	EC50 (M)		Emax (% relaxation of PHE-induced tone)	
	Geometric mean (95% CL)		mean \pm sem	
	arteries	veins	arteries	veins
Hypoxia:				
Control	6.17 (2.48-9.86) x 10 ⁻⁸	33.2 (2.07-4.57) x 10 ⁻⁸	79.82 \pm 4.30	44.58 \pm 7.58
TNF- α (10ng/ml)	4.96 (1.61-8.31) x 10 ⁻⁸		74.40 \pm 6.28	
insulin (1000 μ IU/ml)	4.47 (3.02-5.94) x 10 ⁻⁸		70.95 \pm 6.46	
TNF- α + insulin	5.96 (1.43-10.50) x 10 ⁻⁸	27.6 (2.14-3.38) x 10 ⁻⁸	71.93 \pm 5.42	50.94 \pm 9.22
Hyperoxia:				
Control	10.04 (0.51-1.57) x 10 ⁻⁸	90.4 (0.15-19.50) x 10 ⁻⁸	76.61 \pm 4.99	51.50 \pm 7.53
TNF- α (10ng/ml)	13.8 (0.73-2.03) x 10 ⁻⁸		74.23 \pm 3.80	
insulin (1000 μ IU/ml)	9.91 (5.63-14.20) x 10 ⁻⁸		69.38 \pm 4.09	
TNF- α + insulin	10.4 (0.61-1.42) x 10 ⁻⁸	44.5(3.13-5.77) x 10 ⁻⁸	71.09 \pm 5.02	55.04 \pm 6.43

Table 2. Concentration-response curve parameters for the effect of TNF- α (10ng/ml) and insulin (1000 μ IU/ml) on carbachol-induced relaxation of equine digital arteries and veins, incubated overnight at 37°C under hypoxic or hyperoxic conditions. For graphs, see Figs 3,4 and 5. * and ** indicate significant differences between treatment and control groups (p<0.05 and p<0.01, respectively; paired t-test).

Treatment	EC ₅₀ (M)		Emax (% relaxation of PHE-induced tone)	
	Geometric mean (95% CL)		mean \pm sem	
	arteries	veins	arteries	veins
Hypoxia:				
Control	9.54 (3.42-15.65) x 10 ⁻⁸		68.80 \pm 4.42	
TNF- α (10ng/ml)	7.38 (3.92-10.83) x 10 ⁻⁸		47.81 \pm 6.74**	
Control	14.1 (-2.12-3.03) x 10 ⁻⁸		55.79 \pm 8.93	
Insulin	11.6 (0.35-1.98) x 10 ⁻⁸		56.99 \pm 9.56	
Control	5.75 (2.29-9.21) x 10 ⁻⁸	74.1 (1.29-13.53) x 10 ⁻⁸	68.43 \pm 6.87	33.93 \pm 7.48
TNF- α + insulin	14.5 (0.68-2.22) x 10 ⁻⁸	33.2 (2.73-3.91) x 10 ⁻⁸	44.77 \pm 10.05 *	21.32 \pm 2.11 *
Hyperoxia:				
Control	13.3 (0.82-1.84) x 10 ⁻⁸	46.9 (-0.26-9.64) x 10 ⁻⁸	61.20 \pm 2.80	44.66 \pm 8.71
TNF- α (10ng/ml)	15.1 (0.85-2.18) x 10 ⁻⁸	120.0 (-0.36-2.76) x 10 ⁻⁸	55.23 \pm 5.10	39.89 \pm 8.94
+ insulin (1000 μ IU/ml)				

Figure Legends

Figure 1: The effect of TNF- α (10ng/ml) and insulin (1000 μ IU/ml) on carbachol-induced relaxation of equine digital arteries incubated under hyperoxic (A) or hypoxic (B) conditions for 1½ hours. Each point represents mean \pm sem from 6 separate experiments. No significant differences were observed between treatment and control groups (see Table 1 for curve fitting parameters).

Figure 2: The effect of TNF- α (10ng/ml) and insulin (1000 μ IU/ml) on carbachol-induced relaxation of equine digital veins incubated under hyperoxic (A) or hypoxic (B) conditions for 1½ hours. Each point represents mean \pm sem from 6 (A) or 10 (B) separate experiments. No significant differences were observed between treatment and control groups (see Table 1 for curve fitting parameters).

Figure 3: The effect of TNF- α (10ng/ml) on carbachol-induced relaxation of equine digital arteries incubated overnight under hypoxic conditions. Each point represents mean \pm sem from 12 separate experiments. The maximum relaxation was significantly reduced with TNF- α treatment (see Table 2 for curve fitting parameters).


Figure 4: The effect of (A) insulin (1000 μ IU/ml) and (B) TNF- α (10ng/ml) and insulin (1000 μ IU/ml) on carbachol-induced relaxation of equine digital arteries incubated overnight at 37°C under hypoxic conditions. C: The effect of TNF- α (10ng/ml) and insulin (1000 μ IU/ml) on carbachol-induced relaxation of equine digital arteries incubated overnight at 37°C under hyperoxic conditions. Each point represents mean \pm sem from 6 separate experiments. Under hypoxic conditions, the maximum relaxation was significantly reduced with TNF- α + insulin treatment (Panel B; see Table 2 for curve fitting parameters).

Figure 5: The effect of TNF- α (10ng/ml) and insulin (1000 μ IU/ml) on carbachol-induced relaxation of equine digital veins incubated overnight at 37°C under hyperoxic (A) or hypoxic (B) conditions. Each point represents mean \pm sem from 6 (A) or 7 (B) separate experiments. Under hypoxic conditions, the maximum relaxation was significantly reduced with TNF- α + insulin treatment (Panel B; see Table 2 for curve fitting parameters).

Figure 6: The (A) time course and (B) concentration dependency of the effect of TNF- α on cGMP production by equine digital vein endothelial cells. Data were analysed using a linear mixed effect model. * p <0.05, ** p <0.01, *** p <0.001, **** p <0.0001

Figures

Figure 1



B





Figure 2

A



B




Figure 3






Figure 4

A



B



C





Figure 5

A



B





Figure 6

A



B

