

**RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE**

This is the author's accepted manuscript of the following article:

Meeson, RL and Geddes, AT (2015) Management and long-term outcome of pelvic fractures: a retrospective study of 43 cats. JOURNAL OF FELINE MEDICINE AND SURGERY.

The final publication is available at SAGE Journals via  
<http://dx.doi.org/10.1177/1098612X15606958>.

The full details of the published version of the article are as follows:

TITLE: Management and long-term outcome of pelvic fractures: a retrospective study of 43 cats

AUTHORS: Richard L. Meeson and Alex T. Geddes

JOURNAL TITLE: JOURNAL OF FELINE MEDICINE AND SURGERY

PUBLICATION DATE: 7 October 2015 (online)

PUBLISHER: SAGE Publications

DOI: 10.1177/1098612X15606958

Management and Long-term Outcome of Pelvic Fractures –  
A Retrospective of Forty-Three Cats

Journal:	Journal of Feline Medicine and Surgery
Manuscript ID:	JFMS-15-0152.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Meeson, Richard; Royal Veterinary College, Clinical Sciences and Services Geddes, Alex; Royal Veterinary College, Clinical Sciences and Services
Keywords:	pelvis, constipation, megacolon, surgery, conservative, long-term , feline, fracture, neuropraxia
Abstract:	<p>Objectives: Evaluate the management and long-term outcome of cats with pelvic fractures.</p> <p>Methods: Cats with pelvic fractures had their records and radiographs reviewed. Radiographs were reviewed for fracture configuration, implants and pelvic canal narrowing. Owners were contacted for long-term follow-up using a questionnaire.</p> <p>Results: Forty-three cats met the criteria; mean follow-up of 24 months (range 6-45). The majority (93%) had more than one orthopaedic pelvic injury, with sacro-iliac fracture-luxations seen most commonly. 23% had pre-surgical neurological deficits. Most cats (74%) were managed surgically; 60% of sacroiliac fracture-luxations, 82% ilial fractures, and 50% acetabular fractures received surgery. The complication rate was 22%, most commonly sciatic neuropraxia, (13%). 79% of all neurological deficits resolved and the remainder improved. Mean pelvic canal narrowing after trauma was -15% in surgical and -16% in conservatively managed cats. Canal width was improved postoperatively (-8%), but mildly narrowed further by follow-up (-12%); however these changes were not significant. 19% of cats had had constipation post-surgery; none developed megacolon. There was no clear correlation between the degree of narrowing of the pelvic canal up to -50%, or whether conservative treatment was opted for, and the development of constipation. Long-term mobility was not impaired in 86%, and 84% did not have any lameness detectable.</p> <p>Conclusion and Relevance: The majority were managed surgically, with a 22% complication rate; the most common being transient sciatic neuropraxia. Long-term outcome was generally excellent and most had a full recovery. Constipation/obstipation was very uncommon and no clear relationship to pelvic canal narrowing could be found when considering narrowing of up to 50% in both surgical and conservative groups. As no cats in this cohort had narrowing &gt;50%, the current recommendation of surgery to improve the canal width if narrowing is greater than 45-50% should remain.</p>

1 Title Page

2 Management and Long-term Outcome of Pelvic Fractures

3 – A Retrospective of Forty-Three Cats

4 R.L. Meeson & A.T. Geddes

5 The Royal Veterinary College, Department of Clinical Sciences and Services, Hawkshead Lane, North

6 Mymms, Hatfield, Hertfordshire, AL9 7TA, UK

7 Corresponding Author:

8 Richard L Meeson MA, Vet MB, MVetMed, PGCertVetEd, FHEA, DipECVS, MRCVS

9 [rmeeson@rvc.ac.uk](mailto:rmeeson@rvc.ac.uk)

10 +44(0)1707666366

11 Fax +44(0)1707649384

12 Queen Mother Hospital for Animals, Hawkshead Lane, North Mymms, Hatfield, Hertfordshire, AL9

13 7TA, UK

14

15 Key Words: Pelvis, constipation, megacolon, conservative, fracture, surgery, neuropraxia

16

17

18

19 Abstract

20 Objectives: Evaluate the management and long-term outcome of cats with pelvic fractures.

21 Methods: Cats with pelvic fractures had their records and radiographs reviewed. Radiographs  
22 were reviewed for fracture configuration, implants and pelvic canal narrowing. Owners were  
23 contacted for long-term follow-up using a questionnaire.

24 Results: Forty-three cats met the criteria; mean follow-up of 24 months (range 6-45). The majority  
25 (93%) had more than one orthopaedic pelvic injury, with sacro-iliac fracture-luxations seen most  
26 commonly. 23% had pre-surgical neurological deficits. Most cats (74%) were managed surgically;  
27 60% of sacroiliac fracture-luxations, 82% ilial fractures, and 50% acetabular fractures received  
28 surgery. The complication rate was 22%, most commonly sciatic neuropraxia, (13%). 79% of all  
29 neurological deficits resolved and the remainder improved. Mean pelvic canal narrowing after  
30 trauma was -15% in surgical and -16% in conservatively managed cats. Canal width was improved  
31 postoperatively (-8%), but mildly narrowed further by follow-up (-12%); however these changes  
32 were not significant. 19% of cats had had constipation post-surgery; none developed megacolon.  
33 There was no clear correlation between the degree of narrowing of the pelvic canal up to -50%, or  
34 whether conservative treatment was opted for, and the development of constipation. Long-term  
35 mobility was not impaired in 86%, and 84% did not have any lameness detectable.

36 Conclusion and Relevance: The majority were managed surgically, with a 22% complication rate;  
37 the most common being transient sciatic neuropraxia. Long-term outcome was generally excellent  
38 and most had a full recovery. Constipation/obstipation was very uncommon and no clear  
39 relationship to pelvic canal narrowing could be found when considering narrowing of up to 50% in  
40 both surgical and conservative groups. As no cats in this cohort had narrowing >50%, the current

41 recommendation of surgery to improve the canal width if narrowing is greater than 45-50% should  
42 remain.

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 Introduction

59 Pelvic fractures are common, accounting for 20%-32% of cat fractures<sup>1-3</sup>. In a large retrospective  
60 from the early 1990s of 103 cats with pelvic fractures, 90% of cats had pelvic floor fractures, 60%  
61 had suffered a sacroiliac luxation and 48.5% had ilial fractures<sup>2</sup>. Historically, feline pelvic fractures  
62 were commonly managed conservatively<sup>1,4</sup>, however there has been a shift to surgical management  
63 in recent years, borrowing criteria from canine pelvic fracture management<sup>5</sup>. Indications for  
64 surgery have included pelvic canal narrowing, disruption of the weight bearing axis (acetabular,  
65 ilial body or sacroiliac luxations), nerve impingement, intractable pain, inability to ambulate within  
66 a few days of injury, and bilateral/concomitant orthopaedic injuries<sup>5</sup>. Associated non-orthopaedic  
67 injuries are also common, including urinary tract trauma and neurological deficits being reported  
68 in 59-72% of cases<sup>2,3</sup>. Various techniques have been used to stabilise pelvic fractures in dogs<sup>6-14</sup> and  
69 cats<sup>5,15-20</sup>. Several complications are typically associated with pelvic fractures. Persistent or  
70 subsequent narrowing to the pelvic canal of greater than 45% has been suggested to be a risk factor  
71 for obstipation/constipation<sup>17</sup>. If left unattended, it may progress to megacolon requiring life long  
72 medical treatment or surgical alternatives such as subtotal colectomy and/or pelvic osteotomy<sup>21-24</sup>.  
73 This degree of narrowing therefore has been taken to be an indicator for surgical intervention in  
74 cats<sup>5</sup>. Peripheral nerve damage has also noted to be associated with pelvic fractures, especially ilial  
75 fractures, due to the proximity of the sciatic nerve<sup>3,5,25</sup>. A degree of lameness or decrease in  
76 mobility is also commonly cited post pelvic fracture, however there is sparse evidence to support  
77 this.

78           Currently, there are only limited reports on management of feline pelvic fractures that  
79 include surgical management, and very limited data on their long term outcomes. This study aims  
80 to evaluate the management of feline pelvic fractures, the occurrence of complications, whether

81 there is an association with pelvic canal size and constipation, and what the subsequent long term  
82 outcome is for cats with pelvic fractures.

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98 Materials and methods

99 Medical records (January 2010 to January 2014) of cats admitted with pelvic fractures were  
100 reviewed. Inclusion criteria were presence of any of the following; acetabular, ilial, ischial, pubic  
101 fractures and sacroiliac fracture-luxation, managed either conservatively or surgically, with pre-  
102 operative radiographs available. Surgically managed cats had to have post-operative and follow-up  
103 radiographs. Cats were excluded if follow-up with an owner assessed questionnaire of greater than  
104 six months post fracture was not available. Retrieved data included signalment, fracture  
105 configuration, pre-operative neurological assessment, method of management, post-operative  
106 neurological assessment, and complications. Cats were determined to have cauda-equina signs  
107 when there was a diagnosis recording of the following: 'tail-pull', 'cauda-equina', 'sacrococcygeal  
108 nerve impairment', or the clinical notes recorded a flaccid bladder requiring  
109 expression/catheterization/tube cystostomy; a lack of tail sensation/movement; reduced or absent  
110 anal tone or an absent or decreased perineal reflex. Sciatic neuropraxia was attributed when sciatic  
111 nerve damage was recorded as a diagnosis, or from the clinical notes where a reduced withdrawal  
112 reflex was noted with lack of flexion at the hock, and/or reduced or absent deep pain sensation at  
113 the paw, or knuckling was noted in the absence of other hind limb pathology.

114 Radiographic evaluation included assessment of both lateral and ventrodorsal view  
115 radiographs to determine the fracture configuration, pelvic canal narrowing pre-surgery, post-  
116 surgery and at follow-up using the sacral index (SI)<sup>17</sup>. All measurements were performed in  
117 triplicate and the degree of narrowing was categorised as mild (<10%), moderate (10-30%) and  
118 severe (>30%)<sup>17</sup>. A negative value indicated narrowing and a positive value indicated widening of  
119 the canal above the predicted normal width based on the SI measurement. All radiographic



120 evaluation was performed using DICOM imaging software. (Osirix version 4.1 64-bit open-source  
121 DICOM viewer: Osirix Imaging Software, [http:// www.osirix-viewer.com/OsiriX-64bit.html](http://www.osirix-viewer.com/OsiriX-64bit.html)).

122 Short-term clinical outcome (<3months) and complications were determined from the  
123 patient records at follow-up appointments. Long-term follow-up (>6months) was by postal or  
124 telephone questionnaire to owners using a previously published feline questionnaire<sup>15,17</sup>. Mobility  
125 and lameness were graded from 0-5 with descriptors for each group described to the owners.  
126 Information regarding specific signs of neurological deficits (knuckling, plantigrade stance, low tail  
127 carriage, ataxia) were also requested. Specific questions regarding urination and defecation were  
128 made. Data was gathered, analysed (Microsoft Excel, Microsoft Corp and SPSS v 19.0 IBM Corp), and  
129 assessed for normality and descriptive statistics performed as appropriate. Association of pelvic  
130 narrowing to constipation/obstipation was assessed by Mann-Whitney-U. A P value of <0.05 was  
131 considered significant.

132

133

134

135

136

137

138

139

140 Results

## 141 Cats with pelvic fractures

142 Forty-three cats (mean age 71 months, range 7-219), met the inclusion criteria. Twenty-five cats  
143 were female (23 neutered, two entire), 18 cats were male (17 neutered one entire). Twenty-eight  
144 cats were domestic short hair (65%), seven were domestic long hair (16%), and eight were other  
145 breeds (19%). Fracture configurations and frequency is outlined in Table 1, and sub-classification  
146 of ilial fractures are outlined in Table 2. In summary, when considering bilateral sacroiliac luxation  
147 as more than one fracture, 40/43 (93%) of cats had more than one pelvic injury  
148 (fractures/luxations). Sacroiliac fracture-luxations were most common, being seen in 40/43 (93%);  
149 unilateral or bilateral pubic fractures were present in 31/43 (72%); unilateral or bilateral ischial  
150 fractures were seen in 22/43 (51%) as were ilial fractures, 22/43 (51%). No bilateral ilial fractures  
151 were identified. Acetabular fractures were least common and again were only seen unilaterally, in  
152 11/43 cats (26%).

## 153 Management of fractures

154 The majority of cats (74% 32 cats) underwent surgical stabilisation of their fractures with the  
155 remainder (26% 11 cats) being conservatively managed. More than one surgical  
156 repair/stabilisation was performed in 19/32 cats. Management of fractures were as follows:

- 157 • Sacroiliac fracture-luxations were surgically managed in 24/40 (60%), most commonly  
158 using a unilateral or bilateral 2.0mm or 2.7mm lag screw. Two cats were managed with a  
159 screw and transilial pin, one had a transilial pin alone.
- 160 • Iliac fractures were generally managed surgically in 18/22 (82%) fractures, most  
161 commonly with a single laterally placed 2.0mm DCP, some with a 1.5/2.0 VCP, two cats

162 were double plated, one had a reconstruction plate with K-wires, one had a human radial  
163 2.4mm locking plate, and two were reconstructed using K-wires and lag screws alone.

164 • Acetabular fractures were managed conservatively in 58% (7/12); notably these fractures  
165 tended to be along the caudal acetabular rim or were comminuted, and were combined  
166 with femoral head and neck excision in two cats. Of the surgically managed cats, two had  
167 acetabular plates, one had pins with wire and two were plated using locking or  
168 reconstruction plates.

169 • Pubic fractures were almost exclusively managed conservatively (30/31) other than one  
170 cat who had a pelvic symphyseal separation which had caused bilateral ventroversion of the  
171 hip joints and was managed by pubic symphyseal wiring.

172 • No ischial fractures were managed surgically.

173 The post-operative complication rate was 22%, (7/32). Two cats suffered implant complications  
174 (wire breakage, screw loosening), which did not require any further management; one cat  
175 developed a surgical site swelling suspected to be infection, and the remainder had post-operative  
176 neurological deficits.

#### 177 Neurological injuries

178 Neurological deficits were present in 10/43 cats (23%) on presentation. Sciatic neuropraxia was  
179 most common (7/10), and the remainder (3/10) had cauda-equina signs. No increase in  
180 neurological deficits was seen in the short term in conservatively managed cats, however four cats  
181 surgically managed developed further deficits (sciatic neuropraxia) post surgery (13%). Resolution  
182 of pre-surgical deficits was seen in five cats by follow-up at six to eight weeks, and in the long term  
183 (>6 months), neurological deficits from the trauma or surgery had resolved in 11/14 (79%) of cats,

184 and had improved in a further three. One conservatively managed cat had no detectable  
185 abnormalities at presentation, but went on to develop an unsteady/wobbly gait three months post  
186 fracture.

187 Pelvic canal diameter

188 Mean pre-operative % canal width was not significantly different between surgically managed cats;  
189 -15% (range -43 to +30%) and conservatively managed cats -16% (range -42 to +4%). Post-  
190 surgery, mean canal width had widened to a mean of -8% (range -37 to +26%), however this  
191 increase was not statistically significant. At the six to eight week follow-up, the pelvic canal had  
192 slightly narrowed to -12% (range -51 to +19%), with an average of increased narrowing by 4%,  
193 again not significant. See Table 3 for categorisation of severity of narrowing. Constipation post  
194 fracture was seen in eight cats (19%). Two had problems at least monthly, one only twice a year  
195 and five were very intermittent suffering less than once every year. Half of the cats with  
196 constipation had visited the vet, and 2/8 were medically managed, and 2/8 had no treatment. Cats  
197 that developed constipation had a pelvic canal size range of -27 to +5%. "Severe narrowing" of the  
198 canal of (> 30%)<sup>13,17</sup> was present in six cats managed surgically and conservatively, with a range  
199 from -31 to -51%, however none of these cats developed constipation. Only one cat in this study  
200 had narrowing >45% which has been suggested to be the cut off for increased risk of defecation  
201 problems<sup>17</sup>, however it did not develop any such problems. No cats from this series were reported  
202 to develop megacolon or require any surgical intervention for problems relating to  
203 constipation/obstipation.

204 Long term clinical outcome

205 The mean long term follow-up was 24 months (range 6-45 months). The majority of cats (36/42,  
206 84%) showed no signs of lameness, with only seven (16%) having some degree of permanent  
207 lameness, (see Table 4). The majority of cats were felt to be mobile by their owners, with 86% 'as  
208 expected' to 'very agile for their age'. Only 14% of cats were considered to have impaired mobility  
209 (Table 4).

210

211

212

213

214

215

216

217

218

219

220

221

222

223 Discussion

224 This is the largest group of cats with pelvic fractures that have long-term follow-up of at least six  
225 months post surgery. This cohort was older than previous reports, with a mean age of 71 months,  
226 compared with a mean age of under 17months<sup>3</sup>. The change in demographic may relate to the  
227 increase in motor vehicle traffic since those other cohorts were reviewed or the geographic effect of  
228 living in a metropolitan area.

229 Compared with the largest previously published study<sup>2</sup>, there were significantly higher  
230 levels of sacroiliac fracture-luxations at 93% compared with 60%, and a similar level of ilial body  
231 fractures 51% vs 48.5%. Acetabular fractures were the least common (26% of cats), however this  
232 was still higher than previous reports<sup>2</sup>. The higher levels of individual fracture types, or diagnosis  
233 of them may be attributable to the use of high detail digital radiographs, which were not present  
234 when the previous study was conducted. Furthermore, radiographs were evaluated by board  
235 certified surgical and radiology specialists, potentially increasing the likelihood of detection.  
236 Although not unsurprising, as 93% of pelvic fractures had at least two pelvic orthopaedic injuries  
237 careful evaluation of radiographs needs to be performed if only one fracture is initially identified.

238 Although fixation of the pelvic floor has been described in the literature<sup>26</sup>, this was not  
239 performed routinely in this cohort and did not appear to negatively impact on outcome. Iliac body  
240 fractures are usually an indication for surgical repair<sup>5,15,17</sup>, and surgical stabilisation was performed  
241 in the vast majority. Some combination of lateral plating was most common, usually with a single  
242 DCP plate, and no implant complications were seen other than one cat with screw pull out in the  
243 iliac wing. This cat had a comminuted iliac fracture which was not fully reconstructed, and there was  
244 a conservatively managed sacroiliac luxation which may have contributed to the loads placed upon  
245 the relatively thin cranial iliac wing<sup>15</sup>. Greater consideration may be necessary to stabilising

246 concurrent injuries if there is any compromise in the primarily stabilised fracture. Sacroiliac  
247 fracture luxations were managed surgically in 60% of cats. Several factors are considered when  
248 determining whether to surgically manage these fractures, including whether they are bilateral,  
249 degree of displacement, discomfort and mobility considerations and concurrent injuries<sup>5,18,27</sup>.  
250 Placement of a single or bilateral lag screw<sup>18</sup> remains a popular and successful technique, being  
251 used in most cats here (21/24). Placement of a transilial pin<sup>5,20</sup> in conjunction with a lag screw was  
252 used in two cats that had bilateral sacroiliac luxations, and was also used as sole fixation in one cat.  
253 The transilial pin is a potentially easier technique to perform, and may have particular use for when  
254 sacral wing landmarks are lost, however there are currently no guidelines on placement of  
255 transilial pins in cats. Acetabular fractures were only surgically managed in 50%. This is surprising  
256 as articular fractures are typically treated with reduction and rigid internal fixation, and the  
257 historic opinion that fractures in the caudal 1/3 of the acetabulum did not require surgical  
258 management has been disproved<sup>28</sup>. However, most of the conservatively managed acetabular  
259 fractures in this series had fracture lines which were along the caudal perimeter of the articular  
260 acetabulum and therefore the cost to benefit assessment may have laid in favour of conservative  
261 treatment. The other conservatively managed fractures had degrees of comminution leading to  
262 salvage with a femoral head and neck excision.

263         Neurological deficits were seen in 23% of cats, and therefore careful neurological  
264 evaluation is essential in pelvic fracture cats. Fractures with proximity to other structures, will  
265 inevitably increase the risk of concurrent injuries. The high frequency of sacroiliac fracture-  
266 luxation and ilial fractures seen, could have resulted in damage to the lumbosacral plexus, being  
267 ventral to the sacrum, and feasibly result in a degree of traction or avulsion secondary to sacroiliac  
268 fracture-luxations<sup>29</sup>. Likewise the position of the sciatic nerve medial to the ilial body and then

269 passing over the cranial ischium clearly puts it at risk, and therefore the anatomic proximity would  
270 explain high levels of concurrent neurological impairment. These intimate relationships also  
271 explain the risk of surgically induced nerve impairment<sup>25,30</sup>. During surgery great care is taken to  
272 avoid trapping or stretching nerves, especially the sciatic, however 13% of cats did have post-  
273 operative sciatic neuropraxia. Positively, all of the traumatic and surgically induced neurological  
274 deficits improved, with 79% of cats having complete resolution and the remainder having some  
275 residual impairment, implying that the damage is likely a neuropraxia or axontemesis at worst and  
276 not neurotemesis. Therefore the prognosis for cats with pelvic fractures and hind limb neurological  
277 deficits appears generally good. Only one cat in this cohort developed neurological deficits not  
278 present from the trauma or surgery. This cat was conservatively managed, had bilateral mild  
279 sacroiliac luxations and no neurological deficits on presentation. Although callus healing of bone  
280 fragments have also been suggested to place nerves at risk<sup>3</sup>, the cause of the subsequent weakness  
281 in this cat remains unclear.

282           Post-operative complications occurred in around 1/5 cats, with the majority being post-  
283 surgical sciatic neuropraxia and therefore particular attention should be given to post-operative  
284 neurological deficits when discussing surgical management with owners. Acquired megacolon  
285 secondary to constipation/obstipation is often cited as a potential complication of pelvic fractures,  
286 due to persistent canal narrowing, and is said to account for 25% of megacolon cases.<sup>24,31,32</sup> Pelvic  
287 canal narrowing has become a criteria for surgical management, with narrowing of greater than 45-  
288 50% being reported to increase the risk of megacolon<sup>17</sup> and hence the cut off for surgery. However  
289 there are other causes of megacolon including neurological injury, sacral spinal cord deformity and  
290 most commonly idiopathic<sup>24,31</sup>. This study had a mean follow up of 24months with a minimum of  
291 six months, which was important as clinical signs usually begin shortly after pelvic injury but could



292 take longer than five months<sup>23</sup>. In this follow-up period, only eight cats were reported to have  
293 constipation (19%). No cats were reported to develop megacolon. The cats that developed any  
294 issues with constipation had a pelvic canal size range of -27 to +5%. Severe narrowing of the canal,  
295 when defined as narrowing of greater than 30%<sup>13,17</sup> was present in six cats managed surgically and  
296 conservatively, with a range from -31 to -51%, however none of these cats developed constipation.  
297 Only one cat in this study had narrowing >45% which has been suggested to be the cut off for  
298 increased risk of defecation problems<sup>17</sup>. From the data presented here, it appears that narrowing of  
299 up to 50% does not cause constipation. As no cats in this study had narrowing greater than 50%,  
300 the current recommendation of surgical intervention if the pelvic canal is >45-50% narrowed  
301 should remain, until a cohort of cats with narrowing of greater than 50% has been fully evaluated.

302           Although, it is reassuring to know that the long term outcome of cats with pelvic fractures  
303 is generally excellent, even in those with neurological deficits, there is likely to be some bias in this  
304 population. It is possible that some cats presenting with pelvic fractures may well have had such  
305 severe trauma, including neurological deficits, such as absence of anal tone, perineal reflex, or  
306 bladder function that they may have been euthanised due to the guarded prognosis given. This  
307 study is also unable to determine whether surgical management is superior or not to conservative  
308 management. On the face of it, the outcomes were largely similar, and the pre-operative pelvic canal  
309 narrowing was also similar. However conservative management vs surgical was not randomly  
310 assigned, and usually related to the combinations and configurations of fractures seen. These  
311 populations of cats are therefore not the same. In spite of this, this study shows that cats that  
312 received surgery and those that were intentionally conservatively managed based on current  
313 recommendations<sup>5</sup> can have excellent outcomes.

314

315 Conclusions

316 Current management criteria for feline pelvic fractures appears to work well, with excellent long  
317 term outcomes. Surgical complications are infrequent but are most commonly varying degrees of  
318 sciatic impairment. Positively, neurological deficits from the trauma or surgery resolve in most and  
319 improve in the remainder. No cats developed megacolon however a few did have intermittent  
320 issues with constipation, although the relationship to pelvic injuries is unclear. On balance it  
321 appears that narrowing of up to 45-50% is not a direct risk factor for development of constipation  
322 and megacolon, however narrowing of greater than 50% could potentially still be a risk and  
323 therefore should remain as an indication for surgical intervention.

324 Funding and conflict of interest statement

325 The Authors declare that there is no conflict of interest.

326

327

328

329

330

331

332

333

334

335

336

337 References

- 338 1. Hill FW. A survey of bone fractures in the cat. *J Small Anim Pract.* 1977 Jul;18(7):457-63.
- 339 2. Bookbinder PE, Flanders JA. Characteristics of Pelvic Fracture in the Cat. A 10-Year  
340 Retrospective Study. *Vet Comp Orthop Traumatol.* 1992;5(3):37-42.
- 341 3. Lanz OI, Lanz OI. Lumbosacral and pelvic injuries. *Veterinary Clinics of North America:*  
342 *Small Animal ....* 2002.
- 343 4. Bennett D. Orthopaedic disease affecting the pelvic region of the cat. *J Small Anim Pract.*  
344 1975 Nov;16(11):723-38.
- 345 5. Meeson R, Corr S. Management of Pelvic Trauma. *J. Feline Med. Surg.* 2011 May;13(5):347-  
346 61.
- 347 6. Fitzpatrick N, Lewis D, Cross A. A biomechanical comparison of external skeletal fixation  
348 and plating for the stabilization of ilial osteotomies in dogs. *Vet Comp Orthop Traumatol.*  
349 2008 Jan 1;21(4):349-57.
- 350 7. Miller A, Miller A. Decision making in the management of pelvic fractures in small animals.  
351 *In Practice.* 2002;24(2):54.
- 352 8. Bowlit KL, Fischer A, Friend EJ, Binder E, Delisser P, Reif U, et al. Closed reduction and  
353 percutaneous fixation of sacroiliac luxations in cats using 2.4 mm cannulated screws – a  
354 cadaveric study. *Vet Comp Orthop Traumatol.* 2012;25(1):22-7.
- 355 9. Denny HR. Pelvic fractures in the dog: a review of 123 cases. *Journal of Small Animal*  
356 *Practice.* 1978 Jan;19(1-12):151-66.
- 357 10. Robins G, Dingwall J, Sumner-Smith G. The plating of pelvic fractures in the dog. *Veterinary*  
358 *Record.* 1973 Nov 24;93(21):550-4.
- 359 11. Brown SG, Biggart JF 3rd. Plate fixation of iliac shaft fractures in the dog. *Journal American*  
360 *Vet Med Assoc.* 1975 Sep 1;167(6):472-8.
- 361 12. Hulse D, Van Gundy T, Johnson S et al. Compression Screw Stabilization of Oblique Iliac  
362 Fractures in the Dog. *Vet Comp Orthop Traumatol.* 1989;2(4):32-7.
- 363 13. Breshears LA, Fitch RB, Wallace LJ et al. The radiographic evaluation of repaired canine iliac  
364 fractures (69 cases). *Vet Comp Orthop Traumatol.* 2004;17(2):64.
- 365 14. Gentry SJ, Taylor RA, Dee JF. The use of veterinary cuttable plates: 21 cases. *J Am Anim Hosp*  
366 *Assoc.* 1993.
- 367 15. Langley-Hobbs SJ, Meeson RL, Hamilton MH, Radke H, Lee K. Feline iliac fractures: a  
368 prospective study of dorsal plating and comparison with lateral plating. *Vet Surg.* 2009

- 369 Apr;38(3):334-42.
- 370 16. Langley-Hobbs S, Sissener T, Shales C. Tension band stabilisation of acetabular physeal  
371 fractures in four kittens. *J. Feline Med. Surg* 2007 Jun;9(3):177-87
- 372 17. Hamilton MH, Evans DA, Langley-Hobbs SJ. Feline Iliac Fractures: Assessment of Screw  
373 Loosening and Pelvic Canal Narrowing After Lateral Plating. *Veterinary Surgery*. 2009  
374 Apr;38(3):326-33.
- 375 18. Stabilization of Sacroiliac Luxation in 40 Cats Using Screws Inserted in Lag Fashion. :no-no.
- 376 19. Composite fixation of comminuted iliac wing fractures in cats: three cases. 2011  
377 May;13(5):376-82.
- 378 20. Yap FW, Dunn AL, Farrell M, Calvo I. Trans-iliac pin/bolt/screw internal fixation for  
379 sacroiliac luxation or separation in cats: six cases. *jfm.sagepub.com*.
- 380 21. Muir P, Hardie RJ, Colopy-Poulsen SA, Danova NA. Managing feline obstipation secondary to  
381 pelvic fracture. *Comp Contin Ed*. 2005;27.
- 382 22. Eberhard Rosin DVM. Megacolon in Cats: The Role of Colectomy. *Veterinary Clinics of North  
383 America: Small Animal Practice*. 1993 May;23(3):587-94.
- 384 23. Schrader SC, Schrader SC. Pelvic osteotomy as a treatment for obstipation in cats with  
385 acquired stenosis of the pelvic canal: six cases (1978-1989). *J Am Vet Med Assoc*. 1992 Jan  
386 15;200(2):208-13.
- 387 24. White RN. Surgical management of constipation. *J Feline Med Surg*. 2002 Sep;4(3):129-38
- 388 25. Forterre F, Forterre F, Tomek A, Tomek A, al E, al E. Iatrogenic sciatic nerve injury in  
389 eighteen dogs and nine cats (1997-2006). *Vet Surg*. 2007 Jul 1;36(5):464-71.
- 390 26. Kipfer NM, Kipfer NM, Montavon PM, Montavon PM. Fixation of pelvic floor fractures in cats.  
391 *Veterinary and comparative ....* 2011.
- 392 27. Shales CJ, Shales CJ, White L, White L, Langley-Hobbs SJ, Langley-Hobbs SJ. Sacroiliac  
393 luxation in the cat: defining a safe corridor in the dorsoventral plane for screw insertion in  
394 lag fashion. *Vet Surg*. 2009 Apr;38(3):343-8.
- 395 28. Beck AL, Pead MJ, Draper E. Regional load bearing of the feline acetabulum. *J Biomech*. 2005  
396 Mar;38(3):427-32
- 397 29. Smeak DD, Olmsread M. Fracture/luxations of the sacrococcygeal area in the cat: A  
398 retrospective study of 51 cases. *Vet Surg* 1985; 14: 319-326.
- 399

- 400 30. Bennett D, Vaughan LC. Peroneal nerve paralysis in the cat and dog: an experimental study. J  
401 Small Anim Pract. 1976 Aug;17(8):499-506.
- 402 31. Bertoy RW, Bertoy RW. Megacolon in the cat. Vet Clin North Am Small Anim Pract. 2002  
403 Jul;32(4):901-15.
- 404 32. Hudson EB, Hudson EB, Farrow CS, Farrow CS, Smith SL, Smith SL. Acquired megacolon in a  
405 cat. Mod Vet Pract. 1979 Aug;60(8):625-7.
- 406

Table 1: Fracture classifications, indicating numbers of cats with each fracture type, followed by percentage.

	Acetabular	Iliac	Ischial	Pubic	Sacroiliac
Left	3 (7%)	12 (28%)	9 (21%)	12 (28%)	9 (21%)
Right	8 (18%)	10 (23%)	8 (19%)	13 (30%)	15 (35%)
Bilateral	0 (0%)	0 (0%)	5 (11%)	6 (14%)	16 (37%)
Total	11 (26%)	22 (51%)	22 (51%)	31 (72%)	40 (93%)

Table 2: Sub-classification of ilial fractures, showing ilial fracture configurations and percentage representation. All percentages rounded to nearest whole number.

	Number of Cats	% of ilial fractures
Left comminuted	5	23
Right comminuted	1	5
Total	6	27
Right oblique	4	18
Left oblique	6	27
Total	10	45
Left transverse	1	5
Right transverse	5	23
Total	6	27

Table 3: Classification of pelvic canal narrowing. Widening is pelvic canal diameter greater than the sacral index width. Mild narrowing = 0-10% narrowed, Moderate narrowing = 10-30% and Severe = >30% narrowing.

	Widened	Mild Narrowing	Moderate Narrowing	Severe Narrowing
% Conservative cats	18	9	55	18
% Surgical cats post surgery	22	22	66	0
% Surgical cats follow up	26	30	26	18



Table 4: Lameness and Mobility outcomes from questionnaire. For lameness, grade 0 indicates complete absence of lameness, I indicates barely noticeable lameness, Grade V indicates the lameness could not be worse, and grades II-IV are grades of severity between. For mobility, grade I indicates very agile, grade III indicates mobility consistent with age, grade V indicates poor mobility. Numbers of cats and % out of totals are represented.

Grade	Lameness	Mobility
None	36 (84%)	NA
I	3 (7%)	17 (40%)
II	1 (2%)	7 (16%)
III	2 (5%)	13 (30%)
IV	1 (25%)	6 (14%)
V	0 (0%)	0 (0%)