



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Strategic Trampling to Conserve Soil Moisture in a Namibian Rangeland

I. Zimmermann

Polytechnic of Namibia, Namibia

J. Labuschagne

Farm Weiveld, Namibia

V. K. Namupala

Polytechnic of Namibia, Namibia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/9-3/28>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Strategic trampling to conserve soil moisture in a Namibian rangeland

I. Zimmermann¹, J. Labuschagne² and V.K. Namupala¹

¹ Polytechnic of Namibia, P/Bag 13388, Windhoek, Namibia, E-mail: izimmermann@polytechnic.edu.na

² Farm Weiveld, P.O. Box 53, Gobabis, Namibia

Key words : animal impact, BIOTA, rest, soil moisture, trampling

Introduction There is still a lot of controversy around trampling as a rangeland management tool. Savory (1999) promotes animal impact for a variety of benefits including removal of old plant material, invigoration of existing plants, mulching the soil surface with trampled vegetation, favouring establishment of new plants, returning nutrients to the soil surface and breaking soil crusts that interfere with seed germination and rainfall absorption. On the other hand, Miller (2005) points out that an extensive amount of hydrologic research has failed to support the hypothesis that trampling by large herbivores has beneficial impacts on infiltration, indicating instead that trampling tends to result in lower infiltration rates where it destroys stable soil aggregates. Those who promote trampling all agree that it must be followed by sufficient rest to allow recovery after trampling. This study, falling under the Biodiversity Transect Analysis in Africa (BIOTA) program, investigates trampling applied by three farmers on their low rainfall rangelands.

Materials and methods Strategic trampling is applied on sandy soil (3-8% clay) at Farm Weiveld (23.06° S, 18.88° E) in Namibia's Camelthorn Savanna, with a mean annual rainfall of about 250mm. Initial observations were casual, by digging soil to feel how moist it was at different depths, in relation to trampling by livestock under different conditions. Moisture content of soil augured on two occasions from measured depths on each side of the boundary fence was determined gravimetrically. Gypsum blocks have subsequently been buried at 10, 25, 50 and 80cm depth, both in and outside enclosures, replicated five times on each of three farms, for electronic determination of soil moisture availability. The number of perennial grasses growing within 0.75m of each of 100 sample points were counted on each side of a fence separating two paddocks, one of which had been briefly trampled earlier in the growing season.

Results Casual observations suggested that brief trampling after good rain conserved soil moisture, possibly by breaking the capillary connections that suck moisture from lower layers in untrampled soil. Figure 1 shows slightly more soil moisture at the end of the dry season, in April, between 30 and 80cm depth in the strategically trampled soil, suggesting that infiltration may have been improved. However, by mid dry season, in July, it appeared drier below 30cm, possibly due to higher transpiration through the five times greater density of perennial grasses there. No results are available yet from the gypsum blocks. The perennial grass density was significantly higher where brief trampling had been applied shortly after the first effective rain of the season (Figure 2).




Figure 1 Successive soil moisture profiles under strategic trampling and conventional rotation.




Figure 2 Mean perennial grass density each side of fence that received different trampling treatments (error bars show 95% confidence limits).

Conclusions Initial indications suggest that brief trampling, followed by rest, can be applied strategically both before and after rain to achieve certain rangeland management objectives, provided the conditions are favourable. While Savory (1999) seems to treat animal impact as a simple tool, with variables of type of animal, stocking density and timing, observations at Farm Weiveld suggest that trampling is more complex and should furthermore be differentiated between its variables of season, soil texture, soil moisture profile and organic matter content of the soil.

References

- Miller, M.E. 2005. The structure and function of dryland ecosystems—Conceptual models to inform long-term ecological monitoring. U.S. Geological Survey Scientific Investigations Report 2005-1597, 73pp.
- Savory, A. 1999. *Holistic management*. Island Press, 610pp.