

# Neutrosophic Sets and Systems

---

Volume 38

Article 12

---

11-20-2020

## On New Types of Weakly Neutrosophic Crisp Continuity

Qays Hatem Imran

Riad K. Al-Hamido

Ali Hussein Mahmood Al-Obaidi

Follow this and additional works at: [https://digitalrepository.unm.edu/nss\\_journal](https://digitalrepository.unm.edu/nss_journal)

---

### Recommended Citation

Imran, Qays Hatem; Riad K. Al-Hamido; and Ali Hussein Mahmood Al-Obaidi. "On New Types of Weakly Neutrosophic Crisp Continuity." *Neutrosophic Sets and Systems* 38, 1 (2020).  
[https://digitalrepository.unm.edu/nss\\_journal/vol38/iss1/12](https://digitalrepository.unm.edu/nss_journal/vol38/iss1/12)

This Article is brought to you for free and open access by UNM Digital Repository. It has been accepted for inclusion in *Neutrosophic Sets and Systems* by an authorized editor of UNM Digital Repository. For more information, please contact [amywinter@unm.edu](mailto:amywinter@unm.edu), [lsloane@salud.unm.edu](mailto:lsloane@salud.unm.edu), [sarahrk@unm.edu](mailto:sarahrk@unm.edu).



# On New Types of Weakly Neutrosophic Crisp Continuity

Qays Hatem Imran<sup>1\*</sup>, Riad K. Al-Hamido<sup>2</sup> and Ali Hussein Mahmood Al-Obaidi<sup>3</sup>

<sup>1</sup>Department of Mathematics, College of Education for Pure Science, Al-Muthanna University, Samawah, Iraq.  
E-mail: qays.imran@mu.edu.iq

<sup>2</sup>Department of Mathematics, College of Science, Al-Baath University, Homs, Syria.  
E-mail: riad-hamido1983@hotmail.com

<sup>3</sup>Department of Mathematics, College of Education for Pure Science, University of Babylon, Hillah, Iraq.  
E-mail: aalobaidi@uobabylon.edu.iq

\* Correspondence: qays.imran@mu.edu.iq

**Abstract:** The article processes the conceptualizations of neutrosophic crisp  $\alpha$ -open and neutrosophic crisp semi- $\alpha$ -open sets to define some new types of weakly "neutrosophic crisp continuity" essentially, neutrosophic crisp  $\alpha^*$ -continuous, neutrosophic crisp  $\alpha^{**}$ -continuous, neutrosophic crisp semi- $\alpha$ -continuous, neutrosophic crisp semi- $\alpha^*$ -continuous and neutrosophic crisp semi- $\alpha^{**}$ -continuous functions. Also, we shall explain the relationships between these types of weakly neutrosophic crisp continuity and the concepts of neutrosophic crisp continuity.

**Keywords:** Neutrosophic crisp  $\alpha^*$ -continuous, neutrosophic crisp  $\alpha^{**}$ -continuous, neutrosophic crisp semi- $\alpha$ -continuous, neutrosophic crisp semi- $\alpha^*$ -continuous, and neutrosophic crisp semi- $\alpha^{**}$ -continuous functions.

## 1. Introduction

In 2014, Salama et al. [1] performed the abstraction of neutrosophic crisp topological space (concisely, *NCTS*). Al-Hamido et al. [2] submitted the intellect of neutrosophic crisp semi- $\alpha$ -closed sets in *NCTS*s. Abdel-Basset et al. [3-8] gave a novel neutrosophic approach. Maheswari et al. [9] presented gb-closed sets and gb-continuity in aspects of the neutrosophic theory. Banupriya et al. [10] investigated the notion of  $\alpha$ gs continuity and  $\alpha$ gs irresolute maps in the sense of neutrosophic view. In [11], Dhavaseelan et al. exhibited the theme of neutrosophic  $\alpha^m$ -continuity. Al-Hamido et al. [15] introduced neutrosophic crisp topology via N-topology. Imran et al. [16] introduced and studied the thought of neutrosophic generalized alpha generalized continuity. Hanif PAGE et al. [17] presented neutrosophic generalized homeomorphism. This paper aspires to lay on new types of weakly neutrosophic crisp continuity, for instance, neutrosophic crisp  $\alpha^*$ -continuous, neutrosophic crisp  $\alpha^{**}$ -continuous, neutrosophic crisp semi- $\alpha$ -continuous, neutrosophic crisp semi- $\alpha^*$ -continuous and neutrosophic crisp semi- $\alpha^{**}$ -continuous functions. Likewise, we shall explain the relationships between these types of weakly neutrosophic crisp continuity and the concepts of neutrosophic crisp continuity.

## 2. Preliminaries

For the whole of the disquisition,  $(\mathbb{X}, F_1)$ ,  $(\mathbb{Y}, F_2)$ , and  $(\mathbb{Z}, F_3)$  (merely  $\mathbb{X}$ ,  $\mathbb{Y}$ , and  $\mathbb{Z}$ ) habitually intend *NCTS*s. Let  $\mathcal{C}$  be a neutrosophic crisp set (shortly, *NCS*) in *NCTS*  $(\mathbb{X}, F_1)$  and denote its complement by  $\mathcal{C}^c$ . Indicate the neutrosophic crisp open set as *NC-OS*, and the neutrosophic crisp closed set (its complement) as *NC-CS* in *NCTS*  $(\mathbb{X}, F_1)$ . Additionally, we refer to the neutrosophic crisp closure and neutrosophic crisp interior of  $\mathcal{C}$  via  $NCcl(\mathcal{C})$  and  $NCint(\mathcal{C})$ , correspondingly.

**Definition 2.1 [1]:** Assume that nonempty particular understudy space  $\mathbb{X}$  has mutually disjoint subsets  $\mathcal{C}_1, \mathcal{C}_2$  and  $\mathcal{C}_3$ . A *NCS*  $\mathcal{C}$  with form  $\mathcal{C} = \langle \mathcal{C}_1, \mathcal{C}_2, \mathcal{C}_3 \rangle$  is called an object.

**Definition 2.2:** For any *NCS*  $\mathcal{C}$  in *NCTS*  $(\mathbb{X}, I_1)$ , we have

- (i) if  $\mathcal{C} \subseteq NCint(NCcl(NCint(\mathcal{C})))$ , then it is called a neutrosophic crisp  $\alpha$ -open set and symbolize by *NC $\alpha$ -OS*. Furthermore, its complement is named neutrosophic crisp  $\alpha$ -closed set and signified by *NC $\alpha$ -CS*. Likewise, we reveal the collection consisting of all *NC $\alpha$ -OSs* in  $\mathbb{X}$  with *NC $\alpha$ O*( $\mathbb{X}$ ). [12]
- (ii) if  $\mathcal{C} \subseteq NCcl(NCint(NCcl(NCint(\mathcal{C}))))$ , then it is said to be a neutrosophic crisp semi- $\alpha$ -open set and indicated via *NCS  $\alpha$ -OS*. Moreover, its complement is known as a neutrosophic crisp semi- $\alpha$ -closed set and referred with *NCS $\alpha$ -CS*. Besides, we mentioned the collection of all *NCS $\alpha$ -OSs* in  $\mathbb{X}$  through *NCS $\alpha$ O*( $\mathbb{X}$ ). [2]

**Proposition 2.3 [12]:** For any *NCS*  $\mathcal{C}$  in *NCTS*  $(\mathbb{X}, I_1)$ , then  $\mathcal{C} \in NC\alpha O(\mathbb{X})$  iff we have at least a *NC-OS*  $\mathcal{D}$  satisfying  $\mathcal{D} \subseteq \mathcal{C} \subseteq NCint(NCcl(\mathcal{D}))$ .

**Proposition 2.4 [14]:** Every *NC-OS* is a *NC $\alpha$ -OS*, but the opposite is not valid in general.

**Proposition 2.5 [2]:** In a *NCTS*  $(\mathbb{X}, I_1)$ , the next assertions stand, but not vice versa:

- (i) All *NC-OSs* are *NCS $\alpha$ -OSs*.
- (ii) All *NC $\alpha$ -OSs* are *NCS $\alpha$ -OSs*.

**Definition 2.6 [1]:** Let  $\eta: (\mathbb{X}, I_1) \rightarrow (\mathbb{Y}, I_2)$  be a function, we called it a neutrosophic crisp continuous and denoted by *NC-continuous* iff for all *NC-OSs*  $\mathcal{D}$  from  $\mathbb{Y}$ , then its inverse image  $\eta^{-1}(\mathcal{D})$  is a *NC-OS* from  $\mathbb{X}$ .

**Theorem 2.7 [1]:** A function  $\eta: (\mathbb{X}, I_1) \rightarrow (\mathbb{Y}, I_2)$  is *NC-continuous* iff  $\eta^{-1}(NCint(\mathcal{D})) \subseteq NCint(\eta^{-1}(\mathcal{D}))$  for every  $\mathcal{D} \subseteq \mathbb{Y}$ .

**Definition 2.8 [1]:** Let  $\eta: (\mathbb{X}, I_1) \rightarrow (\mathbb{Y}, I_2)$  be a function, we named it a neutrosophic crisp open and indicated via *NC-open* iff for all *NC-OSs*  $\mathcal{C}$  from  $\mathbb{X}$ , then its image  $\eta(\mathcal{C})$  is a *NC-OS* from  $\mathbb{Y}$ .

**Definition 2.9 [13]:** Let  $\eta: (\mathbb{X}, I_1) \rightarrow (\mathbb{Y}, I_2)$  be a function, we said it a neutrosophic crisp  $\alpha$ -continuous and referred through *NC $\alpha$ -continuous* iff for all *NC-OSs*  $\mathcal{D}$  from  $\mathbb{Y}$ , then its inverse image  $\eta^{-1}(\mathcal{D})$  is a *NC $\alpha$ -OS* from  $\mathbb{X}$ .

**Proposition 2.10 [14]:** Every *NC-continuous* function is a *NC $\alpha$ -continuous*, but the opposite is not valid in general.

### 3. Weakly Neutrosophic Crisp Continuity Functions

**Definition 3.1:** Let  $\eta: (\mathbb{X}, I_1) \rightarrow (\mathbb{Y}, I_2)$  be a function, we call it as

- (i) a neutrosophic crisp  $\alpha^*$ -continuous and denoted by  $NC\alpha^*$ -continuous iff for all  $NC\alpha$ -OSs  $\mathcal{D}$  from  $\mathbb{Y}$ , then its inverse image  $\eta^{-1}(\mathcal{D})$  is a  $NC\alpha$ -OS from  $\mathbb{X}$ .
- (ii) a neutrosophic crisp  $\alpha^{**}$ -continuous and indicated via  $NC\alpha^{**}$ -continuous iff for all  $NC\alpha$ -OS  $\mathcal{D}$  from  $\mathbb{Y}$ , then its inverse image  $\eta^{-1}(\mathcal{D})$  is a  $NC$ -OS from  $\mathbb{X}$ .

**Definition 3.2:** Let  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  be a function, we named it as

- (i) a neutrosophic crisp semi- $\alpha$ -continuous and referred through  $NCS\alpha$ -continuous iff for all  $NC$ -OSs  $\mathcal{D}$  from  $\mathbb{Y}$ , then its inverse image  $\eta^{-1}(\mathcal{D})$  is a  $NCS\alpha$ -OS from  $\mathbb{X}$ .
- (ii) a neutrosophic crisp semi- $\alpha^*$ -continuous and symbolize by  $NCS\alpha^*$ -continuous iff for all  $NCS\alpha$ -OSs  $\mathcal{D}$  from  $\mathbb{Y}$ , then its inverse image  $\eta^{-1}(\mathcal{D})$  is a  $NCS\alpha$ -OS from  $\mathbb{X}$ .
- (iii) a neutrosophic crisp semi- $\alpha^{**}$ -continuous and signified via  $NCS\alpha^{**}$ -continuous iff for all  $NCS\alpha$ -OSs  $\mathcal{D}$  from  $\mathbb{Y}$ , then its inverse image  $\eta^{-1}(\mathcal{D})$  is a  $NC$ -OS from  $\mathbb{X}$ .

**Theorem 3.3:** Let  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  be a function, then the next declarations are same:

- (i)  $\eta$  is a  $NCS\alpha$ -continuous.
- (ii) its inverse image of each  $NC$ -CS from  $\mathbb{Y}$  is  $NCS\alpha$ -CS from  $\mathbb{X}$ .
- (iii)  $\eta(NCint(NCcl(NCint(NCcl(\mathcal{C})))) \subseteq NCcl(\eta(\mathcal{C}))$ , for each  $\mathcal{C} \in \mathbb{X}$ .
- (iv)  $NCint(NCcl(NCint(NCcl(\eta^{-1}(\mathcal{D})))) \subseteq \eta^{-1}(NCcl(\mathcal{D}))$ , for each  $\mathcal{D} \in \mathbb{Y}$ .

**Proof:**

[(i)  $\Rightarrow$  (ii)] Suppose  $\mathcal{D}$  is a  $NC$ -CS from  $\mathbb{Y}$ . This implies that  $\mathcal{D}^c$  stands a  $NC$ -OS. Hence  $\eta^{-1}(\mathcal{D}^c)$  is a  $NCS\alpha$ -OS from  $\mathbb{X}$ . In other words,  $(\eta^{-1}(\mathcal{D}))^c$  stands a  $NCS\alpha$ -OS from  $\mathbb{X}$ . Thus  $\eta^{-1}(\mathcal{D})$  is a  $NCS\alpha$ -CS in  $\mathbb{X}$ .

[(ii)  $\Rightarrow$  (iii)] Let  $\mathcal{C} \in \mathbb{X}$ , then  $NCcl(\eta(\mathcal{C}))$  stays a  $NC$ -CS from  $\mathbb{Y}$ . Hence  $\eta^{-1}(NCcl(\eta(\mathcal{C})))$  is  $NCS\alpha$ -CS in  $\mathbb{X}$ . Thus we have  $\eta^{-1}(NCcl(\eta(\mathcal{C}))) \supseteq NCint(NCcl(NCint(NCcl(\eta^{-1}(NCcl(\eta(\mathcal{C})))))) \supseteq NCint(NCcl(NCint(NCcl(\mathcal{C})))$ .

Or  $NCcl(\eta(\mathcal{C})) \supseteq \eta(NCint(NCcl(NCint(NCcl(\mathcal{C}))))$ .

[(iii)  $\Rightarrow$  (iv)] Since  $\mathcal{D} \in \mathbb{Y}, \eta^{-1}(\mathcal{D}) \in \mathbb{X}$ . So, we have by our hypothesis the corresponding notation  $NCint(NCcl(NCint(NCcl(\eta^{-1}(\mathcal{D})))) \subseteq NCcl(\eta(\eta^{-1}(\mathcal{D}))) \subseteq NCcl(\mathcal{D})$ , and that leads us to this fact  $NCint(NCcl(NCint(NCcl(\eta^{-1}(\mathcal{D})))) \subseteq \eta^{-1}(NCcl(\mathcal{D}))$ .

[(iv)  $\Rightarrow$  (i)] Let  $\mathcal{D}$  be a  $NC$ -OS of  $\mathbb{Y}$ . Let  $\mathcal{C} = \mathcal{D}^c$  and  $\mathcal{D} = \eta^{-1}(\mathcal{C})$  by (iii) we have  $NCint(NCcl(NCint(NCcl(\eta^{-1}(\mathcal{C})))) \subseteq NCcl(\mathcal{C}) = \mathcal{C}$ .

That is  $NCint(NCcl(NCint(NCcl(\eta^{-1}(\mathcal{D}^c)))) \subseteq \eta^{-1}(\mathcal{D}^c)$ . Or  $NCint(NCcl(NCint(NCcl(\eta^{-1}(\mathcal{D})))) \supseteq \eta^{-1}(\mathcal{D})$ . Hence  $\eta^{-1}(\mathcal{D})$  is a  $NCS\alpha$ -OS in  $\mathbb{X}$  and thus  $\eta$  be there a  $NCS\alpha$ -continuous. ■

**Proposition 3.4:**

- (i) all  $NC$ -continuous functions are  $NCS\alpha$ -continuous, but the opposite is not valid in general.
- (ii) all  $NC\alpha$ -continuous functions are  $NCS\alpha$ -continuous, but the opposite is not exact in general.

**Proof:**

(i) Suppose  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  is a  $NC$ -continuous function, and  $\mathcal{D}$  be a  $NC$ -OS from  $\mathbb{Y}$ . Next  $\eta^{-1}(\mathcal{D})$  remains a  $NC$ -OS from  $\mathbb{X}$ . Since any  $NC$ -OS is a  $NCS\alpha$ -OS,  $\eta^{-1}(\mathcal{D})$  stays a  $NCS\alpha$ -OS from  $\mathbb{X}$ . Thus  $\eta$  exists a  $NCS\alpha$ -continuous function.

(ii) Let  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  be a  $NC\alpha$ -continuous function and  $\mathcal{D}$  be a  $NC$ -OS from  $\mathbb{Y}$ . Subsequently  $\eta^{-1}(\mathcal{D})$  happens a  $NC\alpha$ -OS from  $\mathbb{X}$ . Since any  $NC\alpha$ -OS is  $NCS\alpha$ -OS,  $\eta^{-1}(\mathcal{D})$  stays a  $NCS\alpha$ -OS from  $\mathbb{X}$ . Thus  $\eta$  is a  $NCS\alpha$ -continuous function. ■

**Example 3.5:** Suppose  $\mathbb{X} = \{p, q, r, s\}$  and  $\mathbb{Y} = \{u, v, w\}$ . Then  $\Gamma_1 = \{\phi_N, \mathbb{X}_N\} \cup \{\{p\}, \phi, \phi\}$  and  $\Gamma_2 = \{\phi_N, \mathbb{Y}_N\} \cup \{\{u\}, \phi, \phi\}$  be neutrosophic crisp topologies (shortly,  $NCTs$ ) on  $\mathbb{X}$  and  $\mathbb{Y}$ , correspondingly. Define the function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  via  $\eta(\{p\}, \phi, \phi) = \eta(\{q\}, \phi, \phi) = \langle\{u\}, \phi, \phi\rangle$ ,  $\eta(\{r\}, \phi, \phi) = \langle\{v\}, \phi, \phi\rangle$ ,  $\eta(\{s\}, \phi, \phi) = \langle\{w\}, \phi, \phi\rangle$ . Then  $\eta$  is a  $NC \alpha$ -continuous function but not  $NC$ -continuous since  $\langle\{u\}, \phi, \phi\rangle$  is  $NC$ -OS but  $\eta^{-1}(\langle\{u\}, \phi, \phi\rangle) = \langle\{p, q\}, \phi, \phi\rangle$  which is not  $NC$ -OS in  $\mathbb{X}$ . Also,  $\eta$  is a  $NCS\alpha$ -continuous function but not  $NC$ -continuous, since  $\langle\{u\}, \phi, \phi\rangle$  is  $NC$ -OS in  $\mathbb{Y}$  but  $\eta^{-1}(\langle\{u\}, \phi, \phi\rangle) = \langle\{p, q\}, \phi, \phi\rangle$  is not  $NC$ -OS from  $\mathbb{X}$ .

**Example 3.6:** Suppose  $\mathbb{X} = \{p, q, r\}$ . Then  $\Gamma = \{\phi_N, \mathbb{X}_N\} \cup \{\{p\}, \phi, \phi\}, \langle\{q\}, \phi, \phi\rangle, \langle\{p, q\}, \phi, \phi\rangle\}$  be a  $NCT$  on  $\mathbb{X}$ .

Define the function  $\eta: (\mathbb{X}, \Gamma) \rightarrow (\mathbb{X}, \Gamma)$  by  $\eta(\{p\}, \phi, \phi) = \langle\{p\}, \phi, \phi\rangle$ ,  $\eta(\langle\{q\}, \phi, \phi\rangle) = \eta(\langle\{r\}, \phi, \phi\rangle) = \langle\{q\}, \phi, \phi\rangle$ . It is easily seen that  $\eta$  is a  $NCS\alpha$ -continuous function but not  $NC\alpha$ -continuous, since  $\langle\{q\}, \phi, \phi\rangle$  is  $NC$ -OS in  $\mathbb{X}$  but  $\eta^{-1}(\langle\{q\}, \phi, \phi\rangle) = \langle\{q, r\}, \phi, \phi\rangle$  is not  $NC\alpha$ -OS in  $\mathbb{X}$ .

**Remark 3.7:** The concepts of  $NC$ -continuity and  $NC\alpha^*$ -continuity are independent, for examples.

**Example 3.8:** Suppose  $\mathbb{X} = \{p, q, r, s\}$  and  $\mathbb{Y} = \{u, v, w\}$ . Then

$\Gamma_1 = \{\phi_N, \mathbb{X}_N\} \cup \{\{p\}, \phi, \phi\}, \langle\{q, r\}, \phi, \phi\rangle, \langle\{p, q, r\}, \phi, \phi\rangle\}$  and  $\Gamma_2 = \{\phi_N, \mathbb{Y}_N\} \cup \{\{u\}, \phi, \phi\}$  be  $NCTs$  on  $\mathbb{X}$  and  $\mathbb{Y}$ , correspondingly. Define the function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  via  $\eta(\{p\}, \phi, \phi) = \langle\{u\}, \phi, \phi\rangle$ ,  $\eta(\langle\{q\}, \phi, \phi\rangle) = \langle\{v\}, \phi, \phi\rangle$ ,  $\eta(\langle\{r\}, \phi, \phi\rangle) = \eta(\langle\{s\}, \phi, \phi\rangle) = \langle\{w\}, \phi, \phi\rangle$ . Then  $\eta$  is a  $NC$ -continuous function but not  $NC\alpha^*$ -continuous, since  $\langle\{u, v\}, \phi, \phi\rangle$  is  $NC\alpha$ -OS in  $\mathbb{Y}$  but  $\eta^{-1}(\langle\{u, v\}, \phi, \phi\rangle) = \langle\{p, q\}, \phi, \phi\rangle$  is not  $NC\alpha$ -OS in  $\mathbb{X}$ .

**Example 3.9:** Assume  $\mathbb{X} = \{p, q, r, s\}$  and  $\mathbb{Y} = \{u, v, w\}$ . Then  $\Gamma_1 = \{\phi_N, \mathbb{X}_N\} \cup \{\{p\}, \phi, \phi\}$  and  $\Gamma_2 = \{\phi_N, \mathbb{Y}_N\} \cup \{\{u\}, \phi, \phi\}$  be  $NCTs$  on  $\mathbb{X}$  and  $\mathbb{Y}$ , correspondingly. Define the function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  via  $\eta(\{p\}, \phi, \phi) = \eta(\langle\{q\}, \phi, \phi\rangle) = \langle\{u\}, \phi, \phi\rangle$ ,  $\eta(\langle\{r\}, \phi, \phi\rangle) = \langle\{v\}, \phi, \phi\rangle$ ,  $\eta(\langle\{s\}, \phi, \phi\rangle) = \langle\{w\}, \phi, \phi\rangle$ . Then  $\eta$  is a  $NC \alpha^*$ -continuous function but not  $NC$ -continuous, since  $\langle\{u\}, \phi, \phi\rangle$  is  $NC$ -OS in  $\mathbb{Y}$ , but  $\eta^{-1}(\langle\{u\}, \phi, \phi\rangle) = \langle\{p, q\}, \phi, \phi\rangle$  is not  $NC$ -OS in  $\mathbb{X}$ .

**Theorem 3.10:**

- (i) If a function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  is  $NC$ -open,  $NC$ -continuous, and bijective, then  $\eta$  is a  $NC\alpha^*$ -continuous.
- (ii) A function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  is  $NC\alpha^*$ -continuous iff  $\eta: (\mathbb{X}, NC\alpha O(\mathbb{X})) \rightarrow (\mathbb{Y}, NC\alpha O(\mathbb{Y}))$  is a  $NC$ -continuous.

**Proof:**

(i) Let  $\mathcal{D} \in NC\alpha O(\mathbb{Y})$ , to prove that  $\eta^{-1}(\mathcal{D}) \in NC\alpha O(\mathbb{X})$ , i.e.,  $\eta^{-1}(\mathcal{D}) \subseteq NCint(NCcl(NCint(\eta^{-1}(\mathcal{D}))))$ . Let  $r \in \eta^{-1}(\mathcal{D}) \Rightarrow \eta(r) \in \mathcal{D}$ . Hence  $\eta(r) \in NCint(NCcl(NCint(\mathcal{D})))$  (since  $\mathcal{D} \in NC\alpha O(\mathbb{Y})$ ). Therefore, at least  $NC$ -OS  $\mathcal{H}$  from  $\mathbb{Y}$  where  $\eta(r) \in \mathcal{H} \subseteq NCcl(NCint(\mathcal{D}))$ . Then  $r \in \eta^{-1}(\mathcal{H}) \subseteq$

$\eta^{-1}(NCcl(NCint(\mathcal{D})))$  , but  $\eta^{-1}(NCcl(NCint(\mathcal{D}))) \subseteq NCcl(\eta^{-1}(NCint(\mathcal{D})))$  (since  $\eta^{-1}$  is a  $NC$ -continuous, which is equivalent to  $\eta$  is a  $NC$ -open and bijective). Then  $r \in \eta^{-1}(\mathcal{H}) \subseteq NCcl(\eta^{-1}(NCint(\mathcal{D})))$ . Hence  $r \in \eta^{-1}(\mathcal{H}) \subseteq NCcl(\eta^{-1}(NCint(\mathcal{D}))) \subseteq NCcl(NCint(\eta^{-1}(\mathcal{D})))$  (since  $\eta$  is a  $NC$ -continuous). Hence  $r \in \eta^{-1}(\mathcal{H}) \subseteq NCcl(NCint(\eta^{-1}(\mathcal{D})))$ , but  $\eta^{-1}(\mathcal{H})$  remains a  $NC$ -OS from  $\mathbb{X}$  (because  $\eta$  be present a  $NC$ -continuous). Therefore,  $r \in NCint(NCcl(NCint(\eta^{-1}(\mathcal{D}))))$ .

Hence  $\eta^{-1}(\mathcal{D}) \subseteq NCint(NCcl(NCint(\eta^{-1}(\mathcal{D})))) \Rightarrow \eta^{-1}(\mathcal{D}) \in NC\alpha O(\mathbb{X}) \Rightarrow \eta$  is a  $NC\alpha^*$ -continuous.

(ii) The proof of (ii) is easily. ■

**Theorem 3.11:** A function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  is a  $NCS\alpha^*$ -continuous iff  $\eta: (\mathbb{X}, NCS\alpha O(\mathbb{X})) \rightarrow (\mathbb{Y}, NCS\alpha O(\mathbb{Y}))$  is a  $NC$ -continuous.

**Proof:** Obvious. ■

**Remark 3.12:** The concepts of  $NC$ -continuity and  $NCS\alpha^*$ -continuity are independent, for examples.

**Example 3.13:** Suppose  $\mathbb{X} = \{p, q, r, s\}$  and  $\mathbb{Y} = \{u, v, w\}$ .

Then  $\Gamma_1 = \{\phi_N, \mathbb{X}_N\} \cup \{\langle\{p\}, \phi, \phi\rangle, \langle\{q, r\}, \phi, \phi\rangle, \langle\{p, q, r\}, \phi, \phi\rangle\}$  and  $\Gamma_2 = \{\phi_N, \mathbb{Y}_N\} \cup \{\langle\{u\}, \phi, \phi\rangle\}$  be  $NCT$ s on  $\mathbb{X}$  and  $\mathbb{Y}$ , correspondingly. Define the function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  via  $\eta(\langle\{p\}, \phi, \phi\rangle) = \langle\{u\}, \phi, \phi\rangle, \eta(\langle\{q\}, \phi, \phi\rangle) = \langle\{v\}, \phi, \phi\rangle, \eta(\langle\{r\}, \phi, \phi\rangle) = \eta(\langle\{s\}, \phi, \phi\rangle) = \langle\{w\}, \phi, \phi\rangle$ . It is easily seen that  $\eta$  is a  $NC$ -continuous function but not  $NCS\alpha^*$ -continuous, since  $\langle\{u, v\}, \phi, \phi\rangle$  is  $NCS\alpha$ -OS in  $\mathbb{Y}$  but  $\eta^{-1}(\langle\{u, v\}, \phi, \phi\rangle) = \langle\{p, q\}, \phi, \phi\rangle$  is not  $NCS\alpha$ -OS in  $\mathbb{X}$ .

**Example 3.14:** Assume  $\mathbb{X} = \{p, q, r, s\}$  and  $\mathbb{Y} = \{u, v, w\}$ . Then  $\Gamma_1 = \{\phi_N, \mathbb{X}_N\} \cup \{\langle\{p\}, \phi, \phi\rangle\}$  and  $\Gamma_2 = \{\phi_N, \mathbb{Y}_N\} \cup \{\langle\{u\}, \phi, \phi\rangle\}$  be  $NCT$ s on  $\mathbb{X}$  and  $\mathbb{Y}$ , correspondingly.

Define the function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  via  $\eta(\langle\{p\}, \phi, \phi\rangle) = \eta(\langle\{q\}, \phi, \phi\rangle) = \langle\{u\}, \phi, \phi\rangle, \eta(\langle\{r\}, \phi, \phi\rangle) = \langle\{v\}, \phi, \phi\rangle, \eta(\langle\{s\}, \phi, \phi\rangle) = \langle\{w\}, \phi, \phi\rangle$ . Then  $\eta$  is a  $NCS\alpha^*$ -continuous function but not  $NC$ -continuous, since  $\langle\{u\}, \phi, \phi\rangle$  is  $NC$ -OS in  $\mathbb{Y}$ , but  $\eta^{-1}(\langle\{u\}, \phi, \phi\rangle) = \langle\{p, q\}, \phi, \phi\rangle$  is not  $NC$ -OS in  $\mathbb{X}$ .

**Proposition 3.15:** Every  $NC\alpha^*$ -continuous function is a  $NC\alpha$ -continuous and  $NCS\alpha$ -continuous; however, the reverse generally is not valid.

**Proof:** Assume  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  is a  $NC\alpha^*$ -continuous function and let  $\mathcal{D}$  be any  $NC$ -OS from  $\mathbb{Y}$ . Then we have  $\mathcal{D}$  as a  $NC\alpha$ -OS from  $\mathbb{Y}$  [from proposition 2.4]. Since  $\eta$  is a  $NC\alpha^*$ -continuous, then  $\eta^{-1}(\mathcal{D})$  considers a  $NC\alpha$ -OS from  $\mathbb{X}$ . Thus,  $\eta$  stands a  $NC\alpha$ -continuous. Also,  $\eta$  is a  $NCS\alpha$ -continuous. ■

**Example 3.16:** Let  $\mathbb{X} = \{p, q, r, s\}$ .

Then  $\Gamma = \{\phi_N, \mathbb{X}_N\} \cup \{\langle\{p\}, \phi, \phi\rangle, \langle\{q\}, \phi, \phi\rangle, \langle\{p, q\}, \phi, \phi\rangle, \langle\{p, q, r\}, \phi, \phi\rangle\}$  be a  $NCT$  on  $\mathbb{X}$ . Define the function  $\eta: (\mathbb{X}, \Gamma) \rightarrow (\mathbb{X}, \Gamma)$  by  $\eta(\langle\{p\}, \phi, \phi\rangle) = \langle\{p\}, \phi, \phi\rangle, \eta(\langle\{q\}, \phi, \phi\rangle) = \eta(\langle\{r\}, \phi, \phi\rangle) = \langle\{s\}, \phi, \phi\rangle, \eta(\langle\{s\}, \phi, \phi\rangle) = \langle\{r\}, \phi, \phi\rangle$ . It is easily seen that  $\eta$  is a  $NC\alpha$ -continuous function but not  $NC\alpha^*$ -continuous, since  $\langle\{p, q, r\}, \phi, \phi\rangle$  is  $NC\alpha$ -OS in  $\mathbb{X}$ , but  $\eta^{-1}(\langle\{p, q, r\}, \phi, \phi\rangle) = \langle\{p, s\}, \phi, \phi\rangle$  is not  $NC\alpha$ -OS in  $\mathbb{X}$ .

**Example 3.17:** Let  $\mathbb{X} = \{p, q, r\}$ . Then  $\Gamma = \{\phi_N, \mathbb{X}_N\} \cup \{\langle\{p\}, \phi, \phi\rangle, \langle\{q\}, \phi, \phi\rangle, \langle\{p, q\}, \phi, \phi\rangle\}$  be a  $NCT$  on  $\mathbb{X}$ . Define a function  $\eta: (\mathbb{X}, \Gamma) \rightarrow (\mathbb{X}, \Gamma)$  by  $\eta(\langle\{p\}, \phi, \phi\rangle) = \langle\{p\}, \phi, \phi\rangle, \eta(\langle\{q\}, \phi, \phi\rangle) = \eta(\langle\{r\}, \phi, \phi\rangle) = \langle\{q\}, \phi, \phi\rangle$ . It is easily seen that  $\eta$  is a  $NCS\alpha$ -continuous function but not  $NC\alpha^*$ -continuous, since  $\langle\{q\}, \phi, \phi\rangle$  is  $NC\alpha$ -OS in  $\mathbb{X}$ , but  $\eta^{-1}(\langle\{q\}, \phi, \phi\rangle) = \langle\{q, r\}, \phi, \phi\rangle$  is not  $NC\alpha$ -OS in  $\mathbb{X}$ .

**Definition 3.18:** A function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  is called  $\mathcal{M}$ -function iff  $\eta^{-1}(NCint(NCcl(\mathcal{D}))) \subseteq NCint(NCcl(\eta^{-1}(\mathcal{D})))$ , for every  $NC\alpha$ -OS  $\mathcal{D}$  from  $\mathbb{Y}$ .

**Theorem 3.19:** If  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  is a  $NC\alpha$ -continuous function and  $\mathcal{M}$ -function, then  $\eta$  is a  $NC\alpha^*$ -continuous.

**Proof:** Let  $\mathcal{C}$  be any  $NC\alpha$ -OS of  $\mathbb{Y}$ , then we have at least a  $NC$ -OS  $\mathcal{D}$  from  $\mathbb{Y}$  where  $\mathcal{D} \subseteq \mathcal{C} \subseteq NCint(NCcl(\mathcal{D}))$ . Since  $\eta$  is  $\mathcal{M}$ -function, we have  $\eta^{-1}(\mathcal{D}) \subseteq \eta^{-1}(\mathcal{C}) \subseteq \eta^{-1}(NCint(NCcl(\mathcal{D}))) \subseteq NCint(NCcl(\eta^{-1}(\mathcal{D})))$ . By proposition 2.3, we have  $\eta^{-1}(\mathcal{C})$  is a  $NC\alpha$ -OS. Hence,  $\eta$  is a  $NC\alpha^*$ -continuous. ■

**Remark 3.20:** The concepts of  $NC\alpha^*$ -continuity and  $NCS\alpha^*$ -continuity are independent as the following examples show.

**Example 3.21:** Assume  $\mathbb{X} = \{p, q, r, s\}$  and  $\mathbb{Y} = \{u, v, w\}$ .

Then  $\Gamma_1 = \{\phi_N, \mathbb{X}_N\} \cup \{\langle\{p\}, \phi, \phi\rangle, \langle\{q\}, \phi, \phi\rangle, \langle\{p, q\}, \phi, \phi\rangle, \langle\{p, q, r\}, \phi, \phi\rangle\}$  and  $\Gamma_2 = \{\phi_N, \mathbb{Y}_N\} \cup \{\langle\{u\}, \phi, \phi\rangle, \langle\{v\}, \phi, \phi\rangle, \langle\{u, v\}, \phi, \phi\rangle\}$  be  $NCT$ s on  $\mathbb{X}$  and  $\mathbb{Y}$ , correspondingly. Define the function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  via  $\eta(\langle\{p\}, \phi, \phi\rangle) = \eta(\langle\{s\}, \phi, \phi\rangle) = \langle\{v\}, \phi, \phi\rangle, \eta(\langle\{r\}, \phi, \phi\rangle) = \langle\{w\}, \phi, \phi\rangle$  and  $\eta(\langle\{q\}, \phi, \phi\rangle) = \langle\{u\}, \phi, \phi\rangle$ . It is easily seen that  $\eta$  is a  $NCS\alpha^*$ -continuous function but not  $NC\alpha^*$ -continuous, since  $\langle\{v\}, \phi, \phi\rangle$  is  $NC\alpha$ -OS in  $\mathbb{Y}$  but  $\eta^{-1}(\langle\{v\}, \phi, \phi\rangle) = \langle\{p, s\}, \phi, \phi\rangle$  is not  $NC\alpha$ -OS in  $\mathbb{X}$ .

**Example 3.22:** Suppose  $\mathbb{X} = \{p, q, r, s\}$ .

Then  $\Gamma = \{\phi_N, \mathbb{X}_N\} \cup \{\langle\{p\}, \phi, \phi\rangle, \langle\{q\}, \phi, \phi\rangle, \langle\{p, q\}, \phi, \phi\rangle, \langle\{p, q, r\}, \phi, \phi\rangle\}$  be a  $NCT$  on  $\mathbb{X}$ . Define the function  $\eta: (\mathbb{X}, \Gamma) \rightarrow (\mathbb{X}, \Gamma)$  via  $\eta(\langle\{p\}, \phi, \phi\rangle) = \eta(\langle\{q\}, \phi, \phi\rangle) = \langle\{q\}, \phi, \phi\rangle, \eta(\langle\{r\}, \phi, \phi\rangle) = \langle\{s\}, \phi, \phi\rangle, \eta(\langle\{s\}, \phi, \phi\rangle) = \langle\{r\}, \phi, \phi\rangle$ . It is easily seen that  $\eta$  is a  $NC\alpha^*$ -continuous function but not  $NCS\alpha^*$ -continuous, since  $\langle\{p, r\}, \phi, \phi\rangle$  is  $NCS\alpha$ -OS in  $\mathbb{X}$ , but  $\eta^{-1}(\langle\{p, r\}, \phi, \phi\rangle) = \langle\{s\}, \phi, \phi\rangle$  is not  $NCS\alpha$ -OS in  $\mathbb{X}$ .

**Theorem 3.23:** If a function  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  is  $NC\alpha^*$ -continuous,  $NC$ -open and bijective, then it is  $NCS\alpha^*$ -continuous.

**Proof:** Let  $\eta: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  be a  $NC\alpha^*$ -continuous,  $NC$ -open and bijective. Let  $\mathcal{D}$  be a  $NCS\alpha$ -OS in  $\mathbb{Y}$ . Then we have at least a  $NC\alpha$ -OS say  $\mathcal{P}$  where  $\mathcal{P} \subseteq \mathcal{D} \subseteq NCcl(\mathcal{P})$ . Therefore  $\eta^{-1}(\mathcal{P}) \subseteq \eta^{-1}(\mathcal{D}) \subseteq \eta^{-1}(NCcl(\mathcal{P})) \subseteq NCcl(\eta^{-1}(\mathcal{P}))$  (since  $\eta$  is a  $NC$ -open), but  $\eta^{-1}(\mathcal{P}) \in NC\alpha O(\mathbb{X})$  (since  $\eta$  is a  $NC\alpha^*$ -continuous). Hence  $\eta^{-1}(\mathcal{P}) \subseteq \eta^{-1}(\mathcal{D}) \subseteq NCcl(\eta^{-1}(\mathcal{P}))$ . Thus,  $\eta^{-1}(\mathcal{D}) \in NCS\alpha O(\mathbb{X})$ . Therefore,  $\eta$  is a  $NCS\alpha^*$ -continuous. ■

**Remark 3.24:** Let  $\eta_1: (\mathbb{X}, \Gamma_1) \rightarrow (\mathbb{Y}, \Gamma_2)$  and  $\eta_2: (\mathbb{Y}, \Gamma_2) \rightarrow (\mathbb{Z}, \Gamma_3)$  be two functions, then:

- (i) If  $\eta_1$  and  $\eta_2$  are  $NC \alpha$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  need not to be a  $NC\alpha$ -continuous.
- (ii) If  $\eta_1$  and  $\eta_2$  are  $NCS \alpha$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  need not to be a  $NCS\alpha$ -continuous.

**Theorem 3.25:** Let  $\eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Y}, I_2)$  and  $\eta_2: (\mathbb{Y}, I_2) \rightarrow (\mathbb{Z}, I_3)$  be two functions, then:

- (i) If  $\eta_1$  is  $NC \alpha$ -continuous and  $\eta_2$  is  $NC$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha$ -continuous.
- (ii) If  $\eta_1$  is  $NC \alpha^*$ -continuous and  $\eta_2$  is  $NC \alpha$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha$ -continuous.
- (iii) If  $\eta_1$  and  $\eta_2$  are  $NC\alpha^*$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha^*$ -continuous.
- (iv) If  $\eta_1$  and  $\eta_2$  are  $NCS\alpha^*$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NCS\alpha^*$ -continuous.
- (v) If  $\eta_1$  and  $\eta_2$  are  $NC\alpha^{**}$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha^{**}$ -continuous.
- (vi) If  $\eta_1$  and  $\eta_2$  are  $NCS\alpha^{**}$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NCS\alpha^{**}$ -continuous.
- (vii) If  $\eta_1$  is  $NC \alpha^{**}$ -continuous and  $\eta_2$  is  $NC \alpha^*$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha^{**}$ -continuous.
- (viii) If  $\eta_1$  is  $NC \alpha^{**}$ -continuous and  $\eta_2$  is  $NC \alpha$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC$ -continuous.
- (ix) If  $\eta_1$  is  $NC \alpha$ -continuous and  $\eta_2$  is  $NC \alpha^{**}$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha^*$ -continuous.
- (x) If  $\eta_1$  is  $NC$ -continuous and  $\eta_2$  is  $NC \alpha^{**}$ -continuous, then  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha^{**}$ -continuous.

**Proof:**

- (i) Assume  $\mathcal{F}$  considers a  $NC$ -OS from  $\mathbb{Z}$ . Since  $\eta_2$  is a  $NC$ -continuous,  $\eta_2^{-1}(\mathcal{F})$  is a  $NC\alpha$ -OS in  $\mathbb{Y}$ . Since  $\eta_1$  is a  $NC \alpha$ -continuous,  $\eta_1^{-1}(\eta_2^{-1}(\mathcal{F})) = (\eta_2 \circ \eta_1)^{-1}(\mathcal{F})$  is a  $NC \alpha$ -OS in  $\mathbb{X}$ . Thus,  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  exists a  $NC\alpha$ -continuous.
- (ii) Let  $\mathcal{F}$  be a  $NC$ -OS in  $\mathbb{Z}$ . Subsequently  $\eta_2$  stands a  $NC \alpha$ -continuous, and  $\eta_2^{-1}(\mathcal{F})$  stays a  $NC\alpha$ -OS from  $\mathbb{Y}$ . Since  $\eta_1$  is a  $NC\alpha^*$ -continuous,  $\eta_1^{-1}(\eta_2^{-1}(\mathcal{F})) = (\eta_2 \circ \eta_1)^{-1}(\mathcal{F})$  is a  $NC\alpha$ -OS in  $\mathbb{X}$ . Thus,  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha$ -continuous.
- (iii) Let  $\mathcal{F}$  be a  $NC\alpha$ -OS in  $\mathbb{Z}$ . Since  $\eta_2$  is a  $NC\alpha^*$ -continuous,  $\eta_2^{-1}(\mathcal{F})$  is a  $NC\alpha$ -OS in  $\mathbb{Y}$ . Since  $\eta_1$  is a  $NC\alpha^*$ -continuous,  $\eta_1^{-1}(\eta_2^{-1}(\mathcal{F})) = (\eta_2 \circ \eta_1)^{-1}(\mathcal{F})$  is a  $NC\alpha$ -OS in  $\mathbb{X}$ . Thus,  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha^*$ -continuous.
- (iv) Let  $\mathcal{F}$  be a  $NCS\alpha$ -OS in  $\mathbb{Z}$ . Since  $\eta_2$  is a  $NCS\alpha^*$ -continuous,  $\eta_2^{-1}(\mathcal{F})$  is a  $NCS\alpha$ -OS in  $\mathbb{Y}$ . Since  $\eta_1$  is a  $NCS\alpha^*$ -continuous,  $\eta_1^{-1}(\eta_2^{-1}(\mathcal{F})) = (\eta_2 \circ \eta_1)^{-1}(\mathcal{F})$  is a  $NCS\alpha$ -OS in  $\mathbb{X}$ . Thus,  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NCS\alpha^*$ -continuous.
- (v) Let  $\mathcal{F}$  be a  $NC\alpha$ -OS in  $\mathbb{Z}$ . Since  $\eta_2$  is a  $NC\alpha^{**}$ -continuous,  $\eta_2^{-1}(\mathcal{F})$  is a  $NC$ -OS in  $\mathbb{Y}$ . Since any  $NC$ -OS is a  $NC\alpha$ -OS,  $\eta_2^{-1}(\mathcal{F})$  is a  $NC\alpha$ -OS in  $\mathbb{Y}$ . Since  $\eta_1$  is a  $NC\alpha^{**}$ -continuous,  $\eta_1^{-1}(\eta_2^{-1}(\mathcal{F})) = (\eta_2 \circ \eta_1)^{-1}(\mathcal{F})$  is a  $NC$ -OS in  $\mathbb{X}$ . Thus,  $\eta_2 \circ \eta_1: (\mathbb{X}, I_1) \rightarrow (\mathbb{Z}, I_3)$  is a  $NC\alpha^{**}$ -continuous. The proof is obvious for others. ■

**Remark 3.26:** The next figure describes the relationship between various classes of weakly  $NC$ -continuous functions:






Fig. 1

4. Conclusion

We shall use the concepts of  $NC\alpha$ -OS and  $NCS\alpha$ -CS to define several new types of weakly  $NC$ -continuity such as;  $NC\alpha^*$ -continuous,  $NC\alpha^{**}$ -continuous,  $NCS\alpha$ -continuous,  $NCS\alpha^*$ -continuous and  $NCS\alpha^{**}$ -continuous functions. The neutrosophic crisp  $\alpha$ -open and neutrosophic crisp semi- $\alpha$ -open sets can be used to derive some new types of weakly  $NC$ -open ( $NC$ -closed) functions.

References

1. A. A. Salama, F. Smarandache and V. Kroumov, Neutrosophic crisp sets and neutrosophic crisp topological spaces. *Neutrosophic Sets and Systems*, 2(2014), 25-30.
2. R. K. Al-Hamido, Q. H. Imran, K. A. Alghurabi and T. Gharibah, On neutrosophic crisp semi- $\alpha$ -closed sets. *Neutrosophic Sets and Systems*, 21(2018), 28-35.
3. M. Abdel-Basset, R. Mohamed, A.E.N.H. Zaided, & F. Smarandache, A Hybrid Plithogenic Decision-Making Approach with Quality Function Deployment for Selecting Supply Chain Sustainability Metrics. *Symmetry*, 11(7), (2019), 903.
4. M. Abdel-Basset, N.A. Nabeeh, H.A. El-Ghareeb, & A. Aboelfetouh, Utilising neutrosophic theory to solve transition difficulties of IoT-based enterprises. *Enterprise Information Systems*, (2019), 1-21.

5. M. Abdel-Baset, V. Chang, & A. Gamal, Evaluation of the green supply chain management practices: A novel neutrosophic approach. *Computers in Industry*, 108 (2019), 210-220.
6. M. Abdel-Baset, M. Saleh, A. Gamal, & F. Smarandache, An approach of TOPSIS technique for developing supplier selection with group decision making under type-2 neutrosophic number. *Applied Soft Computing*, 77 (2019), 438-452.
7. M. Abdel-Baset, V. Chang, A. Gamal, & F. Smarandache, An integrated neutrosophic ANP and VIKOR method for achieving sustainable supplier selection: A case study in importing field. *Computers in Industry*, 106 (2019), 94-110.
8. M. Abdel-Baset, G. Manogaran, A. Gamal, & F. Smarandache, A group decision making framework based on neutrosophic TOPSIS approach for smart medical device selection. *Journal of medical systems*, 43(2), (2019), 38.
9. C. Maheswari and S. Chandrasekar, Neutrosophic gb-closed Sets and Neutrosophic gb-Continuity. *Neutrosophic Sets and Systems*, 29(2019), 89-100.
10. V. Banupriya and S. Chandrasekar, Neutrosophic  $\alpha$ gs Continuity and Neutrosophic  $\alpha$ gs Irresolute Maps. *Neutrosophic Sets and Systems*, 28(2019), 162-170.
11. R. Dhavaseelan, R. Narmada Devi, S. Jafari and Q. H. Imran, Neutrosophic  $\alpha^m$ -continuity. *Neutrosophic Sets and Systems*, 27(2019), 171-179.
12. A. A. Salama, Basic structure of some classes of neutrosophic crisp nearly open sets & possible application to GIS topology. *Neutrosophic Sets and Systems*, 7(2015), 18-22.
13. A. A. Salama, I. M. Hanafy, H. Elghawalby and M. S. Dabash, Some GIS Topological Concepts via Neutrosophic Crisp Set Theory. *New Trends in Neutrosophic Theory and Applications*, 2016.
14. W. Al-Omeri, Neutrosophic crisp sets via neutrosophic crisp topological spaces. *Neutrosophic Sets and Systems*, 13(2016), 96-104.
15. R. K. Al-Hamido, T. Gharibah, S. Jafari, F. Smarandache, On neutrosophic crisp topology via N-topology. *Neutrosophic Sets and Systems*, 23(2018), 96-109.
16. Q. H. Imran, R. Dhavaseelan, A. H. M. Al-Obaidi and Md. Hanif PAGE, On neutrosophic generalized alpha generalized continuity. *Neutrosophic Sets and Systems*, 35(2020), 511-521.
17. Md. Hanif PAGE and Q. H. Imran, Neutrosophic generalized homeomorphism. *Neutrosophic Sets and Systems*, 35(2020), 340-346.

Received: May 15, 2020. Accepted: Nov, 20, 2020.