

# First report of *Paratylenchus lepidus* Raski, 1975 associated with green tea (*Camellia sinensis* (L.) Kuntze) in Vietnam

Thi Mai Linh Le<sup>1,2</sup>, Huu Tien Nguyen<sup>1,2,3,\*</sup>, Thi Duyen Nguyen<sup>1,2,\*</sup> and Quang Phap Trinh<sup>1,2</sup>

<sup>1</sup>Institute of Ecology and Biological Resources, Vietnam Academy of Sciences and Technology, 18 Hoang Quoc Viet, Cau Giay, 100000, Hanoi, Vietnam.

<sup>2</sup>Graduate University of Science and Technology, Vietnam Academy of Sciences and Technology, 18 Hoang Quoc Viet, Cau Giay, 100000, Hanoi, Vietnam.

<sup>3</sup>Nematology Research Unit, Department of Biology, Ghent University, K.L. Ledeganckstraat 35, 9000, Ghent, Belgium.

\*E-mails: tien.quelampb@gmail.com; tqphap@gmail.com

This paper was edited by Zafar Ahmad Handoo.

Received for publication August 3, 2020.

The genus *Paratylenchus* (Ciobanu et al., 2003) is commonly known as pin nematodes that are ectoparasites and can be frequently found at high density in perennial plants, hop gardens, orchards, or forest trees (Ghaderi et al., 2016; Ghaderi, 2019). Although sometimes plants infected by *Paratylenchus* species show no specific symptoms, large populations of *Paratylenchus* spp. affect the absorption capacity of roots and the general physiology of plants (Ghaderi, 2019). According to Talavera and Navas (2002), *Paratylenchus* is only considered damaging nematodes at a density higher than 500 nematodes per 100 cm<sup>3</sup> of soil. However, several studies reported that the population of *Paratylenchus* can increase from a low number to damaging levels within a short time (Faulkner, 1964; Brzeski et al., 1975). The identification of *Paratylenchus* species was mostly based on morphological characterizations (Ghaderi et al., 2016), but morphological variation can be an obstacle to

## Abstract

The pin nematodes, *Paratylenchus* spp., are relatively small nematodes that can feed on a wide range of host plants. The morphological identification of this nematode is greatly hampered by their small size and variable characters. This study provides the first report of *Paratylenchus lepidus* from Vietnam with a combination of morphological and molecular characterizations. The 28S rDNA phylogenetic tree of the genus and the first *COI* mtDNA barcode of this species are also provided.

## Keywords

28S rDNA, *COI* mtDNA, DNA barcode, Plant-parasitic nematodes, Taxonomy.

the identification process, which make the molecular approach to become more popular in recent studies of pin nematodes. In Vietnam, 16 *Paratylenchus* species have been reported without molecular data, including *Paratylenchus aculentus*, *P. arcuatus*, *P. colbrani*, *P. corbetti*, *P. costatus*, *P. dianthus*, *P. discocephalus*, *P. elachistus*, *P. epicotylus*, *P. laocaiensis*, *P. minusculus*, *P. nawadus*, *P. pandatus*, *P. perlatus*, *P. serricaudatus*, and *P. similis* (Nguyen and Nguyen, 2000; Nguyen et al., 2004). In this study, we provide the first report of *Paratylenchus lepidus* (Raski, 1975) in Vietnam using the combination of morphological and molecular characterizations.

## Material and methods

Soil and root samples were collected from the rhizosphere of green tea (*Camellia sinensis* (L.) Kuntze) in Vietnam. Nematodes were extracted using

the modified Baermann tray method (Whitehead and Hemming, 1965). After that, they were fixed and prepared to make permanent slides following Nguyen et al. (2019a). For morphological characterization, measurements and pictures were taken using Carl Zeiss Axio Lab. A1 light microscope equipped with a Zeiss AxioCam ERc5s digital camera. For molecular characterization, the D2-D3 region of 28S rDNA and *COI* mtDNA gene were amplified using D2A/D3B (5'-ACAAGTACCGTGGGGAAAGTTG-3'/5'-TCGG AAGGAACCAGCTACTA-3') (Subbotin et al., 2006) and JB3/JB4 (5'-TTTTTTGGGCATCCTGAGGTTTAT-3'/5'-TAAAGAAAGAACATAATGAAAATG-3') (Nguyen et al., 2019b) primers. Forward and reverse sequences were assembled using Geneious R11

(www.geneious.com). The best fit model was chosen using Mega 7 and phylogenetic analysis was done following Nguyen et al. (2019c).

## Results and discussion

### Measurements

$n=20$  (♀♀):  $L=340\pm 20$  (307-371)  $\mu\text{m}$ ,  $a=25\pm 1$  (22-27),  $b=4.1\pm 0.3$  (3.7-4.6),  $c=11.5\pm 1.4$  (9.8-13.8),  $c'=3.5\pm 0.4$  (3.0-4.1),  $V\%=82\pm 1$  (81-84), Lip height= $2.7\pm 0.5$  (1.9-3.6)  $\mu\text{m}$ , Lip width= $4.9\pm 0.5$  (4.2-5.9)  $\mu\text{m}$ , Stylet= $25\pm 1$  (24-27)  $\mu\text{m}$ , Median bulb length= $15.2\pm 2$  (12.8-18.0)  $\mu\text{m}$ , Median bulb width= $6.9\pm 0.6$  (5.9-7.7)  $\mu\text{m}$ , SE pore= $75\pm 4$  (67-81)




Figure 1: Female of *P. lepidus* from green tea in Vietnam. A: Entire body; B: Anterior end region; C to F: Variation of the tail region.

$\mu\text{m}$ , Pharynx= $84 \pm 3$  (79-91)  $\mu\text{m}$ , Body width= $13.8 \pm 0.4$  (13.3-14.5)  $\mu\text{m}$ , Vulval body diam.= $12.2 \pm 0.4$  (11.3-12.6)  $\mu\text{m}$ , Anal body diam.= $8.5 \pm 0.4$  (7.9-9.5)  $\mu\text{m}$ , Tail length= $30 \pm 3$  (26-35)  $\mu\text{m}$ .

### Morphological characterization

The female of Vietnamese population of *Paratylenchus lepidus* is characterized by having a slender body, curved ventrally; lateral field with four incisures;

lip region weakly sclerotized, continuous to body contour; median bulb elongate with a distinct valve; isthmus slender, surrounded by nerve ring; basal bulb pyriform; secretory-excretory pore located at level of basal bulb to pharyngo-intestinal junction; hemizonid located just anterior to secretory-excretory pore; gonad monodelphic, post uterine sac absent; vulval lips not protruding but having prominent advulval flap; tail curved ventrally with a finely rounded to bluntly pointed terminus (Fig. 1). Male was not found.




Figure 2: Bayesian inference phylogenetic tree generated from 28S rDNA sequences under HKY + G model (BIC = 2089.633, lnL = -740.031, G = 0.23, R = 3.22, f(A) = 0.226, f(T) = 0.182, f(C) = 0.249, f(G) = 0.343). Bayesian posterior probabilities (in percentage) are given next to each node. Sequences of *P. lepidus* from Vietnam are in bold font.

## Molecular characterization

The 28S rDNA sequence of *P. lepidus* from Vietnam (742bp long, accession number: MT808205) was 99.7% similar (2bp difference) to the sequence of *P. lepidus* from GenBank (accession number: MK886692). The phylogenetic tree based on 28S rDNA sequences showed that the sequence of *P. lepidus* from Vietnam was placed together with the sequence of *P. lepidus* from GenBank (100% PP) (Fig. 2). A *COI* mtDNA sequence of *P. lepidus* from Vietnam (418bp long) was also obtained and submitted to GenBank under the accession number MT828831.

## Remarks

Morphology of *P. lepidus* from green tea in Vietnam is in agreement with the description of the type population (Raski, 1975) with small variations in measurements, however, these variations can be seen from the type population and other populations (Maria et al., 2019). In this study, molecular identification is in agreement with morphological identification to support the presence of *P. lepidus* in Vietnam. The first *COI* mtDNA sequence of *P. lepidus* is also provided to serve as a molecular barcode for molecular identification of *Paratylenchus* species in the future.

## Acknowledgments

This research was supported by a fund from the Institute of Ecology and Biological Resources (code: IEBR.TST.3-20).

## References

Brzeski, M. W., Zepp, A. L. and D'errico, F. P. 1975. Development stages of *Paratylenchus bukowinensis* Micoletzky. *Roczniki Nauk Rolniczych* Seria 5:145–51.

Ciobanu, M., Geraert, E. and Popovici, I. 2003. The genera *Paratylenchus* Micoletzky, 1922 and *Gracilacus* Raski, 1962 in Romania (Nematoda: Tylenchulidae). *Nematologia Mediterranea* 31:55–9.

Faulkner, L. R. 1964. Pathogenicity and population dynamics of *Paratylenchus hamatus* on *Mentha* spp. *Phytopathology* 54:344–8.

Ghaderi, R. 2019. The damage potential of pin nematodes, *Paratylenchus* Micoletzky, 1922 *sensu lato* spp.

(Nematoda: Tylenchulidae). *Journal of Crop Protection* 8:243–57.

Ghaderi, R., Geraert, E. and Karegar, A. 2016. The Tylenchulidae of the world: identification of the family Tylenchulidae (Nematoda: Tylenchida) Academia Press, Belgium.

Maria, M., Miao, W., Ye, W. and Zheng, J. 2019. Updated description of *Paratylenchus lepidus* Raski 1975 and *P. minor* Sharma, Sharma and Khan, 1986 by integrating molecular and ultra-structural observations. *Journal of Nematology* 51:e2019–56.

Nguyen, N. C. and Nguyen, V. T. 2000. Fauna of Vietnam, plant-parasitic nematodes Science and Technics Publishing House, Hanoi, Vietnam.

Nguyen, C. N., Baldwin, J. G. and Choi, Y. E. 2004. New records of *Paratylenchus* Micoletzky, 1922 (Nematoda: Paratylenchinae) from Vietnam with description of *Paratylenchus laocaiensis* sp. n. *Journal of Nematode Morphology and Systematics* 7:51–75.

Nguyen, H. T., Le, T. M. L., Nguyen, T. D., Liebanas, G., Nguyen, T. A. D. and Trinh, Q. P. 2019a. Description of *Geocenamus vietnamensis* sp. n. (Nematoda: Merliniidae) from Vietnam. *Journal of Nematology* 51:e2019–25.

Nguyen, H. T., Trinh, Q. P., Couvreur, M., Singh, P. R., Decraemer, W. and Bert, W. 2019b. Molecular and morphological characterisation of a new root-lesion nematode, *Pratylenchus horti* n. sp. (Tylenchomorpha: Pratylenchidae) from Ghent University Botanical Garden. *Nematology* 21:739–52.

Nguyen, H. T., Trinh, Q. P., Couvreur, M., Singh, P. R., Decraemer, W. and Bert, W. 2019c. Description of *Rotylenchus rhomboides* n. sp. and a Belgian population of *Rotylenchus buxophilus* (Tylenchomorpha: Hoplolaimidae). *Journal of Nematology* 51:1–20.

Raski, D. J. 1975. Revision of the genus *Paratylenchus* Micoletzky, 1922 and descriptions of new species. Part II of three parts. *Journal of Nematology* 3:274–95.

Subbotin, S. A., Sturhan, D., Chizhov, V. N., Vovlas, N. and Baldwin, J. G. 2006. Phylogenetic analysis of Tylenchida Thorne, 1949 as inferred from D2 and D3 expansion fragments of the 28S rRNA gene sequences. *Nematology* 8:455–74.

Talavera, M. and Navas, A. 2002. Incidence of plant-parasitic nematodes in natural and semi-natural mountain grassland the host status of some common grass species. *Nematology* 4:541–52.

Whitehead, A. G. and Hemming, J. R. 1965. A comparison of some quantitative methods of extracting small vermiform nematodes from soil. *Annals of Applied Biology* 55:25–38.