

June 1997

Light scattering from a periodically modulated two-dimensional electron gas with partially filled Landau levels

A. Brataas

Norwegian University of Science and Technology, Trondheim, Norway

C. Zhang

University of Wollongong, czhang@uow.edu.au

K. A. Chao

Norwegian University of Science and Technology, Trondheim, Norway

Follow this and additional works at: <https://ro.uow.edu.au/engpapers>



Part of the [Engineering Commons](#)

<https://ro.uow.edu.au/engpapers/250>

Recommended Citation

Brataas, A.; Zhang, C.; and Chao, K. A.: Light scattering from a periodically modulated two-dimensional electron gas with partially filled Landau levels 1997.

<https://ro.uow.edu.au/engpapers/250>

Light scattering from a periodically modulated two-dimensional electron gas with partially filled Landau levels

Arne Brataas

Department of Physics, Norwegian University of Science and Technology, N-7034 Trondheim, Norway

C. Zhang

Department of Physics, University of Wollongong, New South Wales 2522, Australia

K. A. Chao

*Department of Physics, Norwegian University of Science and Technology, N-7034 Trondheim, Norway
and Department of Theoretical Physics, Lund University, S-223 26 Lund, Sweden*

(Received 19 December 1996)

We study light scattering from a periodically modulated two-dimensional electron gas in a perpendicular magnetic field. If a subband is partially filled, the imaginary part of the dielectric function as a function of frequency contains additional discontinuities to the case of completely filled subbands. The positions of the discontinuities may be determined from the partial filling factor and the height of the discontinuity can be directly related to the modulation potential. The light-scattering cross section contains a peak which is absent for integer filling. [S0163-1829(97)07820-X]

Since the Weiss oscillation was first observed in the magnetoresistivity several years ago,¹ a considerable amount of work, both experimental and theoretical, has been carried out on the electronic and transport properties of a two-dimensional electron system under a periodic potential and a constant magnetic field.²⁻¹⁰ Most recent works include the dc transport in a strong antidot system^{11,12} and the observation of a quantum fractal-like energy spectrum.¹³ Weiss oscillation can be understood as a type of commensurability oscillation originated from the interplay of two different length scales of the system, the periodicity of the modulation potential a , and the radius of the cyclotron motion R_c . The dc resistivity has a set of minima whenever the condition $2R_c = (n - 1/4)a$ is satisfied, where n is any positive integer. Most of the current investigations have been limited to the case where the external field has zero frequency. In a recent paper by Stewart and Zhang,¹⁴ the dielectric response at finite frequency and wave numbers was calculated. Their result indicates that the modulation induced structure at finite frequency is much richer than that of the static case. Most noticeably, the electron-hole pair excitation contains a set of singularities at the excitation band edges.

In this Brief Report, we investigate the density response function within the random-phase approximation for a two-dimensional electron gas under a weak periodic modulation and a constant magnetic field. We shall pay special attention to the case where the Fermi level lies within a Landau band. In what follows, we shall show that this partially filled Landau band at the Fermi level has a profound effect on the dielectric response and in turn alters the light scattering cross section from such a system. Our main results are as follows. (i) Electron-hole pair excitation has a steplike (discontinuous) behavior around $\omega = n\omega_c$. This step can be determined analytically for the case of weak modulation. Thus the amplitude of the modulation potential, which so far has been rather difficult to measure experimentally, can now be deter-

mined. (ii) Also around $\omega = n\omega_c$, the real part of the dielectric function has a logarithm divergence. (iii) When the effects of (i) and (ii) are included in the density response function, a sharp peak can be observed in the light scattering cross section.

We consider a two-dimensional electron gas where a static magnetic field B is perpendicular to the plane. A weak periodic potential is applied in the x direction,

$$V(x) = V_0 \cos(Kx), \quad (1)$$

where $K = 2\pi/a$ and a is the period of the modulation. In the Landau gauge the single-particle wave functions are of the form $\psi_{nk}(x, y) \propto \exp(ik_y y) \phi_{nx_0}(x)$, where $x_0 = k_y l^2$ (magnetic length, $l = [\hbar/(m^* \omega_c)]^{1/2}$) is the center coordinate and $\phi_{nx_0}(x)$ is the eigenfunction of the one-dimensional Hamiltonian,

$$H = -\frac{\hbar^2}{2m^*} \frac{d^2}{dx^2} + \frac{1}{2} m^* \omega_c^2 (x - x_0)^2 + V_0 \cos(Kx), \quad (2)$$

where m^* is the effective mass and $\omega_c = eB/(m^*c)$ is the cyclotron frequency. We assume that the modulation potential is a weak perturbation, $V_0 \ll E_F$, where E_F is the Fermi energy. The energy spectrum to linear order in the modulation potential is³

$$E_n(x_0) = \left(n + \frac{1}{2} \right) \hbar \omega_c + U_n \cos(Kx_0), \quad (3)$$

where $U_n = V_0 \exp(-\mathcal{H}/2) L_n(\mathcal{H})$, $\mathcal{H} = (Kl)^2/2$, and $L_n(\mathcal{H})$ is a Laguerre polynomial.

In order to simplify the discussion of the discontinuity in the imaginary part of the dielectric function as a function of the frequency, let us consider only a vertical transition. The energy difference for the transition from state m', x_0 to state $m' + m, x_0$ is then ($m > 0$)




FIG. 1. Single-particle energies (lines) around the Fermi energy (broken line) as a function of the center coordinate, Kx_0 , when the last subband is half-filled. Electron-hole pair excitations with energies $\omega < \omega_c$, $\omega = \omega_c$, and $\omega > \omega_c$ are shown.

$$E_{m+m'}(x_0) - E_{m'}(x_0) = m\hbar\omega_c + (U_{m+m'} - U_{m'})\cos(Kx_0). \quad (4)$$

For sufficiently small \mathcal{H} we may approximate $L_n(\mathcal{H}) = 1 - n\mathcal{H}$, so that $U_{m+m'} - U_{m'} = -V_0 m\mathcal{H}\exp(-\mathcal{H}/2)$. We also have $U_n(\mathcal{H}) > 0$ for small n and \mathcal{H} . This means that the energy difference between the states $m+m', x_0$ and m', x_0 is less than $m\hbar\omega_c$ for $0 < Kx_0 < \pi/2$ and $3\pi/2 < Kx_0 < 2\pi$. On the other hand, the energy difference is larger than $m\hbar\omega_c$ for $\pi/2 < Kx_0 < 3\pi/2$. Let us now consider a situation where the last Landau band (with index n_F) is half-filled. At $\omega \sim \omega_c$, two transition processes ($n_F - 1 \rightarrow n_F$ and $n_F \rightarrow n_F + 1$) contribute to the electron-hole pair excitation. However, the transition ($n_F - 1 \rightarrow n_F$) only contributes for $\omega < \omega_c$ while for $\omega > \omega_c$ only the transition ($n_F \rightarrow n_F + 1$) contributes (see Fig. 1). At $\omega = \omega_c$, the contributions from these two distinct transitions are different and thus a discontinuity occurs at $\omega = \omega_c$. One can immediately generalize this conclusion for excitations around $\omega \sim m\omega_c$, where a similar discontinuity occurs due to the different contributions from the ($n_F - m \rightarrow n_F$) transition and the ($n_F \rightarrow n_F + m$) transition. Below we shall derive an analytical expression for this discontinuity for a strictly vertical transition.

The case of half-filling is a quite special situation but actually exhibits the general discontinuous behavior of the pair excitation in modulated systems. If the last subband is less than half-filled, the discontinuity in the imaginary part of the dielectric function will appear for frequencies larger than $m\omega_c$. If the last subband is more than half-filled, the discontinuity in the imaginary part of the dielectric function will appear for frequencies lower than $m\omega_c$. In this way the imaginary part of the dielectric function also provides direct information about the filling of the last subband. For integer fillings, such discontinuity will disappear. For noninteger fillings but nonvertical excitations ($q_y \neq 0$), such discontinuity will still be present.

The dielectric function within the random-phase approximation is

$$\begin{aligned} \epsilon(q_x, \omega) = & 1 - 2\pi r_s \frac{k_F}{q_x} \hbar\omega_c \\ & \times \sum_{n \neq n'} \sum_{x_0} C_{nn'} \frac{f_{n,x_0} - f_{n',x_0}}{E_{n,x_0} - E_{n',x_0} + \hbar(\omega + i\delta)}, \end{aligned} \quad (5)$$

where r_s is the plasma parameters and f_{n,x_0} is the Fermi occupation function. The coefficient $C_{nn'}$ ($n > n'$) is given as

$$C_{nn'} = \frac{n!}{n'} X^{n-n'} [L_{n'}^{n-n'}(X)]^2 \exp(-X) \quad (6)$$

with $X = (q_x l)^2/2$ and $L_n^m(X)$ is an associated Laguerre polynomial.

For positive frequencies the imaginary part of the dielectric function is¹⁴

$$\begin{aligned} \text{Im}[\epsilon(q_x, \omega)] = & -2\pi^2 r_s (k_F/q_x) \hbar\omega_c \\ & \times \sum_{m=1}^{\infty} \sum_{m'=0}^{n_F} C_{m+m',m'} (f_{m+m',x_j} - f_{m',x_j}) \\ & \times Q^{mm'}, \end{aligned} \quad (7)$$

where the function

$$Q^{mm'} = \frac{\theta(|U_{m+m'} - U_{m'}| - |m\hbar\omega_c - \hbar\omega|)}{\sqrt{(U_{m+m'} - U_{m'})^2 - (m\hbar\omega_c - \hbar\omega)^2}} \quad (8)$$

gives the square-root singularities. In Eq. (7) we also sum over the simple roots x_j given by

$$\cos Kx_j = -\frac{m\hbar\omega_c - \hbar\omega}{U_{m+m'} - U_{m'}}. \quad (9)$$

For small $(m\hbar\omega_c - \hbar\omega)/(U_{m+m'} - U_{m'})$ we see that in the interval $0 < Kx_0 < 2\pi$ the solutions of Eq. (9) are $Kx_0 = \pi/2 + (m\hbar\omega_c - \hbar\omega)/(U_{m+m'} - U_{m'})$ and $Kx_1 = 3\pi/2 - (m\hbar\omega_c - \hbar\omega)/(U_{m+m'} - U_{m'})$. Let us consider a system where the filling factor is $\nu = 5.5$ and study the excitations around $\omega = 4\omega_c$. We introduce a small energy shift $\delta = \omega - 4\omega_c$ and find from Eqs. (7) and (9) that for $\delta = 0^-$

$$\begin{aligned} \text{Im}[\epsilon(q_x, \delta = 0^-)] = & 4\pi^2 r_s (k_F/q_x) \hbar\omega_c \left[\frac{C_{5,1}}{|U_5 - U_1|} \right. \\ & \left. + \frac{C_{6,2}}{|U_6 - U_2|} + \frac{C_{7,3}}{|U_7 - U_3|} + \frac{C_{8,4}}{|U_8 - U_4|} \right]. \end{aligned} \quad (10)$$




FIG. 2. Imaginary part of the dielectric function, $\text{Im}[\epsilon(q_x, \omega)]$, as a function of ω/ω_c for $\nu=5.5$, $V_0=1$ meV.

On the contrary, for $\delta=0^+$ we find

$$\begin{aligned} \text{Im}[\epsilon(q_x, \delta=0^+)] = & 4\pi^2 r_s (k_F/q_x) \hbar \omega_c \left[\frac{C_{6,2}}{|U_6 - U_2|} \right. \\ & \left. + \frac{C_{7,3}}{|U_7 - U_3|} + \frac{C_{8,4}}{|U_8 - U_4|} + \frac{C_{9,5}}{|U_9 - U_5|} \right]. \end{aligned} \quad (11)$$

Therefore,

$$\begin{aligned} \Delta \text{Im}[\epsilon(q_x)] = & \text{Im}[\epsilon(q_x, \delta=0^+)] - \text{Im}[\epsilon(q_x, \delta=0^-)] \\ = & 4\pi^2 r_s (k_F/q) \hbar \omega_c \left[\frac{C_{5,1}}{|U_5 - U_1|} - \frac{C_{9,5}}{|U_9 - U_5|} \right]. \end{aligned} \quad (12)$$

Similarly the discontinuity around $\omega=3\omega_c$ is given as $C_{5,2}/|U_5 - U_2| - C_{8,5}/|U_8 - U_5|$, etc. Since the first-order potential element U_n is linearly proportional to the modulation potential, we see that apart from numerical constants the discontinuity in the imaginary part of the dielectric function is determined directly by the strength of the modulation potential.

Note also that the imaginary part of the dielectric function should contain i singularities around each frequency band even for partially filled subbands as long as $q_y=0$. For non-zero q_y , in principle one may have $i+1$ singularities as stated in Ref. 14, where i is an integer and counts the resonance frequency.

For our numerical evaluation of the dielectric function we have employed the following parameters for a typical modulated GaAs/Al_xGa_{1-x}As heterostructure: $\kappa=13$, $r_s=0.73$, $E_F=10$ meV, and $m^*=0.067m_e$, giving a filling factor $\nu=5.5$. The amplitude and period of the modulation potential are $V_0=1$ meV and $a=300$ nm, respectively. The calculation was done at zero temperature with $q_x=0.2k_F$, $q_y=0$. The numerical parameters are thus the same as used in Ref. 14 except that we have a strictly vertical transition $q_y=0$ instead of the small $q_y=1 \times 10^6 \text{ m}^{-1}$.

We show in Fig. 2 the imaginary part of the dielectric function around $\omega=4\omega_c$. The discontinuity around

$\omega=4\omega_c$ is clearly resolved together with the four singularities on each side of the frequency band.

If the exchange and correlation effects are neglected, the imaginary part of the dielectric function is proportional to the cross section of spin-density excitations, which may be measured in Raman scattering when the polarization of the incoming and scattered light are perpendicular.¹⁵ Within this approximation the discontinuity step in the imaginary part of the dielectric function is therefore directly measurable.

The real part of the dielectric function is

$$\begin{aligned} \text{Re}[\epsilon(q_x, \omega)] = & 1 - 4\pi^2 r_s \frac{k_F}{q_x} \hbar \omega_c \\ & \times \sum_{m=1}^{\infty} \sum_{m'=0}^{n_F} C_{m+m', m'} (I_1 - I_2 + I_3 - I_4), \end{aligned} \quad (13)$$

where

$$I_1 = I(m\hbar\omega_c + \hbar\omega, U_{m+m'}, E_F - E_{m+m'}, U_{m+m'}), \quad (14a)$$

$$I_2 = I(m\hbar\omega_c + \hbar\omega, U_{m'}, E_F - E_{m'}, U_{m'}), \quad (14b)$$

$$I_3 = I(m\hbar\omega_c - \hbar\omega, U_{m+m'}, E_F - E_{m+m'}, U_{m+m'}), \quad (14c)$$

$$I_4 = I(m\hbar\omega_c - \hbar\omega, U_{m'}, E_F - E_{m'}, U_{m'}) \quad (14d)$$

are given by the Cauchy principal value integral

$$I(a, b, c, d) \equiv \frac{1}{2\pi} \int_0^{2\pi} d\phi \theta(c - d\cos\phi) P \frac{1}{a + b\cos\phi}. \quad (15)$$

The integral may be found by a complex contour integration. For $(c/d)^2 > 1$, which is the case for completely occupied or unoccupied subbands, the integral is simply

$$I = \begin{cases} \theta(c) 2\pi K, & (a/b)^2 > 1 \\ 0, & (a/b)^2 < 1, \end{cases} \quad (16)$$

where the factor K is

$$K = \frac{1}{2\pi a \sqrt{|1 - (b/a)^2|}}. \quad (17)$$

For the partially filled subbands where $(c/d)^2 < 1$ we find

$$I = K \times \begin{cases} 2\pi - 4\arctan\gamma, & (a/b)^2 > 1, \quad d > 0 \\ 4\arctan\gamma, & (a/b)^2 > 1, \quad d < 0 \\ -2\ln \left| \frac{1-\gamma}{1+\gamma} \right|, & (a/b)^2 < 1, \quad d > 0 \\ 2\ln \left| \frac{1-\gamma}{1+\gamma} \right|, & (a/b)^2 < 1, \quad d < 0, \end{cases} \quad (18)$$

where

$$\gamma = \frac{1-b/a}{\sqrt{|1 - (b/a)^2|}} \sqrt{\frac{1-c/d}{1+c/d}}. \quad (19)$$




FIG. 3. Real part of the dielectric function, $\text{Re}[\epsilon(q_x, \omega)]$, as a function of ω/ω_c for $\nu=5.5$, $V_0=1$ meV.

For the partially filled subband $m'=5$ (at filling $\nu=5.5$), the third argument of Eq. (15) is zero so that the real part of the dielectric function is logarithmic divergent around $\omega=4\omega_c$, as can be seen from Eq. (18). In addition, the real part of the dielectric function has four singularities for frequencies slightly larger than $4\omega_c$ and four singularities for frequencies slightly smaller than $4\omega_c$. The real part of the dielectric function is shown in Fig. 3 around $\omega=4\omega_c$.

In a far-infrared absorption or a Raman scattering experiment where the polarizations of the incoming and scattered photon are parallel, the charge-density excitations are measured. That is the scattering cross section which is proportional to $-\text{Im}[1/\epsilon(q, \omega)]$. This function has δ peaks when $\text{Re}[\epsilon(q, \omega)]=0$. For an unmodulated system, the only peaks in the scattering cross section are those due to the magneto-plasmon excitation. For a modulated system, certain spectral




FIG. 4. $-\text{Im}[1/\epsilon(q_x, \omega)]$ as a function of ω/ω_c for $\nu=5.5$, $V_0=1$ meV.

weight is shifted back to the energy corresponding to the electron-hole pair excitation. The peaks due to electron-hole pair excitation are rather sharp but finite even when disorders are non-negligible [14]. Furthermore, we found that the cross section has an additional sharp peak at $\omega=n\omega_c$ due to a logarithmic singularity in the real part of the dielectric function. This is depicted in Fig. 4.

In conclusion we have shown that a partial filling of the last Landau band may be detected in optical spectroscopy either in the spin-density excitation spectra or in the charge-density excitation spectra. The spectra will provide information about the modulation potential and the magnitude of the partial filling.

We would like to thank R.R. Gerhardtts and S. M. Stewart for interesting discussions.

¹D. Weiss, K. v. Klitzing, K. Ploog, and G. Weinmann, *Europhys. Lett.* **8**, 179 (1989).

²R. W. Winkler, J. P. Kotthaus, and K. Ploog, *Phys. Rev. Lett.* **62**, 1177 (1989).

³R. R. Gerhardtts, D. Weiss, and K. v. Klitzing, *Phys. Rev. Lett.* **62**, 1173 (1989).

⁴D. Weiss *et al.*, *Phys. Rev. B* **39**, 13 020 (1989).

⁵D. Pfannkuche and R. R. Gerhardtts, *Phys. Rev. B* **46**, 12 606 (1992).

⁶R. R. Gerhardtts and C. Zhang, *Phys. Rev. Lett.* **64**, 1473 (1990).

⁷C. Zhang and R. R. Gerhardtts, *Phys. Rev. B* **41**, 12 850 (1990).

⁸H. L. Cui, V. Fessatidis, and N. J. M. Horing, *Phys. Rev. Lett.* **63**,

2598 (1989).

⁹C. Zhang, *Phys. Rev. Lett.* **65**, 2207 (1990).

¹⁰A. Manolescu and R. R. Gerhardtts, *Phys. Rev. B* **51**, 1703 (1995).

¹¹D. Weiss *et al.*, *Phys. Rev. Lett.* **70**, 4118 (1993).

¹²P. Rotter, M. Suhrke, and U. Rössler, *Phys. Rev. B* **54**, 4452 (1996).

¹³T. Schlösser, K. Ensslin, J. P. Kotthaus, and M. Holland, *Europhys. Lett.* **33**, 683 (1996).

¹⁴S. M. Stewart and C. Zhang, *Phys. Rev. B* **52**, R17 036 (1995).

¹⁵M. V. Klein, in *Light Scattering in Solids I*, edited by M. Cardona, Topics in Applied Physics Vol. 8 (Springer, Berlin, 1984).