

University of Wollongong
Research Online

Faculty of Informatics - Papers (Archive)

Faculty of Engineering and Information
Sciences

June 1998

Group Divisible Designs, GBRSDS And Generalized Weighing Matrices

D. G. Sarvate

College of Charleston, Charleston, USA

Jennifer Seberry

University of Wollongong, jennie@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/infopapers>

 Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Sarvate, D. G. and Seberry, Jennifer: Group Divisible Designs, GBRSDS And Generalized Weighing Matrices 1998.

<https://ro.uow.edu.au/infopapers/351>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Group Divisible Designs, GBRSDS And Generalized Weighing Matrices

Abstract

We give new constructions for regular group divisible designs, pairwise balanced designs, generalized Bhaskar Rao supplementary difference sets and generalized weighing matrices. In particular if p is a prime power and q divides $p - 1$ we show the following exist:

- (i) $GDD(2(p^2 + p + 1), 2(p^2 + p + 1), rp^2, 2p^2, \lambda_1 = p^2\lambda, \lambda_2 = (p^2 - p)r, m = p^2 + p + 1, n = 2), r = 1, 2$;
- (ii) $GDD(q(p + 1), q(p + 1), p(q - 1), p(q - 1), \lambda_1 = (q - 1)(q - 2), \lambda_2 = (p - 1)(q - 1)^2/q, m = q, n = p + 1)$;
- (iii) $PBD(21, 10; K), K = \{3, 6, 7\}$ and $PBD(78, 38; K), K = \{6, 9, 45\}$;
- (iv) $GW(vk, k^2; EA(k))$ whenever a (v, k, λ) -difference set exists and k is a prime power;
- (v) $PBIBD(vk^2, vk^2, k^2, k^2; \lambda_1 = 0, \lambda_2 = \lambda, \lambda_3 = k)$ whenever a (v, k, λ) -difference set exists and k is a prime power;
- (vi) we give a $GW(21; 9; Z_3)$.

The GDD obtained are not found in W.H. Clatworthy, "Tables of Two-Associate-Class, Partially Balanced Designs", NBS, US Department of Commerce, 1971.

Disciplines

Physical Sciences and Mathematics

Publication Details

This article was originally published as Sarvate, DG and Seberry, J, Group Divisible Designs, GBRSDS And Generalized Weighing Matrices, *Utilitas Mathematica*, 54, 1998, 157-174.

GROUP DIVISIBLE DESIGNS, GBRSDS AND GENERALIZED WEIGHING MATRICES

Dinesh G. Sarvate
Department of Mathematics
College of Charleston
Charleston, S.C. 29424
U.S.A.

and

Jennifer Seberry*
Department of Computer Science
University of Wollongong
Wollongong, NSW, 2500
Australia

June 11, 1998

Abstract

We give new constructions for regular group divisible designs, pairwise balanced designs, generalized Bhaskar Rao supplementary difference sets and generalized weighing matrices. In particular if p is a prime power and q divides $p - 1$ we show the following exist:

- (i) $GDD(2(p^2 + p + 1), 2(p^2 + p + 1), rp^2, 2p^2, \lambda_1 = p^2\lambda, \lambda_2 = (p^2 - p)r, m = p^2 + p + 1, n = 2), r = 1, 2$;
- (ii) $GDD(q(p + 1), q(p + 1), p(q - 1), p(q - 1), \lambda_1 = (q - 1)(q - 2), \lambda_2 = (p - 1)(q - 1)^2/q, m = q, n = p + 1)$;
- (iii) $PBD(21, 10; K), K = \{3, 6, 7\}$ and $PBD(78, 38; K), K = \{6, 9, 45\}$;
- (iv) $GW(vk, k^2; EA(k))$ whenever a (v, k, λ) -difference set exists and k is a prime power;
- (v) $PBIBD(vk^2, vk^2, k^2, k^2; \lambda_1 = 0, \lambda_2 = \lambda, \lambda_3 = k)$ whenever a (v, k, λ) -difference set exists and k is a prime power;
- (vi) we give a $GW(21; 9; Z_3)$.

The GDD obtained are not found in W.H. Clatworthy, *Tables of Two-Associate-Class, Partially Balanced Designs*, NBS, US Department of Commerce, 1971.

1 INTRODUCTION

In this paper we set out to explore the usefulness of Bhaskar Rao designs and generalized matrices in the construction of GDD and found them to be very rich indeed.

A design is a pair (X, B) where X is a finite set of elements and B is a collection of (not necessarily distinct) subsets B_i (called blocks) of X .

A *balanced incomplete block design*, $BIBD(v, b, r, k, \lambda)$, is an arrangement of v elements into b blocks such that:

- (i) each element appears in exactly r blocks;
- (ii) each block contains exactly $k (< v)$ elements; and
- (iii) each pair of distinct elements appear together in exactly λ blocks.

*Research supported by Telecom grant 7027 and ARC grant A48830241

As $r(k-1) = \lambda(v-1)$ and $vr = bk$ are well known necessary conditions for the existence of a $BIBD(v, b, r, k, \lambda)$ we denote this design by $BIBD(v, k, \lambda)$.

Let v and λ be positive integers and K a set of positive integers.

An arrangement of the elements of a set X into blocks is a *pairwise balanced design*, $PBD(v; K; \lambda)$, if:

- (i) X contains exactly v elements;
- (ii) if a block contains k elements then k belongs to K ;
- (iii) each pair of distinct elements appear together in exactly λ blocks.

A pairwise balanced design $PBD(v; \{k\}; \lambda)$, that is where $K = \{k\}$ consists of exactly one integer, is a $BIBD(v, k, \lambda)$. It is well known that a $PBD(v-1; \{k, k-1\}; \lambda)$ can be obtained from the $BIBD(v, b, r, k, \lambda)$.

For the definition of a partially balanced incomplete block design with m associate classes ($PBIBD(m)$) see Raghavarao [34] or Street and Street [53].

A *generalized Bhaskar Rao design*, W is defined as follows. Let W be a $v \times b$ matrix with entries from $G \cup \{0\}$ where $G = \{h_1 = e, h_2, \dots, h_g\}$ is a finite group of order g . W is then expressed as a sum $W = h_1 A_1 + \dots + h_g A_g$, where A_1, \dots, A_g are $v \times b$ $(0, 1)$ matrices such that the Hadamard product $A_i \star A_j = 0$ for any $i \neq j$.

Denote by W^+ the transpose of $h_1^{-1} A_1 + \dots + h_g^{-1} A_g$ and let $N = A_1 + \dots + A_g$. In this paper we are concerned with the special case where W , denoted by $GBRD(v, b, r, k, \lambda; G)$, satisfies

- (i) $WW^+ = rI + \frac{\lambda}{g}(h_1 + \dots + h_g)(J - I)$, and
- (ii) $NN^T = (r - \lambda)I + \lambda J$.

It can be seen that the second condition requires that N be the incidence matrix of a $BIBD(v, b, r, k, \lambda)$ and thus we can use the shorter notation $GBRD(v, k, \lambda; G)$ for a generalized Bhaskar Rao design. A $GBRD(v, k, \lambda; Z_2)$ is also referred to as a $BRD(v, k, \lambda)$.

A $GBRD(v, k, \lambda; G)$ with $v = b$ is a symmetric $GBRD$ or generalized weighing matrix, but a generalized weighing matrix, $W = GW(v, k; G)$ is also used for any square matrix satisfying $WW^+ = kI$ where $h_1 + \dots + h_g = 0$ is used (as in the g th roots of unity). If W has no 0 entries the $GBRD$ is also known as a generalized Hadamard matrix (GH).

A *group divisible design*, $GDD(v, b, r, k, \lambda_1, \lambda_2, m, n)$, on v points is a triple (X, S, A) where

- (i) X is a set (of points), where $|X| = v$,
- (ii) S is a class of non-empty subsets X (called groups), of size n , which partitions X , and $|S| = m$,
- (iii) A is a class of subsets of X (called blocks), each containing at least two points, and $|A| = b$,
- (iv) each pair of distinct points $\{x, y\}$ where x and y are from the same group is contained in precisely λ_1 blocks.
- (v) each pair of distinct points $\{x, y\}$ where x and y are not from the same group is contained in precisely λ_2 blocks.

In general, the number of elements in a group is denoted by n .

Bhaskar Rao designs with elements $0, \pm 1$ have been studied by a number of authors including Bhaskar Rao [3, 4], Seberry [42, 44], Singh [48], Sinha [49], Street [51], Street and Rodger [52] and Vyas [54]. Bhaskar Rao [3] used these designs to construct partially balanced designs and this was improved by Street and Rodger [52] and Seberry [44]. Another technique for studying partially balanced designs has involved looking at generalized orthogonal matrices which have elements from elementary abelian groups and the element 0. Matrices with group elements as entries have been studied by Berman [1, 2], Butson [5, 6], Delsarte and Goethals [13], Drake [16], Rajkundlia [35], Seberry [40, 41], Shrikhande [47] and Street [50].

Generalized Hadamard matrices has been studied by Street [50], Seberry [40, 41], Dawson [8], and de Launey [9, 10].

Bhaskar Rao designs over elementary abelian groups other than Z_2 have been studied by Lam and Seberry [26] and Seberry [45]. de Launey, Sarvate and Seberry [12] considered Bhaskar Rao designs over Z_4 which is an abelian (but not elementary) group. Some Bhaskar Rao designs over the non-abelian groups S_3 and Q_4 have been studied by Gibbons and Mathon [20].

Palmer and Seberry [33] study generalized Bhaskar Rao designs over the non-abelian groups S_3, D_4, Q_4, D_6 and over the small abelian group $Z_2 \times Z_4$. Seberry [46] completed the study of groups of order 8.

We use the following notation for initial blocks of a *GBRD*. We say $(a_\alpha, b_\beta, \dots, c_\gamma)$ is an initial block, when the Latin letters are developed mod v and the Greek subscripts are the elements of the group, which will be placed in the incidence matrix in the position indicated by the Latin letter. For example in the $(i, b - 1 + i)$ th position we place β and so on.

We form the difference table of an initial block $(a_\alpha, b_\beta, \dots, c_\gamma)$ by placing in the position headed by x_δ and by row y_η the element $(x - y)_{\delta\eta^{-1}}$ where $(x - y)$ is mod v and $\delta\eta^{-1}$ is in the group. By the term *totality* of elements we mean that repetitions remain: hence the set union of $\{1,2,3,4,5,6\} \cup \{3,4,7,8\} = \{1,2,3,4,5,6,7,8\}$ while the totality of elements in the two sets $\{1,2,3,4,5,6\} \& \{3,4,7,8\} = \{1,2,3,3,4,4,5,6,7,8\}$. The symbol $\&$ is sometimes written as \uplus .

A set of initial blocks will be said to form a *GBR* difference set (if there is one initial block) or *GBR* supplementary difference sets (if more than one) if in the totality of elements

$$(x - y)_{\delta\eta^{-1}} \pmod{v, G}$$

each non-zero element $a_g, a \pmod{v}, g \in G$, occurs $\lambda / |G|$ times.

Examples of the use of these *GBR* supplementary difference sets (*GBRSDS*) are given in Seberry [42].

2 GROUP DIVISIBLE DESIGNS

Let B be the incidence matrix of a $BIBD(v, b, r, k, \lambda)$. Let A be the matrix formed from a $GBRD(V, B, R, j, tv; G)$, where $|G| = v$, by replacing each zero of the *GBRD* by the $v \times v$ zero matrix and each group element of the *GBRD* by the right regular permutation matrix representation from the group $EA(v)$.

Then A is a $GDD(vV, vB, R, j, \lambda_1 = 0, \lambda_2 = t, m = V, n = v)$.

Lemma 1 *Suppose there exists a $BIBD(v, b, r, k, \lambda)$, Y , and a $GBRD(V, B, R, j, tv; G)$, A , with $|G| = v$. Then there exists a $GDD(vV, bB, rR, jk, \lambda_1 = R\lambda, \lambda_2 = trk, m = V, n = v)$.*

Proof. Let $C = A \times Y$, where the group element g_i of G with matrix representation G_i is replaced by $G_i Y$ and zero by the $v \times b$ zero matrix. Then all the parameters of C except λ_1 and λ_2 are immediate. The inner product of any two rows of Y is λ and $G_i Y$ also has inner product of rows λ . $G_i Y$'s occur R times in each row of C so $\lambda_1 = R\lambda$.

The inner product of rows of the $GBRD$ gives t copies of the group so the contribution to the inner product of rows of different GDD groups is

$$\& \quad tG_i Y Y^T G_j^{-1} = \quad \& \quad tG_i(r - \lambda)I + \lambda J)G_j^{-1} = t(r - \lambda + \lambda v)J.$$

$$g_i, g_j \in G \quad \quad \quad g_i, g_j \in G$$

Hence $\lambda_2 = trk$. Another way to check that $\lambda_2 = trk$ is to observe that we will have as inner product

$$t(G_1 + \cdots + G_v)Y Y^T = t(J)((r - \lambda)I + \lambda J) = t(r - \lambda)J + t(\lambda v)J,$$

Now use $\lambda(v - 1) = r(k - 1)$ to get the result. □

Example 1 Let the $Y = SBIBD(3, 2, 1)$ and A be the $GBRD(6, 6, 6; Z_3)$. Then the Lemma gives us $GDD(18, 18, 12, 12, \lambda_1 = 6, \lambda_2 = 8, m = 6, n = 3)$ which is given below and which is not found in Clatworthy's tables [7].

Example 2 Let A be formed from the $GBRD(7, 21, 9, 3, 3; Z_3) = D$ which may be written as:

$$D = \left[\begin{array}{cccccc|cccccc|cccccc} 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 0 & 0 & 1 & \omega^2 & 0 & w & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 0 & 0 & 1 & \omega^2 & 0 & w & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 0 & 1 & \omega^2 & 0 & w & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 1 & 0 & \omega^2 & 0 & 0 & 0 & 0 & 1 & \omega & 0 & w & 0 & 0 & 0 & 0 & 1 & \omega^2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \omega^2 & 0 & 0 & 0 & 0 & 1 & \omega & 0 & w & 0 & 0 & 0 & 0 & 1 & \omega^2 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 0 & 1 & \omega^2 & 0 & w & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 0 & 1 & \omega^2 & 0 & w & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

$$\text{Let } B = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}, \quad T = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \quad \text{and } \mathbf{O} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

We form C from A by replacing 0 by \mathbf{O} and ω^i by $T^i B$. Then C is a $GDD(21, 63, 18, 6, \lambda_1 = 9, \lambda_2 = 4, m = 7, n = 3)$.

Corollary 2 A $GBRD(p^2 + p + 1, p^2 + p + 1, p^2, p^2, p^2 - p; Z_2)$ always exists for p a prime power so a $GDD(2(p^2 + p + 1), 2(p^2 + p + 1), rp^2, kp^2, \lambda_1 = p^2\lambda, \lambda_2 = (p^2 - p)r)$ exists where $(k, r, \lambda) = (2, 2, 2)$ or $(1, 1, 0)$.

Corollary 3 A $GBRD(p + 1, p, p - 1; Z_q)$ exists for every prime power p , if q divides $p - 1$. An $SBIBD(q, q - 1, q - 2)$ exists. Hence there exists a $GDD(q(p + 1), q(p + 1), p(q - 1), p(q - 1), \lambda_1 = (q - 1)(q - 2), \lambda_2 = (p - 1)(q - 1)^2/q, m = p + 1, n = q)$.

Corollary 4 If a $GBRD(V, B, R, j, tv; EA(v))$ exists then a $GDD(vV, vB, R(v - 1), j(v - 1), \lambda_1 = R(v - 2), \lambda_2 = t(v - 1)^2, m = V, n = v)$ exists.

Proof. An $SBIBD(v, v - 1, v - 2)$ always exists. □

Example 2 (continued). There exists a $GH(3t, Z_3) = H, 3t > 7$, and write $SBIBD(7, 3, 1) = S$. Take seven rows of H and replace its elements by 0 and T^i to give,

$$H' = GDD(21, 9t, 3t, 7, \lambda_1 = 0, \lambda_2 = t, m = 7, n = 3).$$

Replace the zeros and ones of S by the 3×1 matrix of zeros and ones respectively to form,

$$S' = GDD(21, 7, 3, 9, \lambda_1 = 3, \lambda_2 = 1, m = 7, n = 3).$$

We note $L = I_7 \times J_{3,1}$ is a $GDD(21, 7, 1, 3, \lambda_1 = 1, \lambda_3 = 0, m = 7, n = 3)$.

Then

$$C_1 = [C : H'(t = 3) : H'(t = 4) : S']$$

is a $PBD(21, 12; K)$, where $K = \{6, 7, 9\}$, and

$$C_2 = [C : H'(t = 6) : L]$$

is a $PBD(21, 10; K)$, where $K = \{6, 7, 3\}$.

Table 1 gives some GDD on 21 varieties which exist.

v	k	m	n	λ_1	λ_2	Comment
21	12	7	3	4	2	from $SBIBD(7, 4, 2)$
21	7	7	3	0	t	(a $GH(3t, Z_3)$ exists, $3t > 7$)
21	18	7	3	6	5	$(J - I)_7 \times J_{3,1}$
21	3	7	3	1	0	$I_7 \times J_{3,1}$
21	9	7	3	3	1	from $SBIBD(7, 3, 1)$
21	k	7	3	r	λ	from $BIBD(7, b, r, k, \lambda)$
21	K	7	3	0	Δ	from $GBRD(7, B, R, K, 3\Delta; Z_3)$

Table 1

Remark 1 The GDD s with $\lambda_1 = r$ in Table 1 and in Table 2 can be constructed from a $BIBD$ and are singular but we have listed these parameters for easy reference so as to be able to apply them in the following Lemma and the Table 2 parameters in Lemma 6.

Clatworthy's tables [7] give $R188$ with $v = b = 21, k = r = 8, m = 7, n = 3, \lambda_1 = 7, \lambda_2 = 1$ but no GDD with $v = b = 21$, and the designs of Table 1 appear to be new.

Lemma 5 *Combinations from Table 1 can be used to give $PBD(21, \mu; K)$ for many μ and $k_i \in K$.*

Glynn [21] has found a $GW(13, 9, 6; S_3)$ which is circulant with the following first row: $[o e a d o a f e f o d]$ where $a = (1), b = (123)(456), c = (132)(465), d = (14)(26)(35), e = (15)$.

Example 7 Using Lemma 1 with the $SBIBD(6, 5, 4)$ we get

$$C = GDD(78, 78, 45, 45, \lambda_1 = 36, \lambda_2 = 20, m = 13, n = 6).$$

No.	v	k	m	n	λ_1	λ_2	Comment
B_1	78	54	13	6	9	6	from $SBIBD(13, 9, 6)$
B_2	78	45	13	6	36	20	above
B_3	78	13	13	6	0	t	(if a $GH(6t, G)$ exists $ G = b$, $6t > 13$: none are known),
B_4	78	72	13	6	12	11	$(J - I)13 \times J_{6,1}$
B_5	78	6	13	6	1	0	$I_{13} \times J_{6,1}$
B_6	78	24	13	6	4	1	from $SBIBD(13, 4, 1)$
B_7	78	k	13	6	r	λ	from $BIBD(13, b, r, k, \lambda)$
B_8	78	K	13	6	0	Δ	from $GBRD(13, B, R, K, 6\Delta; S_3)$
B_9	78	9	13	6	0	6	from $GW(13, 9, 6; S_3)$

Table 2

Table 2 gives some GDD on 78 varieties which exist.

We note that a $PBD(78, 38, K)$ with $K = \{6, 9, 45\}$ can be formed by taking

$$[B_2 : 2 \text{ copies } B_5 : 3 \text{ copies } B_9].$$

Clatworthy's tables list $R201$ which has $v = b = 78$, $r = k = 9$, $m = 13$, $n = 6$ but all the other designs in Table 2 appear to be new.

Lemma 6 *Combinations from Table 2 can be used to give $PBD(78, \mu; K)$ for many μ and K .*

3 GENERALIZED SUPPLEMENTARY DIFFERENCE SETS

We slightly extend a Lemma of de Launey and Seberry [11, Lemma 6.1.1] to get a new result.

Theorem 7 *Suppose there exist $n - \{v; k; \lambda\}$ supplementary difference sets and a square $GBRD(k, j, tg; G)$, $Y = (y_{su})$, where $|G| = g$. Then there exist $nk - \{v; j; t\lambda g; G\} - GBRSDS$.*

Proof. Let the $n - \{v; k; \lambda\}$ SDS , D_i , $i = 1, \dots, n$, have elements $d_1^i, d_2^i, \dots, d_k^i$.

Using the $GBRD(y_{su})$ we form nk $GBRSDS$ by choosing the initial blocks

$$d_{1_{y_{1u}}}^i, d_{2_{y_{2u}}}^i, \dots, d_{k_{y_{ku}}}^i, i = 1, 2, \dots, n; u = 1, 2, \dots, k,$$

where if y_{su} is 0, then we remove $d_{s_{y_{su}}}^i$ from the block (see Example 8).

These blocks are developed modulo v so that in a block, developed from an initial block with y_{au} in position $(1, a)$, position $(1 + b, a + b)$ is also y_{au} . Note that $1 + b$ and $a + b$ are both reduced modulo v .

Because the initial sets, D_i , had each element $1, 2, \dots, v - 1$ occurring as the solution of the equation

$$d_a^i - d_b^i, i \in \{1, \dots, n\}, a, b \in \{1, \dots, k\}$$

exactly λ times, the new design will have

$$g_{aj}g_{bj}^{-1}, j = 1, \dots, k, a, b \in \{1, \dots, k\}$$

occurring λtg times. Hence we have the starting blocks of an $nk - \{v; j; t\lambda g; G\} - GBRSDS$.

□

Corollary 8 Let $p \equiv 1 \pmod{4}$ be a prime power. Then there exist $2 - \{p; \frac{1}{2}(p-1); \frac{1}{2}(p-3)\}$ SDS. Suppose there exists a $GBRD((p-1)/2, k, tg; G)$ where $|G| = g$. Then there exist $(p-1) - \{p; k; tg(p-3)/2; G\} - GBRSDS$.

Example 8 We use the $GBRD(5, 5, 4, 4, 3; Z_3)$

$$Y = (y_{iu}) = \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & \omega & \omega^2 \\ 1 & 1 & 0 & \omega^2 & \omega \\ 1 & \omega & \omega^2 & 0 & 1 \\ 1 & \omega^2 & \omega & 1 & 0 \end{bmatrix} \text{ written as } \begin{bmatrix} \star & 0 & 0 & 0 & 0 \\ 0 & \star & 0 & 1 & 2 \\ 0 & 0 & \star & 2 & 1 \\ 0 & 1 & 2 & \star & 0 \\ 0 & 2 & 1 & 0 & \star \end{bmatrix}.$$

Attaching \star to an element is the same as multiplying it by zero in multiplicative notation and so removes that element from the starting block.

Now there are $3 - \{7; 5; 10\}$ SDS namely $\{0, 1, 2, 3, 4\}$, $\{0, 1, 2, 4, 5\}$ and $\{0, 1, 2, 3, 5\}$. So we make 15 starting blocks

$$\begin{aligned} & \{1_0, 2_0, 3_0, 4_0\}, \quad \{0_0, 2_0, 3_1, 4_2\}, \quad \{0_0, 1_0, 3_2, 4_1\}, \quad \{0_0, 1_1, 2_2, 4_0\}, \quad \{0_0, 1_2, 2_1, 3_0\}, \\ & \{1_0, 2_0, 4_0, 5_0\}, \quad \{0_0, 2_0, 4_1, 5_2\}, \quad \{0_0, 1_0, 4_2, 5_1\}, \quad \{0_0, 1_1, 2_2, 5_0\}, \quad \{0_0, 1_2, 2_1, 4_0\}, \\ & \{1_0, 2_0, 3_0, 5_0\}, \quad \{0_0, 2_0, 3_1, 5_2\}, \quad \{0_0, 1_0, 3_2, 5_1\}, \quad \{0_0, 1_1, 2_2, 5_0\}, \quad \{0_0, 1_2, 2_1, 3_0\}, \end{aligned}$$

which give a $15 - \{7; 4; 30; Z_3\} - GBRSDS$.

Applying the same method to the $(11, 5, 2)$ -difference set $\{1, 3, 4, 5, 9\}$ gives 5 starting blocks, $D_i^1, i = 1, \dots, 5$, namely $\{3_0, 4_0, 5_0, 9_0\}$, $\{1_0, 4_0, 5_1, 9_2\}$, $\{1_0, 3_0, 5_2, 9_1\}$, $\{1_0, 3_1, 4_2, 9_0\}$ and $\{1_0, 3_2, 4_1, 5_0\}$ which give a $5 - \{11; 4; 6; Z_3\} - GBRSDS$. (Note: superscript 1 in D_i^1 is not necessary in this example.)

Corollary 9 Let $p \equiv 3 \pmod{4}$ be a prime power. Suppose there exists a $GBRD(v, b, r, k, tp; G)$, $|G| = p$. Then there exist a $GDD(vp, bp, \frac{1}{2}r(p-1), \frac{1}{2}k(p-1), \lambda_1 = \frac{1}{4}r(p-3), \lambda_2 = \frac{1}{4}t(p-1)^2, m = v, n = p)$ and a $GDD(vp, bp, \frac{1}{2}r(p+1), \frac{1}{2}k(p+1), \lambda_1 = \frac{1}{4}r(p+1), \lambda_2 = \frac{1}{4}(p+1)^2, m = v, n = p)$.

Proof. Use the $(p, \frac{1}{2}(p-1), \frac{1}{4}(p-3))$ -difference set in the theorem or the $(p, \frac{1}{2}(p+1), \frac{1}{4}(p+1))$ difference set. \square

Corollary 10 Let $p \equiv 3 \pmod{4}$ and $p+1$ both be prime powers. Then there exist a $GDD(p(p+2), p(p+2), \frac{1}{2}(p^2-1), \frac{1}{2}(p^2-1), \lambda_1 = \frac{1}{4}(p+1)(p-3), \lambda_2 = \frac{1}{4}(p-1)^2, m = p+2, n = p)$ and an $SBIBD(p(p+2), \frac{1}{2}(p+1)^2, \frac{1}{4}(p+1)^2)$.

Proof. Use the previous corollary and the $GBRD(p+2, p+1, p; Z_p)$. \square

Example 9 Over $GF(2^3)$ with the primitive equation $\gamma^3 = \gamma + 1$ we have

$$\gamma, \gamma^2, \gamma^3 = \gamma + 1, \gamma^4 = \gamma^2 + \gamma, \gamma^5 = \gamma^3 = \gamma + 1, \gamma^6 = \gamma^2 + 1, \gamma^7 = 1$$

and choosing $m_{00} = m_{ii} = 0, m_{0i} = m_{i0} = 1, i = 1, \dots, 8$ and $m_{ij} = a^k$ if $\gamma^k = \gamma^j + \gamma^i$

	0	1	γ	γ^2	γ^3	γ^4	γ^5	γ^6
0	1	1	1	1	1	1	1	1
1	1	0	1	a	a^2	a^3	a^4	a^5
γ	1	1	0	a^3	a^2	a	a^5	a^4
γ^2	1	a	a^3	0	a^4	1	a^2	a^6
γ^3	1	a^2	a^2	a^4	0	a^5	a	a^3
γ^4	1	a^3	a	1	a^5	0	a^6	a^2
γ^5	1	a^4	a^2	a^2	a	a^6	0	1
γ^6	1	a^5	a^4	a^6	a^3	a^2	1	0
	1	a^6	a^2	a^5	1	a^4	a^3	a

We map $0 \rightarrow 0_7, a^i \rightarrow T^i B$. If B is an $SBIBD(7, k, \lambda)$ the new matrix has order 63, row and column sum $8k$, $\lambda_1 = 8\lambda$ and $\lambda_2 = k^2$. So we have an $SBIBD(63, 63, 32, 32, 16)$ or a $GDD(63, 63, 24, 24; \lambda_1 = 8, \lambda_2 = 9, m = 9, n = 7)$.

In general this construction takes a $GBRD(p+1, p, p-1)$ where p is a prime power and an $SBIBD(p-1, k, \lambda)$ and makes a $GDD(p^2-1, p^2-1, pk, pk, \lambda_1 = p\lambda, \lambda_2 = k^2)$.

Corollary 11 *Let $p \equiv 3 \pmod{4}$ and $q = p-1$ both be prime powers. Then there exists a $GDD(p(q^2+q+1), p^2(q^2+q+1), \frac{1}{2}p^2(p-1), \frac{1}{2}p(p-1); \lambda_1 = \frac{1}{4}p^2(p-3), \lambda_2 = \frac{1}{4}(p-1)^2)$ and a $GDD(p(q^2+q+1), p^2(q^2+q+1), \frac{1}{2}p^2(p+1), \frac{1}{2}p(p-1); \lambda_1 = \frac{1}{4}p^2(p+1), \lambda_2 = \frac{1}{4}(p+1)^2, m = q^2+q+1, n = p)$.*

Proof. Use the $p - \{q^2+q+1; q+1; q+1; G\} - GBRSDS, |G| = p$, to make a $GBRD(q^2+q+1, p(q^2+q+1), p^2, p, p; G)$. Then use the $SBIBD(p, \frac{1}{2}(p-1), \frac{1}{4}(p-3))$ and $SBIBD(p, \frac{1}{2}(p+1), \frac{1}{4}(p+1))$ to obtain the second GDD of the enunciation. \square

For example, we know that there exist a difference set for $SBIBD(7, 3, 1)$ and a $GBRD(3, 3, 3; Z_3)$. We apply Theorem 7 to get $3 - \{7, 3, 3; Z_3\} - GBRSDS$.

Now we use Lemma 1 and the trivial $BIBD(3, 1, 0) = I_3$ to obtain a $GDD(21, 63, 9, 3; 0, 1)$ and Lemma 1 and the $BIBD(3, 2, 1)$ to obtain a $GDD(21, 63, 9, 3; 9, 4)$.

4 GENERALIZED WEIGHING MATRICES

Write $G = (g_{ij})$ for a symmetric $GH(k, G), |G| = k$, where G comprises the k th roots of unity, $1, \gamma, \dots, \gamma^{k-1}$ with the relation $1 + \gamma + \gamma^2 + \dots + \gamma^{k-1} = 0$. G is in normalized form so $g_{0i} = g_{i0} = 1, i = 0, \dots, k-1$ and $g_{ij} = \gamma^{ij}$.

Let $D = \{d_1, \dots, d_k\}$ be a (v, k, λ) -difference set. Form the $k - \{v; k; k\lambda\} - GBRSDS, D_i = \{g_{i1}d_1, g_{i2}d_2, \dots, g_{ik}d_k\}, i = 1, \dots, k$. Call the matrices developed from D_i, A_i . Now we form a matrix, W , of order k^2 by choosing the circulant matrix with first row

$$[A_1 : A_2 : \dots : A_k].$$

We claim W is a generalized weighing matrix.

Theorem 12 *If k is a prime power and there exists a (v, k, λ) -difference set then there exists a $GW(vk, k^2; EA(k))$.*

Proof. We use the normalized $GH(k, EA(k)), G = (g_{ij})$ whose elements are the k th roots of unity as above. We form W as above.

There are three products to check: the inner product of row x and row $x + yk$, $y \neq 0$; the inner product of row x and row y where $x, y \in S_i = \{ik, ik + 1, \dots, ik + k - 1, i = 1, \dots, k\}$; the inner product of row x and row y where $x \in S_i, y \in S_j, i \neq j$.

The first row of A_i has $a_{1,d_j} = g_{ij}$, $a_{1,n} = 0$ otherwise. Hence the x th row of A_i has $a_{x,j} = a_{1,j-x+1} = g_{1m}$ if $j - x + 1 = d_m$ and $a_{xj} = 0$ otherwise.

Case 1: The inner product of the x th row and the $x + yk$ th row of W is

$$\begin{aligned} & \sum_{z=0}^{k-1} \sum_{j=1}^v a_{x,j+zk} a_{x+yk,j+zk}^{-1}, \quad y \neq 0 \\ &= \sum_{z=0}^{k-1} \sum_{j=1}^v a_{1,j+zk-x+1} a_{1,j+zk-x-yk+1}^{-1}, \quad y \neq 0 \\ &= \sum_{z=0}^{k-1} \sum_{d_m \in D} g_{zm} g_{(z-y),m}^{-1}, \quad y \neq 0, \text{ if } d_m = j - x + 1, d_m \in D \\ &= 0 \end{aligned}$$

since $\sum_{d \in D} g_{zm} g_{(z-y),m}^{-1}$ is the inner product of two rows of the GH, G , for which $1 + \gamma + \gamma^2 + \dots + \gamma^{k-1} = 0$.

Case 2: The inner product of the x th row and the y th row, $x, y \in S_i$ is

$$\begin{aligned} & \sum_{z=0}^{k-1} \sum_{j=1}^v a_{x,j+zk} a_{y,j+zk}^{-1}, \quad y \neq x \\ &= \sum_{z=0}^{k-1} \sum_{d_m, d_n \in D} g_{zm} g_{zn}^{-1}, \quad d_m = j - x + 1, d_n = j - y + 1, d_m, d_n \in D \\ &= 0 \end{aligned}$$

since $\sum_{z=0}^{k-1} g_{zm} g_{zn}^{-1}$ is the inner product of two rows of the GH .

Case 3: The inner product of x th and y th rows, $x \in S_i, y \in S_j, i \neq j$ is

$$\begin{aligned} & \sum_{z=0}^{k-1} \sum_{j=1}^v a_{x,j+zk} a_{y,j+zk}^{-1} \\ &= \sum_{z=0}^{k-1} \sum_{d_m, d_n \in D} g_{zm} g_{z+w,n}^{-1}, \quad \text{some } w \neq 0, \end{aligned}$$

$$d_m = j - x + 1, d_n = j - y + 1, d_m, d_n \in D.$$

(The w reflects that where x, y come from different S_i , the elements of row y have all been incremented by the same fixed constant (w) due to the block cyclic structure of W)

$$\begin{aligned} &= \sum_{z=0}^{k-1} \sum_{d_m, d_n \in D} \gamma^{zm} \gamma^{-zn - wn} \\ &= \sum_{z=0}^{k-1} \sum_{d_m, d_n \in D} \gamma^{z(m-n) - wn} \end{aligned}$$

$$\begin{aligned}
&= \sum_{d_m, d_n \in D} \sum_{z=0}^{k-1} \gamma^{z(m-n)-\omega n} \\
&= 0 \text{ as } \sum_{z=0}^{k-1} \gamma^{z(m-n)-\omega n} = 0.
\end{aligned}$$

Thus we have the result. \square

Example 10 The $GW(21, 9; Z_3)$ is given. Similarly one can construct a $GW(55, 25; Z_5)$.

$$\begin{bmatrix}
0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 \\
0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 \\
0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega \\
1 & 0 & 0 & 0 & 1 & 1 & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega & 0 & \omega & 0 & 0 & 0 & 1 & \omega^2 & 0 \\
0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega & 0 & \omega & 0 & 0 & 0 & 1 & \omega^2 \\
1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 \\
1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 \\
\\
0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 \\
0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 \\
0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 \\
\omega & 0 & 0 & 0 & 1 & \omega^2 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega & 0 \\
0 & \omega & 0 & 0 & 0 & 1 & \omega^2 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega \\
\omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 \\
1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 \\
\\
0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 \\
0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\
0 & 0 & 0 & 1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\
\omega^2 & 0 & 0 & 0 & 1 & \omega & 0 & \omega & 0 & 0 & 0 & 1 & \omega^2 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 \\
0 & \omega^2 & 0 & 0 & 0 & 1 & \omega & 0 & \omega & 0 & 0 & 0 & 1 & \omega^2 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \\
\omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 1 \\
1 & \omega & 0 & \omega^2 & 0 & 0 & 0 & 1 & \omega^2 & 0 & \omega & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 & 0 & 0
\end{bmatrix}$$

Lemma 13 *If k is a prime power and there exists a (v, k, λ) -difference set then there exists a $PBIBD(vk^2, vk^2, k^2, k^2; \lambda_1 = 0, \lambda_2 = \lambda, \lambda_3 = k)$.*

Proof. We replace the elements of the $GW(vk, k^2; EA(k))$ by their matrix representation as before. This gives $\lambda_1 = 0$.

The set of x th and $(x + yk)$ th rows, $y = 0, \dots, k - 1$ of the GW give the third association which has $\lambda_3 = k$.

The set of rows corresponding to the product of the x th rows and the y th rows, $x \in S_i, y \in S_j, i \neq j$ give the second association class with $\lambda_2 = \lambda$. \square

Table 3 gives some of the generalized weighing matrices and $PBIBDs$ parameters obtained by using Theorem 12 and Lemma 13.

Example 11 From the $GW(21, 9; Z_3)$ with ω^i replaced by T^i we have the classes comprising rows $3j + 1, 3j + 2, 3j + 3, j = 0, 1, \dots, 20$ with inner product zero.

Rows $3j + 1, 3j + 2, 3j + 3$ with any of $21 + 3j + 1, 21 + 3j + 2, 21 + 3j + 3$ (and vice versa) and with any of $42 + 3j + 1, 42 + 3j + 2, 42 + 3j + 3, j = 0, 1, \dots, 7$ (and vice versa) have inner product 3.

All other pairs of rows have inner product 1.

Difference set (v, k, λ)	GW ($vk, k^2; EA(k)$)	$PBIBD$ ($vk^2, vk^2, k^2, k^2; 0, \lambda, k$)
(4, 3, 2)	(12, 9; $EA(3)$)	(36, 36, 9, 9; 0, 2, 3)
(7, 3, 1)	(21, 9; $EA(3)$)	(63, 63, 9, 9; 0, 1, 3)
(5, 4, 3)	(20, 16; $EA(4)$)	(80, 80, 16, 16; 0, 3, 4)
(7, 4, 2)	(28, 16; $EA(4)$)	(112, 112, 16, 16; 0, 2, 4)
(13, 4, 1)	(52, 16; $EA(4)$)	(208, 208, 16, 16; 0, 1, 4)
(6, 5, 4)	(30, 25; $EA(5)$)	(150, 150, 25, 25; 0, 4, 5)
(11, 5, 2)	(55, 25; $EA(5)$)	(275, 275, 25, 25; 0, 2, 5)
(21, 5, 1)	(105, 25; $EA(5)$)	(525, 525, 25, 25; 0, 1, 5)
(8, 7, 6)	(56, 49; $EA(7)$)	(392, 392, 49, 49; 0, 6, 7)
(15, 7, 3)	(105, 49; $EA(7)$)	(735, 735, 49, 49; 0, 3, 7)
(9, 8, 7)	(72, 64; $EA(8)$)	(576, 576, 64, 64; 0, 7, 8)
(15, 8, 4)	(120, 64; $EA(8)$)	(960, 960, 64, 64; 0, 4, 8)
(57, 8, 1)	(456, 64; $EA(64)$)	(3648, 3648, 64, 64; 0, 1, 8)

Table 3

So the $PBIBD$, X , satisfies $XJ = JX = 9$,

$$XX^T = 9I_3 \times I_7 \times I_3 + (J - I)_3 \times J_7 \times J_3 + (J - I)_3 \times I_7 \times 2J.$$

Hence we have a $PBIBD(63, 63, 9, 9; \lambda_1 = 0, \lambda_2 = 1, \lambda_3 = 3)$.

Acknowledgment: We wish to thank Dr. Kishore Sinha for his helpful advice and comments.

References

- [1] Gerald Berman (1977), Weighing matrices and group divisible designs determined by $EG(t, pn)$, $t > 2$, *Utilitas Math.*, **12**, 183-192.
- [2] Gerald Berman (1978), Families of generalised weighing matrices, *Canad. J. Math.*, **30**, 1016-1028.
- [3] M. Bhaskar Rao (1966), Group divisible family of PBIB designs. *J. Indian Stat. Assoc.*, **4**, 14-28.
- [4] M. Bhaskar Rao (1970), Balanced orthogonal designs and their applications in the construction of some BIB and group divisible designs. *Sankhya Ser. A*, **32**, 439-448.
- [5] A.T. Butson (1962), Generalized Hadamard matrices, *Proc. Amer. Math. Soc.*, **13**, 894-898.
- [6] A.T. Butson (1963), Relations among generalized Hadamard matrices, relative difference sets and maximal length recurring sequences, *Canad. J. Math.*, **15**, 42-48.
- [7] Clatworthy W. H., *Tables of Two-Associate-Class Partially Balanced Designs*, National Bureau of Standards, US Commerce Department, 1971.

- [8] Jeremy E. Dawson (1985), A construction for the generalized Hadamard matrices $GH(4q, EA(q))$, *J. Statist. Plann. and Inference*, **11**, 103-110.
- [9] Warwick de Launey (1984), On the non-existence of generalised Hadamard matrices, *J. Statist. Plann. and Inference*, **10**, 385-396.
- [10] Warwick de Launey (1986), A survey of generalised Hadamard matrices and difference matrices $D(k, \lambda; G)$ with large k , *Utilitas Math.*, **30**, 5-29.
- [11] Warwick de Launey and Jennifer Seberry (1984), Generalised Bhaskar Rao designs of block size four, *Congressus Numerantium*, **41**, 229-294.
- [12] Warwick de Launey, D.G. Sarvate, Jennifer Seberry (1985). Generalised Bhaskar Rao Designs with block size 3 over Z_4 , *Ars Combinatoria*, **19A**, 273-286.
- [13] P. Delsarte and J.M. Goethals (1969), Tri-weight codes and generalized Hadamard matrices, *Information and Control*, **15**, 196-206.
- [14] P. Delsarte and J.M. Goethals (1971), On quadratic residue-like sequences in Abelian groups, Report R168, MBLE Research Laboratory, Brussels.
- [15] A. Dey and C.K. Midha (1976), Generalised balance matrices and their applications, *Utilitas Math.*, **10**, 139-149.
- [16] D.A. Drake (1979), Partial λ -geometries and generalized matrices over groups, *Canad. J. Math.*, **31**, 617-627.
- [17] P. Eades (1980), On circulant (v, k, λ) -designs, *Combinatorial Mathematics VII*, Lecture Notes in Mathematics, Vol **829**, Springer-Verlag, Berlin-Heidelberg-New York, 83-93.
- [18] A.V. Geramita and J. Seberry (1979), *Orthogonal Designs: Quadratic Forms and Hadamard Matrices*, Marcel Dekker, New York.
- [19] A.V. Geramita, N.J. Pullman and J. Seberry Wallis, Families of weighing matrices, *Bull. Austral. Math. Soc.*, **10**, 119-122.
- [20] Peter Gibbons and Rudolf Mathon (1987), Construction Methods for Bhaskar Rao and related designs, *J. Austral. Math. Soc. Ser A*, **42**, 5-30.
- [21] D.G. Glynn (1978), *Finite Projective Planes and Related Combinatorial Systems*, Ph.D. Thesis, University of Adelaide, 1978.
- [22] Marshall Hall Jr. (1967), *Combinatorial Mathematics*, Blaisdell, Waltham, Mass.
- [23] H. Hanani (1961), The existence and construction of balanced incomplete block designs, *Ann. Math. Stat.*, **32**, 361-386.
- [24] H. Hanani, (1975), Balanced incomplete block designs and related designs, *Discrete Math.*, **11**, 255-369.
- [25] Dieter Jungnickel (1979), On difference matrices, resolvable TD's and generalized Hadamard matrices, *Math. Z.*, **167**, 49-60.
- [26] Clement Lam and Jennifer Seberry (1984), Generalized Bhaskar Rao designs, *J. Statist. Plann. and Inference*, **10**, 83-95.
- [27] R.C. Mullin (1974), Normal affine resolvable designs and orthogonal matrices, *Utilitas Math.*, **6**, 195-208.

- [28] R.C. Mullin (1975), A note on balanced weighting matrices, *Combinatorial Mathematics III*, Lecture Notes in Mathematics, Vol. **452**, Springer-Verlag, Berlin-Heidelberg-New York, 28-41.
- [29] R.C. Mullin and R.G. Stanton (1975), Group matrices and balanced weighing designs, *Utilitas Math.*, **8**, 303-310.
- [30] R.C. Mullin and R.G. Stanton (1975), Balanced weighing matrices and group divisible designs, *Utilitas Math.*, **8**, 277-301.
- [31] R.C. Mullin and R.G. Stanton (1976), Corrigenda to: Balanced weighing matrices and group divisible designs, *Utilitas Math.*, **9**, 347.
- [32] William D. Palmer, (1990), Generalized Bhaskar Rao designs with two association classes, *Australas. J. Combin.*, **1**, 161-180.
- [33] William D. Palmer and Jennifer Seberry, Bhaskar Rao designs over small groups, *Ars Combinatoria*, **26A**, (1988) 125-148
- [34] D. Raghavarao (1971), *Construction and Combinatorial Problems in Design of Experiments*, Wiley, New York.
- [35] Dinesh Rajkundlia (1983), Some techniques for constructing infinite families of BIBDs, *Discrete Math.*, **44**, 61-96.
- [36] G.M. Saha and A.D. Dab(1978), An infinite class of PBIB designs, *Canad. J. Statist.*, **6**, 25-32.
- [37] G.M. Saha and Gauri Shankar (1976), On a Generalized Group Divisible family of association schemes and PBIB designs based on the schemes, *Sankhya*, **38B**, 393-403.
- [38] Jennifer Seberry Wallis (1972), Hadamard matrices. Part IV of *Combinatorics: Room squares, sum free sets and Hadamard matrices*, Lecture Notes in Mathematics, Vol **292**, Springer-Verlag, Berlin-Heidelberg-New York, 273-489.
- [39] Jennifer Seberry (1978), A class of group divisible designs, *Ars Combinatoria*, **6**, 151-152.
- [40] Jennifer Seberry (1979), Some remarks on generalized Hadamard matrices and theorems of Rajkundlia on SBIBDs, *Combinatorial Mathematics IV*, Lecture Notes in Mathematics, Vol **748**, Springer Verlag, Berlin-Heidelberg-New York, 154-164.
- [41] Jennifer Seberry (1980), A construction for generalized Hadamard matrices, *J. Statist. Plann. and Inference*, **4**, 365-368.
- [42] Jennifer Seberry (1982), Some families of partially balanced incomplete block designs. *Combinatorial Mathematics IX*, Lecture Notes in Mathematics, Vol **952**, Springer, Berlin-Heidelberg-New York, 378-386.
- [43] Jennifer Seberry (1982), The skew-weighing matrix conjecture, *Uni. of Indore Research J. Science*, **7**, 1-7.
- [44] Jennifer Seberry (1984), Regular group divisible designs and Bhaskar Rao designs with block size 3, *J. Statist. Plann. and Inference*, **10**, 69-82.
- [45] Jennifer Seberry (1985), Generalized Bhaskar Rao designs of block size three, *J. Statist. Plann. and Inference*, **11**, 373-379.

- [46] Jennifer Seberry (1988), Bhaskar Rao designs of block size 3 over group of order 8, University College, UNSW, Technical report CS88/4.
- [47] S.S. Shrikhande (1964), Generalized Hadamard matrices and orthogonal arrays of strength 2, *Canad. J. Math.*, **16**, 736-740.
- [48] S.J. Singh (1982), Some Bhaskar Rao designs and applications for $k = 3, \lambda = 2$, *University of Indore J. Science*, **7**, 8-15.
- [49] Kishore Sinha (1978), Partially balanced incomplete block designs and partially balanced weighing designs, *Ars Combinatoria*, **6**, 91-96.
- [50] Deborah J. Street (1979), Generalized Hadamard matrices, orthogonal arrays and F -squares, *Ars Combinatoria*, **8**, 131-141.
- [51] Deborah J. Street (1981), Bhaskar Rao designs from cyclotomy, *J. Austral. Math. Soc. Ser. A*, **29**, 425-430.
- [52] D.J. Street and C.A. Rodger (1980), Some results on Bhaskar Rao designs, *Combinatorial Mathematics VII*, Lecture Notes in Mathematics, Vol **829**, Springer-Verlag, Berlin-Heidelberg-New York, 238-245.
- [53] Anne P. Street and Deborah J. Street, *Combinatorics of Experimental Design*, Oxford University Press, Oxford 1987.
- [54] R. Vyas (1982), Some Bhaskar Rao Designs and applications for $k = 3, \lambda = 4$, *University of Indore J. Science*, **7**, 16-25.
- [55] R.M. Wilson (1974), A few more squares, Proc. Fifth South Eastern Conf. on Combinatorics, Graph Theory and Computing, *Congressus Numerantium XXI*, Utilitas Mathematica, Winnipeg, 675-680.
- [56] R.M. Wilson (1975), Construction and uses and pairwise balanced designs, *Combinatorics*, Edited by M. Hall Jr. and J.H. van Lint, (Mathematisch Centrum, Amsterdam), 19-42
- [57] R.M. Wilson (1975), An existence theory of pairwise balanced designs III; proof of the existence conjectures, *J. Combinatorial Theory Series A*, **18**, 71-79.