

1-1-2009

Dietary fish oil is anti-hypertrophic but does not enhance post-ischemic myocardial function in female mice

Catherine E. Huggins
University of Melbourne

Claire L. Curl
University of Melbourne

Ruchi Patel
Monash University

Peter L. McLennan
University of Wollongong, petermcl@uow.edu.au

Mandy L. Theiss
University of Wollongong, mlt01@uow.edu.au

See next page for additional authors

Follow this and additional works at: <https://ro.uow.edu.au/medpapers>



Part of the [Medicine and Health Sciences Commons](#)

Citation

Huggins, Catherine E.; Curl, Claire L.; Patel, Ruchi; McLennan, Peter L.; Theiss, Mandy L.; Pedrazzini, Thierry; Pepe, Salvatore; and Delbridge, Lea M.D., 2009, Dietary fish oil is anti-hypertrophic but does not enhance post-ischemic myocardial function in female mice, *American Journal of Physiology*, 296, H957-H966.

<https://ro.uow.edu.au/medpapers/7>

Dietary fish oil is anti-hypertrophic but does not enhance post-ischemic myocardial function in female mice

Abstract

Clinically and experimentally, a case for omega-3 polyunsaturated fatty acid (PUFA) cardioprotection in females has not been clearly established. The goal of this study was to investigate whether dietary omega-3 PUFA supplementation could provide ischemic protection in female mice with an underlying genetic predisposition to cardiac hypertrophy. Mature female transgenic mice (TG) with cardiac-specific overexpression of angiotensinogen that develop normotensive cardiac hypertrophy and littermate wild-type (WT) mice were fed a fish oil-derived diet (FO) or PUFA-matched control diet (CTR) for 4 wk. Myocardial membrane lipids, ex vivo cardiac performance (intraventricular balloon) after global no-flow ischemia and reperfusion (15/30 min), and reperfusion arrhythmia incidence were assessed. FO diet suppressed cardiac growth by 5% and 10% in WT and TG, respectively ($P < 0.001$). The extent of mechanical recovery [ratepressure product (RPP) = beats/min \times mmHg] of FO-fed WT and TG hearts was similar ($50 \pm 7\%$ vs. $45 \pm 12\%$, 30 min reperfusion), and this was not significantly different from CTR-fed WT or TG. To evaluate whether systemic estrogen was masking a protective effect of the FO diet, the responses of ovariectomized (OVX) WT and TG mice to FO dietary intervention were assessed. The extent of mechanical recovery of FO-fed OVX WT and TG (RPP, $50 \pm 4\%$ vs. $64 \pm 8\%$) was not enhanced compared with CTR-fed mice (RPP, $60 \pm 11\%$ vs. $80 \pm 8\%$, $P = 0.335$). Dietary FO did not suppress the incidence of reperfusion arrhythmias in WT or TG hearts (ovary-intact mice or OVX). Our findings indicate a lack of cardioprotective effect of dietary FO in females, determined by assessment of mechanical and arrhythmic activity postischemia in a murine ex vivo heart model.

Keywords

enhance, myocardial, post, mice, dietary, female, fish, oil, anti, function, hypertrophic, but, ischemic, does, not

Disciplines

Medicine and Health Sciences

Authors

Catherine E. Huggins, Claire L. Curl, Ruchi Patel, Peter L. McLennan, Mandy L. Theiss, Thierry Pedrazzini, Salvatore Pepe, and Lea M.D. Delbridge

Catherine E. Huggins, Claire L. Curl, Ruchi Patel, Peter L. McLennan, Mandy L. Theiss, Thierry Pedrazzini, Salvatore Pepe and Lea M. D. Delbridge
Am J Physiol Heart Circ Physiol 296:957-966, 2009. First published Jan 30, 2009;
doi:10.1152/ajpheart.01151.2008

You might find this additional information useful...

This article cites 49 articles, 27 of which you can access free at:

<http://ajpheart.physiology.org/cgi/content/full/296/4/H957#BIBL>

Updated information and services including high-resolution figures, can be found at:

<http://ajpheart.physiology.org/cgi/content/full/296/4/H957>

Additional material and information about *AJP - Heart and Circulatory Physiology* can be found at:

<http://www.the-aps.org/publications/ajpheart>

This information is current as of March 18, 2010 .

Dietary fish oil is antihypertrophic but does not enhance postischemic myocardial function in female mice

Catherine E. Huggins,¹ Claire L. Curl,¹ Ruchi Patel,² Peter L. McLennan,³ Mandy L. Theiss,³ Thierry Pedrazzini,⁴ Salvatore Pepe,^{1,5} and Lea M. D. Delbridge¹

¹Cardiac Phenomics Laboratory, Department of Physiology and ⁵Murdoch Children's Research Institute, Department of Paediatrics, University of Melbourne and Department of Cardiology, Royal Children's Hospital, Parkville, Victoria; ²Department of Physiology, Monash University, Victoria; ³Division of Medical Sciences, Graduate School of Medicine, School of Health Sciences, University of Wollongong, New South Wales, Australia; ⁴Department of Medicine, University of Lausanne Medical School, Lausanne, Switzerland

Submitted 30 October 2008; accepted in final form 26 January 2009

Huggins CE, Curl CL, Patel R, McLennan PL, Theiss ML, Pedrazzini T, Pepe S, Delbridge LM. Dietary fish oil is antihypertrophic but does not enhance postischemic myocardial function in female mice. *Am J Physiol Heart Circ Physiol* 296: H957–H966, 2009. First published January 30, 2009; doi:10.1152/ajpheart.01151.2008.—Clinically and experimentally, a case for omega-3 polyunsaturated fatty acid (PUFA) cardioprotection in females has not been clearly established. The goal of this study was to investigate whether dietary omega-3 PUFA supplementation could provide ischemic protection in female mice with an underlying genetic predisposition to cardiac hypertrophy. Mature female transgenic mice (TG) with cardiac-specific overexpression of angiotensinogen that develop normotensive cardiac hypertrophy and littermate wild-type (WT) mice were fed a fish oil-derived diet (FO) or PUFA-matched control diet (CTR) for 4 wk. Myocardial membrane lipids, ex vivo cardiac performance (intraventricular balloon) after global no-flow ischemia and reperfusion (15/30 min), and reperfusion arrhythmia incidence were assessed. FO diet suppressed cardiac growth by 5% and 10% in WT and TG, respectively ($P < 0.001$). The extent of mechanical recovery [rate-pressure product (RPP) = beats/min \times mmHg] of FO-fed WT and TG hearts was similar ($50 \pm 7\%$ vs. $45 \pm 12\%$, 30 min reperfusion), and this was not significantly different from CTR-fed WT or TG. To evaluate whether systemic estrogen was masking a protective effect of the FO diet, the responses of ovariectomized (OVX) WT and TG mice to FO dietary intervention were assessed. The extent of mechanical recovery of FO-fed OVX WT and TG (RPP, $50 \pm 4\%$ vs. $64 \pm 8\%$) was not enhanced compared with CTR-fed mice (RPP, $60 \pm 11\%$ vs. $80 \pm 8\%$, $P = 0.335$). Dietary FO did not suppress the incidence of reperfusion arrhythmias in WT or TG hearts (ovary-intact mice or OVX). Our findings indicate a lack of cardioprotective effect of dietary FO in females, determined by assessment of mechanical and arrhythmic activity postischemia in a murine ex vivo heart model.

polyunsaturated fatty acids; ischemia-reperfusion; Langendorff perfused hearts

LARGE CLINICAL TRIALS HAVE demonstrated that when omega-3 polyunsaturated fatty acid (PUFA) from fish oil (FO) [docosahexaenoic acid (DHA) + eicosapentaenoic acid (EPA)] are used in secondary prevention following myocardial infarction, there is a marked reduction in cardiac-related death and particularly in sudden cardiac death (3, 17). More recently omega-3 PUFA supplementation (DHA + EPA) has been demonstrated to be a beneficial adjunct to present therapies for

treatment of heart failure (16). Although these findings have been of landmark significance, a limitation has been the underrepresentation of women, with women either not included in the cohort (3) or comprising only a small proportion (8–20%) of participants (16, 17). This limited female representation is significant because sex differences in cardiac function and in the incidence and progression of cardiovascular diseases are well documented (10, 25).

One recent exception is the Japan EPA Lipid Intervention Study (JELIS), which was a large study (18,645 participants) of predominantly postmenopausal women (~70%). Interestingly, this study found no significant association between EPA supplementation and sudden cardiac death (49). It is plausible that this reflects a sex specificity of omega-3 PUFA not previously identified (34). In a subanalysis of the JELIS cohort, a risk reduction for major coronary events (combining sudden cardiac death, fatal myocardial infarction, and nonfatal myocardial infarction) could be identified but only in the quartile comprising a largest ratio of men to women (40).

Experimentally there is also a dearth of information on the cardiac effects of omega-3 PUFA in the female heart. There is some evidence to suggest that the balance between omega-3 and omega-6 PUFA affects males and females differently. Korotkova et al. (26) demonstrated that the supply of dietary PUFA (of varying ratios of n3:n6 PUFA) during the perinatal period in rats had long-term effects on blood pressure in adulthood that were different for males and females (associated with the specific n3:n6 PUFA ratio of the diet). Thus the possibility that dietary omega-3 PUFA supplementation may have distinct cardiovascular actions in females requires evaluation.

There is strong evidence from experimental studies (of males) that dietary omega-3 PUFA have antiarrhythmic properties in the setting of ischemia and reperfusion and postmyocardial infarction (29, 31, 39) although the cellular mechanisms are incompletely understood and controversial (4, 29). Omega-3-mediated improvement of postischemic contractile recovery independent of vascular function (39) has been attributed at least in part to the incorporation of these PUFA into membrane phospholipids. Systemic estrogen has also been shown to afford protection against arrhythmogenesis, ischemia-reperfusion injury, and myocardial hypertrophy (22, 35, 43). Various in vitro studies suggest that the biological actions and cellular

Address for reprint requests and other correspondence: L. Delbridge, Cardiac Phenomics Laboratory, Dept. of Physiology, Univ. of Melbourne, Parkville, Victoria, Australia 3010 (e-mail: lmd@unimelb.edu.au).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

targets of omega-3 PUFA and estrogen may be similar, suggesting that omega-3 PUFA may exert a greater effect in females after a period of estrogen withdrawal; albeit a possible convergence of action has not been directly investigated experimentally (7, 18, 27, 33, 47).

Given the lack of available female-specific experimental data, and in the absence of convincing clinical findings relating to the cardiovascular efficacy of omega-3 PUFA in women, a case for omega-3 cardioprotection in females has not been clearly established. Therefore, the aim of this study was to determine whether dietary omega-3 PUFA supplementation could provide cardioprotection in female mice, specifically by enhancing the recovery of ex vivo hearts to an ischemic insult. Cardiac hypertrophy is associated with increased risk of arrhythmia and vulnerability to ischemia and reperfusion injury (14, 24) and hence is a pathological condition that would be expected to be favorably modulated by the protective actions of omega-3 PUFA supplementation. Thus our study included a cohort of female mice additionally compromised by an underlying genetic predisposition for cardiac hypertrophy. Our experiments utilized the angiotensinogen-over-expressing mouse, a model of load-independent cardiac hypertrophy resulting from amplified cardiac-specific angiotensin II (Ang II) production (30). We hypothesized that dietary FO supplementation would be most efficacious (in enhancing postischemic recovery and suppressing arrhythmias) in females where there is an underlying predisposition for cardiac hypertrophy and additionally that a protective effect of dietary FO would be more prominent in the context of systemic estrogen depletion.

MATERIALS AND METHODS

The project was approved by the University of Melbourne Animal Ethics Committee (AEC Project 05173).

Experimental model. All animals were handled in the manner specified by the Prevention of Cruelty to Animals Act 1986 and NHMRC/CSIRO/ACC Australian Code of Practice for the Care and Use of Animals for Scientific Purposes (1997). The generation and characterization of a transgenic heterozygous mouse (TG1306/1R) harboring multiple copies of a transgene expressing the rat angiotensinogen gene under the α -myosin heavy chain promoter have been reported previously (9, 30). A colony of these mice was established in the Biological Research Facility (Department of Physiology, University of Melbourne). In this study, female age-matched transgenic mice (TG) and littermate wild-type (WT) controls were investigated with the use of a 4-wk treatment (the reported time required to achieve maximum membrane omega-3 incorporation) (37) terminating at age 34 wk. At the conclusion of the 4-wk treatment period, mice were anesthetized with pentobarbital sodium (100 mg/kg), and hearts were excised and perfused ex vivo (described below). In a separate cohort of mice, hearts were excised, adipose tissue and lungs removed, and hearts blotted dry and weighed. Atria and ventricle were then separated and weighed. Tissues were snap frozen in liquid and stored at -80°C until determination of membrane fatty acid composition.

Diets. During the treatment period, mice were fed ad libitum a fully fabricated phytoestrogen-free diet (based on the American Institute of Nutrition standard diet, 1993; AIN93), prepared at the Smart Foods Center, University of Wollongong (P. L. McLennan). The control diet (CTR) was specified to be representative of a "standard" commercial chow diet with minimum nutritional level omega-3 content and was fabricated (% of dry weight) from 50.5% cornstarch, 10% sucrose, 15% casein, 5% gelatin, 5% fiber, 3.5% mineral mix, 1% vitamin mix, and 10% oil (5% sunflower seed oil and 5% olive oil). The FO omega-3-rich diet comprised 10% oil content consisting of 7% tuna fish oil (Nu-Mega Ingredients, Brisbane, QLD, Australia) and 3%

olive oil (FO diet). The FO diet was matched for total level of fatty acid polyunsaturation and caloric content to the CTR diet (1).

Cardiac membrane phospholipid fatty acid analysis. Total nonfractionated ventricular membrane phospholipid fatty acids were extracted from 70–100 mg of ventricular tissue using a modification of the Folch method as previously described (31). Phospholipids were isolated by solid-phase extraction, using silica Sep-pak cartridges




Fig. 1. Body weight and cardiac growth of female angiotensinogen transgenic mice (TG, hatched bars) at diet treatment completion (4 wk), age 34 wk, compared with littermate controls [wild-type (WT), open bars]. Data are presented as means \pm SE ($n = 8/\text{group}$). Two-way ANOVA (factors, genetic type, and diet). A: body weight: WT vs. TG, $P = 0.305$; control diet (CTR) vs. fish oil diet (FO), $P = 0.483$; interaction factor $P = 0.507$. B: heart weight: WT vs. TG, $*P < 0.001$; CTR vs. FO, $P = 0.081$; interaction factor $P = 0.272$. C: heart weight normalized to body weight (mg/g): WT vs. TG, $*P < 0.001$; CTR vs. FO, $\#P = 0.014$; interaction factor, $P = 0.422$.

Table 1. Membrane PUFA levels at diet completion

		LA	Arachidonic	EPA	DHA	Omega-3:Omega-6
		18:2 n6	20:4 n6	20:5 n3	22:6 n3	
WT	CTR	16.3±0.5	10.6±0.1	ND	24.2±0.3	0.8±0.1
	FO	2.8±0.1	3.2±0.1	0.5±0.1	47.5±0.2	5.9±0.1
	CTR	16.5±0.5	10.5±0.1	ND	24.0±0.3	0.8±0.1
TG	FO	2.9±0.1	2.9±0.1	0.4±0.1	48.8±0.1	6.2±0.1
	WT vs. TG	0.763	0.240	0.292	0.244	0.033
P	CTR vs. FO	<0.001	<0.001		<0.001	<0.001
	interaction	0.904	0.630		0.010	0.014

Data are presented as a percentage of total membrane phospholipid fatty acids (means ± SE; $n = 3$ -5/group). 2-way ANOVA (factors, genetic type, and diet), P , probability. Eicosapentaenoic acid (EPA) was analyzed by unpaired t -test. PUFA, polyunsaturated fatty acid; CTR, control diet; FO, fish oil; LA, linoleic acid; DHA, docosahexaenoic acid; WT, wild-type mice; TG, transgenic mice.

(Waters, Milford, MA). Fatty acids were methylated using methanol: toluene (4:1 vol/vol) plus 200 μ l of acetyl chloride and placed on a dry heat block at 100°C for 1 h. Fatty acid methyl esters were analyzed by gas chromatography using a Shimadzu GC-17A with flame ionization detection. The column used was a FAMEWax Crossbond-PEG 30 $m \times 0.25$ mm $\times 0.25$ μ m (Restek, Bellefonte, PA) with hydrogen used as the carrier gas (temperature rise 150–230°C over 28 min). Fatty acids were identified from authentic fatty acid methyl ester standards (Sigma-Aldrich, San Leandro, CA) and expressed as a percentage of total fatty acids. All solvents used during lipid extraction and transesterification of phospholipids were of analytical grade (Sigma-Aldrich) with 0.01% butylated hydroxy toluene added as an antioxidant.

Heart perfusion and ventricular pressure measurement. Hearts were excised, cannulated, and perfused at a constant pressure of 80 mmHg in Langendorff mode and allowed to spontaneously beat at 37°C as previously described (20). Hearts were perfused with a filtered (0.45 μ m) modified Krebs-Henseleit buffer containing the following: 119 mmol/l NaCl, 22 mmol/l NaHCO₃, 4 mmol/l KCl, 1.2 mmol/l MgCl₂, 1.2 mmol/l KH₂PO₄, 0.5 mmol/l EDTA, 2.5 mmol/l CaCl₂, 5 mmol/l glucose (BDH, AnalaR), and 100 μ U/ml insulin (Humulin, 100 IU/ml; Eli Lilly, West Ryde NSW, Australia). Perfusate was equilibrated with 95% O₂ and 5% CO₂ (pH 7.4). Baseline contractile function was assessed after a 20-min normoxic equilibration period. Global no-flow ischemia was induced for 15 min followed by 30-min reperfusion.

For assessment of isovolumic contractile function, a small fluid-filled plastic-wrap balloon was inserted into the left ventricle via the mitral valve and inflated to a set diastolic pressure (5 mmHg), and this pressure was maintained throughout the equilibration period (4.0 ± 0.4 mmHg at end of equilibration, i.e. 20-min normoxic perfusion). The parameters measured included intrinsic heart rate (HR, beats/min), systolic pressure (SP, mmHg), diastolic pressure (DP, mmHg), left ventricular developed pressure (LVDevP = SP-DP), rate-pressure product (LVDevP \times HR, RPP mmHg/min), and the maximum positive and negative pressure derivatives (+dP/dt and -dP/dt, mmHg/ms). Coronary flow was measured by timed collection of the coronary effluent and normalized to wet heart weight (ml·min⁻¹·g⁻¹). Myo-

cardial oxygen consumption (M \dot{V} O₂) was determined by measuring the coronary flow rate and the partial pressure of oxygen (Po₂) (ABL 615; Radiometer, Brønshøj, Denmark) of both the perfusate and the coronary effluent at 5-min reperfusion. M \dot{V} O₂ was expressed as mol O₂/min per gram wet weight using the equation: M \dot{V} O₂ = Po₂ (arterial-venous) \times solubility of O₂/mmHg \times coronary flow rate/wet heart weight (39). Hearts were excluded from the study where the baseline cardiac function was unstable, as previously defined in detail (20).

Arrhythmia analysis. The extent of ectopy during the first 10 min of reperfusion was assessed from the ventricular pressure traces, and given by the equation: number of ectopic beats/total beats \times 100. Arrhythmias were also classified by type according to the following designations: ventricular premature beats (VPB; early contraction before relaxation), bigeminy (a variant of a VPB: paired beats occurring in repetition), potentiated contraction (normal sinus rhythm, slight delay, subsequent increased single contraction, resumption of sinus rhythm), ventricular tachycardia (VT; 4 or more consecutive VPBs), and ventricular fibrillation (VF; barely discernable beat, developed pressure <5 mmHg), as previously described (19).

Bilateral ovariectomy. To withdraw gonadal estrogen, mice from each experimental group (i.e., CRT WT and TG and FO WT and TG) underwent bilateral ovariectomy or sham surgery. Mice were anesthetized with isoflurane (2.4% and 200–220 ml airflow; Univentor U400, Zejtun, Malta). A small transverse incision was made to expose the uterus, and the uterine horns were ligated just distal to the ovarian pedicle (5–0 silk; Dynek, London, UK). Ovaries were removed (OVX) and the incision closed by suture. After surgery, mice were monitored daily for 4-wk diet treatment period until euthanized for isolated heart perfusion experiments.

A period of 4 wk of estrogen withdrawal was employed on the basis of our previous findings in rats that this duration is sufficient to induce significant changes in cardiomyocyte Ca²⁺ kinetics (7). At the conclusion of the 4-wk intervention, uterine weight was used to confirm ovarian estrogen withdrawal. In all mice, ovariectomy was associated with significant uterine atrophy compared with intact mice (109 ± 8 vs. 24 ± 1 mg), and uterine weights were comparable for all OVX groups.

Table 2. Baseline ex vivo functional parameters

		LVDevP, mmHg	+dP/dt, mmHg/s	-dP/dt, mmHg/s	HR, beats/min	Flow, ml·min ⁻¹ ·g ⁻¹
WT	CTR	163±6	5089±537	-3689±194	347±18	15.6±1.5
	FO	153±11	5554±422	-3694±164	348±17	16.3±1.5
	CTR	143±7	4871±417	-3723±255	371±23	15.0±1.6
TG	FO	150±10	4643±365	-3630±214	353±22	16.3±0.9
	WT vs. TG	0.199	0.408	0.700	0.454	0.828
P	CTR vs. FO	0.839	0.706	0.892	0.672	0.484
	interaction	0.344	0.553	0.876	0.639	0.868

Data are presented as means ± SE ($n = 8$ /group). All parameters were measured after 20-min normoxic perfusion (equilibration). 2-way ANOVA, P , probability. LVDevP, left ventricular developed pressure; +dP/dt, rate of pressure development; -dP/dt, rate of pressure decline. HR, heart rate.

Statistical analyses. Data are expressed as means \pm SE. Group sizes for cardiac weight index determinations and for functional analyses ranged from $n = 8-12$, and for membrane analyses, $n = 3-4$ (these data exhibit low variance). Data were analyzed using two-way ANOVA and with repeated measures where appropriate. Multiple-comparisons post hoc analysis was performed using Tukey HSD. Arrhythmia duration data were analyzed with a nonparametric Kruskal-Wallis test and incidence analyzed with the Fisher's exact

test. All data were considered significant at $P < 0.05$. Statistical calculations were performed using the SPSS V.15.0 (SPSS, Chicago, IL).

RESULTS

Somatic growth, cardiac growth, and ventricular membrane fatty acid composition. Body mass and food consumption were tracked throughout the 4-wk diet treatment period. The mean




Fig. 2. Effects of 4-wk dietary FO supplementation compared with CTR diet in WT (circles) and TG (triangles) on ex vivo contractile recovery following 15-min global no-flow ischemia. Time course shows basal (time point B), which was measured after 20-min equilibration and time points at 5-min intervals during 30-min reperfusion. Data are presented as means \pm SE ($n = 8$ /group), analyzed by 2-way ANOVA (factors, genetic type, and diet) with repeated measures. A: developed pressure: WT vs. TG, $P = 0.581$; CTR vs. FO, $P = 0.146$; interaction factor, $P = 0.596$. B: baseline normalized developed pressure: WT vs. TG, $P = 0.304$; CTR vs. FO, $P = 0.230$; interaction factor, $P = 0.394$. C: rate pressure product (RPP, developed pressure \times heart rate): WT vs. TG, $P = 0.494$; CTR vs. FO, $P = 0.155$; interaction factor, $P = 0.215$. D: baseline normalized RPP: WT vs. TG, $P = 0.353$; CTR vs. FO, $P = 0.366$; interaction factor, $P = 0.278$. E: rate of contraction (+dP/dt): WT vs. TG, $P = 0.518$; CTR vs. FO, $P = 0.129$; interaction factor, $P = 0.434$. F: baseline normalized rate of contraction: WT vs. TG, $P = 0.283$; CTR vs. FO, $P = 0.179$; interaction factor, $P = 0.221$. G: heart rate: WT vs. TG, $P = 0.503$; CTR vs. FO, $P = 0.358$; interaction factor, $\dagger P = 0.026$. H: baseline normalized heart rate: WT vs. TG, $P = 0.918$; CTR vs. FO, $P = 0.721$; interaction factor, $P = 0.072$.

food intake in g/mouse per day was not different across diet group (CTR diet 3.1 ± 0.3 g, FO diet 2.8 ± 0.3 g, $P = 0.459$). At the end of the diet treatment period, body mass was similar between all groups (Fig. 1A). Heart mass and the normalized cardiac weight index (heart mass normalized to body mass) were significantly greater for TG hearts compared with WT hearts in both dietary groups ($P < 0.001$, $n = 8$ /group; Fig. 1, B and C). FO diet was associated with smaller normalized heart mass for both TG (by 10%) and WT hearts (by 5%) compared with strain-matched CTR diet-fed mice ($P = 0.014$, Fig. 1C) but no significant diet and genetic type interaction was detected (Fig. 1C).

Four weeks of dietary FO markedly elevated the omega-3:omega-6 PUFA ratio of both WT and TG (Table 1). This effect was primarily achieved through decreased ventricular membrane linoleic and arachidonic acid and elevated EPA and DHA. For FO-fed TG mice there was a small but selectively greater increase in DHA and in the omega-3:omega-6 PUFA ratio compared with diet-matched WT (Table 1).

Effects of dietary FO on cardiac performance pre- and postischemia. Ex vivo isovolumic mechanical function, coronary flow, and $\dot{M}\dot{V}O_2$ were measured pre- and postischemia in hearts of WT and TG mice ($n = 8$ /group). Under basal conditions (normoxia), there were no significant differences between any of the groups (Table 2). Global ischemia (15 min) rapidly reduced contractile function in all of the groups, and all hearts had resumed mechanical function by the end of reperfusion (30 min), albeit at a reduced level compared with baseline (Fig. 2). The time course of postischemic recovery throughout reperfusion is shown in Fig. 2 and depicted as nonnormalized and baseline-normalized values. For the non-normalized data, postischemic developed pressure (Fig. 2A), the RPP (Fig. 2C), rate of contraction (+dP/dt, Fig. 2E), and rate of pressure decline (-dP/dt, data not shown) were all not significantly different between WT and TG hearts or between the CTR- and FO-fed hearts. For intrinsic HR (Fig. 2G), a significant interaction between genetic type and diet ($P = 0.026$) was detected, indicating that the FO diet had a differential effect in WT (increased HR) and TG (lowered HR) hearts.

Normalization of postischemic function to preischemic "baseline" values (Fig. 2, right) did not reveal any further significant differences across any of the groups. Hearts of CTR-fed TG appeared to recover a slightly higher level of developed pressure, RPP, rate of contraction (+dP/dt) (Fig. 2, B, D, and F), and rate of pressure decline (-dP/dt, data not shown) compared with all other groups, but this effect was not significant for any parameters (interaction $P = 0.394$, $P = 0.366$, $P = 0.221$, $P = 0.209$ for each parameter, respectively). Baseline-normalized intrinsic HR followed a similar pattern to the nonnormalized data, but the interaction effect (differential effect of FO diet in TG and WT) did not quite attain statistical significance ($P = 0.072$, Fig. 2H).

Coronary flow and myocardial oxygen consumption were markedly reduced postischemia (compared with baseline), by ~10–30% (Fig. 3 vs. Table 2). Dietary FO did not alter coronary regulation for either WT or TG mice, compared with genetically matched CTR-fed mice (Fig. 3A). Myocardial oxygen consumption paralleled coronary flow (Fig. 3B), indicating a direct relationship between oxygen delivery and extraction for hearts of all groups.





Fig. 3. Effects of 4-wk dietary FO supplementation (compared with CTR) in WT and TG on ex vivo cardiac performance during early reperfusion following 15-min global no-flow ischemia. A: coronary flow: WT vs. TG, $P = 0.791$; CTR vs. FO, $P = 0.754$; interaction factor, $P = 0.361$. B: myocardial oxygen consumption ($\dot{M}\dot{V}O_2$): WT vs. TG, $P = 0.661$; CTR vs. FO, $P = 0.681$; interaction factor, $P = 0.479$.

Effects of dietary FO on the performance of the estrogen-withdrawn heart pre- and postischemia. To determine whether circulating ovarian hormones were masking a protective effect of dietary FO, a separate cohort of WT and TG mice underwent ovariectomy to eliminate gonadal supply of sex steroids during the 4-wk diet treatment period (surgery success was confirmed by uterine atrophy). Hearts were excised and cannulated, and mechanical function was measured ex vivo. Hearts were weighed at the end of the perfusion protocol (CTR WT/TG, 9.7 ± 0.2 , 10.5 ± 0.3 mg/g; FO WT/TG, 9.4 ± 0.4 , 10.1 ± 0.6 mg/g, 2-way ANOVA $P > 0.05$). Postischemia, a variable and marked extent of myocardial tissue edema, contributes to the error in determining cardiac weight index. Although mean cardiac weight index was higher in TG (vs. WT) and lower in FO-fed mice (vs. CTR-fed), significant differences could not be resolved in postperfused hearts of OVX groups.

Under basal conditions, mechanical function of hearts from OVX mice was similar between WT and TG and between the CTR and FO diet groups for the range of parameters investigated: developed pressure CTR WT, 138 ± 7 vs. CTR TG, 139 ± 7 mmHg; FO WT, 163 ± 9 vs. FO TG, 149 ± 10 mmHg; rate pressure product; and rate of contraction (Fig. 4, A and C, at time point B, $P > 0.05$, $n = 8$ –12/group). Overall, OVX TG hearts (i.e., both diet groups) exhibited lower basal coronary flow and myocardial oxygen consumption compared

Fig. 4. Effects of 4-wk dietary FO supplementation (compared with CTR) in ovariectomized (OVX) WT (circles) and TG (triangles) on ex vivo contractile recovery and coronary flow following 15-min global no-flow ischemia. Time course shows basal (*time point B*), which was measured after 20-min equilibration and time points at 5-min intervals during 30-min reperfusion. Data are presented as means \pm SE ($n = 8-12$ /group). Two-way ANOVA (factors, genetic type, and diet) with repeated measures. A: RPP (developed pressure \times heart rate): WT vs. TG, $*P = 0.016$; CTR vs. FO, $P = 0.335$; interaction factor, $P = 0.615$. B: baseline normalized rate pressure product: WT vs. TG, $*P = 0.014$; CTR vs. FO, $P = 0.175$; interaction factor, $P = 0.873$. C: rate of contraction (+dP/dt): WT vs. TG, $*P = 0.005$; CTR vs. FO, $P = 0.341$; interaction factor, $P = 0.417$. D: baseline normalized rate pressure product: WT vs. TG, $*P = 0.003$; CTR vs. FO, $P = 0.207$; interaction, $P = 0.813$.



with hearts of OVX WT (CTR 13.8 ± 1.1 vs. 16.5 ± 1.0 $\text{ml} \cdot \text{min}^{-1} \cdot \text{g}^{-1}$ wet wt, $P = 0.009$; 7.4 ± 0.5 vs. 8.8 ± 0.4 $\mu\text{mol O}_2 \cdot \text{min}^{-1} \cdot \text{g}^{-1}$ wet wt, $P = 0.006$).

The time course of postischemic recovery throughout reperfusion (of OVX hearts) is shown in Fig. 4 and depicted as nonnormalized and baseline-normalized values. Overall throughout reperfusion, OVX TG hearts (from both diet groups) exhibited enhanced recovery of RPP product and rate of pressure development (+dP/dt), by $\sim 25\%$ (Fig. 4, A and C), compared with OVX WT hearts. Dietary FO did not enhance mechanical recovery of OVX WT or TG hearts for any of the parameters measured, including RPP (Fig. 4A), rate of pressure development (+dP/dt, Fig. 4C), and developed pressure or rate of pressure decline ($-dP/dt$) (data not shown). In the baseline-normalized data (Fig. 4, right), a similar pattern to the nonnormalized values was observed. In contrast to the ovary-intact mice (Fig. 2), FO diet did not significantly alter intrinsic HR of WT or TG hearts. Similar to the hearts of ovary-intact mice (Fig. 3), dietary FO did not significantly alter coronary regulation for either WT or TG mice, compared with genetic-matched CTR-fed mice (Fig. 5A). Myocardial oxygen consumption paralleled coronary flow (Fig. 5B).

Interestingly, in general, ovariectomy was associated with an apparent increase in the propensity for mechanical alternans (beat-to-beat alterations in twitch amplitude ≥ 5 mmHg at a constant HR) (19). Thirty-one percent of hearts ($n = 13/41$) from the OVX groups displayed alternans compared with only 6% of hearts ($n = 2/32$) from the ovary-intact groups, and notably this vulnerability was not detected at all in the ovary-intact WT groups ($P = 0.001$). Furthermore, separation of the WT hearts by diet groups revealed that the incidence of alternans in CTR WT was markedly greater in the OVX hearts ($n = 6/10$) than in the ovary-intact group, where there was no occurrence ($n = 0/8$, $P = 0.01$). This difference was not

significant in the FO-fed WT OVX (OVX $n = 3/8$ vs. intact $n = 0/8$).

Dietary FO supplementation and postischemic arrhythmia. Arrhythmia incidence was evaluated during the first 10 min of reperfusion from the ventricular pressure record and classified as previously described (19). In the ovary-intact and OVX groups, no significant differences in the incidence of VPB, VT (4 or more consecutive premature beats), potentiated contraction, or the duration of bigeminy and VF were detected between WT and TG hearts. Hearts of FO-fed mice did not exhibit an altered incidence of these arrhythmia types in either the ovary-intact or OVX groups (Table 3).

To obtain an aggregated "snap shot" of arrhythmia incidence, the percentage ectopy was determined by expressing the number of arrhythmic beats as a percentage of total beats. Overall (i.e., from both diet groups), TG hearts of both ovary-intact (Fig. 5A) and OVX groups (Fig. 5B) exhibited a smaller percentage ectopy compared with WT hearts. Compared with mice fed the CTR diet, dietary FO had no effect on the extent of ectopy exhibited in the reperfused hearts in both ovary-intact (Fig. 6A) and OVX groups (Fig. 6B).

DISCUSSION

This is the first experimental study to address the question of whether dietary omega-3 PUFA intervention confers cardioprotection in the ischemic female heart. The major findings of this study are that, in the female murine heart, dietary FO 1) modestly suppresses hypertrophic growth 2) does not enhance postischemic recovery in the normal or the hypertrophic heart, 3) does not enhance postischemic recovery in the normal or hypertrophic hearts of estrogen-depleted mice, and 4) does not lower the incidence of reperfusion arrhythmia in the normal or hypertrophic heart. Thus contrary to our hypothesis, we find




Fig. 5. Effects of 4-wk dietary FO compared with CTR diet in OVX WT (circles) and TG (triangles) on ex vivo cardiac performance during early reperfusion following 15-min global no-flow ischemia. A: coronary flow: WT vs. TG, $P = 0.386$; CTR vs. FO, $P = 0.232$; interaction factor, $P = 0.843$. B: MVO₂: WT vs. TG, $P = 0.129$; CTR vs. FO, $P = 0.134$; interaction factor, $P = 0.763$.

no evidence of a protective effect of dietary FO on myocardial postischemic recovery or arrhythmia suppression in females in the normal heart, the hypertrophic heart, or the estrogen-withdrawn heart. An important finding of this study is that sex should be considered in future investigations of the cardiac mechanisms of action of omega-3 PUFA.

Dietary FO modulates myocardial membrane composition and growth. The finding that dietary FO suppresses heart growth is important. FO-fed TG hearts had a greater proportion of membrane-incorporated omega-3 DHA (22:6 omega-3) compared with WT hearts, which may be linked to hypertrophic growth suppression. There have been only two other (published) experimental studies that have investigated the effects of dietary omega-3 PUFA on cardiac hypertrophic growth, and these studies have been in male rodents. Duda et al. (11) reported that, in a rat model of pressure overload, the increase in left ventricular mass was less marked in the FO-fed group compared with rats fed the contrast diet (rich in saturated fat). A FO diet-induced reduction in ventricular mass in a murine model of systemic carnitine deficiency (JVS mice) where cardiac enlargement occurs secondary to lipotoxicity has also been reported (45). The mechanisms leading to regression of cardiac hypertrophy in vivo remain to be elucidated. Takahashi et al. (45) observed that dietary FO prevented the membrane translocation of several cytosolic protein kinase C isozymes and suggested that this may be the mechanism for the hypertrophy suppression in the JVS mice. In vitro studies in isolated neonatal rat cardiac myocytes (sex of rats not reported) have demonstrated that DHA and EPA inhibit the growth response (i.e., protein synthesis) induced by phenylephrine and endothelin-1 stimulation via the Ras/Raf/Erk1/2 and JNK signaling pathways (41, 42). In the present study, the growth reduction effect of FO diet was observed without a prior hemodynamic loading stimulus and therefore likely reflects a direct modulation of intrinsic myocardial trophic maintenance signaling. It has been proposed that in females adaptive hypertrophic growth plays a more important role in systolic functional maintenance in compromised situations (36). A detailed molecular understanding is required to discern whether growth suppression associated with omega-3 intervention in females is cardioprotective.

Postischemic recovery in TG females is not undermined. The hypertrophic TG female hearts did not exhibit contractile dysfunction under basal conditions (when compared with WT),

Table 3. Type and incidence of reperfusion arrhythmias in hearts of CTR-fed and FO-fed intact and ovariectomized WT and TG mice

	WT		TG	
	CTR	FO	CTR	FO
<i>Intact</i>	<i>n</i> = 8	<i>n</i> = 8	<i>n</i> = 8	<i>n</i> = 8
VPB (incidence)	104 ± 16 (8)	66 ± 14 (8)	77 ± 29 (8)	62 ± 18 (8)
VT (incidence)	2 ± 1 (3)	ND	ND	ND
potentiated (incidence)	5 ± 2 (8)	18 ± 5 (7)	13 ± 3 (7)	29 ± 25 (5)
bigeminy (seconds)	82 ± 21 (8)	61 ± 22 (6)	39 ± 8 (7)	46 ± 26 (5)
VF (seconds)	51 (1)	108 (1)	ND	ND
<i>Ovariectomized</i>	<i>n</i> = 10	<i>n</i> = 8	<i>n</i> = 12	<i>n</i> = 11
VPB (incidence)	70 ± 9 (10)	89 ± 24 (8)	62 ± 11 (12)	67 ± 13 (11)
VT (incidence)	2 ± 1 (3)	9 (1)	ND	ND
potentiated (incidence)	25 ± 11 (6)	20 ± 8 (6)	24 ± 16 (7)	21 ± 13 (6)
bigeminy (seconds)	56 ± 18 (8)	36 ± 16 (5)	55 ± 17 (8)	55 ± 15 (7)
VF (seconds)	344 ± 120 (2)	174 (1)	ND	ND

Assessment of ex vivo arrhythmia during the first 10 min of reperfusion. Arrhythmias were identified from the ventricular pressure record, using the criteria described in MATERIALS AND METHODS. Numbers in parentheses indicate the number of hearts exhibiting the type of arrhythmia (SE is omitted where this number is 1 only). Ventricular premature beat (VPB), ventricular tachycardia (VT), and potentiated contraction are expressed as number of events (incidence) occurring during the first 10 min of reperfusion and shown as means ± SE for the number of hearts in parentheses. Bigeminy and ventricular fibrillation (VF) are presented as duration (seconds), occurring during the first 10 min of reperfusion and shown as means ± SE for the number of hearts in parentheses. No significant differences were detected. ND, not detected.




Fig. 6. Assessment of ex vivo arrhythmia incidence during the first 10 min of reperfusion in WT and TG mice 4 wk after dietary FO supplementation or CTR. Data (means \pm SE; $n = 8$ – 12 /group) are expressed as a percentage ectopy (arrhythmic beats/total beats \times 100). ANOVA (factors, genetic type, and diet). A: hearts of ovary-intact WT and TG mice: WT vs. TG, $*P = 0.030$; CTR vs. FO, $P = 0.299$; interaction factor, $P = 0.947$. B: hearts of OVX WT and TG mice: WT vs. TG, $*P = 0.015$; CTR vs. FO, $P = 0.529$; interaction factor, $P = 0.245$.

which contrasts with our previous observations in male TG of similar age (20). Although our earlier study did not involve a specified diet and comparison, the diet used most closely resembles the CTR diet described here. The present finding accords with other reports in the literature that have shown that the occurrence, onset, and progression of cardiac hypertrophy are frequently different in females compared with males (10). In addition, this study found that the TG female hearts exhibit an intrinsic functional resistance to ischemia and reperfusion (equivalent to WT). Hypertrophic exacerbation of ischemia and reperfusion damage has been previously observed in some experimental models where cardiac growth and function are modified by systemic loading conditions. This study shows that in the female, under these experimental conditions, primary chronic hypertrophy occurring without hemodynamic complication does not impair postischemic mechanical recovery or promote postischemic arrhythmia. This finding is consistent with previous studies in the rat that have shown that female hearts are protected from ischemia-reperfusion, exhibiting lower incidence of arrhythmia, improved recovery of contractility, and decreased necrosis when compared with males (2, 22, 35). These differences are largely attributed to sex steroids although the cellular and molecular mechanisms underlying sex differences in cardiac function remain unresolved (25). The sarcolemmal ATP-sensitive potassium channel may play a role because expression of this channel is estrogen modulated and

pharmacological block has been shown to abolish sex differences in ischemia-reperfusion damage in ex vivo hearts (23).

FO may not confer protection in the female heart. Contrary to our primary hypothesis, the extent of postischemic mechanical recovery was found not to be enhanced in the hearts of FO-fed TG (or WT), when compared with CTR-fed mice. Dietary FO, which is generally considered to be cardioprotective (29, 31, 39), albeit not always (4), had no effect on postischemic mechanical recovery or arrhythmia suppression (VPB, VT, VF) in WT hearts. In younger animals fed diets identical to those used in the present study, we have previously identified a postischemic protective effect of FO in WT and TG males (21). In the present study, hearts of FO-fed TG hearts did exhibit a lower intrinsic HR during reperfusion compared with TG CTR-fed mice. This finding is consistent with previous studies in the rat and human (1) and has been suggested to be beneficial since a lower resting HR reduces risk of arrhythmia. However, surprisingly in this study we did not find an antiarrhythmic effect of dietary FO on postischemia arrhythmia incidence in the TG (or WT) hearts when compared with CTR-fed TG.

Review of the literature provides a basis for inferring that omega-3 may act at similar cellular targets to estrogen (7, 18, 27, 33, 47). Dietary omega-3 PUFA and estrogen have been both observed to modulate cardiomyocyte ion homeostasis, particularly the ionic fluxes involved in excitation-contraction coupling, and have been linked with intracellular Ca^{2+} -dependent downstream signaling. In addition, omega-3 PUFA and estrogen may confer benefit by directly targeting the genome (i.e., bind to nuclear transcription factors). In the estrogen-replete female, activation and transcription of genes involved with cardiac remodeling, cellular repair, and mitochondrial biogenesis may already be invoked (by estrogen), hence precluding the potential for further omega-3 PUFA action. Thus we advanced a second hypothesis that the effects of dietary FO may be masked in the WT and TG hearts attributable to the already “protected” state of the estrogen-replete female.

Withdrawal of ovarian sex hormones does not unmask a cardioprotective action of dietary FO. To evaluate whether systemic estrogen was “preempting” a protective effect of omega-3 PUFA (i.e., the effects of both estrogen and omega-3 PUFA are not additive), we compared the responses of OVX WT and TG mice to FO dietary intervention. It has been previously established that a 4-wk period of experimental estrogen depletion by ovariectomy is sufficient to reverse the functional cardiomyocyte modeling conferred by endogenous estrogen (7). OVX CTR-fed WT hearts exhibited a distinct vulnerability to ischemia, displaying a markedly higher incidence of mechanical alternans suggestive of abnormal calcium handling (5, 12). Data subanalysis suggests that FO may selectively influence alternans occurrence in WT-OVX. Although mechanical alternans occurrence cannot be strictly considered of arrhythmogenic origin without additional characterization, this observation does provide a possible avenue for further cellular investigations. Previous dietary and acute (in vitro) studies (in males) have indicated that omega-3 PUFA confers arrhythmia protection through favorably modulating intracellular calcium flux (18, 32, 47). Our finding was that dietary FO did not enhance postischemic mechanical recovery in the estrogen-depleted WT or TG hearts. Moreover, there was no beneficial effect of dietary FO on the incidence of reperfusion arrhythmia.

These findings, in particular the lack of effect on reperfusion arrhythmias, represent intriguing experimental outcomes given the consistent (but not universal) reports in the literature that dietary FO-derived omega-3 PUFA are antiarrhythmic in male rodents following ischemia and reperfusion (29, 31, 39). This unanticipated finding adds to the emerging evidence that omega-3 PUFA may not be antiarrhythmic in all contexts. Given that the diets fabricated for this study were matched for total PUFA content, these findings may suggest that, in females, both omega-3 and omega-6 PUFA classes exert similar effects. The absence of a cardioprotective FO effect in the estrogen-deplete female in this study prompts consideration of the proposition that testosterone is important in males in mediating the actions of omega-3 intervention. There is evidence to suggest that testosterone may reduce the activities of the enzymes involved in the desaturation process of the 18-carbon chain PUFA to the longer more unsaturated PUFA (28). Thus supplementation studies with long-chain omega-3 PUFA (e.g., DHA and EPA) may not achieve the same outcome in females because of relatively greater intrinsic capacity to produce these PUFA endogenously.

An additional finding of this study was that postischemia differences in the extent of mechanical recovery were evident in OVX TG (compared with intact) that were not apparent in the OVX WT (Fig. 4C vs. Fig. 2F). In these TG hearts, ovariectomy was associated with a higher level of intrinsic contractility (i.e., $+dP/dt$) postischemia compared with genotype-matched intact mice. Thus our findings suggest that the absence of circulating sex hormones impacts differently on the myocardium when there is a chronic local activation of RAS. Previous studies have reported an Ang II-estrogen interaction (38, 48). Chronic infusion with Ang II was associated with a greater pressor effect in OVX females and males compared with intact females, indicating a protective role of ovarian hormones against Ang II-induced hypertension (48). The finding that postischemic excitation-contraction coupling is altered suggests that in vivo the local cardiac growth mediator milieu may interact with systemic estrogen to modulate excitation-contraction coupling.

Previous studies have shown that Ang II stimulates increases in myocyte cytosolic free Ca^{2+} by activating voltage-sensitive Ca^{2+} channels, producing a subsequent increase in contractility (13). Ang II modulates beat-to-beat Ca^{2+} involved in excitation-contraction coupling, and an Ang II-mediated positive inotropic effect on isolated rat myocytes has been demonstrated (8). The positive inotropic effects of Ang II have been attributed to an increase in the Ca^{2+} transient with (15) or without (46) an increase in myofilament sensitivity to Ca^{2+} . In contrast, it has been shown that estrogen suppresses cellular Ca^{2+} flux (6, 7, 33). Thus, in the TG, the removal of ovarian estrogen would be expected to enhance intracellular Ca^{2+} flux, and, in a relatively low workload environment (as in this study, with a modest ischemic insult), an augmented Ca^{2+} flux may improve inotropic state—conferring relative “hypercontractility”. However, more severe in vivo ischemic or workload demands could be expected to unmask mechanical deficits associated with higher resting intracellular Ca^{2+} . In such circumstances, it might be predicted that Ang II hypercontractility postischemia undermines long-term mechanical stability and recovery. Interestingly, in a different pathological setting of a disease model of familial hypertrophic cardiomyopathy, dietary phytoestrogen enrichment (soy intake) has been shown

to exacerbate mechanical dysfunction (44). In this context, it may be that high levels of ingested exogenous estrogens undermine the supply of activator Ca^{2+} for excitation-contraction coupling and impair the adaptive response to counter genetic abnormality. Further work is required to understand the effects of exogenous and endogenous estrogens on Ca^{2+} modulation in hypertrophic disease states. The importance of using fully specified diets rather than diets of regionally and seasonally variable phytoestrogen content is also increasingly apparent.

In summary, this study provides the first experimental evaluation of the effects of dietary FO in modulating the myocardial responses of female hearts to ischemic challenge. Our findings indicate a lack of selective cardioprotective effect of FO dietary intervention determined by assessment of mechanical and arrhythmic activity in the murine ex vivo heart model. Furthermore, relative to the CTR diet, FO intervention was without functional benefit even in the presence of cardiac hypertrophy (induced by cardiac Ang II overproduction) or in the context of chronic systemic estrogen withdrawal (surgically simulated “menopause”). Interestingly, FO diet exerted a significant (modest) cardiac growth suppression effect, which was not selective for hypertrophic hearts. These novel experimental findings highlight the importance of gaining a more complete understanding of the role of omega-3 PUFA in regulating growth and function specifically in the female heart. The assumed benefits of omega-3 intervention/supplementation which have been largely identified in male cohort studies may not necessarily translate to females.

ACKNOWLEDGMENTS

The tuna fish oil was a gift from Nu-Mega Ingredients (Australia).

GRANTS

This work was supported by research and career (C. Huggins and C. Curl) support provided by the National Heart Foundation Australia and the National Health and Medical Research Council of Australia.

REFERENCES

1. **Abdukeyum GG, Owen AJ, McLennan PL.** Dietary (n-3) long-chain polyunsaturated fatty acids inhibit ischemia and reperfusion arrhythmias and infarction in rat heart not enhanced by ischemic preconditioning. *J Nutr* 138: 1902–1909, 2008.
2. **Bell JR, Porrello ER, Huggins CE, Harrap SB, Delbridge LM.** The intrinsic resistance of female hearts to an ischemic insult is abrogated in primary cardiac hypertrophy. *Am J Physiol Heart Circ Physiol* 294: H1514–H1522, 2008.
3. **Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PM, Elwood PC, Deadman NM.** Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 2: 757–761, 1989.
4. **Coronel R, Wilms-Schopman FJ, Den Ruijter HM, Belterman CN, Schumacher CA, Opthof T, Hovenier R, Lemmens AG, Terpstra AH, Katan MB, Zock P.** Dietary n-3 fatty acids promote arrhythmias during acute regional myocardial ischemia in isolated pig hearts. *Cardiovasc Res* 73: 386–394, 2007.
5. **Cross HR, Lu L, Steenbergen C, Philipson KD, Murphy E.** Overexpression of the cardiac Na^{+}/Ca^{2+} exchanger increases susceptibility to ischemia/reperfusion injury in male, but not female, transgenic mice. *Circ Res* 83: 1215–1223, 1998.
6. **Curl CL, Delbridge LM, Wendt IR.** Sex differences in cardiac muscle responsiveness to Ca^{2+} and L-type Ca^{2+} channel modulation. *Eur J Pharmacol* 586: 288–292, 2008.
7. **Curl CL, Wendt IR, Canny BJ, Kotsanas G.** Effects of ovariectomy and 17 beta-oestradiol replacement on $[Ca^{2+}]_i$ in female rat cardiac myocytes. *Clin Exp Pharmacol Physiol* 30: 489–494, 2003.
8. **Delbridge LM, Morgan TO, Harris PJ.** Effects of endothelin-1 on the contractility of cardiomyocytes from the spontaneously hypertensive rat. *Clin Exp Pharmacol Physiol* 22: 755–762, 1995.

9. Domenighetti AA, Wang Q, Egger M, Richards SM, Pedrazzini T, Delbridge LM. Angiotensin II-mediated phenotypic cardiomyocyte remodeling leads to age-dependent cardiac dysfunction and failure. *Hypertension* 46: 426–432, 2005.
10. Du XJ, Fang L, Kiriazis H. Sex dimorphism in cardiac pathophysiology: experimental findings, hormonal mechanisms, and molecular mechanisms. *Pharmacol Ther* 111: 434–475, 2006.
11. Duda MK, O'Shea KM, Lei B, Barrows BR, Azimzadeh AM, McElfresh TE, Hoyt BD, Kop WJ, Stanley WC. Dietary supplementation with omega-3 PUFA increases adiponectin and attenuates ventricular remodeling and dysfunction with pressure overload. *Cardiovasc Res* 76: 303–310, 2007.
12. Dumitrescu C, Narayan P, Efimov IR, Cheng Y, Radin MJ, McCune SA, Altschuld RA. Mechanical alternans and restitution in failing SHHF rat left ventricles. *Am J Physiol Heart Circ Physiol* 282: H1320–H1326, 2002.
13. Freer RJ, Pappano AJ, Peach MJ, Bing KT, McLean MJ, Vogel S, Sperelakis N. Mechanism for the positive inotropic effect of angiotensin II on isolated cardiac muscle. *Circ Res* 39: 178–183, 1976.
14. Friehs I, del Nido PJ. Increased susceptibility of hypertrophied hearts to ischemic injury. *Ann Thorac Surg* 75: S678–S684, 2003.
15. Fujita S, Endoh M. Influence of a Na⁺-H⁺ exchange inhibitor ethylisopropylamiloride, a Na⁺-Ca²⁺ exchange inhibitor KB-R7943 and their combination on the increases in contractility and Ca²⁺ transient induced by angiotensin II in isolated adult rabbit ventricular myocytes. *Naunyn-Schmiedeberg's Arch Pharmacol* 360: 575–584, 1999.
16. GISSI-HF Investigators, Tavazzi L, Maggioni AP, Marchioli R, Barlera S, Franzosi MG, Latini R, Lucci D, Nicolosi GL, Porcu M, Tognoni G. Effect of n-3 polyunsaturated fatty acids in patients with chronic heart failure (the GISSI-HF trial): a randomised, double-blind, placebo-controlled trial. *Lancet* 372: 1223–1230, 2008.
17. GISSI Prevenzione Investigators. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. *Lancet* 354: 447–455, 1999.
18. Honen BN, Saint DA, Laver DR. Suppression of calcium sparks in rat ventricular myocytes and direct inhibition of sheep cardiac RyR channels by EPA, DHA and oleic acid. *J Membr Biol* 196: 95–103, 2003.
19. Huggins CE, Bell JR, Pepe S, Delbridge LM. Bench-marking ventricular arrhythmias in the mouse—revisiting the 'Lambeth Conventions' 20 years on. *Heart Lung Circ* 17: 445–450, 2008.
20. Huggins CE, Domenighetti AA, Pedrazzini T, Pepe S, Delbridge LM. Elevated intracardiac angiotensin II leads to cardiac hypertrophy and mechanical dysfunction in normotensive mice. *J Renin Angiotensin Aldosterone Syst* 4: 186–190, 2003.
21. Huggins CE, McLennan PL, Theiss ML, Pedrazzini T, Pepe S, Delbridge LMD. Sex differences in murine myocardial membrane lipid composition modulate growth and postischemic arrhythmia (Abstract). *J Mol Cell Cardiol* 42: S4, 2007.
22. Humphreys RA, Kane KA, Parratt JR. The influence of maturation and gender on the anti-arrhythmic effect of ischaemic preconditioning in rats. *Basic Res Cardiol* 94: 1–8, 1999.
23. Johnson MS, Moore RL, Brown DA. Sex differences in myocardial infarct size are abolished by sarcolemmal KATP channel blockade in rat. *Am J Physiol Heart Circ Physiol* 290: H2644–H2647, 2006.
24. Kahan T, Bergfeldt L. Left ventricular hypertrophy in hypertension: its arrhythmogenic potential. *Heart* 91: 250–256, 2005.
25. Konhilas JP, Leinwand LA. The effects of biological sex and diet on the development of heart failure. *Circulation* 116: 2747–2759, 2007.
26. Korotkova M, Gabriellsson BG, Holmang A, Larsson BM, Hanson LA, Strandvik B. Gender-related long-term effects in adult rats by perinatal dietary ratio of n-6/n-3 fatty acids. *Am J Physiol Regul Integr Comp Physiol* 288: R575–R579, 2005.
27. Kravtsov GM, Kam KW, Liu J, Wu S, Wong TM. Altered Ca²⁺ handling by ryanodine receptor and Na⁺-Ca²⁺ exchange in the heart from ovariectomized rats: role of protein kinase A. *Am J Physiol Heart Circ Physiol* 292: C1625–C1635, 2007.
28. Marra CA, de Alaniz MJ. Influence of testosterone administration on the biosynthesis of unsaturated fatty acids in male and female rats. *Lipids* 24: 1014–1019, 1989.
29. Mathan NR, Jordan H, Chung M, Lichtenstein AH, Lathrop DA, Lau J. A systematic review and meta-analysis of the impact of omega-3 fatty acids on selected arrhythmia outcomes in animal models. *Metabolism* 54: 1557–1565, 2005.
30. Mazzolai L, Nussberger J, Aubert JF, Brunner DB, Gabbiani G, Brunner HR, Pedrazzini T. Blood pressure-independent cardiac hypertrophy induced by locally activated renin-angiotensin system. *Hypertension* 31: 1324–1330, 1998.
31. McLennan PL. Relative effects of dietary saturated, monounsaturated, and polyunsaturated fatty acids on cardiac arrhythmias in rats. *Am J Clin Nutr* 57: 207–212, 1993.
32. McLennan PL, Abeywardena MY. Membrane basis for fish oil effects on the heart: linking natural hibernators to prevention of human sudden cardiac death. *J Membr Biol* 206: 85–102, 2005.
33. Meyer R, Linz KW, Surges R, Meinardus S, Vees J, Hoffmann A, Windholz O, Grohe C. Rapid modulation of L-type calcium current by acutely applied estrogens in isolated cardiac myocytes from human, guinea-pig and rat. *Exp Physiol* 83: 305–321, 1998.
34. Mozaffarian D. Fish and n-3 fatty acids for the prevention of fatal coronary heart disease and sudden cardiac death. *Am J Clin Nutr* 87: 1991S–1996S, 2008.
35. Nikolic I, Liu D, Bell JA, Collins J, Steenbergen C, Murphy E. Treatment with an estrogen receptor-beta-selective agonist is cardioprotective. *J Mol Cell Cardiol* 42: 769–780, 2007.
36. Olsson MC, Palmer BM, Leinwand LA, Moore RL. Gender and aging in a transgenic mouse model of hypertrophic cardiomyopathy. *Am J Physiol Heart Circ Physiol* 280: H1136–H1144, 2001.
37. Owen AJ, Peter-Przyborowska BA, Hoy AJ, McLennan PL. Dietary fish oil dose- and time-response effects on cardiac phospholipid fatty acid composition. *Lipids* 39: 955–961, 2004.
38. Pedram A, Razandi M, Aitkenhead M, Levin ER. Estrogen inhibits cardiomyocyte hypertrophy in vitro. Antagonism of calcineurin-related hypertrophy through induction of MCIPI. *J Biol Chem* 280: 26339–26348, 2005.
39. Pepe S, McLennan PL. Cardiac membrane fatty acid composition modulates myocardial oxygen consumption and postischemic recovery of contractile function. *Circulation* 105: 2303–2308, 2002.
40. Saito Y, Yokoyama M, Origasa H, Matsuzaki M, Matsuzawa Y, Ishikawa Y, Oikawa S, Sasaki J, Hishida H, Itakura H, Kita T, Kitabatake A, Nakaya N, Sakata T, Shimada K, Shirato K. Effects of EPA on coronary artery disease in hypercholesterolemic patients with multiple risk factors: sub-analysis of primary prevention cases from the Japan EPA Lipid Intervention Study (JELIS). *Atherosclerosis* 200: 135–140, 2008.
41. Shimamoto N, Jesmin S, Zaedi S, Maeda S, Soma M, Aonuma K, Yamaguchi I, Miyauchi T. Eicosapentaenoic acid prevents endothelin-1-induced cardiomyocyte hypertrophy in vitro through the suppression of TGF-β 1 and phosphorylated JNK. *Am J Physiol Heart Circ Physiol* 291: H835–H845, 2006.
42. Siddiqui RA, Shaikh SR, Kovacs R, Stillwell W, Zaloga G. Inhibition of phenylephrine-induced cardiac hypertrophy by docosahexaenoic acid. *J Cell Biochem* 92: 1141–1159, 2004.
43. Skavdahl M, Steenbergen C, Clark J, Myers P, Demianenko T, Mao L, Rockman HA, Korach KS, Murphy E. Estrogen receptor-β mediates male-female differences in the development of pressure overload hypertrophy. *Am J Physiol Heart Circ Physiol* 288: H469–H476, 2005.
44. Stauffer BL, Konhilas JP, Luczak ED, Leinwand LA. Soy diet worsens heart disease in mice. *J Clin Invest* 116: 209–216, 2006.
45. Takahashi R, Okumura K, Asai T, Hirai T, Murakami H, Murakami R, Numaguchi Y, Matsui H, Ito M, Murohara T. Dietary fish oil attenuates cardiac hypertrophy in lipotoxic cardiomyopathy due to systemic carnitine deficiency. *Cardiovasc Res* 68: 213–223, 2005.
46. Vila Petroff MG, Mattiazzi AR. Angiotensin II and cardiac excitation-contraction coupling: questions and controversies. *Heart Lung Circ* 10: 90–98, 2001.
47. Xiao YF, Gomez AM, Morgan JP, Lederer WJ, Leaf A. Suppression of voltage-gated L-type Ca²⁺ currents by polyunsaturated fatty acids in adult and neonatal rat ventricular myocytes. *Proc Natl Acad Sci USA* 94: 4182–4187, 1997.
48. Xue B, Pamidimukkala J, Hay M. Sex differences in the development of angiotensin II-induced hypertension in conscious mice. *Am J Physiol Heart Circ Physiol* 288: H2177–H2184, 2005.
49. Yokoyama M, Origasa H, Matsuzaki M, Matsuzawa Y, Saito Y, Ishikawa Y, Oikawa S, Sasaki J, Hishida H, Itakura H, Kita T, Kitabatake A, Nakaya N, Sakata T, Shimada K, Shirato K. Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomised open-label, blinded endpoint analysis. *Lancet* 369: 1090–1098, 2007.