

2012

## The human foveal confluence and high resolution fMRI

Mark M. Schira

*University of Wollongong*, [mschira@uow.edu.au](mailto:mschira@uow.edu.au)

Follow this and additional works at: <https://ro.uow.edu.au/sspapers>



Part of the [Education Commons](#), and the [Social and Behavioral Sciences Commons](#)

---

### Recommended Citation

Schira, Mark M., "The human foveal confluence and high resolution fMRI" (2012). *Faculty of Social Sciences - Papers*. 254.

<https://ro.uow.edu.au/sspapers/254>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: [research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

## The human foveal confluence and high resolution fMRI

### Abstract

Abstract presented at the Vision Sciences Society meeting, 11-16 May 2012, Naples FL.

### Keywords

foveal, human, confluence, high, fmri, resolution

### Disciplines

Education | Social and Behavioral Sciences

### Publication Details

Schira, M. M. (2012). The human foveal confluence and high resolution fMRI. *Journal of Vision*, 12 (9), 1396-1-1396-1.

After remaining terra incognita for 40 years, the detailed organization of the foveal confluence has just recently been described in humans. I will present recent high resolution mapping results in human subjects and introduce current concepts of its organization in human and other primates (Schira et al., J. Nsci, 2009). I will then introduce a new algebraic retino-cortical projection function that accurately models the V1-V3 complex to the level of our knowledge about the actual organization (Schira et al. PLoS Comp. Biol. 2010). Informed by this model I will discuss important properties of foveal cortex in primates. These considerations demonstrate that the observed organization though surprising at first hand is in fact a good compromise with respect to cortical surface and local isotropy, proving a potential explanation for this organization. Finally, I will discuss recent advances such as multi-channel head coils and parallel imaging which have greatly improved the quality and possibilities of MRI. Unfortunately, most fMRI research is still essentially performed in the same old 3 by 3 by 3 mm style - which was adequate when using a 1.5T scanner and a birdcage head coil. I will introduce simple high resolution techniques that allow fairly accurate estimates of the foveal organization in research subjects within a reasonable timeframe of approximately 20 minutes, providing a powerful tool for research of foveal vision.

Meeting abstract presented at VSS 2012