



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Science, Medicine and Health - Papers

Faculty of Science, Medicine and Health

2013

An appraisal of the extent and geomorphological diversity of the coral reefs of the United Kingdom Dependent Territories

S Hamylton

University of Wollongong, shamylto@uow.edu.au

S Andrefouet

Institut Pour Recherche Et Developpement, serge.andrefouet@gmail.com

Publication Details

Hamylton, S. & Andrefouet, S. (2013). An appraisal of the extent and geomorphological diversity of the coral reefs of the United Kingdom Dependent Territories. In C. R.C. Sheppard (Eds.), *Coral Reefs of the United Kingdom Overseas Territories* (pp. 1-11). Dordrecht: Springer.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

An appraisal of the extent and geomorphological diversity of the coral reefs of the United Kingdom Dependent Territories

Abstract

The 14 Dependent Territories governed by the United Kingdom (UK) Foreign and Commonwealth Office include in alphabetic order Anguilla, Ascension Island and Tristan da Cunha, British Antarctic Territory, Bermuda, British Indian Ocean Territory, British Virgin Islands, Cayman Islands, Falkland Islands, Gibraltar, Montserrat, Pitcairn Island, St Helena, South Georgia and South Sandwich Islands, Sovereign Base Areas on Cyprus and the Turks and Caicos Islands (Fig. 1.1). UK governance responsibilities for these territories include the strengthening of democracy, environmental protection, improvement of public services and law enforcement (Oldfield and Sheppard 1997)

Keywords

kingdom, diversity, reefs, geomorphological, territories, dependent, extent, coral, appraisal, united

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Hamylton, S. & Andrefouet, S. (2013). An appraisal of the extent and geomorphological diversity of the coral reefs of the United Kingdom Dependent Territories. In C. R.C. Sheppard (Eds.), *Coral Reefs of the United Kingdom Overseas Territories* (pp. 1-11). Dordrecht: Springer.

An appraisal of the extent and geomorphological diversity of the coral reefs of the United Kingdom Dependent Territories

S. Hamylton¹ and S. Andréfouët²

1. *School of Earth and Environmental Sciences, University of Wollongong, Australia.*

2. *Institut de Recherche pour le Développement, U227 Coreus Noumea, New Caledonia.*

The United Kingdom (UK) governs, through its Dependent Territories, a total coral reef area of 4712 km² making it, in jurisdictional terms, approximately the twelfth reef nation of the world, allowing for uncertainty in reef area estimates. Of the UK Overseas Territories, the reef nations include Anguilla, Turks and Caicos, Bermuda, the British Virgin Islands, the Cayman Islands, the British Indian Ocean Territories and the Pitcairn Islands. This national value is inferred from geomorphological inventory of the reef systems falling within these Territories, conducted using an internally consistent typology generated as part of the Millennium Coral Reef Mapping Project using Landsat 7+ ETM satellite images from the Caribbean, Indian Ocean and South Pacific regions. These Territories host the world's largest atoll, the northernmost reefs of the Atlantic and the most easterly atoll of the Indo-Pacific reef province. In terms of main morphological units, atolls, banks, barrier reefs, fringing reefs and patch reefs are represented.

Keywords: Remote sensing, Millennium Mapping, Landsat

Introduction

The fourteen Dependent Territories governed the United Kingdom (UK) Foreign and Commonwealth Office include in alphabetic order Anguilla, British Antarctic Territory, Bermuda, British Indian Ocean Territory, British Virgin Islands, Cayman Islands, Falkland Islands, Gibraltar, Montserrat, Pitcairn Island, St Helena, Ascension Island and Tristan da Cunha, South Georgia and South Sandwich Islands, Sovereign Base Areas on Cyprus and the Turks and Caicos Islands (Figure 1). UK governance responsibilities for these territories include the strengthening of democracy, environmental protection, improvement of public services and law enforcement (Oldfield and Sheppard, 1997)

Seven of these territories incorporate substantial reef systems, including Anguilla, Bermuda, the British Indian Ocean Territory (also known as the Chagos Islands), the British Virgin Islands, the Cayman Islands, the Pitcairn Islands and the Turks and Caicos Islands. These territories are composed of small remote islands that support a disproportionately large area of reef and associated marine biodiversity. The total reef area inside the Dependent Territories mapped by the Millennium Mapping Project is 4712km², which makes the UK approximately the twelfth reef nation of the World.

At the global scale, current estimates of national or regional reef areas derive from a variety of sources including marine charts, remote sensing satellite images. They provide estimates that vary widely in accuracy. Basic information (reef or non-reef) on the location, extent and geomorphological nature of reefs systems has been compiled in the *World Atlas of Coral Reefs* published by the United Nations Environment Programme World Conservation Monitoring Centre (Spalding et al., 2001). This atlas provided the first global overview of reef distribution, however, the level of detail was largely determined by the variable availability and specification (scale, accuracy, precision) of marine charts across different

reef regions. As a consequence, the inventory is often inconsistent from one area to another. Other areas have since benefited from specific mapping projects that have provided accurate estimates of reef areas, itemized in different categories that follow a predefined typology of reef types and habitats (for instance, for Hawaii see Rohmann, et al. 2005).

Starting in 2004, a global inventory of reef geomorphological diversity and units has produced consistent maps of reef areas worldwide. The data source is made of Landsat 7 Enhanced Thematic Mapper Plus (ETM+) images, completed by occasional Landsat 5 and Aster satellite images. The project, named Coral Reef Millennium Mapping Project (MCRMP) has produced GIS files and electronic atlases for different part of the world. Regional atlases include Islands of the Central and Western Indian Ocean (Andréfouët et al. 2009), Papua New Guinea (Andréfouët et al. 2006) and the French Territories (Andréfouët et al. 2008). Several projects have distributed degraded (both in spatial and thematic resolutions) versions of the MCRMP products, for the Caribbean (Burke et al. 2004) and globally (Burke et al. 2010). The principles used to design the MCRMP typology, as well as the main hierarchical structure are described elsewhere (Andréfouët et al., 2006; Andréfouët, 2011). The typology and products have proved to be relevant in various contexts: geological appraisals (Andréfouët al. 2009), conservation planning (Green et al. 2009), fisheries and food security (Bell et al. 2009) and to enhance the mapping of specific habitats such a seagrass beds (Wabnitz et al. 2008).

This chapter uses MCRMP products to conduct a geomorphological appraisal of the reef systems of the UK Dependent Territories with the following three objectives:

- i.* to identify where the coral reefs of the Dependent Territories are,
- ii.* to generate a consistent and accurate measurement of reef areas, and
- iii.* to delineate the morphological zones associated with these reef systems.




Figure 1. Location of the UK Dependent Territories. Asterisk denotes reef territories.

From a geomorphic perspective, coral reefs are three-dimensional structures that have evolved over geological timescales according to local sea level variations, subsidence, tectonics, hydrodynamic and climate forcing, and dominant living community types. They range in area from 1 – 100 km² in extent (Hopley, 2011). As a result, reefs display a myriad of shapes and structures at a scale that can be resolved by high spatial resolution (1-30 meters) optical spaceborne sensors down to a water depth of about 40 meters in very clear waters. Typical depth penetration limit is around 20-30 meters.

The UK Dependent Territories assessment was carried out using GIS layers generated as part of the *Millennium Coral Reef Mapping Project* (MCRCP). High resolution Landsat 7 ETM+ satellite images of coral reefs were interpreted using segmentation and photo-interpretation techniques to delineate regions belonging to different morphometric groups within a globally applicable typology of 800 classes (Andréfouët et al. 2006). Reef typology refers to the definition of categories of reef objects according to a series of characteristics relevant for a given purpose (Andréfouët, 2011).

The MCRCP hierarchical typology employs 5 hierarchical levels (Andréfouët et al. 2009a):

- Level 1: discriminates between oceanic and continental reefs;
- Level 2: discriminates the main reef complexes. Atolls, banks, uplifted atolls and islands can be both oceanic and continental. Then, patch reefs, barrier reefs, fringing reefs and marginal structures are defined for continental reefs.
- Level 3: discriminates further details within each of the Level 2 blocks that are too numerous to cite here, but include barrier, fringing and patch reefs of islands for instance, as well as different categories of these types: outer barrier, coastal barrier, multiple barrier, faro barrier, etc.; lagoon exposed fringing, ocean-exposed fringing, etc.
- Level 4: defines the geomorphological units discernable on Landsat imagery within each of the previous blocks, including forereef, reef flat, pass, enclosed lagoon, reef island, etc.
- Level 5: combines categories for Levels 1-4 to provide a final typology of 800 classes worldwide, although any single reef complex is likely to include between 1 and 20 classes at most.

This appraisal was primarily conducted at Level 3 of the typological hierarchy, which provided an optimal level of detail to facilitate comparison between the different territories.

In addition to the various geomorphologic attributes, each of the MCRCP polygon shapefiles has an associated “Reef” attribute that denoted whether the geomorphic unit supports significant coral communities (with a 1 value assigned to reefs and 0 assigned to non-reefs). For instance, forereef, reef flat, subtidal reef flat, pass, pinnacle, etc. are considered as Reef, whereas, terrace (i.e., sedimentary areas), lagoon, etc. are not considered part of the coral reef per se. To calculate the overall reef areas in the present assessment, we considered all polygons with a Reef attribute of value 1, for which the the geometry calculator was employed to calculate the area, followed by the summary statistics tool to sum the areas of all the reef polygons.

Reef extent will clearly depend on the definition used: the “Reef” definition used here was largely consistent and compatible with classical definition of coral reefs. For instance, it was compatible with the definition that state that coral reefs are “physical structure which has been built up and continues to grow over decadal time scales, as a result of the accumulation of calcium carbonate laid down by hermatypic corals and other organisms” (Spalding et al. 2001). Other definitions as provided by *The Encyclopedia of Modern Coral Reefs* elaborate further on the geomorphological components of coral reefs, including coral tracts (large areas of indefinite extent) and massive






structures (in basal area and thickness and wave resistance) (Done, 2011). These definitions were also consistent with the MCRMP definition.

Reef area

The reef systems of the Dependent Territories included 19 Level 3 reef classes, covering a total reef area of 4712 km² (this figure relates to the areas that were identified as reef only). These included atolls, barrier reefs, banks, fringing reefs and patch reefs (Table 1).

TERRITORY	AREA MAPPED/ KM²	REEF AREA/ KM²	# CLASSES	FIVE DOMINANT CLASSES
Anguilla	2537	43	7	<i>Shelf slope (89%) Main land (8%) Ocean exposed fringing (2%) Shelf terrace (1%) Bank lagoon (0.4%)</i>
Bermuda	733	340	7	<i>Island lagoon (34%) Outer Barrier complex (32%) Ocean exposed fringing (13%) Intra-lagoon patch reef complex (11%) Main land (8%)</i>
British Indian Ocean Territory	15639	2859	10	<i>Drowned atoll (77%) Atoll lagoon (15%) Atoll rim (4%) Drowned Bank (3%) Bank lagoon (1%)</i>
British Virgin Islands	4500	138	9	<i>Shelf slope (91%) Lagoon exposed fringing (3.5%) Outer barrier reef complex (2.8%) Intra-lagoon patch reef complex (1%) Shelf patch reef complex (0.8%)</i>
Cayman Islands	471	471	9	<i>Main Land (56%) Island lagoon (16%) Coastal Barrier reef complex (7%) Ocean exposed fringing (6%) Shelf terrace (5%)</i>
Turks and Caicos Islands	6885	822	11	<i>Shelf terrace (60%) Main land (14%) Shelf slope (12%) Shelf structure (8%) Coastal barrier reef complex (3%)</i>
Pitcairn Islands	89	39	9	<i>Main land (52%) Shelf slope (20%) Bank barrier (10%) Atoll rim (7%) Ocean exposed fringing (7%)</i>

Table 1 A morphometric summary of the reef areas of the Dependent Territories (N.B. The right hand side column quotes % cover of the five most dominant geomorphic classes and therefore does not sum to 100%).






Figure 2 Geomorphological maps of the reef systems of the UK Dependent Territories. Pie charts indicate % coverage of the geomorphic units for each territory (see key for units).

It can be seen from Figure 3 that in many cases (5 out of 7 nations), the UNEP study delineated more extensive reef areas than the MCRMP. The differences are significant in several instances (reaching 139% in the case of the British Virgin Islands). Similar discrepancies have been reported elsewhere (Wabnitz et al., 2010), illustrating the inherent variability associated with the different mapping approaches.




Figure 3. Reef areas for each of the Dependent Territories compared between the previous reference (World Atlas of Coral Reefs, in blue) and the Millennium Coral Reef Mapping Project (red).

Characteristics of each Territory

For each Territory, we provide the Level 1 and Level 2 MCRMP label, and some information from the literature combined with the new inventory at Level 3.

Turks and Caicos (*Oceanic/Island*)

The limestone islands of the Turks and Caicos Islands group stretch across the northern extent of the relatively small Turks Bank and the much larger Caicos Bank (area 3933 km²). The margins of these banks slope down to a deeper shelf structure at a depth of 20-30m that descends into oceanic water exceeding 4 km depth (Sullivan et al. 1994). Across the northern shore of the Caicos Islands is a coastal barrier reef complex with exposed fringing reefs on the oceanward aspect. Geologically, the subaerial islands that have developed in the Turks and Caicos Islands (area 924 km²) consist of oolitic limestone sediments, with eolianite hills that have developed on the windward shores reaching up to 75 m above sea-level and karst limestone cliffs (Wanless and Dravis, 1989). Of the overall typology, the Turks and Caicos Islands consisted of 11 different classes, with the shelf terrace occupying 60% of the mapped shallow platform.

Anguilla (*Oceanic/Island*)

Anguilla is a low coralline island (area 192 km²), which has developed on a volcanic base, as part of the Lesser Antilles arc, which stretches 800 km across the eastern margin of the Caribbean Sea (Stein et al. 1982). The total area of reef systems mapped for Anguilla was

2537 km² comprised of submerged banks and terraces, a shelf slope and fringing reefs. The fringing reefs have developed along the north and south coast of the island together with a number of offshore cays that support smaller reef platforms. The dominant morphological feature mapped for Anguilla was the submarine shelf it shares with St Martin to the southeast, which occupied 89% of the area of the reef system. Along the northern extent of this Bank are exposed linear segments of fringing reef along the shelf edge. This 17 km-long reef along the southeast coast is considered to be one of the most important unbroken reefs in the eastern Caribbean (Putney 1982).

British Virgin Islands (*Oceanic/Continental island*)

The Virgin Islands constitute the eastern extremity of the Greater Antilles arc and, in administrative terms, the shallow shelf on which the Virgin Islands sit can be subdivided into the US Virgin islands in the lower south western portion of the shelf and the British Virgin Islands on the upper north eastern portion. There are 40 uplifted volcanic islands, small cays, reef platforms and rocks in the group, the largest of which is Tortola (54km²). These rise from the Puerto Rican shelf, which sits at 65 m below sea level. The small shelf patch reef complex has largely developed within a matrix of volcanic uplift around the larger islands across the central shelf area. On the eastern windward side an outer barrier reef complex has developed in association with Anegada, a relatively flat emergent coral limestone platform (altitude 8 m) (Oldfield et al., 1999).

Cayman Islands (*Oceanic/Island*)

The flat, low lying Cayman Islands consist of three islands: Grand Cayman, Cayman Brac and Little Cayman, which sit at the western end of the Greater Antilles group. The islands sit along the Cayman Ridge, which forms the northern margin of the east-west aligned Oriente Transform Fault (Brunt & Davies, 1994). The Cayman Islands have a collective land area of 261km², which supports a series of terraced fringing and barrier reefs upon which spur and groove formations have developed that display considerable variability in structural form in relation to local wave power dynamics around the islands (Roberts, 1974). A total reef area of 126km² is supported by these islands in the form of an outer and coastal barrier reef complex, exposed fringing reef and shelf terrace.

The British Indian Ocean Territory (*Oceanic/Island*)

The British Indian Ocean Territory lies at the southernmost extension of the north-south aligned Chagos-Laccadive Ridge and is composed of a limestone cap a few hundred kilometres thick that has developed over the volcanic remains of the Reunion hotspot (Sheppard & Wells, 1988; Parson & Evans, 2005). The reef systems of the British Indian Ocean Territories are comprised of 2859 km² reef area, which fall into 10 classes related to the major morphological units (Andréfouët et al., 2009). These include five atolls (Blenheim Reef, Diego Garcia, Egmont, Peros Banhos and Salomon), a drowned atoll (Great Chagos Bank, the largest atoll structure in the world at 9210 km²) and three drowned banks (Speakers Bank, Pitt Bank and Centurion Bank). Each of the atolls have substantial lagoons, ranging in size from 11 - 940 km² with carbonate rims of varying degrees of subaerial exposure around their perimeter. All atolls and submerged banks appear to be actively growing reefs (Sheppard & Wells, 1988).

Pitcairn Islands (*Oceanic/Continental island*)

The Pitcairn group is comprised of four widely spaced atolls and islands in the South Pacific Ocean that fall along two geological structural lineations associated with hotspot activity of the clockwise-spreading Pacific plate (Spencer, 1995). These four structures are

Pitcairn Island (a volcanic island), Henderson Island (an uplifted atoll) and two small atolls, Oeno and Ducie. Collectively the islands can be classified into 9 geomorphic units, dominated by emergent volcanic and reef islands, which represent 52% of the area mapped. Pitcairn is a volcanic island that rises 3.5km from the seafloor with a peak that stands 347 m above sea level with continuous fringing reef around it (Benton and Spencer, 1995). The atoll of Henderson Island is a reef-capped volcano that was uplifted as a result of crustal loading by the adjacent Pitcairn volcano (Fosberg et al. 1983; Wells & Jenkins, 1988), giving rise to several unique biodiversity characteristics for which Henderson has been designated a World Heritage Site. Oeno atoll has a marked outer reef rim perimeter, with an island of area 0.7km² that has developed at the centre of the lagoon. Ducie atoll (area 6 km²) is the most easterly atoll of the Indo-Pacific reef province and the southern most atoll of the world, thought to be the surface expression of a field of seamounts (Spencer, 1995).

Bermuda (*Oceanic/Island*)

Bermuda is comprised of 150 isolated coral limestone islands in the Sargasso Sea, western Atlantic Ocean, that have formed along the rim of an extinct submarine volcano approximately 1000 km east of the North Carolina on the continental USA coastline. The extinct volcano sits on top of the Bermuda Platform, a topographic high of the Bermuda Pedestal, a basement that lies in water depths around 75 m (Vacher & Rowe, 1997). The land area (56km²) is predominantly comprised of a network of 10 main islands that are joined by causeways. The extinct volcano rim surrounds a substantial island lagoon (246km²). The Bermuda reef system (total area 677km²) is formed by the most northerly coral reefs in the world, which form a large outer barrier reef structure that encompasses an island lagoon of area 246km².

Further refinements to the reef inventory

Figures 4 and 5 demonstrate the full detail (Level 5) contained in the MCRCP products for the Chagos Islands and Anguila. The additional level of detail is apparent from the number of classes represented at this level, which are 24 and 71 for the Chagos islands and Anguila respectively (as opposed to 10 and 22 at Level 3).




Figure 4. The Chagos Islands (or British Indian Ocean Territory), illustrating the detail included at Level 5 of the MCRCP Geomorphological map.




Figure 5. Anguilla, illustrating the detail included at Level 5 of the MCRCP Geomorphological map.

The identification of 4712km² of reef within the UK Dependent Territories was possible using a remote sensing dataset of the requisite accuracy, resolution, consistency and completeness for consistently delineating shallow reef morphological units. Such consistency is important for regions that span the Atlantic, Indian and Pacific Oceans yet fall under a common governance framework. All of the major (Level 2) morphological reef units identified in the *MCRCP* global typology including atolls, banks, uplifted atolls, islands, patch reefs barrier reefs, fringing reefs and marginal structures are represented in the UK Dependent Territories.

As satellite remote sensing images have become increasingly available at a resolution commensurate with reef landform morphological variability (1 – 100 km²), morphometrics derived from them represent an important source of information for managing both global environmental change and anthropogenic influences on reefs. Assessment of reefs according to morphogenetic phenomena, such as tectonic activity, sea level rise, sediment and hydrodynamics, provides a fundamental basis on which ecological dynamics and the impacts of human activities can be superimposed and understood. To this end, this geomorphological assessment, generated for the first time from consistent images of the UK Dependent territories, presents a useful foundation for the incorporation of morphodynamic information into marine environmental management decisions and policies.

While this study draws on the best available data for the time being, this assessment could be further improved by combining higher specification remote sensing data and targeted ground referencing field campaigns in a coordinated manner across the UK Dependent Territories.

References

- Andréfouët, S, Muller-Karger, F.E. Robinson, J.A. Kranenburg, C.J., Torres-Pulliza, D. Spraggins, S.A and Murch, B. (2006a) Global assessment of modern coral reef extent and diversity for regional science and management applications: a view from space. *Proceedings of 10th International Coral Reef Symposium*, 1732-1745.
- Andréfouët S, Chagnaud N, Chauvin, C., Kranenburg C (2006b) *Atlas of the Papua New Guinea Coral Reefs*, Centre IRD de Nouméa, Nouméa, Nouvelle-Calédonie. CD-ROM.
- Andréfouët S, Chagnaud N, Chauvin, C., Kranenburg C (2008) *Atlas of the French Overseas Coral Reefs*, Centre IRD de Nouméa, Nouméa, Nouvelle-Calédonie. CD-ROM.
- Andréfouët S, Cabioch G, Flamand B, Pelletier B (2009a) A reappraisal of the diversity of geomorphological and genetic processes of New Caledonian coral reefs: a synthesis from optical remote sensing, coring and acoustic multibeam observations. *Coral Reefs* 28:691-707.
- Andréfouët S, Chagnaud N, Kranenburg C (2009b) *Atlas of Western and Central Indian Ocean Coral Reefs*, Centre IRD de Nouméa, Nouméa, Nouvelle-Calédonie. CD-ROM.
- Andréfouët S (2011) Reef typology. In: Hopley D (ed) *Encyclopedia of Modern Coral Reefs*. Springer, Dordrecht, The Netherlands, p 906.
- Bell, J.D., Kronen, M., Vunisea, A., Nash, W., Keeble, G., Demmke, A., Andréfouët S (2009) Planning the use of fish for food security in the Pacific, *Marine Policy* 33: 64-76.
- Benton, T & Spencer, T. (1995) *The Pitcairn Islands: Biogeography, Ecology and Prehistory*, Academic Press.
- Brunt, MA & Davies, JE (1994) *The Cayman Islands: Natural History and Biogeography*, Kluwer Academic publishers, The Netherlands.
- Burke, L., Maidens, J., Spalding, M., Kramer, P., Green, E., Greenhalgh, S., Nobles, H. and Kool, J. (2004) *Reefs at Risk in the Caribbean*, publication of the World Resources Institute.
- Burke, L., Reyntar, K., Spalding, M. and Perry, A (2010) *Reefs at Risk Revisited*, publication of the World Resources Institute.
- Done, T (2011) Coral Reef, Definition. In: Hopley D (ed) *Encyclopedia of Modern Coral Reefs*. Springer, Dordrecht, The Netherlands, p 261-266.
- Fosberg, F.R., Sacht, M.H. & Stoddart, D.R. (1983) Henderson Island (southeastern Polynesia): summary of current knowledge. *Atoll Research Bulletin* 329: 1- 18.
- Green, A., S.E. Smith, G. Lipsett-Moore, C. Groves, N. Peterson, S. Sheppard, P. Lokani, R. Hamilton, J. Almany, J. Aitsi, and L. Bualia (2009) Designing a resilient network of marine protected areas for Kimbe Bay, Papua New Guinea. *Oryx* 43: 488-498.
- Hopley D (ed) (2011) *Encyclopedia of Modern Coral Reefs*. Springer, Dordrecht, The Netherlands.
- Oldfield, S & Sheppard, S. (1997) Conservation of Biodiversity and Research Needs in the UK Dependent Territories, *Journal of Applied Ecology*, 34:1111-1121.
- Oldfield, S., Procter, D. & Fleming, L.V. (1999) *Biodiversity: The UK Overseas Territories*, Report published by the Joint Nature Conservation Committee, ISBN: 1 86107 502 2.
- Parson, L. & Evans, A. 2005 Seafloor topography and tectonic elements of the Western Indian Ocean. *Phil. Trans. R. Soc*, 363, 1826: 15-24.
- Putney, A.D. (1982) Survey of conservation priorities in the Lesser Antilles. Final Report. Caribbean Environment Technical Report. Caribbean Conservation Association.
- Roberts, H.H. (1974) Variability of Reefs with regard to changes in wave power around an island, *Proceedings of the Second International Coral Reef Symposium*, 2:497 – 512.
- Rohmann, S.O., Hayes, J.J., Newhall, R.C., Monaco, M.E., and Grigg, R.W. (2005) The area of potential shallow water tropical and sub-tropical coral ecosystems in the United States, *Coral Reefs*, 24: 370-383.
- Sheppard, C. And Wells, S. (1988) *Coral Reefs of the World Volume 2: Indian Ocean, Red Sea and Gulf*. UNEP Regional Seas Directories and Bibliographies. IUCN, Glad, Switzerland and Cambridge, UK. pp. 1-389.

- Wells, S. And Jenkins, M (1988) Coral Reefs of the World Volume 3: Central and Western Pacific. UNEP Regional Seas Directories and Bibliographies. IUCN, Glad, Switzerland and Cambridge, UK. pp. 1-329.
- Spalding, M., Ravilious, C. & Green, E. (2001) World Atlas of Coral Reefs, University of California Press.
- Spencer, T. (1995) The Pitcairn Islands, South Pacific Ocean: plate tectonics and climatic contexts, pp 13-32. In: The Pitcairn Islands: Biogeography, Ecology and Prehistory, Benton, T & Spencer, T. (1995) Academic Press.
- Stein, S., Engela, J. & Weins, D.A. (1982) Subduction seismicity and Tectonics in the Lesser Antilles Arc, Journal of Geophysical Research, 87: 8642 – 8664.
- Sullivan, K.M., Chiappone, M., & Lott, C. (1994) Abundance patterns of stony corals on platform margin reefs of the Caicos Bank. Bahamas Journal of Science, 1: 2-12.
- Vacher, H.L. & Rowe, M.P. (1997) Geology and Hydrogeology of Bermuda pp35-91. In: Vacher, H.L. & Quinn, T.M. (Eds) Geology and Hydrogeology of Carbonate Islands, Elsevier, New York.
- Wabnitz CC, Andréfouët S, Torres-Pulliza D, Müller-Karger FE, Kramer PA (2008) Regional-scale seagrass habitat mapping in the Wider Caribbean Region using Landsat sensors: applications to conservation and ecology. Remote Sens Environ 112: 3455-3467
- Wanless, H.R. & Dravis, J.J. (1989) Carbonate Environments and sequences of the Caicos Platform: field trip guidebook T374. Washington DC, American Geophysical Union.