

2013

An investigation of mismatch negativity in current and ex-cannabis users using a feature controlled method

Felicity Webster

University of Wollongong, faw122@uowmail.edu.au

Samantha Broyd

University of Wollongong, sbroyd@uow.edu.au

Lisa-Marie Greenwood

University of Wollongong, lmg001@uowmail.edu.au

Rodney J. Croft

University of Wollongong, rcroft@uow.edu.au

Juanita Todd

University Of Newcastle

See next page for additional authors

Follow this and additional works at: <https://ro.uow.edu.au/sspapers>



Part of the [Education Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Webster, Felicity; Broyd, Samantha; Greenwood, Lisa-Marie; Croft, Rodney J.; Todd, Juanita; Michie, Patricia; Johnstone, Stuart; Lee-Bates, Benjamin; Coyle, Hannah; and Solowij, Nadia, "An investigation of mismatch negativity in current and ex-cannabis users using a feature controlled method" (2013). *Faculty of Social Sciences - Papers*. 879.

<https://ro.uow.edu.au/sspapers/879>

An investigation of mismatch negativity in current and ex-cannabis users using a feature controlled method

Abstract

Abstract presented at the 23rd Australasian Society for Psychophysiology Conference, 20-22 Nov 2013, Wollongong, Australia

Keywords

cannabis, ex, controlled, current, method, negativity, feature, mismatch, investigation, users

Disciplines

Education | Social and Behavioral Sciences

Publication Details

Webster, F., Broyd, S., Greenwood, L., Croft, R., Todd, J., Michie, P. T., Johnstone, S., Lee-Bates, B., Coyle, H. & Solowij, N. (2013). An investigation of mismatch negativity in current and ex-cannabis users using a feature controlled method. 23rd Australasian Society for Psychophysiology Conference (p. 65). Wollongong, Australia: Australian Society for Psychophysiology Inc.

Authors

Felicity Webster, Samantha Broyd, Lisa-Marie Greenwood, Rodney J. Croft, Juanita Todd, Patricia Michie, Stuart Johnstone, Benjamin Lee-Bates, Hannah Coyle, and Nadia Solowij

An investigation of mismatch negativity in current and ex- cannabis users using a feature controlled method

Felicity Webster^{1*}, Samantha Broyd¹, Lisa-marie Greenwood¹, Rodney Croft¹, Juanita Todd^{2,3}, Patricia T. Michie^{2,3}, Stuart Johnstone¹, Ben Lee-Bates¹, Hannah Coyle¹ and Nadia Solowij^{1,3}

¹School of Psychology and Centre for Psychophysics, Psychophysiology and Psychopharmacology, University of Wollongong, Australia

²School of Psychology and Priority Research Centre for Translational Neuroscience and Mental Health, Australia

³Schizophrenia Research Institute, Australia

Aims: The Mismatch Negativity (MMN) is a brain event-related potential marker of sensory memory and prediction error. Studies have found reduced MMN amplitude in long-term cannabis users, and in ex-cannabis users, relative to non-user controls. These groups have not been directly compared. A criticism of previous research is a lack of control for perceptual differences between the deviant and standard tone within the oddball sequence of a multifeature paradigm, which may enhance N1 and therefore overestimate MMN. The current study investigated the use of a novel, feature-controlled extraction method to further explore MMN in chronic users, ex-users and controls. **Method:** 39 chronic users, 16 ex-users and 44 non-user controls completed a multi-feature MMN paradigm with duration (100 ms), frequency (1200 Hz) and intensity (90 dB) deviants (deviants 6%; standards 82%, 50 ms, 1000 Hz, 80 dB), with runs preceded by trains of deviants presented as standards. MMN was extracted using (i) the traditional method (deviant – oddball standard) and (ii) a feature-controlled method (deviant – perceptually identical stimuli presented as standards prior to the oddball sequence). **Results:** A main effect of Method type indicated the traditional method produced larger MMN amplitude estimates for all groups and deviant conditions. A main effect of Group was identified for frequency MMN indicating reduced MMN in chronic users compared to controls. In ex-users, frequency MMN was reduced relative to controls using the traditional method, but only at trend level for the feature-controlled method. No differences between chronic and ex-users were identified for any deviant condition with either method. **Conclusions:** Reduced frequency MMN in chronic and 33-month abstinent users suggests chronic use may lead to early sensory information processing deficits that persist after cessation of use. These data demonstrate the utility of a feature-controlled method of examining MMN, and suggest the traditional method may overestimate MMN due to stimulus perceptual differences enhancing N1.

Acknowledgements: This study was funded by the National Health and Medical Research Council (Project Grant 1007593) and the Australian Research Council (FT110100752).

Keywords: mismatch negativity, Cannabis, feature-controlled method, chronic users, ex-users

doi: 10.3389/conf.fnhum.2013.213.00021

* Correspondence: Ms. Felicity Webster, School of Psychology and Centre for Psychophysics, Psychophysiology and Psychopharmacology, University of Wollongong, Wollongong, NSW, Australia, faw122@uowmail.edu.au