

University of Wollongong
Research Online

Faculty of Science - Papers (Archive)

Faculty of Science, Medicine and Health

1-1-2005

The effects of continuous and fluctuating copper exposures on the marine alga *Phaeodactylum tricornutum*


Brad Angel
University of Wollongong, bma06@uow.edu.au

Stuart L. Simpson
CSIRO, stuart.simpson@csiro.au

Jenny L. Stauber
CSIRO, jenny.stauber@csiro.au

Dianne Jolley
University of Wollongong, djolley@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/scipapers>

 Part of the [Life Sciences Commons](#), [Physical Sciences and Mathematics Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Angel, Brad; Simpson, Stuart L.; Stauber, Jenny L.; and Jolley, Dianne: The effects of continuous and fluctuating copper exposures on the marine alga *Phaeodactylum tricornutum* 2005.
<https://ro.uow.edu.au/scipapers/4767>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

The effects of continuous and fluctuating copper exposures on the marine alga *Phaeodactylum tricornutum*

Abstract

Contaminant concentrations in aquatic systems are seldom constant. Erratic inputs such as industrial discharges, rain water flushing, and random spills may cause concentrations to increase rapidly. Environmental processes may contribute through the dispersive actions of tides and currents, adsorptive losses to or release from resuspended sediments, and contaminant losses due to photo-degradation and volatilization. Despite such variability in contaminant concentrations, environmental guidelines are derived from toxicity test data using continuous exposure, where contaminant concentrations at the beginning of the exposure are assumed to remain relatively constant over the test duration. Responses of organisms exposed to fluctuating contaminant concentrations may differ from those exposed continuously to contaminants, even for equal contaminant loads. The current knowledge gap regarding the differing responses of organisms to contaminants from continuous and pulsed exposures is impeding decision making processes of both regulatory bodies and discharging industries. We have investigated the effects of continuous and pulsed copper exposure on the growth of the copper-sensitive microalga *Phaeodactylum tricornutum* as an indicator of ecosystem health.

Keywords

phaeodactylum, marine, alga, tricornutum, effects, exposures, copper, fluctuating, continuous

Disciplines

Life Sciences | Physical Sciences and Mathematics | Social and Behavioral Sciences

Publication Details

Angel, B., Simpson, S. L., Stauber, J. L. & Jolley, D. (2005). The effects of continuous and fluctuating copper exposures on the marine alga *Phaeodactylum tricornutum*. Book of Abstracts: ICOBTE 8th International Conference on the Biogeochemistry of Trace Elements Australia: CSIRO Land and Water.

The Effects of Continuous and Fluctuating Copper Exposures on the Marine Alga *Phaeodactylum tricornutum*

Brad Angel^{1,2}, Stuart Simpson¹, Jenny Stauber¹ and Dianne Jolley²

¹Centre for Environmental Contaminants Research, CSIRO Energy Technology, Private Mailbag 7, Bangor, NSW 2234, AUSTRALIA (Brad.Angel@csiro.au)

²GEOQUEST, Department of Chemistry, University of Wollongong, NSW 2522, AUSTRALIA

INTRODUCTION

Contaminant concentrations in aquatic systems are seldom constant. Erratic inputs such as industrial discharges, rain water flushing, and random spills may cause concentrations to increase rapidly. Environmental processes may contribute through the dispersive actions of tides and currents, adsorptive losses to or release from resuspended sediments, and contaminant losses due to photo-degradation and volatilization. Despite such variability in contaminant concentrations, environmental guidelines are derived from toxicity test data using continuous exposure, where contaminant concentrations at the beginning of the exposure are assumed to remain relatively constant over the test duration. Responses of organisms exposed to fluctuating contaminant concentrations may differ from those exposed continuously to contaminants, even for equal contaminant loads. The current knowledge gap regarding the differing responses of organisms to contaminants from continuous and pulsed exposures is impeding decision making processes of both regulatory bodies and discharging industries.

We have investigated the effects of continuous and pulsed copper exposure on the growth of the copper-sensitive microalga *Phaeodactylum tricornutum* as an indicator of ecosystem health.

METHODS

Algae were exposed to equal loads of copper either continuously or in different pulse exposures (varying pulse concentration and duration) and algal growth inhibition and changes in cell size/morphology were examined. Growth of *P. tricornutum* was investigated in experiments using copper spiking to create pulses. Algae were separated by centrifugation so that contaminated seawater could be replaced with clean seawater without loss of algal cells. Algal growth in controls was not affected in preliminary experiments using this technique. Rates of copper adsorption, internalization and elimination of copper were determined during and following exposure to different copper concentrations. A 30-min wash in 0.01 M EDTA (in 3.5 % NaCl) removed extracellular copper without disrupting cell integrity. An acid digest of the remaining algal pellet using concentrated HNO₃ gave the intracellular copper. Intra- and extracellular copper were measured using combinations of graphite furnace-atomic emission spectroscopy and anodic stripping voltammetry, respectively. Scanning electron microscopy (SEM), transmission electron microscopy (TEM) and optical microscope techniques were used to investigate changes in cell size/morphology.

Bioassays used inhibition of growth (cell yield) as the test endpoint after 72 h. Initial cell densities of $4-8 \times 10^3$ cells/mL were chosen to reflect cell densities in natural waters in order to achieve environmentally relevant partitioning of copper in solution (Franklyn *et al.*, 2002). Copper exposures included 1, 4, and 8-h pulses and continuous exposures (over 72 h) all equivalent to a time-averaged concentration near the 72-h continuous exposure IC₅₀ of 7 µg Cu/L (Figure 1). Clean filtered sea water with 15 mg NO₃⁻/L and 1.5 mg PO₄³⁻/L was used as

the test medium. Incubations were undertaken at 21°C, using a 12:12 h light:dark cycle of cool white fluorescent light at 140 $\mu\text{mol photons/m}^2/\text{s}$.

MODELLING THE EFFECT OF COPPER ON ALGAL GROWTH

Algal growth is exponential when nutrients are not limiting. Normal exponential growth in control cells can be described by $N_t = N_0 e^{\mu t}$, where, N_t is the cell biomass at any time, N_0 is the initial cell biomass, μ_t is the growth rate, and t is time. As the concentration of copper increases the algal growth rate decreases resulting in a decrease in algal cell biomass at the end of the test. Continuous exposure bioassays run at the same time as pulse exposures were used to calculate model constants. A three-parameter equation was used to predict copper inhibition of growth according to $\mu_t = \mu_i - C/(1+\exp(A+B\log(C_t)))$, where μ_t is the growth rate at time t , μ_i is the initial growth rate, C_t is the bioassay dissolved concentration at time t , and A , B , and C are constants chosen to minimize the sum of the differences between the real and model inhibitions (Simpson et al., 2003).

RESULTS AND DISCUSSION

Algal growth inhibition following copper exposure to equivalent time-averaged copper concentrations was dependent on the exposure scenario (pulse concentration and duration). Intra- and extracellular copper also varied with the different exposure scenarios (pulse and continuous) of equivalent time-averaged concentrations. The 1-h pulse scenario and the equivalent continuous exposure resulted in the highest intracellular copper concentrations, and the greatest algal growth inhibition.

A model was developed that described the response of *P. tricornutum* to both continuous and pulsed copper exposures. This model used the uptake and elimination rates to account for the delay (lag) between copper exposure and the toxic response. A linear relationship was found between intra- and extracellular copper (Figure 2).




Figure 1. Exposure concentrations used to achieve the time-averaged IC50




Figure 2. Intracellular vs extracellular copper

ACKNOWLEDGEMENTS

The authors would like to thank Dr Simon Apte, Leigh Hales, Karyn Wilde, and Nicola Creighton for their help with method development. This project was partially funded by the CRC for Coastal Zone, Estuary and Waterway Management

REFERENCES

- Franklin N.M., Stauber J.L., Apte S. C., Lim R.P. (2002) Effect of initial cell density on the bioavailability and toxicity of copper in microalgal bioassays. *Environ. Toxicol. Chem.* 21:742-751.
- Simpson S.L., Roland M.G.E., Stauber J.L., Batley G.E. (2003) Effect of declining toxicant concentrations on algal bioassay endpoints. *Environ. Toxicol. Chem.* 22:2073-2079.