

University of Wollongong
Research Online

Faculty of Engineering and Information
Sciences - Papers: Part A

Faculty of Engineering and Information
Sciences

1-1-2014

Equilibrium states on the Cuntz-Pimsner algebras of self-similar actions


Marcelo Laca
University of Victoria

Iain Raeburn
University of Otago, raeburn@uow.edu.au

Jacqui Ramagge
University of Wollongong, ramagge@uow.edu.au

Michael Whittaker
University of Wollongong, mwhittak@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/eispapers>

 Part of the [Engineering Commons](#), and the [Science and Technology Studies Commons](#)

Recommended Citation

Laca, Marcelo; Raeburn, Iain; Ramagge, Jacqui; and Whittaker, Michael, "Equilibrium states on the Cuntz-Pimsner algebras of self-similar actions" (2014). *Faculty of Engineering and Information Sciences - Papers: Part A*. 2302.
<https://ro.uow.edu.au/eispapers/2302>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Equilibrium states on the Cuntz-Pimsner algebras of self-similar actions

Abstract

We consider a family of Cuntz-Pimsner algebras associated to self-similar group actions, and their Toeplitz analogues. Both families carry natural dynamics implemented by automorphic actions of the real line, and we investigate the equilibrium states (the KMS states) for these dynamical systems. We find that for all inverse temperatures above a critical value, the KMS states on the Toeplitz algebra are given, in a very concrete way, by traces on the full group algebra of the group. At the critical inverse temperature, the KMS states factor through states of the Cuntz-Pimsner algebra; if the self-similar group is contracting, then the Cuntz-Pimsner algebra has only one KMS state. We apply these results to a number of examples, including the self-similar group actions associated to integer dilation matrices, and the canonical self-similar actions of the basilica group and the Grigorchuk group.

Keywords

states, cuntz, pimsner, equilibrium, algebras, actions, self, similar

Disciplines

Engineering | Science and Technology Studies

Publication Details

Laca, M., Raeburn, I., Ramagge, J. & Whittaker, M. (2014). Equilibrium states on the Cuntz-Pimsner algebras of self-similar actions. *Journal of Functional Analysis*, 266 (11), 6619-6661.

EQUILIBRIUM STATES ON THE CUNTZ-PIMSNER ALGEBRAS OF SELF-SIMILAR ACTIONS

MARCELO LACA, IAIN RAEBURN, JACQUI RAMAGGE, AND MICHAEL F. WHITTAKER

ABSTRACT. We consider a family of Cuntz-Pimsner algebras associated to self-similar group actions, and their Toeplitz analogues. Both families carry natural dynamics implemented by automorphic actions of the real line, and we investigate the equilibrium states (the KMS states) for these dynamical systems.

We find that for all inverse temperatures above a critical value, the KMS states on the Toeplitz algebra are given, in a very concrete way, by traces on the full group algebra of the group. At the critical inverse temperature, the KMS states factor through states of the Cuntz-Pimsner algebra; if the self-similar group is contracting, then the Cuntz-Pimsner algebra has only one KMS state. We apply these results to a number of examples, including the self-similar group actions associated to integer dilation matrices, and the canonical self-similar actions of the basilica group and the Grigorchuk group.

1. INTRODUCTION

We study operator-algebraic dynamical systems consisting of an action σ of the real line \mathbb{R} on a C^* -algebra B . Such systems have been used to model time evolution in physics, and there the states are positive functionals on B . In models from statistical mechanics, the equilibrium states are time-invariant states which satisfy a commutation relation called the KMS_β condition, where β is a real parameter called the inverse temperature [3]. However, the KMS condition is purely C^* -algebraic, and there is a great deal of evidence that the KMS states can be very interesting even when the system (B, σ) is not physical. A famous example is the number-theoretic system studied by Bost and Connes [2], which exhibits a phase transition like that of a freezing liquid. Their work generated enormous interest in the computation of KMS states for systems of purely mathematical origin (see, for example, [11, 13, 16, 15, 5]).

In [17], we analysed the KMS states on a family of Exel crossed products associated to self-coverings of the torus \mathbb{T}^d . For an integer matrix A , the covering map $e^{2\pi i x} \mapsto e^{2\pi i Ax}$ induces an endomorphism α_A of $C(\mathbb{T}^d)$ for which there is a natural transfer operator L ; the Exel crossed product is then, almost by definition [6, 4], the Cuntz-Pimsner algebra of a Hilbert bimodule M_L over $C(\mathbb{T}^d)$ defined using α_A and L . Both the Cuntz-Pimsner algebra $\mathcal{O}(M_L)$ and the Toeplitz algebra $\mathcal{T}(M_L)$ carry natural actions σ of \mathbb{R} . We showed in [17] that the system $(\mathcal{T}(M_L), \sigma)$ has no KMS states for β less than a critical value $\beta_c := \log |\det A|$, and a large simplex of KMS_β states for β greater than β_c ; when A is a dilation matrix, there is only one KMS state with inverse temperature β_c , and this state factors through a state of $(\mathcal{O}(M_L), \sigma)$.

Our analysis in [17] exploited the existence of an orthonormal basis for the right Hilbert module M_L [22, 18], which gives a Cuntz family of isometries $\{s_i\}$ in $\mathcal{O}(M_L)$. The canonical embedding of $C(\mathbb{T}^d)$ gives a unitary representation u of \mathbb{Z}^d in $\mathcal{O}(M_L)$, and Proposition 3.3 of [17] describes a presentation of $\mathcal{O}(M_L)$ in terms of the u_n and s_i .

Date: January 22, 2013.

This research was supported by the Natural Sciences and Engineering Research Council of Canada, the Marsden Fund of the Royal Society of New Zealand, and the Australian Research Council.

Our present project started when we noticed that Nekrashevych had defined “Cuntz-Pimsner algebras” for self-similar groups by specifying a similar presentation [19, 21]. In this paper we extend the analysis in [17] to cover quite general self-similar groups, with uniqueness at the critical inverse temperature for a class of self-similar actions that includes the contracting ones.

A self-similar group consists of a group G , a finite set X , and an action of G on the set X^* of finite words in the alphabet X for which there is a map $(g, x) \mapsto g|_x$ satisfying $g \cdot (xw) = (g \cdot x)(g|_x \cdot w)$ for $w \in X^*$ (see §2). Each integer matrix A gives a self-similar group (\mathbb{Z}^d, Σ) in which Σ is a set of coset representatives for $\mathbb{Z}^d/A^t\mathbb{Z}^d$ (see §2.2), but there are many more: indeed, self-similar groups have been a fertile source of interesting examples for infinite group theory (see [20], for example).

For each self-similar group (G, X) , we construct a Hilbert bimodule M over the group C^* -algebra $C^*(G)$ such that the right module has an orthonormal basis $\{e_x : x \in X\}$ and the left action of $C^*(G) = \overline{\text{span}}\{\delta_g\}$ satisfies $\delta_g \cdot e_x = e_{g \cdot x} \cdot \delta_{g|_x}$. This bimodule has a Toeplitz algebra $\mathcal{T}(M)$ and a Cuntz-Pimsner algebra $\mathcal{O}(M)$, and both carry canonical actions σ of \mathbb{R} . The Cuntz-Pimsner algebra is the same as that of Nekrashevych [21], but the Toeplitz algebra appears to be new. As previous studies in this general area have consistently showed [8, 14, 16, 17, 12], the Toeplitz system $(\mathcal{T}(M), \sigma)$ has a much richer supply of KMS states.

As in [17], there is a critical inverse temperature $\beta_c := \log |X|$ such that $(\mathcal{T}(M), \sigma)$ has no KMS states for β less than β_c . For β larger than β_c , we show that the KMS_β states $(\mathcal{T}(M), \sigma)$ are parametrised by the normalised traces on $C^*(G)$, and we give a formula for the values of these states on a set of elements which span a dense subalgebra of $\mathcal{T}(M)$ (Theorems 6.1 and 5.1). When the restrictions $g|_v$ of each fixed g form a finite set (see §2 for details), there is a unique KMS_{β_c} state on $(\mathcal{T}(M), \sigma)$, and it is the only KMS state of $(\mathcal{T}(M), \sigma)$ which factors through a state of $(\mathcal{O}(M), \sigma)$ (Theorem 7.3). We do not have an explicit formula for the values of this last state, but we describe a combinatorial procedure for computing its value on a particular generator, and illustrate this procedure in some examples (see §8.2).

Since we suspect that many operator algebraists are not familiar with self-similar group actions, we begin in §2 with a review of their basic properties. We then discuss some key examples, including odometers, actions of \mathbb{Z}^d associated to integer matrices, and two nonabelian groups called the basilica group and the Grigorchuk group. We then construct our Hilbert bimodule M over $C^*(G)$, and describe presentations of the Toeplitz algebra $\mathcal{T}(M)$ (Proposition 3.2) and the Cuntz-Pimsner algebra $\mathcal{O}(M)$ (Corollary 3.5).

Our computation of KMS states for $\beta > \beta_c$ follows the general program developed in [16], [17] and [12]. We first find an easily verified relation which allows us to recognise KMS states (Proposition 4.1). We then prove existence of KMS states using representation-theoretic methods (Theorem 5.1). As in [17], our construction uses induced representations, but in the setting of self-similar groups, we can use the bimodule M and ideas from [14] involving Rieffel induction to get a more systematic approach. We prove surjectivity of our parametrisation in §6, by showing that KMS states are characterised by their conditioning to a small corner in $\mathcal{T}(M)$. In §7, we discuss KMS states on the Cuntz-Pimsner algebra, and then we close with a section on examples.

2. SELF-SIMILAR ACTIONS

If X is a set, we write X^n for the set of words of length n in X , with $X^0 = \{\emptyset\}$, and $X^* := \bigcup_{n=0}^{\infty} X^n$. A *self-similar action* (G, X) consists of a finite set X and a faithful action

of a group G on X^* such that, for all $g \in G$ and $x \in X$, there exist unique $y \in X$ and $h \in G$ such that

$$g \cdot (xw) = y(h \cdot w) \quad \text{for all } w \in X^*. \quad (2.1)$$

We also assume that $g \cdot \emptyset = \emptyset$, and then taking $w = \emptyset$ shows that $y = g \cdot x$. We call h the *restriction* of g to x and denote it by $g|_x$. Thus (2.1) becomes

$$g \cdot (xw) = (g \cdot x)(g|_x \cdot w) \quad \text{for all } w \in X^*.$$

Then for $g \in G$ and $w = w_1 w_2 \cdots w_n$ in X^n , we have

$$g \cdot w = (g \cdot w_1)(g|_{w_1} \cdot (w_2 \cdots w_n)) = \cdots = (g \cdot w_1)(g|_{w_1} \cdot w_2) \cdots (g|_{w_1 w_2 \cdots w_{n-1}} \cdot w_n),$$

and in particular $g \cdot w \in X^n$.

Lemma 2.1 ([20, §1.3]). *Suppose (G, X) is a self-similar action.*

(1) *For each $(g, v) \in G \times X^n$, there exists $g|_v \in G$ satisfying*

$$g \cdot (vw) = (g \cdot v)(g|_v \cdot w) \quad \text{for all } w \in X^*. \quad (2.2)$$

(2) *For $g, h \in G$ and $v, w \in X^*$, we have*

$$g|_{vw} = (g|_v)|_w, \quad gh|_v = g|_{h \cdot v} h|_v, \quad \text{and} \quad g|_v^{-1} = g^{-1}|_{g \cdot v}.$$

(3) *For every $g \in G$, the map $g : X^n \rightarrow X^n$ is bijective.*

Suppose that (G, X) is a self-similar action. We can view X^* as the vertices of a rooted tree T_X with root \emptyset and edges from $w \rightarrow wx$, and then (2.2) implies that G acts on T_X by graph automorphisms. Indeed, since the action is faithful, the action gives an embedding of G in the automorphism group $\text{Aut } T_X$. Many of the important examples are constructed by specifying X and the subgroup of $\text{Aut } T_X$.

A self-similar action (G, X) is *finite-state* if for every $g \in G \setminus \{e\}$, the set $\{g|_v : v \in X^*\}$ is finite [20, page 11]. As in [20, §2.11], (G, X) is *contracting* if there is a finite subset S of G such that for every $g \in G$ there exists n with $\{g|_v : v \in X^*, |v| \geq n\} \subset S$; the smallest such set

$$\mathcal{N} := \bigcup_{g \in G} \bigcap_{n=0}^{\infty} \{g|_v : v \in X^*, |v| \geq n\} \quad (2.3)$$

is then called the *nucleus* of (G, X) .

Suppose that (G, X) is a self-similar action and S is a subset of G that is closed under restriction. The *Moore diagram* of S is the labelled directed graph with vertex set $E^0 = S$ and a directed edge from g to $g|_x$ labelled $(x, g \cdot x)$ for each $x \in X$. So an edge

$$g \xrightarrow{(x,y)} h$$

in the Moore diagram encodes the self-similar relation $g \cdot (xw) = y(h \cdot w)$.

We are particularly interested in the Moore diagram of the nucleus, and will use Moore diagrams to help find the nucleus. Later, we will use larger Moore diagrams to compute values of KMS states.

Proposition 2.2. *Suppose (G, X) is a self-similar action and S is a subset of G that is closed under restriction. Every vertex in the Moore diagram of S that can be reached from a cycle belongs to the nucleus.*

Proof. Suppose $g \in G$ is a vertex in the Moore diagram of S , and there is a cycle of length $n \geq 1$ consisting of edges labelled $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with $s(x_1, y_1) =$




FIGURE 1. The Moore diagram for the nucleus of the odometer with $N = 4$.

g , $r(x_i, y_i) = s(x_{i+1}, y_{i+1})$, and $r(x_n, y_n) = g$. By definition of the Moore diagram we have $g \cdot (x_1 \cdots x_n) = y_1 \cdots y_n$ and $g|_{x_1 \cdots x_n} = g$. Thus $g = g|_{(x_1 \cdots x_n)^m}$ for all $m \in \mathbb{N}$ and

$$g \in \bigcap_{n \geq 0} \{g|_v : v \in X^*, |v| \geq n\} \implies g \in \bigcup_{h \in G} \bigcap_{n \geq 0} \{h|_v : v \in X^*, |v| \geq n\} = \mathcal{N}.$$

A similar argument shows that if g can be reached from a cycle, then there are arbitrarily long paths ending at g . \square

In the rest of this section, we discuss some key examples of self-similar actions.

2.1. Odometers. Fix an integer $N > 1$, and let $X_N = \{0, 1, \dots, N-1\}$. We consider the multiplicative free abelian group G with generator g , so that $G = \{g^k : k \in \mathbb{Z}\}$. We define an action of G on X_N^* by

$$g \cdot v = \begin{cases} (v_1 + 1)v_2 \cdots v_{|v|} & \text{if } v_1 < N - 1 \\ 0 \cdots 0(v_k + 1)v_{k+1} \cdots v_{|v|} & \text{if } v_1 = \cdots = v_{k-1} = N - 1 \text{ and } v_k < N - 1. \end{cases}$$

Then (G, X_N) is a self-similar action with $g|_i = e$ for $i < N - 1$ and $g|_{N-1} = g$. This action is contracting with nucleus $\mathcal{N} = \{e, g, g^{-1}\}$: indeed, if $k > 0$ and $|w| > \log_N k$, then $g^k|_w$ is either e or g , and if $k < 0$ and $|w| > \log_N |k|$, then $g^k|_w$ is either e or g^{-1} . The Moore diagram for \mathcal{N} for $N = 4$ is shown in Figure 1.

The self-similar action (G, X_N) is called an *odometer*. To see why, identify X_N^n with $\{0, 1, \dots, N^n - 1\}$ by sending v to $\sum_{i=1}^n v_i N^{i-1}$, and then the action of g on X_N^n adds 1 (mod N^n).

2.2. Integer matrices. Suppose that $A \in M_d(\mathbb{Z})$ has $N := |\det A| > 1$, and write $B := A^t$ for its transpose. We choose a set Σ of coset representatives for the quotient $\mathbb{Z}^d/B\mathbb{Z}^d$, and we assume that $0 \in \Sigma$. For $n \in \mathbb{Z}^d$, we write $c(n)$ for the representative of $n + B\mathbb{Z}^d$ in Σ . We note that $\det B = \det A = N$, and hence Σ has cardinality N .

We now fix an integer $k \geq 1$. Then the set Σ^k gives a parametrisation

$$\{b_k(w) = w_1 + Bw_2 + \cdots + B^{k-1}w_k + B^k\mathbb{Z}^d : w \in \Sigma^k\}$$

of $\mathbb{Z}^d/B^k\mathbb{Z}^d$. The following straightforward lemma tells us how the different bijections b_k combine.

Lemma 2.3. *Write B for the injective homomorphism $B : \mathbb{Z}^d/B^k\mathbb{Z}^d \rightarrow \mathbb{Z}^d/B^{k+1}\mathbb{Z}^d$ which takes $m + B^k\mathbb{Z}^d$ to $Bm + B^{k+1}\mathbb{Z}^d$. Then for $x \in \Sigma$ and $w \in \Sigma^k$ we have $b_{k+1}(xw) = x + B(b_k(w))$.*

Proposition 2.4. *The actions of the additive abelian group \mathbb{Z}^d on its quotients $\mathbb{Z}^d/B^k\mathbb{Z}^d$ combine to give an action of \mathbb{Z}^d on Σ^* such that*

$$n \cdot w = b_k^{-1}(n \cdot b_k(w)) \quad \text{for } k \geq 1 \text{ and } w \in \Sigma^k. \quad (2.4)$$

The pair (\mathbb{Z}^d, Σ) is a self-similar action, and for $n \in \mathbb{Z}^d$, $x \in \Sigma$ we have

$$n \cdot x = c(n + x) \quad \text{and} \quad n|_x = B^{-1}(n + x - c(n + x)). \quad (2.5)$$




FIGURE 2. The Moore diagram for the nucleus of the basilica group.

If A is a dilation matrix (in the sense that all its complex eigenvalues λ satisfy $|\lambda| > 1$), then (\mathbb{Z}^d, Σ) is contracting.

Proof. Since

$$\begin{aligned} (m+n) \cdot (p + B^k \mathbb{Z}^d) &= (m+n) + p + B^k \mathbb{Z}^d = m \cdot (n + p + B^k \mathbb{Z}^d) \\ &= m \cdot (n \cdot (p + B^k \mathbb{Z}^d)), \end{aligned}$$

the formula (2.4) gives an action of the additive group \mathbb{Z}^d on Σ^k . To establish (2.5), we take $x \in \Sigma$, $w \in \Sigma^k$, and compute:

$$\begin{aligned} b_{k+1}(n \cdot (xw)) &= n + x + Bw_1 \cdots + B^k w_k + B^{k+1} \mathbb{Z}^d \\ &= c(n+x) + (n+x - c(n+x)) + Bw_1 \cdots + B^k w_k + B^{k+1} \mathbb{Z}^d \\ &= c(n+x) + B(B^{-1}(n+x - c(n+x)) + w_1 \cdots + B^{k-1} w_k + B^k \mathbb{Z}^d), \\ &= c(n+x) + B(b_k(B^{-1}(n+x - c(n+x)) \cdot w)), \end{aligned}$$

which by Lemma 2.3 is $b_{k+1}(c(n+x)(B^{-1}(n+x - c(n+x)) \cdot w))$. Thus

$$n \cdot (xw) = c(n+x)(B^{-1}(n+x - c(n+x)) \cdot w),$$

which implies that (\mathbb{Z}^d, Σ) is self-similar, and gives (2.5).

Now we suppose that A is a dilation matrix. For $x \in \Sigma$, the virtual endomorphism ϕ_x associated to $x \in X$ (as in [20, §2.5]) is the map $n \mapsto n|_x$ from the stabiliser of x into \mathbb{Z}^d ; the stabiliser is $B\mathbb{Z}^d$, and for $n \in B\mathbb{Z}^d$, $\phi_x(n) = B^{-1}n$. Thus the linear transformation $\mathbb{Q} \otimes \phi : \mathbb{Q}^d \rightarrow \mathbb{Q}^d$ considered in [20, Theorem 2.12.1] has matrix B^{-1} . Since $\det B = \det A \neq 0$, the eigenvalues of B^{-1} are the inverses λ^{-1} of the eigenvalues of B . Since A is a dilation matrix, we have $|\lambda^{-1}| < 1$ for all such λ , and B^{-1} has spectral radius $\rho(B^{-1}) < 1$. Thus [20, Theorem 2.12.1] implies that (\mathbb{Z}^d, Σ) is contracting. \square

2.3. The basilica group. Let X be the set $\{x, y\}$ with $|X| = 2$, and consider the rooted homogeneous tree T_X with vertex set X^* . We recursively define two automorphisms a and b of T_X by

$$\begin{aligned} a \cdot (xw) &= y(b \cdot w) & a \cdot (yw) &= xw \\ b \cdot (xw) &= x(a \cdot w) & b \cdot (yw) &= yw \end{aligned} \tag{2.6}$$

for $w \in X^*$. Then the *basilica group* B is the subgroup of $\text{Aut } T_X$ generated by $\{a, b\}$. The pair (B, X) is then a self-similar action.

We now show that the basilica action (B, X) is contracting and compute the nucleus. This is probably well-known, but since our answer seems to contradict an assertion in [20, page 111], we give a detailed proof.

Proposition 2.5. *The basilica group action (B, X) is contracting, with nucleus*

$$\mathcal{N} = \{e, a, b, a^{-1}, b^{-1}, ab^{-1}, ba^{-1}\};$$

the Moore diagram of \mathcal{N} is in Figure 2.

Proof. Let $\mathcal{S} = \{e, a, b, a^{-1}, b^{-1}, ab^{-1}, ba^{-1}\}$, for which we have the Moore diagram in Figure 2. Since this Moore diagram has a cycle at every vertex, Proposition 2.2 implies that $\mathcal{S} \subset \mathcal{N}$. So we have to prove $\mathcal{N} \subset \mathcal{S}$,

We claim that every $g \in B \setminus \{ab^{-1}, ba^{-1}\}$ which can be written as a reduced product of two elements of $T := \{a, b, a^{-1}, b^{-1}\}$ has the property that $g|_v \in T \cup \{e\}$ for every word $v \in X^2$. There are 12 non-trivial length-two words in G ; we delete ab^{-1}, ba^{-1} from the list, and compute the other restrictions to x and y . We find that

$$\begin{array}{llll} a^2|_x = b & a^2|_y = b & ab|_x = ba & ab|_y = e \\ a^{-2}|_x = b^{-1} & a^{-2}|_y = b^{-1} & a^{-1}b|_x = a & a^{-1}b|_y = b^{-1} \\ a^{-1}b^{-1}|_x = a^{-1} & a^{-1}b^{-1}|_y = b^{-1} & ba|_x = b & ba|_y = a \\ b^2|_x = a^2 & b^2|_y = e & b^{-1}a|_x = b & b^{-1}a|_y = a^{-1} \\ b^{-1}a^{-1}|_x = e & b^{-1}a^{-1}|_y = a^{-1}b^{-1} & b^{-2}|_x = a^{-2} & b^{-2}|_y = e. \end{array}$$

Now we observe that further restrictions of $b^{-1}a^{-1}, ab, b^2$ and b^{-2} are all in $T \cup \{e\}$, and we have justified our claim.

Next we suppose that $n \geq 3$ and that g can be written as a reduced product of n elements of T , and take $v \in X^2$. We claim that $g|_v$ can be written as a product of at most $n - 1$ elements of T . We factor off the last two elements of T , say $g = g'h$. Since $g|_v = g'|_{h \cdot v} h|_v$, the claim in the previous paragraph implies that, unless $(h, v) = (ab^{-1}, xy)$ or (ba^{-1}, yx) , we have $h|_v \in T$. Since $g'|_{h \cdot v}$ is a product of at most $n - 2$ elements of T , we have $g|_v = g'|_{h \cdot v} h|_v$ written as a product of $n - 1$ elements of T . So we have to deal with $(h, v) = (ab^{-1}, xy)$ and $h = (ba^{-1}, yx)$. Now, since g' is a product of $n - 2$ elements of T and $n \geq 3$, we can pull one element t out of g' , and it suffices for us to prove that $(tab^{-1})|_{xy}$ and $(tba^{-1})|_{yx}$ can be written as products of two elements of T . We compute:

$$\begin{aligned} (a^2b^{-1})|_x &= a|_{(ab^{-1}) \cdot x} (ab^{-1})|_x = a|_y ba^{-1} = ba^{-1} \\ (b^{-1}ab^{-1})|_x &= b^{-1}|_y (ba^{-1}) = ba^{-1} \\ (bab^{-1})|_x &= b|_y ba^{-1} = ba^{-1} \\ (a^{-1}ba^{-1})|_y &= a^{-1}|_x ab^{-1} = ab^{-1} \\ (b^2a^{-1})|_{yx} &= (b|_x (ab^{-1}))|_x = (a(ab^{-1}))|_x = a|_y ba^{-1} = ba^{-1} \\ (aba^{-1})|_{yx} &= (a|_x ab^{-1})|_x = (bab^{-1})|_x = b|_y ba^{-1} = ba^{-1}. \end{aligned}$$

This completes the proof of the claim.

Successive applications of the claim in the previous paragraph show that if g is a product of n elements of T and $n \geq 3$, then $g|_v$ is a product of at most 2 elements for every v with $|v| \geq 2(n - 2)$. Now the calculations in the first paragraph show that a further restriction to a word in X^2 gets us into \mathcal{S} . Thus for $|v| \geq 2(n - 1)$, we have $g|_v \in \mathcal{S}$. So the inside intersection in (2.3) is contained in \mathcal{S} , and so is \mathcal{N} . \square

Our next proposition says that B has a large abelian quotient. We believe this is known, but we do not know where a proof has been published.

We want to use the presentation of B found by Bartholdi and Virág [1, Lemma 11], building on work of Grigorchuk and Żuk [10]. We consider the collection Y^* of all nonempty words in $Y := \{a, b, a^{-1}, b^{-1}\}$, and the transformation $\sigma : Y^* \rightarrow Y^*$ which




FIGURE 3. The Moore diagram for the nucleus of the Grigorchuk group.

replaces every appearance of a by bb , every a^{-1} by $b^{-1}b^{-1}$, every b by a , and every b^{-1} by a^{-1} . For $c, d \in Y^*$, we write c^{-1} for the word obtained by formally inverting c , and $[c, d]$ for the word $c^{-1}d^{-1}cd$. Then [1, Lemma 11] says that B has the presentation

$$B = \langle a, b : \sigma^n([a, b^{-1}ab]) = e \text{ for all } n \in \mathbb{N} \rangle. \quad (2.7)$$

Proposition 2.6. *Let $[B, B]$ be the commutator subgroup of the basilica group B , and let $q : B \rightarrow B/[B, B]$ be the quotient map. Then there is an isomorphism ϕ of $B/[B, B]$ onto \mathbb{Z}^2 such that $\phi(q(a)) = (1, 0)$ and $\phi(q(b)) = (0, 1)$.*

Proof. Since $a' := (1, 0)$ and $b' := (0, 1)$ commute, we have $\sigma^n([a', (b')^{-1}a'b']) = e$ for all n . Thus there is a homomorphism of B into \mathbb{Z}^2 taking $\{a, b\}$ to $\{a', b'\}$, and this factors through a homomorphism $\phi : B/[B, B] \rightarrow \mathbb{Z}^2$.

Since $[a, b]$ and $[a, b^{-1}]$ belong to the commutator subgroup, $q(a)$ commutes with $q(b)$ and $q(b^{-1})$, and hence for every $g \in B = \langle a, b \rangle$, there are $k, l \in \mathbb{Z}$ such that $q(g) = q(a^k b^l)$. Since $\phi(q(a^k b^l)) = (k, l)$, we deduce both that ϕ is surjective and that ϕ is injective. \square

2.4. The Grigorchuk group. We again consider the set $X = \{x, y\}$ and the associated rooted tree T_X with vertex set X^* . We define automorphisms a, b, c , and d of T_X recursively by

$$\begin{aligned} a \cdot (xw) &= yw & a \cdot (yw) &= xw & (2.8) \\ b \cdot (xw) &= x(a \cdot w) & b \cdot (yw) &= y(c \cdot w) \\ c \cdot (xw) &= x(a \cdot w) & c \cdot (yw) &= y(d \cdot w) \\ d \cdot (xw) &= xw & d \cdot (yw) &= y(b \cdot w). \end{aligned}$$

Then the Grigorchuk group G is the subgroup of $\text{Aut } T_X$ generated by $\{a, b, c, d\}$.

The first assertions of the next proposition are also in the proof of [20, Theorem 1.6.1]; the assertion about the nucleus is stated without proof on page 57 of [20].

Proposition 2.7. *The generators a, b, c, d of G all have order two, and satisfy $cd = b = dc$, $db = c = bd$ and $bc = d = cb$. The self-similar action (G, X) is contracting with nucleus $\mathcal{N} = \{e, a, b, c, d\}$.*

Proof. The first two relations in (2.8) imply that $a^2 = e$. Now the other relations imply that

$$\begin{aligned} b^2 \cdot (xw) &= x(a^2 \cdot w) = xw & b^2 \cdot (yw) &= y(c^2 \cdot w) \\ c^2 \cdot (xw) &= x(a^2 \cdot w) = xw & c^2 \cdot (yw) &= y(d^2 \cdot w) \end{aligned}$$

$$d^2 \cdot (xw) = xw \qquad d^2 \cdot (yw) = y(b^2 \cdot w),$$

and we can prove by induction on $n = |v|$ that $b^2 \cdot v = c^2 \cdot v = d^2 \cdot v = v$ for all $v \in X^*$. Thus $b^2 = c^2 = d^2 = e$ in $G \subset \text{Aut } T_X$. In particular, every element of G is a product of generators $\{a, b, c, d\}$.

Next we note that a is determined by the first two relations in (2.8), and then the other six determine (b, c, d) . A computation shows that $(b', c', d') = (cd, db, bc)$ satisfies the same six recurrence relations, and hence we have $cd = b$, $db = c$ and $bc = d$. Since the generators all have order two, inverting gives $dc = b$, $bd = c$ and $cb = d$. Thus the only elements of G which are products of two generators are the elements of

$$R := \{ab, ba, ac, ca, ad, da\}.$$

Twelve calculations show that for every $g \in R$, both $g|_x$ and $g|_y$ belong to $\{e, a, b, c, d\}$. Thus if g is a product of n generators, we have $g|_v \in \{e, a, b, c, d\}$ for every word v with $|v| \geq n - 1$. This proves that (G, X) is contracting, and that the nucleus is contained in $\{e, a, b, c, d\}$.

Since every vertex in the Moore diagram of $\{e, a, b, c, d\}$ in Figure 3 can be reached from a cycle, Proposition 2.2 implies that $\{e, a, b, c, d\}$ is contained in the nucleus. \square

3. UNIVERSAL ALGEBRAS ASSOCIATED TO A SELF-SIMILAR ACTION

Suppose that (G, X) is a self-similar action, and let $C^*(G)$ be the full group C^* -algebra of G generated by the unitary representation $\{\delta_g : g \in G\}$. We are interested in two C^* -algebras associated to (G, X) , which we construct as the Toeplitz algebra and the Cuntz-Pimsner algebra of a Hilbert bimodule M over $C^*(G)$.

As a right Hilbert $C^*(G)$ -module, M is the direct sum $M = \bigoplus_{x \in X} C^*(G)$; thus $M = \{m = (m_x)_{x \in X} : m_x \in C^*(G)\}$, with module action $(m_x) \cdot a = (m_x a)$ and inner product

$$\langle m, n \rangle = \sum_{x \in X} m_x^* n_x.$$

For $y \in X$ we define $e_y \in M$ by

$$(e_y)_x = \begin{cases} 1_{C^*(G)} = \delta_e & \text{if } x = y \\ 0 & \text{otherwise,} \end{cases}$$

and then $\{e_x : x \in X\}$ is an orthonormal basis for M with reconstruction formula

$$m = \sum_{x \in X} e_x \cdot \langle e_x, m \rangle \quad \text{for } m \in M. \quad (3.1)$$

The left action of $C^*(G)$ on M will be the integrated form of the unitary representation T in the next proposition.

Proposition 3.1. *Let (G, X) be a self-similar action, and let $g \in G$. Then there is an adjointable operator T_g on M such that*

$$T_g(e_x \cdot a) = e_{g \cdot x} \cdot (\delta_{g|x} a) \quad \text{for } x \in X \text{ and } a \in C^*(G), \quad (3.2)$$

and $T : g \mapsto T_g$ is a unitary representation of G in $\mathcal{L}(M)$.

Proof. We define $T_g : M \rightarrow M$ by

$$T_g(m) = \sum_{y \in X} e_{g \cdot y} \cdot (\delta_{g|y} \langle e_y, m \rangle).$$

For $\mathfrak{a} \in C^*(G)$ and $\mathfrak{m} \in M$, we have

$$T_g(\mathfrak{m} \cdot \mathfrak{a}) = \sum_{y \in X} e_{g \cdot y} \cdot (\delta_{g|_y} \langle e_y, \mathfrak{m} \cdot \mathfrak{a} \rangle) = \sum_{y \in X} e_{g \cdot y} \cdot (\delta_{g|_y} \langle e_y, \mathfrak{m} \rangle \mathfrak{a}) = T_g(\mathfrak{m}) \cdot \mathfrak{a},$$

and hence T_g is $C^*(G)$ linear. The computation

$$T_g(e_x \cdot \mathfrak{a}) = \sum_{y \in X} e_{g \cdot y} \cdot (\delta_{g|_y} \langle e_y, e_x \cdot \mathfrak{a} \rangle) = e_{g \cdot x} \cdot (\delta_{g|x} \mathfrak{a})$$

shows that T_g satisfies Equation (3.2).

We next show that T_g is adjointable with $T_g^* = T_{g^{-1}}$. Let $g \in G$, $x, y \in X$ and $\mathfrak{a}, \mathfrak{b} \in C^*(G)$. Then

$$\begin{aligned} \langle T_g(e_x \cdot \mathfrak{a}), e_y \cdot \mathfrak{b} \rangle &= \langle e_{g \cdot x} \cdot (\delta_{g|x} \mathfrak{a}), e_y \cdot \mathfrak{b} \rangle \\ &= \begin{cases} (\delta_{g|x} \mathfrak{a})^* \mathfrak{b} & \text{if } y = g \cdot x \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} \mathfrak{a}^* \delta_{(g|x)^{-1}} \mathfrak{b} & \text{if } y = g \cdot x \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} \mathfrak{a}^* \delta_{g^{-1}|_{g \cdot x}} \mathfrak{b} & \text{if } y = g \cdot x \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} \mathfrak{a}^* \delta_{g^{-1}|_y} \mathfrak{b} & \text{if } x = g^{-1} \cdot y \\ 0 & \text{otherwise} \end{cases} \\ &= \langle e_x \cdot \mathfrak{a}, e_{g^{-1} \cdot y} \cdot \delta_{g^{-1}|_y} \mathfrak{b} \rangle \\ &= \langle e_x \cdot \mathfrak{a}, T_{g^{-1}}(e_y \cdot \mathfrak{b}) \rangle. \end{aligned}$$

which implies that T_g is adjointable with $T_g^* = T_{g^{-1}}$. Next we let $g, h \in G$, and the calculation

$$\begin{aligned} T_{gh}(e_x \cdot \mathfrak{a}) &= e_{(gh) \cdot x} \cdot (\delta_{gh|x} \mathfrak{a}) \\ &= e_{g \cdot (h \cdot x)} \cdot (\delta_{g|h \cdot x} \mathfrak{a}) \quad \text{by Lemma 2.1} \\ &= T_g(e_{h \cdot x} \cdot (\delta_{h|x} \mathfrak{a})) \\ &= T_g T_h(e_x \cdot \mathfrak{a}). \end{aligned}$$

shows that $T_{gh} = T_g T_h$. Since $T_g^* = T_{g^{-1}}$, this implies that each T_g is unitary, and that T is a homomorphism of G into the unitary group $\mathcal{UL}(M)$, or, in other words, a unitary representation in $\mathcal{L}(M)$. \square

By [26, Proposition C.17], the unitary representation $T: G \rightarrow \mathcal{UL}(M)$ has an integrated form $\pi_T: C^*(G) \rightarrow \mathcal{L}(M)$ satisfying $\pi_T(\delta_g) = T_g$, and with the left action defined by $\mathfrak{a} \cdot \mathfrak{m} = \pi_T(\mathfrak{a})\mathfrak{m}$, M becomes a Hilbert bimodule over $C^*(G)$.

A representation of M in a C^* -algebra B consists of a linear map $\psi: M \rightarrow B$ and a homomorphism $\pi: C^*(G) \rightarrow B$ satisfying

$$\begin{aligned} \psi(x \cdot \mathfrak{a}) &= \psi(x)\pi(\mathfrak{a}), \\ \psi(x)^* \psi(y) &= \pi(\langle x, y \rangle_{C^*(G)}), \text{ and} \\ \psi(\mathfrak{a} \cdot x) &= \pi(\mathfrak{a})\psi(x) \end{aligned}$$

for all $x, y \in X$ and $\mathfrak{a} \in C^*(G)$ (see [9, Section 1]). A representation (ψ, π) of M in B induces a homomorphism $(\psi, \pi)^{(1)}: \mathcal{K}(M) \rightarrow B$ such that $(\psi, \pi)^{(1)}(\Theta_{\mathfrak{m}, \mathfrak{n}}) = \psi(\mathfrak{m})\psi(\mathfrak{n})^*$

for $m, n \in M$ [9, Proposition 1.6], and (ψ, π) is *Cuntz-Pimsner covariant* if¹

$$(\psi, \pi)^{(1)}(\pi_{\mathcal{T}}(a)) = \pi(a) \quad \text{whenever } \pi_{\mathcal{T}}(a) \in \mathcal{K}(M).$$

By [9, Proposition 1.3], the Hilbert bimodule M has a Toeplitz algebra $\mathcal{T}(M)$ generated by a universal representation $(i_M, i_{C^*(G)}) : M \rightarrow \mathcal{T}(M)$; if (ψ, π) is a representation of M in B , we write $\psi \times \pi$ for the homomorphism of $\mathcal{T}(M)$ into B such that $(\psi \times \pi) \circ i_M = \psi$ and $(\psi \times \pi) \circ i_{C^*(G)} = \pi$. The Cuntz-Pimsner algebra $\mathcal{O}(M)$ is the quotient of $\mathcal{T}(M)$ which is generated by a universal Cuntz-Pimsner covariant representation $(j_M, j_{C^*(G)})$. We call $\mathcal{T}(G, X) := \mathcal{T}(M)$ and $\mathcal{O}(G, X) := \mathcal{O}(M)$ the *Toeplitz algebra* and *Cuntz-Pimsner algebra* of the self-similar action (G, X) . It will follow from Corollary 3.5 below that $\mathcal{O}(G, X)$ is the same as the universal Cuntz-Pimsner algebra \mathcal{O}_G in [21, Definition 3.1].

We will use the following presentation of $\mathcal{T}(G, X)$.

Proposition 3.2. *Let (G, X) be a self-similar action, and set $u_g := i_{C^*(G)}(\delta_g)$ for $g \in G$, and $s_x := i_M(e_x)$ for $x \in X$. Then*

- (1) $u : G \rightarrow \mathcal{T}(G, X)$ is a unitary representation of G ,
- (2) $\{s_x : x \in X\}$ is a Toeplitz-Cuntz family of isometries in $\mathcal{T}(G, X)$, and
- (3) $u_g s_x = s_{g \cdot x} u_{g|x}$ for $g \in G$ and $x \in X$.

The set $\{u_g : g \in G\} \cup \{s_x : x \in X\}$ generates $\mathcal{T}(G, X)$, and $(\mathcal{T}(G, X), (u, s))$ is universal for families $\{U_g : g \in G\}$ and $\{S_x : x \in X\}$ satisfying (1), (2) and (3).

Proof. The map u is a unitary representation because $\delta : G \rightarrow \text{UC}^*(G)$ is, and $i_{C^*(G)}$ is a unital homomorphism (which follows from [4, Corollary 3.3]). We have

$$s_x^* s_y = i_M(e_x)^* i_M(e_y) = i_{C^*(G)}(\langle e_x, e_y \rangle) = \begin{cases} 1 & \text{if } x = y \\ 0 & \text{if } x \neq y, \end{cases}$$

which implies that $\{s_x : x \in X\}$ is a Toeplitz-Cuntz family. For (3), we compute

$$\begin{aligned} u_g s_x &= i_{C^*(G)}(\delta_g) i_M(e_x) = i_M(\delta_g \cdot e_x) = i_M(T_g(e_x)) \\ &= i_M(e_{g \cdot x} \cdot \delta_{g|x}) = i_M(e_{g \cdot x}) i_{C^*(G)}(\delta_{g|x}) = s_{g \cdot x} u_{g|x}. \end{aligned}$$

The u_g generate $i_{C^*(G)}(C^*(G))$, and for $m \in M$, the reconstruction formula (3.1) gives

$$i_M(m) = i_M\left(\sum_{x \in X} e_x \cdot \langle e_x, m \rangle\right) = \sum_{x \in X} i_M(e_x) i_{C^*(G)}(\langle e_x, m \rangle).$$

Thus $C^*(u_g, s_x)$ contains all the generators of $\mathcal{T}(G, X) = \mathcal{T}(M)$, and must be all of $\mathcal{T}(G, X)$.

To see the universal property, suppose D is a C^* -algebra, and $\{U_g\} \subset D$ and $\{S_x\} \subset D$ satisfy (1), (2) and (3). We have to find a homomorphism $\pi_{U,S} : \mathcal{T}(G, X) \rightarrow D$ such that $\pi_{U,S}(u_g) = U_g$ and $\pi_{U,S}(s_x) = S_x$. Let $\pi_U : C^*(G) \rightarrow D$ be the integrated form of U . Since each element of M has a unique expansion $\sum_{x \in X} e_x \cdot a_x$, there is a well-defined linear function $\psi : M \rightarrow M$ such that $\psi(e_x \cdot a) = S_x \pi_U(a)$ for $x \in X$ and $a \in C^*(G)$.

We claim that (ψ, π_U) is a representation of M . Let $a \in C^*(G)$ and $x \in X$. Then

$$\psi((e_x \cdot a) \cdot b) = \psi(e_x \cdot (ab)) = S_x \pi_U(ab) = (S_x \pi_U(a)) \pi_U(b) = \psi(e_x \cdot a) \pi_U(b).$$

Next we consider the left action of $b = \delta_g$, which is implemented by the operator T_g of Proposition 3.1. We calculate using relation (3):

$$\psi(\delta_g \cdot (e_x \cdot a)) = \psi(e_{g \cdot x} \cdot (\delta_{g|x} a)) = S_{g \cdot x} \pi_U(\delta_{g|x} a)$$

¹This is Pimsner's original definition [24]; many authors use a slightly different definition due to Katsura, but the two definitions give the same algebras for the bimodules we consider.

$$= S_{g \cdot x} U_{g|x} \pi_U(a) = U_g S_x \pi(a) = \pi_U(\delta_g) \psi(e_x \cdot a),$$

which implies that $\psi(a \cdot m) = \pi(a) \psi(m)$ for $a \in C^*(G)$ and $m \in M$. For the inner product, we have

$$\begin{aligned} \pi_U(\langle e_x \cdot a, e_y \cdot b \rangle) &= \begin{cases} \pi_U(a^* b) & \text{if } x = y \\ 0 & \text{if } x \neq y \end{cases} \\ &= \pi_U(a)^* S_x^* S_y \pi_U(b) = (S_x \pi_U(a))^* (S_y \pi_U(b)) \\ &= \psi(e_x \cdot a)^* \psi(e_y \cdot b). \end{aligned}$$

So (ψ, π_U) is a Toeplitz representation of M , as claimed. Thus there is a homomorphism $\psi \times \pi_U : \mathcal{T}(G, X) = \mathcal{T}(M) \rightarrow D$. We have

$$(\psi \times \pi_U)(u_g) = (\psi \times \pi_U)(i_{C^*(G)}(\delta_g)) = \pi_U(\delta_g) = U_g,$$

and similarly $(\psi \times \pi_U)(s_x) = S_x$. Thus $\pi_{U,S} := \psi \times \pi_U$ has the required properties. \square

We now recall some standard notation for working with the Toeplitz-Cuntz family $\{s_x : x \in X\}$. For $v \in X^n$, we write $s_v := s_{v_1} s_{v_2} \cdots s_{v_n}$. Then for each n , $\{s_v : v \in X^n\}$ is a Toeplitz-Cuntz family, so we have $1 \geq \sum_{v \in X^n} s_v s_v^*$. For $v, w \in X^*$, the product $s_v^* s_w$ vanishes unless either $v = wv'$ or $w = vv'$, and then collapses down to $s_{v'}^*$ or $s_{w'}$. The relation (3) in Proposition 3.2 extends to $u_g s_v = s_{g \cdot v} u_{g|v}$ for $v \in X^*$.

Corollary 3.3. *Let (G, X) be a self-similar action, and take (u, s) as in Proposition 3.2. Then*

$$\mathcal{T}(G, X) = \overline{\text{span}}\{s_v u_g s_w^* : v, w \in X^*, g \in G\}.$$

As usual, we prove that $A_0 := \text{span}\{s_v u_g s_w^*\}$ is a $*$ -subalgebra of $\mathcal{T}(G, X)$, and then since A_0 contains all the generators $\{u_g\} \cup \{s_x\}$, its closure has to be all of $\mathcal{T}(G, X)$. Since A_0 is closed under taking adjoints, it remains to show that $\{s_v u_g s_w^*\}$ is closed under multiplication. Since we will need the result of the computation, we state it separately:

Lemma 3.4. *For $v, w, y, z \in X^*$ and $g, h \in G$, we have*

$$(s_v u_g s_w^*)(s_y u_h s_z^*) = \begin{cases} s_{v(g \cdot y')} u_{g|y'} h s_z^* & \text{if } y = wy' \\ s_v u_{g(h|_{h^{-1} \cdot w'})} s_{z(h^{-1} \cdot w')}^* & \text{if } w = yw' \\ 0 & \text{otherwise.} \end{cases} \quad (3.3)$$

Proof. We have $s_w^* s_y = 0$ unless either $y = wy'$ or $w = yw'$, and hence a computation using the relations $u_g s_w = s_{g \cdot w} u_{g|w}$ gives

$$\begin{aligned} s_v u_g s_w^* s_y u_h s_z^* &= \begin{cases} s_v u_g s_{y'} u_h s_z^* & \text{if } y = wy' \\ s_v u_g s_{w'}^* u_h s_z^* & \text{if } w = yw' \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} s_v s_{g \cdot y'} u_{g|y'} u_h s_z^* & \text{if } y = wy' \\ s_v u_g (s_{h^{-1} \cdot w'} u_{h^{-1}|w'})^* s_z^* & \text{if } w = yw' \\ 0 & \text{otherwise.} \end{cases} \\ &= \begin{cases} s_{v(g \cdot y')} u_{g|y'} h s_z^* & \text{if } y = wy' \\ s_v u_{g(h|_{h^{-1} \cdot w'})} s_{z(h^{-1} \cdot w')}^* & \text{if } w = yw' \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

as required. \square

Corollary 3.5. *Let (G, X) be a self-similar action, and take (u, s) as in Proposition 3.2. Then $\mathcal{O}(G, X)$ is the quotient of $\mathcal{T}(G, X)$ by the ideal I generated by $1 - \sum_{x \in X} s_x s_x^*$.*

Proof. Since $\{e_x : x \in X\}$ is an orthonormal basis for M , it follows from [7, Lemma 2.5] that a Toeplitz representation (ψ, π) is Cuntz-Pimsner covariant if and only if

$$1 = \sum_{x \in X} \psi(e_x) \psi(e_x)^* = \sum_{x \in X} (\psi \times \pi)(i_M(e_x) i_M(e_x)^*) = (\psi \times \pi) \left(\sum_{x \in X} s_x s_x^* \right),$$

and hence if and only if $\psi \times \pi$ vanishes on I . \square

We write u_g and s_x also for the images of the generators in $\mathcal{O}(G, X)$. Since $1 = \sum_{x \in X} s_x s_x^*$ in $\mathcal{O}(G, X)$, the image of the Toeplitz-Cuntz family $\{s_x : x \in X\}$ in $\mathcal{O}(G, X)$ is a Cuntz family. The same is true of the Toeplitz-Cuntz families $\{s_v : v \in X^n\}$, so for every $n \in \mathbb{N}$ we have $1 = \sum_{v \in X^n} s_v s_v^*$ in $\mathcal{O}(G, X)$.

Remark 3.6. As we observed earlier, Corollary 3.5 implies that $\mathcal{O}(G, X)$ is the universal Cuntz-Pimsner algebra \mathcal{O}_G in [21, Definition 3.1]. It is not necessarily the same as the Cuntz-Pimsner algebra in [19], which is generated by a Cuntz family $\{s_x : x \in X\}$ and a unitary representation u of G which factors through a particular ‘‘permutation representation’’ of $C^*(G)$.

Corollary 3.7. *Let (G, X) be a self-similar action with nucleus \mathcal{N} . Then*

$$\mathcal{O}(G, X) = \overline{\text{span}}\{s_v u_g s_w^* : v, w \in X^*, g \in \mathcal{N}\}. \quad (3.4)$$

Proof. Since $\mathcal{O}(G, X)$ is a quotient of $\mathcal{T}(G, X)$, Corollary 3.3 implies that the elements $\{s_v u_h s_w^* : v, w \in X^*, h \in G\}$ span a dense subspace of $\mathcal{O}(G, X)$. We will show that each $s_v u_h s_w^*$ belongs to the right-hand side of (3.4). Since (G, X) has nucleus \mathcal{N} , there exists $n \in \mathbb{N}$ such that $h|_y \in \mathcal{N}$ for all $y \in X^n$. But then the Cuntz relation $1 = \sum_{y \in X^n} s_y s_y^*$ gives

$$s_v u_h s_w^* = s_v u_h \left(\sum_{y \in X^n} s_y s_y^* \right) s_w^* = \sum_{y \in X^n} s_v (h \cdot y) u_{h|_y} s_w^*,$$

which belongs to the right-hand side of (3.4). \square

Corollary 3.8. *If (G, X) is contracting with trivial nucleus $\mathcal{N} = \{e\}$, then $\mathcal{O}(G, X)$ is the Cuntz algebra $\mathcal{O}_{|X|}$.*

Proof. If $\mathcal{N} = \{e\}$, then Corollary 3.7 implies that $\mathcal{O}(G, X)$ is generated by the Cuntz family $\{s_x : x \in X\}$, and hence by the uniqueness theorem for the Cuntz algebra is canonically isomorphic to $\mathcal{O}_{|X|}$. \square

3.1. Universal algebras associated with integer matrices. We consider a matrix $A \in M_d(\mathbb{Z})$ with $N = |\det A| > 1$, and the associated self-similar group (\mathbb{Z}^d, Σ) of §2.2. We want to show that $\mathcal{T}(\mathbb{Z}^d, \Sigma)$ and $\mathcal{O}(\mathbb{Z}^d, \Sigma)$ are the Toeplitz algebra $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L)$ and Exel crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ studied in [17].

As in [17] and [7], we consider the the N -to-1 covering map $\sigma_A : \mathbb{T}^d \rightarrow \mathbb{T}^d$ such that $\sigma_A(e^{2\pi i x}) = e^{2\pi i A x}$, and the endomorphism $\alpha_A : f \rightarrow f \circ \sigma_A$ of $C(\mathbb{T}^d)$. The function $L : C(\mathbb{T}^d) \rightarrow C(\mathbb{T}^d)$ defined by

$$L(f)(z) = \frac{1}{N} \sum_{\sigma_A(w)=z} f(w).$$

is a transfer operator for α_A , and $(C(\mathbb{T}^d), \alpha_A, L)$ is the Exel system studied in [7] and [17]. Following [4], we write M_L for the associated Hilbert bimodule over $C(\mathbb{T}^d)$, with inner product and operations given by

$$\langle m, n \rangle := L(m^* n), \quad f \cdot m := f m \quad \text{and} \quad m \cdot f := m \alpha_A(f)$$

for $m, n \in M_L$ and $f \in C(\mathbb{T}^d)$. It is shown in [18, Lemma 3.3] that M_L is complete in the norm defined by the inner product. The Toeplitz algebra $\mathcal{T}(C(\mathbb{T}^d), \alpha_A, L)$ in [17] is by definition the Toeplitz algebra $\mathcal{T}(M_L)$, and the Exel crossed product $C(\mathbb{T}^d) \rtimes_{\alpha_A, L} \mathbb{N}$ is the quotient $\mathcal{O}(M_L)$ (by [4]).

In [17, Proposition 3.1], we showed that $\mathcal{T}(M_L)$ is the universal algebra generated by a unitary representation $u : \mathbb{Z}^d \rightarrow \mathcal{U}\mathcal{T}(M_L)$ and an isometry v satisfying

$$(E1) \quad vu_m = u_{Bm}v,$$

$$(E2) \quad v^*u_mv = \begin{cases} u_{B^{-1}m} & \text{if } m \in B\mathbb{Z}^d \\ 0 & \text{otherwise.} \end{cases}$$

We will use this presentation and that of Proposition 3.2 to identify $\mathcal{T}(\mathbb{Z}^d, \Sigma)$ with $\mathcal{T}(M_L)$. (The use of the same letter u for the unitary representation of \mathbb{Z}^d in both presentations should not cause problems because our isomorphism takes one u_n to the other u_n .)

Proposition 3.9. *Suppose that $A \in M_d(\mathbb{Z})$ has $|\det A| > 1$, write $B := A^t$, and consider the bimodule M_L constructed above. Define $b : \Sigma^* \rightarrow \mathbb{Z}^d$ by $b(w) = w_1 + Bw_2 + \dots + B^{k-1}w_k$ for $w \in \Sigma^k$. Then there is an isomorphism θ of $\mathcal{T}(\mathbb{Z}^d, \Sigma) = C^*(u, s)$ onto $\mathcal{T}(M_L) = C^*(u, v)$ such that*

$$\theta(s_y u_n s_w^*) = u_{b(y)+B^k n} v^k v^{*l} u_{b(w)}^* \quad (3.5)$$

$$= u_{b(y)} v^k v^{*l} u_{b(w)+B^l n}^* \quad (3.6)$$

for $y \in \Sigma^k, w \in \Sigma^l, n \in \mathbb{Z}^d$.

Proof. We begin by building a representation of $\mathcal{T}(\mathbb{Z}^d, \Sigma)$ in $\mathcal{T}(M_L) = C^*(u, v)$. We have the u_n , and they satisfy condition (a) of Proposition 3.2 because $u : \mathbb{Z}^d \rightarrow \mathcal{T}(M_L)$ is a unitary representation. For $x \in \Sigma$, we define $S_x = u_x v$. Then for $x, y \in \Sigma$, property (E2) gives

$$S_x^* S_y = v^* u_{-x} u_y v = v^* u_{y-x} v = \begin{cases} u_{B^{-1}(y-x)} & \text{if } y-x \in B\mathbb{Z}^d \\ 0 & \text{otherwise;} \end{cases}$$

since both x and y are in Σ , $y-x \in B\mathbb{Z}^d$ if and only if $x = y$. Thus the $\{S_x\}$ are isometries with orthogonal ranges, and form a Toeplitz-Cuntz family, as required in Proposition 3.2 (b). Next we use (E1):

$$\begin{aligned} u_n S_x &= u_{n+x} v = u_{c(n+x)} u_{n+x-c(n+x)} v \\ &= u_{c(n+x)} v u_{B^{-1}(n+x-c(n+x))} \\ &= S_{c(n+x)} u_{B^{-1}(n+x-c(n+x))}, \end{aligned}$$

which is $S_{n \cdot x} u_{n|x}$ by (2.5). Now the universal property of $(\mathcal{T}(\mathbb{Z}^d, \Sigma), u, s)$ gives us a homomorphism $\theta = \theta_{u, S} : \mathcal{T}(\mathbb{Z}^d, \Sigma) \rightarrow \mathcal{T}(M_L)$ such that $\theta(s_x) = u_x v$ and $\theta \circ u = u$. The range contains all the generators u_n and $v = S_0$, and hence θ is onto.

To see that θ is injective, we build an inverse. We define $V := s_0$. Then V is certainly an isometry. For $m \in \mathbb{Z}^d$, an application of (2.5) gives

$$u_{Bm} V = u_{Bm} s_0 = s_{c(0+Bm)} u_{Bm|0} = s_0 u_{B^{-1}(Bm+0-c(Bm+0))} = V u_m,$$

which is (E1). For (E2), we use (2.5) again:

$$V^* u_m V = s_0^* u_m s_0 = s_0^* s_{c(m)} u_{B^{-1}(m-c(m))};$$

since the s_x have mutually orthogonal ranges, this last term vanishes unless $c(m) = 0$, or equivalently $m \in B\mathbb{Z}^d$, in which case it is $s_0^* s_0 u_{B^{-1}m} = u_{B^{-1}m}$. Now the universal property of $\mathcal{T}(M_L)$ gives a homomorphism $\theta' : \mathcal{T}(M_L) \rightarrow \mathcal{T}(\mathbb{Z}^d, \Sigma)$ such that $\theta'(u_m) =$

u_m and $\theta'(v) = V$. Then $\theta' \circ \theta(s_x) = \theta'(u_x v) = u_x s_0 = s_x$, so $\theta' \circ \theta$ is the identity, and θ is injective.

To check the formulas for θ on spanning elements, we take $y \in \Sigma^k$, $w \in \Sigma^l$ and $n \in \mathbb{Z}^d$. Then

$$\begin{aligned} \theta(s_y u_n s_w^*) &= \theta(s_{y_1} s_{y_2} \cdots s_{y_k} u_n s_w^*) = u_{y_1} v u_{y_2} v \cdots u_{y_k} v u_n s_w^* \\ &= u_{y_1} u_{B y_2} v^2 \cdots u_{y_k} v u_n s_w^* \\ &= u_{y_1 + B y_2 + \cdots + B^{k-1} y_k} v^k u_n v^{*l} u_{w_1 + B w_2 + \cdots + B^{l-1} w_l}^* \\ &= u_{y_1 + B y_2 + \cdots + B^{k-1} y_k + B^k n} v^k v^{*l} u_{w_1 + B w_2 + \cdots + B^{l-1} w_l}^* \end{aligned}$$

which is the first formula (3.5). For (3.5), instead of pulling u_n past v^k using (E1) at the last step, pull it past v^{*l} using the adjoint of (E1). \square

Corollary 3.10. *Suppose that A and M_L are as above. Then the isomorphism θ of Proposition 3.9 induces an isomorphism $\bar{\theta}$ of $\mathcal{O}(\mathbb{Z}^d, \Sigma) = C^*(u_n, s_x)$ onto $\mathcal{O}(M_L) = C^*(\bar{u}_n, \bar{v})$ such that*

$$\bar{\theta}(s_v u_n s_w^*) = \bar{u}_{b(v) + B^k n} \bar{v}^k \bar{v}^{*l} \bar{u}_{b(w)}^*.$$

Proof. We know from Corollary 3.5 that $\mathcal{O}(\mathbb{Z}^d, \Sigma)$ is the quotient of $\mathcal{T}(\mathbb{Z}^d, \Sigma)$ by the ideal I generated by $1 - \sum_{x \in \Sigma} s_x s_x^*$, and from [17, Proposition 3.3] that $\mathcal{O}(M_L)$ is the quotient of $\mathcal{T}(M_L)$ by the ideal J generated by $1 - \sum_{x \in \Sigma} (u_x v)(u_x v)^*$. Since $\theta(s_x) = u_{b(x)} v = u_x v$, we have

$$\theta\left(1 - \sum_{x \in \Sigma} s_x s_x^*\right) = 1 - \sum_{x \in \Sigma} (u_x v)(u_x v)^*,$$

so $\theta(I) = J$, and the result follows. \square

4. A CHARACTERISATION OF KMS STATES

Let (G, X) be a self-similar action. The Toeplitz algebra $\mathcal{T}(G, X) = \mathcal{T}(M)$ carries a strongly continuous gauge action $\gamma : \mathbb{T} \rightarrow \text{Aut}(\mathcal{T}(G, X))$ such that $\gamma_z(i_{C^*(G)}(a)) = i_{C^*(G)}(a)$ for $a \in C^*(G)$ and $\gamma_z(i_M(m)) = z i_M(m)$ for $m \in M$. We define $\sigma : \mathbb{R} \rightarrow \text{Aut}(\mathcal{T}(G, X))$ by $\sigma_t = \gamma_{e^{it}}$. In terms of the presentation of Proposition 3.2, we have

$$\sigma_t(u_g) = u_g \quad \text{and} \quad \sigma_t(s_v) = e^{it|v|} s_v.$$

We also write σ for the induced action of \mathbb{R} on $\mathcal{O}(G, X)$. Our main goal is to find the KMS states of $(\mathcal{T}(G, X), \sigma)$ and $(\mathcal{O}(G, X), \sigma)$. In this section, we give a characterisation of KMS states which will make them easier to identify.

Our conventions for KMS states are the same as those of [16] and [17], and are explained at the beginning of [16, §7], for example. For our purposes, a state ϕ of a system (B, \mathbb{R}, α) is a *KMS state with inverse temperature β* (a KMS_β state) if $\phi(ab) = \phi(b\alpha_{i\beta}(a))$ for all a, b in a family \mathcal{F} of analytic elements which span a dense subspace of B . We distinguish between KMS_∞ states, which are by definition limits of KMS_β states as $\beta \rightarrow \infty$, and ground states, for which $z \mapsto \phi(\alpha_z(b))$ is bounded in the upper-half plane for all $a, b \in \mathcal{F}$. (This distinction is not made in the standard references [3, 23].)

The spanning elements $s_v u_g s_w^* \in \mathcal{T}(G, X)$ are analytic for σ since

$$\sigma_t(s_v u_g s_w^*) = e^{it(|v|-|w|)} s_v u_g s_w^* \tag{4.1}$$

and the function $z \mapsto e^{iz(|v|-|w|)}$ is entire. Thus a state ϕ is KMS_β for σ if and only if

$$\begin{aligned} \phi((s_v u_g s_w^*)(s_y u_h s_z^*)) &= \phi((s_y u_h s_z^*)\sigma_{i\beta}(s_v u_g s_w^*)) \\ &= e^{-\beta(|v|-|w|)} \phi((s_y u_h s_z^*)(s_v u_g s_w^*)). \end{aligned} \tag{4.2}$$

We now have the following analogue of [16, Lemma 8.3] and [17, Proposition 4.1].

Proposition 4.1. *Let (G, X) be a self-similar action and suppose that $\sigma : \mathbb{R} \rightarrow \text{Aut } \mathcal{T}(G, X)$ satisfies (4.1).*

- (1) *For $\beta < \log |X|$, there are no KMS_β -states for σ .*
(2) *For $\beta \geq \log |X|$, a state ϕ is a KMS_β -state for σ if and only if*

$$\phi(\mathbf{u}_g \mathbf{u}_h) = \phi(\mathbf{u}_h \mathbf{u}_g) \quad \text{for } g, h \in G \quad (4.3)$$

and

$$\phi(s_v \mathbf{u}_g s_w^*) = \begin{cases} e^{-\beta|v|} \phi(\mathbf{u}_g) & \text{if } v = w \\ 0 & \text{otherwise} \end{cases} \quad (4.4)$$

Proof. Suppose that ϕ is a KMS_β -state. First, for $g, h \in G$ we have

$$\phi(\mathbf{u}_g \mathbf{u}_h) = \phi(\mathbf{u}_h \sigma_{i\beta}(\mathbf{u}_g)) = \phi(\mathbf{u}_h \mathbf{u}_g).$$

Next, we take $v, w \in X^*$ and calculate

$$\begin{aligned} \phi(s_v \mathbf{u}_g s_w^*) &= \phi(\mathbf{u}_g s_w^* \sigma_{i\beta}(s_v)) \\ &= \begin{cases} e^{-\beta|v|} \phi(\mathbf{u}_g s_w^* s_v) & \text{if } v = wv' \text{ or } w = vw' \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} e^{-\beta|v|} \phi(s_v \sigma_{i\beta}(\mathbf{u}_g s_w^*)) & \text{if } v = wv' \text{ or } w = vw' \\ 0 & \text{otherwise} \end{cases} \\ &= \begin{cases} e^{-\beta(|v|-|w|)} \phi(s_v \mathbf{u}_g s_w^*) & \text{if } v = wv' \text{ or } w = vw' \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Thus

$$\phi(s_v \mathbf{u}_g s_w^*) \neq 0 \implies |v| = |w| \text{ and either } v = wv' \text{ or } w = vw' \iff v = w.$$

If $v = w$, then

$$\phi(s_v \mathbf{u}_g s_v^*) = \phi(\mathbf{u}_g s_v^* \sigma_{i\beta}(s_v)) = e^{-\beta|v|} \phi(\mathbf{u}_g s_v^* s_v) = e^{-\beta|v|} \phi(\mathbf{u}_g).$$

Since $\{s_x : x \in X\}$ is a Toeplitz-Cuntz family, we have

$$1 = \phi(1) \geq \phi\left(\sum_{x \in X} s_x s_x^*\right) = \sum_{x \in X} \phi(s_x s_x^*) = \sum_{x \in X} e^{-\beta} \phi(s_x^* s_x) = \sum_{x \in X} e^{-\beta} = |X| e^{-\beta},$$

so that $\beta \geq \log |X|$. This completes the proof of (1) and the forward implication in (2).

For the backward implication in (2), suppose ϕ is a tracial state on $C^*(G)$ satisfying (4.3) and (4.4). We aim to show that ϕ satisfies (4.2). We first suppose that $|y| \geq |w|$. Since y is longer than w , the product $s_w^* s_y$ in the middle of the left-hand side of (4.2) vanishes unless $y = wy'$. If $y = wy'$, then Lemma 3.4 implies that

$$\phi((s_v \mathbf{u}_g s_w^*)(s_y \mathbf{u}_h s_z^*)) = \phi(s_{v(g \cdot y')} \mathbf{u}_{g|_y, h} s_z^*),$$

which by (4.4) vanishes unless $z = v(g \cdot y')$. For $y = wy'$ and $z = v(g \cdot y')$, we compute

$$\begin{aligned} \phi((s_v \mathbf{u}_g s_w^*)(s_y \mathbf{u}_h s_z^*)) &= \phi(s_{v(g \cdot y')} \mathbf{u}_{g|_y, h} s_z^*) \\ &= e^{-\beta|v(g \cdot y')|} \phi(\mathbf{u}_{g|_y, h}) \\ &= e^{-\beta(|v|+|y'|)} \phi(\mathbf{u}_h \mathbf{u}_{g|_y, h}) \quad (\text{using (4.3)}) \\ &= e^{-\beta(|v|+|y'|)} e^{\beta|y|} \phi(s_y \mathbf{u}_h \mathbf{u}_{g|_y, h} s_y^*) \quad (\text{using (4.4)}) \\ &= e^{-\beta(|v|+|y'|-|y|)} \phi(s_y \mathbf{u}_h \mathbf{u}_{g|_y, h} s_y^* s_w^*) \\ &= e^{-\beta(|v|-|w|)} \phi(s_y \mathbf{u}_h (s_y' \mathbf{u}_{g^{-1}|_{g \cdot y}})^* s_w^*) \end{aligned}$$

$$\begin{aligned}
&= e^{-\beta(|v|-|w|)} \phi(s_y \mathbf{u}_h (s_{g^{-1}(g \cdot y')} \mathbf{u}_{g^{-1}|_{g \cdot y'}})^* s_w^*) \\
&= e^{-\beta(|v|-|w|)} \phi(s_y \mathbf{u}_h (\mathbf{u}_{g^{-1} s_{g \cdot y'}})^* s_w^*) \\
&= e^{-\beta(|v|-|w|)} \phi(s_y \mathbf{u}_h s_{g \cdot y'}^* \mathbf{u}_g s_w^*) \\
&= e^{-\beta(|v|-|w|)} \phi(s_y \mathbf{u}_h s_{v(g \cdot y')}^* s_v \mathbf{u}_g s_w^*) \\
&= e^{-\beta(|v|-|w|)} \phi((s_y \mathbf{u}_h s_z^*)(s_v \mathbf{u}_g s_w^*)). \tag{4.5}
\end{aligned}$$

To complete the proof of (2), we observe that

$$\begin{aligned}
\phi((s_y \mathbf{u}_h s_z^*)(s_v \mathbf{u}_g s_w^*)) &\neq 0 \\
&\implies \text{either } v = zv' \text{ and } w = y(h \cdot v'), \text{ or } z = vz' \text{ and } y = w(g^{-1} \cdot z') \\
&\implies z = vz' \text{ and } y = w(g^{-1} \cdot z') \quad (\text{because } |y| \geq |w|) \\
&\implies z = v(g \cdot y') \text{ and } y = wy' \quad (\text{with } y' = g^{-1} \cdot z').
\end{aligned}$$

Thus if there is no y' satisfying $y = wy'$ and $z = v(g \cdot y')$, we have

$$e^{-\beta(|v|-|w|)} \phi((s_y \mathbf{u}_h s_z^*)(s_v \mathbf{u}_g s_w^*)) = 0 = \phi((s_v \mathbf{u}_g s_w^*)(s_y \mathbf{u}_h s_z^*)). \tag{4.6}$$

Together, (4.5), which holds when there exists y' such that $y = wy'$ and $z = v(g \cdot y')$, and (4.6), which holds otherwise, imply (4.2) for $|y| \geq |w|$.

We now suppose that $|y| < |w|$, and take adjoints to reduce to the case in the previous paragraph. Since $\phi(a) = \overline{\phi(a^*)}$, we have

$$\begin{aligned}
\phi((s_v \mathbf{u}_g s_w^*)(s_y \mathbf{u}_h s_z^*)) &= \overline{\phi((s_z \mathbf{u}_{h^{-1}} s_y^*)(s_w \mathbf{u}_{g^{-1}} s_v^*))} \\
&= e^{-\beta(|z|-|y|)} \overline{\phi((s_w \mathbf{u}_{g^{-1}} s_v^*)(s_z \mathbf{u}_{h^{-1}} s_y^*))} \\
&= e^{-\beta(|z|-|y|)} \phi((s_y \mathbf{u}_h s_z^*)(s_v \mathbf{u}_g s_w^*)). \tag{4.7}
\end{aligned}$$

The calculation in the previous paragraph shows that the right-hand side of (4.7) vanishes unless $w = yw'$ and $v = z(g^{-1} \cdot w')$, in which case $|v| - |z| = |w'| = |w| - |y|$ and $|z| - |y| = |v| - |w|$. Thus

$$\phi((s_v \mathbf{u}_g s_w^*)(s_y \mathbf{u}_h s_z^*)) = e^{-\beta(|v|-|w|)} \phi((s_y \mathbf{u}_h s_z^*)(s_v \mathbf{u}_g s_w^*)),$$

and we have proved (4.2) in the remaining case $|y| < |w|$. \square

5. EXISTENCE OF KMS STATES ABOVE THE CRITICAL INVERSE TEMPERATURE

Theorem 5.1. *Let (G, X) be a self-similar action, and let τ be a normalised trace on $C^*(G)$. Then for every $\beta > \log |X|$, there is a KMS_β state $\psi_{\beta, \tau}$ satisfying*

$$\psi_{\beta, \tau}(s_v \mathbf{u}_g s_w^*) = \begin{cases} (1 - |X|e^{-\beta}) \sum_{k=0}^{\infty} e^{-\beta(k+|v|)} \left(\sum_{\{y \in X^k : g \cdot y = v\}} \tau(\delta_{g|_y}) \right) & \text{if } v = w \\ 0 & \text{otherwise.} \end{cases} \tag{5.1}$$

To prove Theorem 5.1, we adapt ideas from the proofs of [14, Theorem 2.1] and [17, Proposition 6.1]. Both involve induced representations; as in [14] rather than [17], we apply Rieffel's Hilbert-bimodule formulation of induced representations to the Fock bimodule $\mathcal{F}(M) := \bigoplus_{j=0}^{\infty} M^{\otimes j}$. We take $\pi_\tau : C^*(G) \rightarrow B(\mathcal{K}_\tau)$ to be the GNS-representation of $C^*(G)$, and then Rieffel induction gives a representation

$$\pi := \bigoplus_{j=0}^{\infty} M^{\otimes j}\text{-Ind } \pi_\tau \quad \text{on the Hilbert space} \quad \mathcal{H}_\pi := \bigoplus_{j=0}^{\infty} M^{\otimes j} \otimes_{C^*(G)} \mathcal{K}_\tau. \tag{5.2}$$

To calculate with the induced representations $M^{\otimes j}$ -Ind π_τ , we need to understand the bimodules $M^{\otimes j}$.

Lemma 5.2. *Suppose that (G, X) is a self-similar action and $\{e_x : x \in X\}$ is the orthonormal basis for the Hilbert bimodule M constructed in §3. Fix $j \geq 1$. Then*

$$\{e_v := e_{v_1} \otimes \cdots \otimes e_{v_j} : v \in X^j\}$$

is an orthonormal basis for $M^{\otimes j}$ with reconstruction formula

$$m = \sum_{v \in X^j} e_v \cdot \langle e_v, m \rangle \quad \text{for } m \in M^{\otimes j}. \quad (5.3)$$

The left action of $C^*(G)$ on $M^{\otimes j}$ satisfies $\delta_g \cdot e_v = e_{g \cdot v} \cdot \delta_{g|v}$.

Proof. We prove this result by induction on j . Proposition 3.1 and the surrounding discussion give the result for $j = 1$. Suppose it is true for $j = k$. For two words $w = w_1 w'$ and $v = v_1 v'$ in X^{k+1} , we have $|w'| = |v'| = k$, and

$$\begin{aligned} \langle e_w, e_v \rangle &= \langle e_{w_1} \otimes e_{w'}, e_{v_1} \otimes e_{v'} \rangle = \langle e_{w'}, \langle e_{w_1}, e_{v_1} \rangle \cdot e_{v'} \rangle \\ &= \delta_{w_1, v_1} \langle e_{w'}, e_{v'} \rangle = \delta_{w_1, v_1} \delta_{w', v'} 1_{C^*(G)} = \delta_{w, v} 1_{C^*(G)}, \end{aligned}$$

giving orthonormality. For $m = m_1 \otimes m' \in M^{\otimes(k+1)} = M \otimes_{C^*(G)} M^{\otimes k}$, we have

$$\begin{aligned} m &= \left(\sum_{x \in X} e_x \cdot \langle e_x, m_1 \rangle \right) \otimes m' = \sum_{x \in X} e_x \otimes \langle e_x, m_1 \rangle \cdot m' \\ &= \sum_{x \in X} e_x \otimes \left(\sum_{v \in X^k} e_v \cdot \langle e_v, \langle e_x, m_1 \rangle \cdot m' \rangle \right) \\ &= \sum_{x \in X, v \in X^k} e_x \otimes e_v \cdot \langle e_x \otimes e_v, m_1 \otimes m' \rangle, \\ &= \sum_{x \in X, v \in X^k} e_{xv} \cdot \langle e_{xv}, m \rangle, \end{aligned}$$

which is the right-hand side of (5.3) for $j = k + 1$. This formula extends by linearity and continuity of the inner product to $m \in M^{\otimes(k+1)}$. Finally, let $w = w_1 w' \in X^{k+1}$ and $g \in G$. Then because the tensor product is balanced over $C^*(G)$, the inductive hypothesis gives

$$\begin{aligned} \delta_g \cdot e_w &= (\delta_g \cdot e_{w_1}) \otimes e_{w'} = (e_{g \cdot w_1} \cdot \delta_{g|w_1}) \otimes e_{w'} \\ &= e_{g \cdot w_1} \otimes (\delta_{g|w_1} \cdot e_{w'}) = e_{g \cdot w_1} \otimes (e_{g|w_1 \cdot w'} \cdot \delta_{(g|w_1)|w'}) \\ &= (e_{g \cdot w_1} \otimes e_{g|w_1 \cdot w'}) \cdot \delta_{g|w} = e_{g \cdot w} \cdot \delta_{g|w}, \end{aligned}$$

and we now have the whole inductive hypothesis for $j = k + 1$. \square

Proof of Theorem 5.1. As promised, we take the GNS representation π_τ of $C^*(G)$ on \mathcal{K}_τ , and consider the representation π of (5.2). Lemma 5.2 implies that every vector in $M^{\otimes j} \otimes_{C^*(G)} \mathcal{K}_\tau$ is a finite sum $\sum_{v \in X^j} e_v \otimes k_v$, and that the representation $M^{\otimes j}$ -Ind π_τ of $C^*(G)$ is the integrated form of the unitary representation U^j of G on $M^{\otimes j} \otimes_{C^*(G)} \mathcal{K}_\tau$ characterised by

$$U_g^j(e_v \otimes k) = T_g(e_v) \otimes k = (e_{g \cdot v} \cdot \delta_{g|v}) \otimes k = e_{g \cdot v} \otimes \pi_\tau(\delta_{g|v})k; \quad (5.4)$$

we set $U_g = \bigoplus U_g^j$. Since the $\{e_v \otimes k : v \in X^j\}$ are mutually orthogonal, there are isometries S_x on \mathcal{H}_π such that $S_x(e_v \otimes k) = e_{xv} \otimes k$, and these isometries form a Toeplitz-Cuntz family. The following calculation using (5.4) shows that U and S satisfy (SSR):

$$S_{g \cdot x} U_{g|x}(e_v \otimes k) = S_{g \cdot x}(e_{g|x \cdot v} \otimes \pi_\tau(\delta_{(g|x)|v})k) = e_{(g \cdot x)(g|x \cdot v)} \otimes \pi_\tau(\delta_{g|x \cdot v})k$$

$$= e_{g \cdot (xv)} \otimes \pi_\tau(\delta_{g|_{xv}})k = U_g(e_{xv} \otimes k) = U_g S_x(e_v \otimes k).$$

Now Proposition 3.2 gives us a representation $\pi_{U,S} : \mathcal{T}(G, X) \rightarrow B(\mathcal{H}_\pi)$ such that $\pi_{U,S}(s_v u_g s_w^*) = S_v U_g S_w^*$.

We now take ξ_τ to be the canonical cyclic vector for the GNS representation π_τ (so that ξ_τ is the image in \mathcal{K}_τ of the identity $1_{C^*(G)}$), and define $\psi_{\beta,\tau} : \mathcal{T}(G, X) \rightarrow \mathbb{C}$ by

$$\psi_{\beta,\tau}(a) := (1 - |X|e^{-\beta}) \sum_{j=0}^{\infty} e^{-\beta j} \left(\sum_{z \in X^j} (\pi_{U,S}(a)(e_z \otimes \xi_\tau) | e_z \otimes \xi_\tau) \right). \quad (5.5)$$

Since $\psi_{\beta,\tau}$ is a norm-convergent sum of vector states with non-negative coefficients, it is a positive functional; since $|X^j| = |X|^j$, summing the geometric series $\sum_j (|X|e^{-\beta})^j$ shows that $\psi_{\beta,\tau}(1) = 1$, and $\psi_{\beta,\tau}$ is a state.

To verify (5.1), we take $a = s_v u_g s_w^*$. Then

$$\begin{aligned} (\pi_{\beta,\tau}(a)(e_z \otimes \xi_\tau) | e_z \otimes \xi_\tau) &= (S_v U_g S_w^*(e_z \otimes \xi_\tau) | e_z \otimes \xi_\tau) \\ &= (U_g S_w^*(e_z \otimes \xi_\tau) | S_v^*(e_z \otimes \xi_\tau)). \end{aligned} \quad (5.6)$$

We have $S_w^*(e_z \otimes \xi_\tau) = 0$ unless $z = wz'$, and hence (5.6) vanishes unless $z = wz' = vz''$, in which case

$$\begin{aligned} (\pi_{\beta,\tau}(a)(e_z \otimes \xi_\tau) | e_z \otimes \xi_\tau) &= (U_g(e_{z'} \otimes \xi_\tau) | e_{z''} \otimes \xi_\tau) \\ &= (e_{g \cdot z'} \otimes \pi_\tau(\delta_{g|_{z'}}) \xi_\tau | e_{z''} \otimes \xi_\tau). \end{aligned}$$

This last inner product vanishes unless $g \cdot z' = z''$, which implies $|z'| = |z''|$ and $|v| = |z| - |z'| = |z| - |z''| = |w|$; now $z = wz' = vz''$ forces $v = w$ and $z' = z''$. Thus the inner product vanishes unless $z = vz'$ and $g \cdot z' = z'$. Noticing that $z = vz'$ implies $|z| \geq |v|$ and writing y for z' , we find that

$$\psi_{\beta,\tau}(s_v u_g s_w^*) = \begin{cases} (1 - |X|e^{-\beta}) \sum_{j=|v|}^{\infty} e^{-\beta j} \left(\sum_{\{y \in X^{j-|v|} : g \cdot y = y\}} (\pi_\tau(\delta_{g|_y}) \xi_\tau | \xi_\tau) \right) & \text{if } v = w \\ 0 & \text{otherwise.} \end{cases}$$

Since $(\pi_\tau(\delta_{g|_y}) \xi_\tau | \xi_\tau) = \tau(1_{C^*(G)}^* \delta_{g|_y} 1_{C^*(G)}) = \tau(\delta_{g|_y})$, taking $k = j - |v|$ gives (5.1).

We show that $\psi_{\beta,\tau}$ is a KMS_β state by checking properties (4.4) and (4.3) of Proposition 4.1. The first is straightforward. We trivially have $\psi_{\beta,\tau}(s_v u_g s_w^*) = 0$ if $v \neq w$. For $g \in G$ and $v \in X^j$, we have

$$e^{-\beta|v|} \psi_{\beta,\tau}(u_g) = (1 - |X|e^{-\beta}) e^{-\beta|v|} \sum_{k=0}^{\infty} e^{-\beta k} \left(\sum_{\{y \in X^k : g \cdot y = y\}} \tau(\delta_{g|_y}) \right),$$

which on pulling $e^{-\beta|v|}$ inside the sum becomes the right-hand side of the formula (5.1) for $\psi_{\beta,\tau}(s_v u_g s_v^*)$. For the second, we need to take $g, h \in G$ and compare

$$\psi_{\beta,\tau}(u_g u_h) = \psi_{\beta,\tau}(u_{gh}) = (1 - |X|e^{-\beta}) \sum_{k=0}^{\infty} e^{-\beta k} \left(\sum_{\{y \in X^k : (gh) \cdot y = y\}} \tau(\delta_{(gh)|_y}) \right) \quad (5.7)$$

with

$$\psi_{\beta,\tau}(u_h u_g) = \psi_{\beta,\tau}(u_{hg}) = (1 - |X|e^{-\beta}) \sum_{k=0}^{\infty} e^{-\beta k} \left(\sum_{\{z \in X^k : (hg) \cdot z = z\}} \tau(\delta_{(hg)|_z}) \right). \quad (5.8)$$

The function $f : X^k \rightarrow X^k$ defined by $f(y) = h \cdot y$ is a bijection, and

$$(gh) \cdot y = y \iff g \cdot (h \cdot y) = h^{-1} \cdot (h \cdot y) \iff (hg) \cdot (h \cdot y) = h \cdot y,$$

so f maps the index set $\{y \in X^k : (gh) \cdot y = y\}$ in (5.7) onto the one $\{z \in X^k : (hg) \cdot z = z\}$ in (5.8). We claim that the function f also matches up the corresponding summands. To see this, suppose $(gh) \cdot y = y$. Then because τ is a trace, we have

$$\tau(\delta_{(gh)|y}) = \tau(\delta_{g|h \cdot y}) = \tau(\delta_{h|y} \delta_{g|h \cdot y}) = \tau(\delta_{h|h^{-1} \cdot (h \cdot y)} \delta_{g|h \cdot y}).$$

The identity $(gh) \cdot y = y$ implies that $h^{-1} \cdot (h \cdot y) = g \cdot (h \cdot y)$, so

$$\tau(\delta_{(gh)|y}) = \tau(\delta_{h|g \cdot (h \cdot y)} \delta_{g|h \cdot y}) = \tau(\delta_{(hg)|h \cdot y}) = \tau(\delta_{(hg)|f(y)}),$$

as claimed. We deduce that $\psi_{\beta, \tau}(u_g u_h) = \psi_{\beta, \tau}(u_h u_g)$, and now Proposition 5.1 implies that $\psi_{\beta, \tau}$ is a KMS_β state. \square

While we have the formulas for the induced representations handy, we describe the ground states and KMS_∞ states of our system.

Proposition 5.3. *Suppose that (G, X) is a self-similar action. Then for every state ω of $C^*(G)$, there is a ground state ϕ_ω on $(\mathcal{T}(G, X), \sigma)$ such that*

$$\phi_\omega(s_v u_g s_w^*) = \begin{cases} \omega(\delta_g) & \text{if } v = w = \emptyset \\ 0 & \text{otherwise.} \end{cases} \quad (5.9)$$

The map $\omega \mapsto \phi_\omega$ is an affine homeomorphism of the state space $S(C^*(G))$ onto the ground states of $(\mathcal{T}(G, X), \sigma)$. For $\omega \in S(C^*(G))$, ϕ_ω is a KMS_∞ state if and only if ω is a trace.

That states on $C^*(G)$ give ground states is proved in greater generality in [14, Theorem 2.2]. However, as in Theorem 5.1, we can use the special features of our situation to give specific formulas.

We begin with an analogue of [16, Lemma 8.4] which will allow us to recognise ground states. The proof of that lemma carries over almost verbatim to this situation.

Lemma 5.4. *Suppose that (G, X) is a self-similar action. A state ϕ of $\mathcal{T}(G, X)$ is a ground state of $(\mathcal{T}(G, X), \sigma)$ if and only if*

$$\phi(s_v u_g s_w^*) = \begin{cases} \phi(u_g) & \text{if } v = w = \emptyset \\ 0 & \text{otherwise.} \end{cases} \quad (5.10)$$

Proof of Proposition 5.3. Given a state f of $C^*(G)$, we take the GNS representation π_f of $C^*(G)$ on \mathcal{K}_f with cyclic vector ξ_f , and consider the representation $\pi_{u, s}$ of $\mathcal{T}(G, X)$ on $\bigoplus_{j=0}^{\infty} M^{\otimes j} \otimes_{C^*(G)} \mathcal{K}_f$, as in the proof of Theorem 5.1. Then we define

$$\phi_f(a) = (\pi_{u, s}(a)(e_\emptyset \otimes \xi_f) | e_\emptyset \otimes \xi_f) \quad \text{for } a \in \mathcal{T}(G, X).$$

Then ϕ_ω is a state, and

$$\begin{aligned} \phi_f(s_v u_g s_w^*) &= (S_v U_g S_w^*(e_\emptyset \otimes \xi_f) | e_\emptyset \otimes \xi_f) \\ &= \begin{cases} 0 & \text{unless } v = w = \emptyset \\ (\pi_f(\delta_g) \xi_f | \xi_f) = f(\delta_g) & \text{if } v = w = \emptyset. \end{cases} \end{aligned}$$

Lemma 5.4 implies that ϕ_f is a ground state. The map $\omega \mapsto \phi_f$ is continuous, affine and injective, and it is onto because $\phi = \phi_f$ for $f = \phi|_{C^*(G)}$.

If ϕ is a KMS_∞ state, then ϕ is the limit of a sequence of KMS_β states, and equation (4.3) in Proposition 4.1 implies that $\tau := \phi|_{C^*(G)}$ is a trace. For the converse, suppose that τ is a trace on $C^*(G)$. Then we can use weak* compactness to get a sequence $\{\psi_{\beta_n, \tau}\}$

which converges to a KMS_∞ state ϕ . Since (4.4) gives $\psi_{\beta,\tau}(s_\nu u_g s_\nu^*) = e^{-\beta|\nu|} \psi_{\beta,\tau}(u_g)$, we have $\psi_{\beta,\tau}(s_\nu u_g s_\nu^*) \rightarrow 0$ as $\beta \rightarrow \infty$ whenever $|\nu| > 0$. On the other hand, (5.1) gives

$$\psi_{\beta,\tau}(u_g) = (1 - |X|e^{-\beta}) \left(\tau(\delta_g) + \sum_{k=1}^{\infty} e^{-k\beta} \left(\sum_{\{y \in X^k : g \cdot y = y\}} \tau(\delta_{g|_y}) \right) \right).$$

Now

$$\left| \sum_{k=1}^{\infty} e^{-k\beta} \left(\sum_{\{y \in X^k : g \cdot y = y\}} \tau(\delta_{g|_y}) \right) \right| \leq |X|e^{-\beta} \sum_{j=0}^{\infty} |X|^j e^{-\beta j} = \frac{|X|e^{-\beta}}{1 - |X|e^{-\beta}}$$

converges to 0 as $\beta \rightarrow \infty$, and hence $\psi_{\beta,\tau}(u_g) \rightarrow \tau(\delta_g)$. Thus the limit ϕ is the state ϕ_τ described in (5.9), and ϕ_τ is KMS_∞ . \square

In the situation of [17], where the group $G = \mathbb{Z}^d$ is abelian, every state on $C^*(G)$ is a trace, and we recover [17, Proposition 8.1]: every ground state of $(\mathcal{T}(\mathbb{Z}^d, \Sigma), \sigma)$ is a KMS_∞ state. For nonabelian G , though, there are many states of $C^*(G)$ which are not traces, and $(\mathcal{T}(G, X), \sigma)$ has many ground states which are not KMS_∞ states.

6. PARAMETERISATION OF KMS_β STATES ON THE TOEPLITZ ALGEBRA

Theorem 6.1. *Suppose that (G, X) is a self-similar action and $\beta > \log |X|$. The map $\tau \mapsto \psi_{\beta,\tau}$ in Theorem 5.1 is an affine homeomorphism from the simplex of normalised traces on the full group C^* -algebra $C^*(G)$ onto the simplex of KMS_β states on $(\mathcal{T}(G, X), \sigma)$.*

For the proof, we need some lemmas. As in [16, §10] and [17, §7], the idea is to show that a KMS_β state can be reconstructed from its conditioning to a corner $P\mathcal{T}(G, X)P$. Here we take

$$P := 1 - \sum_{x \in X} s_x s_x^* \in \mathcal{T}(G, X).$$

Lemma 6.2. *Suppose that ϕ is a KMS_β state, and define $\phi_P : \mathcal{T}(G, X) \rightarrow \mathbb{C}$ by*

$$\phi_P(a) = \frac{1}{1 - |X|e^{-\beta}} \phi(PaP).$$

Then $\phi_P|_{C^(G)}$ is a normalised trace.*

Proof. The function ϕ_P is a positive linear functional because ϕ is, and the computation

$$\begin{aligned} \phi(1) &= \frac{1}{1 - |X|e^{-\beta}} \phi\left(1 - \sum_{x \in X} s_x s_x^*\right) = \frac{1}{1 - |X|e^{-\beta}} \left(1 - e^{-\beta} \sum_{x \in X} \phi(s_x^* s_x)\right) \\ &= \frac{1 - |X|e^{-\beta}}{1 - |X|e^{-\beta}} = 1 \end{aligned}$$

shows that ϕ_P is a state.

With a view to proving that ϕ_P is tracial on $C^*(G)$, we claim that $u_g P = P u_g$. Indeed, for $x \in X$ and $g \in G$ we have

$$u_g s_x s_x^* = u_g s_x s_x^* u_g^* u_g = (u_g s_x)(u_g s_x)^* u_g = (s_{g \cdot x} u_{g|_x})(s_{g \cdot x} u_{g|_x})^* u_g = s_{g \cdot x} s_{g \cdot x}^* u_g.$$

Thus for $g \in G$, we have

$$\begin{aligned} u_g P &= u_g \left(1 - \sum_{x \in X} s_x s_x^*\right) = u_g - \sum_{x \in X} u_g s_x s_x^* \\ &= u_g - \sum_{x \in X} s_{g \cdot x} s_{g \cdot x}^* u_g = \left(1 - \sum_{x \in X} s_{g \cdot x} s_{g \cdot x}^*\right) u_g, \end{aligned}$$

and since $1 - \sum_{x \in X} s_{g \cdot x} s_{g \cdot x}^* = P$ we get $u_g P = P u_g$, as claimed. Now since ϕ is a KMS_β state, we have

$$\phi(P u_g u_n P) = \phi(u_g P u_n) = \phi(P u_n \sigma_{i\beta}(u_g)) = \phi(P u_n u_g) = \phi(P u_n u_g P),$$

which implies that $\phi|_{C^*(G)}$ is a trace. \square

Lemma 6.3. *Let (G, X) be a self-similar action. For each $n \in \mathbb{N}$, the element*

$$p_n := \sum_{j=0}^n \sum_{v \in X^j} s_v P s_v^*$$

is a projection in $\mathcal{T}(G, X)$, and if ϕ is a KMS_β state of $(\mathcal{T}(G, X), \sigma)$ and $a \in \mathcal{T}(G, X)$, then $\phi(p_n a p_n) \rightarrow \phi(a)$ as $n \rightarrow \infty$.

Proof. Each $s_v P s_v^*$ is a projection, so we need to show that $s_v P s_v^*$ and $s_w P s_w^*$ are mutually orthogonal when $v \neq w$. Since $\{s_z : z \in X^m\}$ is a Toeplitz-Cuntz family for each m , this is trivially true for $|v| = |w|$. So suppose $|v| \neq |w|$. The product $P s_v^* s_w P$ vanishes unless $v = wv'$ or $w = vv'$; since $(P s_v^* s_w P) = P s_w s_v^* P$, we may as well assume that $|w| > |v|$ and $w = vv'$. Then, writing w'_1 for the first letter in w' , we have

$$P s_v^* s_w P = P s_{w'} P = \left(1 - \sum_{x \in X} s_x s_x^*\right) s_{w'} P = s_{w'} P - s_{w'_1} s_{w'_1}^* s_{w'} P = 0. \quad (6.1)$$

Thus each p_n is a projection.

Lemma 7.3 of [17] says that if ϕ is a state of a unital C^* -algebra A , and $\{p_n\}$ is a sequence of projections in A such that $\phi(p_n) \rightarrow 1$, then $\phi(p_n a p_n) \rightarrow \phi(a)$ for every $a \in A$. So we aim to show that $\phi(p_n) \rightarrow 1$ as $n \rightarrow \infty$. The KMS condition gives

$$\begin{aligned} \phi(p_n) &= \sum_{j=0}^n \sum_{v \in X^j} \phi(s_v P s_v^*) = \sum_{j=0}^n \sum_{v \in X^j} e^{-\beta j} \phi(P) \\ &= \phi(P) \sum_{j=0}^n (|X| e^{-\beta})^j = (1 - |X| e^{-\beta}) \sum_{j=0}^n (|X| e^{-\beta})^j, \end{aligned}$$

which converges to 1 as $n \rightarrow \infty$. Thus the result follows from [17, Lemma 7.3] \square

The following reconstruction formula is an analogue of [17, Proposition 7.2].

Lemma 6.4. *Suppose $\beta > \log |X|$ and ϕ is a KMS_β state on $\mathcal{T}(G, X)$. Then for $a \in \mathcal{T}(G, X)$,*

$$\phi(a) = (1 - |X| e^{-\beta}) \sum_{j=0}^{\infty} \sum_{v \in X^j} e^{-\beta j} \phi_P(s_v^* a s_v). \quad (6.2)$$

Proof. Lemma 6.3 gives

$$\begin{aligned} \phi(a) &= \lim_{n \rightarrow \infty} \phi(p_n a p_n) = \lim_{n \rightarrow \infty} \sum_{j=0}^n \sum_{l=0}^n \sum_{v \in X^j} \sum_{w \in X^l} \phi(s_v P s_v^* a s_w P s_w^*) \\ &= \lim_{n \rightarrow \infty} \sum_{j=0}^n \sum_{l=0}^n \sum_{v \in X^j} \sum_{w \in X^l} e^{-\beta j} \phi(P s_v^* a s_w P s_w^* s_v P) \\ &= \lim_{n \rightarrow \infty} \sum_{j=0}^n \sum_{v \in X^j} e^{-\beta j} \phi(P s_v^* a s_v P) \quad (\text{using (6.1)}) \\ &= (1 - |X| e^{-\beta}) \sum_{j=0}^{\infty} \sum_{v \in X^j} e^{-\beta j} \phi_P(s_v^* a s_v). \quad \square \end{aligned}$$

Proof of Theorem 6.1. By an application of the monotone convergence theorem, we can deduce from (5.1) that $\tau \mapsto \psi_{\beta,\tau}$ is affine and weak* continuous. Since both sets of states are weak* compact, it suffices to show that $\tau \mapsto \psi_{\beta,\tau}$ is bijective.

To see injectivity, suppose that $\psi_{\beta,\tau} = \psi_{\beta,\rho}$, and take $g \in G$. Then the formula (5.1) gives

$$\begin{aligned} \psi_{\beta,\tau}(u_g) &= (1 - |\mathcal{X}|e^{-\beta}) \sum_{j=0}^{\infty} \sum_{\{y \in \mathcal{X}^j : g \cdot y = y\}} e^{-\beta j} \tau(\delta_{g|_y}) \\ &= (1 - |\mathcal{X}|e^{-\beta}) \tau(\delta_g) + (1 - |\mathcal{X}|e^{-\beta}) \sum_{k=0}^{\infty} \sum_{\{y \in \mathcal{X}^{k+1} : g \cdot y = y\}} e^{-\beta(k+1)} \tau(\delta_{g|_y}). \end{aligned}$$

We can write the index set for the last sum as

$$\{y \in \mathcal{X}^{k+1} : g \cdot y = y\} = \{xy' : x \in \mathcal{X}, y' \in \mathcal{X}^k, g \cdot x = x, g|_x \cdot y' = y'\},$$

and then another application of (5.1) gives

$$\begin{aligned} \sum_{k=0}^{\infty} \sum_{\{y \in \mathcal{X}^{k+1} : g \cdot y = y\}} e^{-\beta(k+1)} \tau(\delta_{g|_y}) &= e^{-\beta} \sum_{k=0}^{\infty} \sum_{\{x \in \mathcal{X} : g \cdot x = x\}} \sum_{\{y' \in \mathcal{X}^k : g|_x \cdot y' = y'\}} e^{-\beta k} \tau(\delta_{(g|_x)|_{y'}}) \\ &= \frac{e^{-\beta}}{1 - |\mathcal{X}|e^{-\beta}} \left(\sum_{\{x \in \mathcal{X} : g \cdot x = x\}} \psi_{\beta,\tau}(u_{g|_x}) \right). \end{aligned}$$

Thus

$$\psi_{\beta,\tau}(u_g) = (1 - |\mathcal{X}|e^{-\beta}) \tau(\delta_g) + e^{-\beta} \left(\sum_{\{x \in \mathcal{X} : g \cdot x = x\}} \psi_{\beta,\tau}(u_{g|_x}) \right). \quad (6.3)$$

Similarly, we have

$$\psi_{\beta,\rho}(u_g) = (1 - |\mathcal{X}|e^{-\beta}) \rho(\delta_g) + e^{-\beta} \left(\sum_{\{x \in \mathcal{X} : g \cdot x = x\}} \psi_{\beta,\rho}(u_{g|_x}) \right). \quad (6.4)$$

Since $\psi_{\beta,\tau} = \psi_{\beta,\rho}$, subtracting (6.3) from (6.4) shows that $\tau(\delta_g) = \rho(\delta_g)$. Thus $\tau = \rho$, and $\tau \mapsto \psi_{\beta,\tau}$ is injective.

To see surjectivity, suppose that ϕ is a KMS_{β} state on $\mathcal{T}(G, \mathcal{X})$. Lemma 6.2 implies that $\tau := \phi|_{C^*(G)}$ is a normalised trace, and we aim to show that $\phi = \psi_{\beta,\tau}$. By (5.1), it suffices to show that $\phi(u_g) = \psi_{\beta,\tau}(u_g)$ for all $g \in G$. Fix $g \in G$. Then the reconstruction formula (6.2) gives

$$\begin{aligned} \phi(u_g) &= (1 - |\mathcal{X}|e^{-\beta}) \sum_{j=0}^{\infty} \sum_{y \in \mathcal{X}^j} e^{-\beta j} \phi_P(s_y^* u_g s_y) \\ &= (1 - |\mathcal{X}|e^{-\beta}) \sum_{j=0}^{\infty} \sum_{y \in \mathcal{X}^j} e^{-\beta j} \tau(s_y^* s_{g \cdot y} u_{g|_y}) \\ &= \lim_{n \rightarrow \infty} (1 - |\mathcal{X}|e^{-\beta}) \sum_{j=0}^n \sum_{\{y \in \mathcal{X}^j : g \cdot y = y\}} e^{-\beta j} \tau(u_{g|_y}), \end{aligned}$$

which by (5.1) is precisely $\psi_{\beta,\tau}(u_g)$. Thus $\phi = \psi_{\beta,\tau}$, and $\tau \mapsto \psi_{\beta,\tau}$ is surjective. \square

For every discrete group G , there are at least two normalised traces on $C^*(G)$. The usual trace τ_e on $C^*(G)$ satisfies

$$\tau_e(\delta_g) = \begin{cases} 1 & \text{if } g = e \\ 0 & \text{otherwise.} \end{cases}$$

To see that there is such a trace, consider the left-regular representation λ of G on $\ell^2(G)$, and define $\tau_e : C^*(G) \rightarrow \mathbb{C}$ in terms of the usual orthonormal basis $\{\xi_g : g \in G\}$ by $\tau_e(a) = (\lambda(a)\xi_e | \xi_e)$. Then it is easy to check on $\text{span}\{\delta_g\}$ that τ_e has the required properties, and continuity does the rest. The other trace is the integrated form $\tau_1 : C^*(G) \rightarrow \mathbb{C}$ of the trivial representation $g \mapsto 1$, which is a scalar-valued homomorphism, and hence is trivially a trace.

Since τ_e and τ_1 do not agree on the δ_g with $g \neq e$, they are distinct traces, and hence by Theorem 6.1 give distinct KMS states. We look at these states.

Corollary 6.5. *Suppose that (G, X) is a self-similar action and $\beta > \log |X|$. For $g \in G$ and $k \geq 0$, we set*

$$F_g^k := \{v \in X^k : g \cdot v = v \text{ and } g|_v = e\}. \quad (6.5)$$

Then there is a KMS_β state ψ_{β, τ_e} on $(\mathcal{T}(G, X), \sigma)$ such that

$$\psi_{\beta, \tau_e}(s_v u_g s_w^*) = \begin{cases} e^{-\beta|v|} (1 - |X|e^{-\beta}) \sum_{k=0}^{\infty} e^{-\beta k} |F_g^k| & \text{if } v = w \\ 0 & \text{otherwise,} \end{cases}$$

where we interpret $|\emptyset|$ as 0.

Proof. The state ψ_{β, τ_e} is the one given by Theorem 5.1, so we just need to check the formula for $\psi_{\beta, \tau_e}(s_v u_g s_w^*)$. It is certainly 0 if $v \neq w$, so we suppose $v = w$. Then since $\tau_e(\delta_e) = 1$ and $\tau_e(\delta_h) = 0$ for $h \neq e$, the sum on the right-hand side of (5.1) collapses to give

$$\psi_{\beta, \tau_e}(s_v u_g s_v^*) = (1 - |X|e^{-\beta}) \sum_{k=0}^{\infty} e^{-\beta(k+|v|)} \left(\sum_{y \in F_g^k} 1 \right),$$

which on pulling out $e^{-\beta|v|}$ gives the required formula. \square

Corollary 6.6. *Suppose that (G, X) is a self-similar action and $\beta > \log |X|$. For $g \in G$ and $k \geq 0$, we set*

$$G_g^k := \{v \in X^k : g \cdot v = v\}. \quad (6.6)$$

Then there is a KMS_β state ψ_{β, τ_1} on $(\mathcal{T}(G, X), \sigma)$ such that

$$\psi_{\beta, \tau_1}(s_v u_g s_w^*) = \begin{cases} e^{-\beta|v|} (1 - |X|e^{-\beta}) \sum_{k=0}^{\infty} e^{-\beta k} |G_g^k| & \text{if } v = w \\ 0 & \text{otherwise.} \end{cases}$$

Proof. As in the proof of the previous corollary, the second sum on the right-hand side of (5.1) counts the number of elements of G_g^k , and hence this follows from Theorem 5.1. \square

Although the formulas in the last two corollaries look a bit messy, they are quite computable, and we will later discuss ways of doing these computations using Moore diagrams. But it is easy to give a quick example now.

Example 6.7. Consider the basilica group $(B, X = \{x, y\})$ of §2.3. The first two relations in (2.6) imply that the generator a changes the first letter of every word, so $F_a^k = G_a^k = \emptyset$ for every $k \geq 1$, and $\psi_{\beta, \tau_e}(\delta_a) = \psi_{\beta, \tau_1}(\delta_a) = 0$. On the other hand, b fixes x with $b|_x = a$, and hence satisfies $b \cdot (xw) \neq xw$ for every longer word xw . Thus $F_b^k = \{yw : w \in X^*\}$ and $G_b^k = \{yw : w \in X^*\} \cup \{x\}$. We deduce that

$$\psi_{\beta, \tau_e}(\delta_b) = (1 - 2e^{-\beta}) \sum_{k=0}^{\infty} e^{-\beta k} 2^{k-1} = \frac{1}{2}(1 - 2e^{-\beta}) \sum_{k=0}^{\infty} (2e^{-\beta})^k = \frac{1}{2}, \text{ and}$$

$$\psi_{\beta, \tau_1}(\delta_b) = (1 - 2e^{-\beta}) \sum_{k=0}^{\infty} e^{-\beta k} (2^{k-1} + 1) = \frac{1}{2} + \frac{1 - 2e^{-\beta}}{1 - e^{-\beta}}.$$

Remark 6.8. When the group G is abelian, the normalised traces on $C^*(G) \cong C(\hat{G})$ are given by probability measures on the compact dual group \hat{G} . Thus in [17] (see also §8.1 below), the KMS states with inverse temperature $\beta > \beta_c$ on $(\mathcal{T}(\mathbb{Z}^d, \Sigma), \sigma)$ are parametrised by the probability measures on \mathbb{T}^d .

When G has an abelian quotient Q , $C^*(Q)$ is a quotient of $C^*(G)$, and the probability measures on \hat{Q} give traces on $C^*(G)$ and KMS states on $(\mathcal{T}(G, X), \sigma)$. This applies in particular to the self-similar action (B, X) associated to the basilica group in §2.3, since Proposition 2.6 implies that B has a quotient isomorphic to \mathbb{Z}^2 . Thus for each $\beta > \log |X|$, Theorem 6.1 gives a simplex S_Q of KMS_β states of $(\mathcal{T}(B, X), \sigma)$ parametrised by the probability measures on $\hat{Q} = \mathbb{T}^2$. The simplex S_Q includes the state ψ_{β, τ_1} of Corollary 6.6, which corresponds to the point mass at $1 \in \mathbb{T}^2$. However, since the trace τ_e does not factor through the quotient map, Theorem 6.1 implies that S_Q does not include the state ψ_{β, τ_e} of Corollary 6.5.

7. KMS STATES AT THE CRITICAL INVERSE TEMPERATURE

In this section we describe the KMS states on $\mathcal{T}(G, X)$ at the critical inverse temperature $\beta_c = \log |X|$. We start by showing that we are effectively dealing with the KMS states on the Cuntz-Pimsner algebra $\mathcal{O}(G, X)$.

Proposition 7.1. *Let (G, X) be a self-similar action. Every $\text{KMS}_{\log |X|}$ state of $(\mathcal{T}(G, X), \sigma)$ factors through a $\text{KMS}_{\log |X|}$ state on $\mathcal{O}(G, X)$.*

Proof. Suppose that ϕ is a $\text{KMS}_{\log |X|}$ state of $(\mathcal{T}(G, X), \sigma)$. Then Proposition 4.1 implies that $\phi(s_x s_x^*) = |X|^{-1}$, and hence

$$\phi\left(1 - \sum_{x \in X} s_x s_x^*\right) = 1 - |X||X|^{-1} = 0.$$

Now the argument in [16, Lemma 10.3] implies that ϕ vanishes on the ideal I generated by $1 - \sum_{x \in X} s_x s_x^*$. (Or one could apply the more general result in [12, Lemma 2.2] to the family $\mathcal{F} = \{s_v u_g s_w^*\}$ of analytic elements.) Corollary 3.5 says that I is the kernel of the quotient map of $\mathcal{T}(G, X)$ onto $\mathcal{O}(G, X)$, and hence ϕ factors through this quotient map. \square

To state our main results about states of $\mathcal{O}(G, X)$, we need some information about the sets F_g^k in Corollary 6.5.

Proposition 7.2. *Suppose that (G, X) is a self-similar action. For $g \in G \setminus \{e\}$ and $k \geq 0$, we consider again*

$$F_g^k = \{v \in X^k : g \cdot v = v \text{ and } g|_v = e\}.$$

The sequence $\{|X|^{-k} |F_g^k|\}$ is increasing and converges with limit $c_g \in [0, 1)$.

Proof. If $v \in F_g^k$ and $x \in X$, then

$$g \cdot (vx) = v(g|_v \cdot x) = vx \quad \text{and} \quad g|_{vx} = (g|_v)|_x = e|_x = e,$$

so $vx \in F_g^{k+1}$. Thus $|F_g^{k+1}| \geq |X| |F_g^k|$, and multiplying by $|X|^{-k-1}$ shows that $\{|X|^{-k} |F_g^k|\}$ is increasing.

Since the action of G on X^* is faithful, g acts non-trivially on some X^j , say $g \cdot v \neq v$. Then v is not in F_g^j , and no word of the form vw is in any F_g^l . So for $k > j$,

$$|F_g^k| \leq |X|^k - |X|^{k-j} = |X|^k (1 - |X|^{-j}).$$

Thus $|X|^{-k}|F_g^k| \leq 1 - |X|^{-j} < 1$ for $k > j$, and the sequence converges to a limit c_g satisfying $c_g < 1$. \square

We can now state our main theorem about $\mathcal{O}(G, X)$. Notice that part (3) applies in particular when (G, X) is contracting.

Theorem 7.3. *Suppose that (G, X) is a self-similar action.*

- (1) *Every KMS state of $(\mathcal{O}(G, X), \sigma)$ has inverse temperature $\log |X|$.*
- (2) *Take c_g as in Proposition 7.2. Then there is a $\text{KMS}_{\log |X|}$ state on $\mathcal{O}(G, X)$ such that*

$$\psi(s_v u_g s_w^*) = \begin{cases} |X|^{-|v|} c_g & \text{if } v = w \\ 0 & \text{otherwise.} \end{cases} \quad (7.1)$$

- (3) *Suppose that for every $g \in G \setminus \{e\}$, the set $\{g|_v : v \in X^*\}$ is finite. Then the state in part (2) is the only KMS state of $(\mathcal{O}(G, X), \sigma)$.*

Proof of Theorem 7.3 (1). Suppose that ϕ is a KMS state of $(\mathcal{O}(G, X), \sigma)$ with inverse temperature β . Then the Cuntz relation $\sum_{x \in X} s_x s_x^* = 1$ and the KMS condition give

$$\begin{aligned} 1 = \phi(1) &= \phi\left(\sum_{x \in X} s_x s_x^*\right) = \sum_{x \in X} \phi(s_x s_x^*) = \sum_{x \in X} \phi(s_x^* \sigma_{i\beta}(s_x)) \\ &= \sum_{x \in X} e^{-\beta} \phi(s_x^* s_x) = \sum_{x \in X} e^{-\beta} = |X| e^{-\beta}, \end{aligned}$$

and hence $\beta = \log |X|$. \square

We will prove existence of the $\text{KMS}_{\log |X|}$ state ψ by taking a limit of KMS_β states as $\beta \rightarrow \beta_c = \log |X|$. To evaluate the limit, we need the following analytic lemma.

Lemma 7.4. *Suppose that $\{c_k\}$ is an increasing sequence of real numbers with $c_k \rightarrow c$. Then*

$$\sum_{k=0}^{\infty} (1-r)c_k r^k \rightarrow c \quad \text{as } r \rightarrow 1-.$$

Proof. Fix $\epsilon > 0$, and choose K such that $k \geq K \implies 0 \leq c - c_k < \frac{\epsilon}{2}$. Choose $\delta > 0$ such that

$$0 < 1-r < \delta \implies \sum_{k=0}^K (1-r)c_k r^k < \frac{\epsilon}{2}.$$

Then for r satisfying $0 < 1-r < \delta$, we have $\sum_{k=0}^{\infty} (1-r)r^k = 1$, so

$$\begin{aligned} \left| c - \sum_{k=0}^{\infty} (1-r)c_k r^k \right| &= \left| \sum_{k=0}^{\infty} (1-r)(c - c_k) r^k \right| \\ &\leq \sum_{k=0}^K (1-r)(c - c_j) r^k + (1-r)(c - c_K) \left(\sum_{k=K+1}^{\infty} r^k \right) \\ &= \sum_{k=0}^K (1-r)(c - c_k) r^k + (1-r)(c - c_K) r^K (1-r)^{-1}, \end{aligned}$$

which is less than ϵ by choice of K and δ (and because $r^K < 1$). \square

Proof of Theorem 7.3 (2). We choose a decreasing sequence $\{\beta_n\}$ such that $\beta_n \rightarrow \log |X|$, and consider the KMS_{β_n} states $\psi_{\beta_n} := \psi_{\beta_n, \tau_e}$ of Corollary 6.5. By weak* compactness of the state space, we can by passing to a subsequence assume that $\{\psi_{\beta_n}\}$ converges

weak* to a state ψ . Proposition 5.3.23 of [3] implies that ψ is a $\text{KMS}_{\log|X|}$ state. (Or we could wait till we have the formula (7.1), and apply Proposition 4.1.)

We now compute the limit of $\psi_{\beta_n}(s_v u_g s_v^*)$. We know from (5.1) that $\psi_{\beta_n}(s_v u_g s_v^*) = 0$ unless $v = w$, and satisfies

$$\begin{aligned} \psi_{\beta_n}(s_v u_g s_v^*) &= e^{-\beta_n|v|} (1 - |X|e^{-\beta_n}) \sum_{k=0}^{\infty} e^{-\beta_n k} |F_g^k| \\ &= e^{-\beta_n|v|} \left(\sum_{k=0}^{\infty} (1 - |X|e^{-\beta_n}) |X|^{-k} |F_g^k| (|X|e^{-\beta_n})^k \right). \end{aligned}$$

Now we are in the situation of Lemma 7.4 with $r = |X|e^{-\beta_n}$ and $c_k = |X|^{-k} |F_g^k| \rightarrow c_g$. Since $r_n := |X|e^{-\beta_n}$ converges to 1 from below as $n \rightarrow \infty$, Lemma 7.4 implies that

$$\sum_{k=0}^{\infty} (1 - r_n) c_k r_n^k \rightarrow c_g \quad \text{as } n \rightarrow \infty.$$

Thus

$$\psi(s_v u_g s_v^*) = \lim_{n \rightarrow \infty} \psi_{\beta_n}(s_v u_g s_v^*) = \lim_{n \rightarrow \infty} e^{-\beta_n|v|} \left(\sum_{k=0}^{\infty} (1 - r_n) c_k r_n^k \right) = |X|^{-|v|} c_g,$$

as required. \square

Proof of Theorem 7.3 (3). Suppose that ϕ is a KMS state on $\mathcal{O}(G, X)$. We need to show that ϕ is the state ψ in (2). Part (1) implies that ϕ has inverse temperature $\log|X|$. Now Proposition 4.1 implies that it suffices for us to prove that $\phi(u_g) = \psi(u_g)$ whenever $g \neq e$.

Suppose that $g \in G \setminus \{e\}$. Since $\{g|_v : v \in X^*\}$ is finite and the action of G on X^* is faithful, there exists j such that for each $v \in X^*$ with $g|_v \neq e$, there exists $u \in X^j$ with $g|_v \cdot u \neq u$. We will show that

$$|X|^{-nj} |G_g^{nj} \setminus F_g^{nj}| = |X|^{-nj} |\{w \in X^{nj} : g \cdot w = w\} \setminus F_g^{nj}| \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (7.2)$$

and use this to show that $\phi(u_g) = c_g = \psi(u_g)$.

We prove by induction that

$$|G_g^{nj} \setminus F_g^{nj}| \leq (|X|^j - 1)^n \quad (7.3)$$

for all $n \geq 1$. Our choice of j ensures that, for every $v \in G_g^j$, the set $\{w \in X^j : g|_v \cdot w = w\}$ is not all of X^j ; thus we have (7.3) for $n = 1$. Assume that (7.3) holds for n . Then

$$|G_g^{(n+1)j} \setminus F_g^{(n+1)j}| = |\{vv' : v \in X^{nj}, v' \in X^j, g \cdot vv' = vv'\} \setminus F_g^{(n+1)j}|,$$

and we have

$$vv' \in G_g^{(n+1)j} \setminus F_g^{(n+1)j} \implies v \in G_g^{nj} \setminus F_g^{nj} \text{ and } g|_v \cdot v' = v'.$$

On the other hand, for each $v \in G_g^{nj} \setminus F_g^{nj}$, we have $g|_v \neq e$, and thus there exists $v' \in X^j$ such that $g|_v \cdot v' \neq v'$. Thus for each $v \in G_g^{nj} \setminus F_g^{nj}$,

$$|\{v' : vv' \in G_g^{(n+1)j} \setminus F_g^{(n+1)j}\}| \leq |X|^j - 1,$$

and the inductive hypothesis gives

$$|G_g^{(n+1)j} \setminus F_g^{(n+1)j}| \leq |G_g^{nj} \setminus F_g^{nj}| (|X|^j - 1) \leq (|X|^j - 1)^{n+1}.$$

Thus (7.3) holds for all $n \geq 1$. Now we have

$$0 \leq |X|^{-nj} |G_g^{nj} \setminus F_g^{nj}| \leq |X|^{-nj} (|X|^j - 1)^n = \left(1 - \frac{1}{|X|^j}\right)^n \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

which gives (7.2).

To complete the proof we show that $\phi(u_g) = c_g$. For every $n \in \mathbb{N}$, we use the Cuntz relation $1 = \sum_{w \in X^{nj}} s_w s_w^*$ and Proposition 4.1 to compute

$$\begin{aligned}
\phi(u_g) &= \phi\left(u_g \sum_{w \in X^{nj}} s_w s_w^*\right) \\
&= \sum_{w \in X^{nj}} \phi(s_{g \cdot w} u_{g|_w} s_w^*) \\
&= \sum_{\{w \in X^{nj} : g \cdot w = w\}} |X|^{-nj} \phi(u_{g|_w}) \\
&= \sum_{w \in G_g^{nj} \setminus F_g^{nj}} |X|^{-nj} \phi(u_{g|_w}) + \sum_{w \in F_g^{nj}} |X|^{-nj} \phi(u_e) \\
&= \sum_{w \in G_g^{nj} \setminus F_g^{nj}} |X|^{-nj} \phi(u_{g|_w}) + |X|^{-nj} |F_g^{nj}|. \tag{7.4}
\end{aligned}$$

Let $\varepsilon > 0$. By Proposition 7.2, there exists $N \in \mathbb{N}$ such that

$$n \geq N \implies |c_g - |X|^{-nj} |F_g^{nj}| < \varepsilon/2 \quad \text{and} \quad \left(1 - \frac{1}{|X|^j}\right)^n < \varepsilon/2.$$

Then for $n \geq N$, (7.4) gives

$$\begin{aligned}
|\phi(u_g) - c_g| &< \sum_{w \in G_g^{nj} \setminus F_g^{nj}} |X|^{-nj} |\phi(u_{g|_w})| + \varepsilon/2 \\
&\leq |G_g^{nj} \setminus F_g^{nj}| |X|^{-nj} + \varepsilon/2 \\
&\leq \left(1 - \frac{1}{|X|^j}\right)^n + \varepsilon/2 < \varepsilon,
\end{aligned}$$

which implies that $\phi(u_g) = c_g$. □

Somewhat surprisingly, our construction of KMS states at the critical inverse temperature gives a third trace on $C^*(G)$.

Corollary 7.5. *Suppose that (G, X) is a self-similar action, and take $\{c_g\}$ as in Proposition 7.2. Then there is a trace τ on $C^*(G)$ such that $\tau(\delta_g) = c_g$ for $g \neq e$.*

Proof. Proposition 4.1 (2) implies that $\phi \circ \pi_u$ is a trace on $C^*(G)$ for every KMS state ϕ of $\mathcal{T}(G, X)$ or $\mathcal{O}(G, X)$, and taking ϕ to be the $\text{KMS}_{\log|X|}$ state of $\mathcal{O}(G, X)$ in Theorem 7.3 (2) gives the required trace $\tau := \phi \circ \pi_u$. □

It will follow from Propositions 8.2 and 8.4 below that, for the self-similar actions of the basilica and Grigorchuk groups, the trace of Corollary 7.5 is distinct from the traces τ_e and τ_1 considered in §6.

Remark 7.6. In [25, §3.4], Planchat constructs a trace Tr on a quotient $C_\rho^*(G)$ of $C^*(G)$, and the value $\text{Tr}(\rho_g)$ at a unitary generator is (in our notation) the limit $\lim_{k \rightarrow \infty} |X|^{-k} |G_g^k|$ of the decreasing sequence $\{|X|^{-k} |G_g^k|\}$. When (G, X) has the finite-state property of Theorem 7.3 (3), the calculation (7.2) implies that $|X|^{-k} |G_g^k| \rightarrow c_g$ also, and hence our trace coincides with the lift of Planchat's trace to $C^*(G)$. For the groups generated by automata studied in [25], the pair (G, X) always has this finite-state property. (To see this, note that G is generated by a finite set S which is closed under restriction. This generating family induces a length function l on G , and then the properties of

restriction imply that $l(g|_v) \leq l(g)$ for all $g \in G$ and $v \in X^*$. Since there are finitely many words of a fixed length, it follows that each $\{g|_v : v \in X^*\}$ is finite.)

Our calculations in the next section suggest that it may be easier to compute the values of this trace using the formula $c_g = \lim_{k \rightarrow \infty} |X|^{-k} |\mathbb{F}_g^k|$.

We finish by showing that for a contracting self-similar action, the values of c_g on the nucleus determine the function c , and hence the KMS state at critical inverse temperature. For convenience, we define $c_e := 1$.

Corollary 7.7. *Suppose that (G, X) is a contracting self-similar action with nucleus \mathcal{N} . For $g \in G$, choose $k \in \mathbb{N}$ such that $g|_w \in \mathcal{N}$ for every $w \in X^k$. Then*

$$c_g = \sum_{\{w \in X^k : g \cdot w = w\}} |X|^{-k} c_{g|_w}.$$

Proof. We let ϕ be the unique $\text{KMS}_{\log|X|}$ state of $(\mathcal{O}(G, X), \sigma)$, so that in particular $\phi(u_g) = c_g$ for all g (see Theorem 7.3 (2)). Now the result follows from the calculation in the first three lines of (7.4). \square

8. EXAMPLES

8.1. Dilation matrices. Suppose that $A \in M_d(\mathbb{Z})$ has $|\det A| > 1$, and consider the associated self-similar action (\mathbb{Z}^d, Σ) of §2.2. We first check that the states constructed in Theorem 5.1 are the same as the ones in [17, Proposition 6.1].

The Fourier transform gives an isomorphism of $C^*(G) = C^*(\mathbb{Z}^d)$ onto $C(\mathbb{T}^d)$; we choose the one which takes δ_n to the function $z \mapsto z^n$. Traces on $C^*(\mathbb{Z}^d)$ are given by probability measures on \mathbb{T}^d ; given such a measure μ , we consider the trace τ_μ such that $\tau_\mu(\delta_n) = \int_{\mathbb{T}^d} z^n d\mu(z)$. We want to compute the values of the state ψ_{β, τ_μ} of Theorem 5.1 on an element $s_w u_n s_w^*$ (it vanishes on the other spanning elements). For $j \geq 0$ and $u \in \Sigma^j$, we have

$$n \cdot u = u \iff n + b_j(u) = b_j(u) \iff n \in B^j \mathbb{Z}^d,$$

so $\{u \in \Sigma^j : n \cdot u = u\}$ is either Σ^j (when $n \in B^j \mathbb{Z}^d$) or empty. If $n \cdot u = u$, then $n|_u = B^{-j}n$ by (2.5), so the right-hand side of (5.1) is

$$(1 - |\det A| e^{-\beta}) \sum_{\{j \geq 0 : n \in B^j \mathbb{Z}^d\}} |\det A|^j e^{-\beta(|w|+j)} \tau_\mu(\delta_{B^{-j}n}). \quad (8.1)$$

Thus we have $n \in B^j \mathbb{Z}^d \iff B^{|w|}n \in B^{|w|+j} \mathbb{Z}^d$, and writing $j' = |w| + j$ in (8.1) gives

$$\psi_{\beta, \tau_\mu}(s_w u_n s_w^*) = (1 - |\det A| e^{-\beta}) \sum_{\{j' \geq |w| : B^{|w|}n \in B^{j'} \mathbb{Z}^d\}} |\det A|^{j'-|w|} e^{-\beta j'} \int_{\mathbb{T}^d} z^{B^{(|w|-j')}n} d\mu(z). \quad (8.2)$$

The isomorphism $\theta : \mathcal{T}(\mathbb{Z}^d, \Sigma) \rightarrow \mathcal{T}(M_L)$ of Proposition 3.9 carries an element $s_w u_n s_w^*$ into $u_{b(w)+B^{|w|}n} v^{|w|} v^{*|w|} u_{b(w)}^*$, and we can check that the right-hand side of (8.2) is the same as the value of the state $\psi_{\beta, \mu}$ of [17, Proposition 6.1] on the spanning element $u_{b(w)+B^{|w|}n} v^{|w|} v^{*|w|} u_{b(w)}^*$.

Proposition 8.1 ([17, Theorem 5.3]). *Suppose that $A \in M_d(\mathbb{Z})$ has $N := |\det A| \neq 0$. Then there is a $\text{KMS}_{\log N}$ state ϕ of $(\mathcal{O}(M_L), \sigma) = C^*(u, v)$ such that*

$$\phi(u_m v^k v^{*l} u_n^*) = \begin{cases} 0 & \text{unless } k = l \text{ and } m = n \\ N^{-k} & \text{if } k = l \text{ and } m = n. \end{cases} \quad (8.3)$$

If A is a dilation matrix, then this is the only KMS state of $(\mathcal{O}(M_L), \sigma)$.

Proof. To apply Theorem 7.3 (2) to the associated self-similar group (\mathbb{Z}^d, Σ) , we need to compute the numbers $|F_n^j|$. For $u \in \Sigma^j$, we have $n \cdot u = u \iff n \in B^j \mathbb{Z}^d$, and then $n|_u = B^{-j}n$, so $n|_u = 0 \iff n = 0$. Thus $F_n^j = \emptyset$ for all $n \neq 0$, and the state ψ of Theorem 7.3 (2) satisfies

$$\psi(s_v u_n s_w^*) = \begin{cases} 0 & \text{unless } v = w \text{ and } n = 0 \\ |\Sigma|^{-|w|} = N^{-|w|} & \text{if } v = w \text{ and } n = 0. \end{cases} \quad (8.4)$$

We take $\phi := \psi \circ \theta^{-1}$. Then the elements of the form $\theta(s_w s_w^*)$ are the $u_m v^k v^{*l} u_n^*$ with $k = l = |w|$ and $m = n = b(w)$, and (8.4) reduces to the formula (8.3) for ϕ .

Now suppose that A is a dilation matrix. Then Proposition 2.4 implies that (\mathbb{Z}^d, Σ) is a contracting self-similar action, and the uniqueness follows from Theorem 7.3 (3). \square

8.2. Computing using the Moore diagram. To calculate values of the KMS states explicitly, we need to compute the sizes of the sets F_g^k and G_g^k defined in (6.5) and (6.6). We begin with G_g^k .

For each $v \in G_g^k$ we get the following path μ_v in the Moore diagram:

$$\mu_v := g \xrightarrow{(v_1, v_1)} g|_{v_1} \xrightarrow{(v_2, v_2)} g|_{v_1 v_2} \xrightarrow{(v_3, v_3)} \cdots \xrightarrow{(v_k, v_k)} g|_v$$

Notice that all the labels have the form (x, x) . Every path with labels (x, x) arises this way: given

$$\mu := g \xrightarrow{(x_1, x_1)} h_1 \xrightarrow{(x_2, x_2)} h_2 \xrightarrow{(x_3, x_3)} \cdots \xrightarrow{(x_k, x_k)} h_k,$$

we have $h_i = (\cdots ((g|_{x_1})|_{x_2}) \cdots)|_{x_i} = g|_{x_1 \cdots x_{i-1}}$, and $v = x_1 x_2 \cdots x_k$ belongs to G_g^k with $\mu_v = \mu$. We call paths μ of this form *stationary*, because they give elements v of X^* such that $s(\mu) \cdot v = v$, where $s(\mu) \in G$ is the source of the path μ . Thus G_g^k is in one-to-one correspondence with the set of stationary paths in the Moore diagram starting at g .

For $v \in F_g^k$, we have $g \cdot v = v$ and $g|_v = e$, so the last vertex on μ_v is e . Thus the elements of F_g^k are in one-to-one correspondence with the stationary paths starting at g and ending at e .

Thus we can compute $|G_g^k|$ and $|F_g^k|$ by counting stationary paths in the Moore diagram. Notice that for a given g , we only need to draw the part of the Moore diagram which consists of the stationary edges reachable by stationary paths from g . For examples of such computations, see Examples 8.3 and 8.5 below.

8.3. The basilica group. We now consider the self-similar action (B, X) which defines the basilica group (see §2.3). In §6.7, we discussed KMS states on the Toeplitz system $(\mathcal{T}(B, X), \sigma)$ at inverse temperatures greater than the critical value $\beta_c = \log |X| = \log 2$. At the critical inverse temperature, Proposition 7.1 implies that every $\text{KMS}_{\log 2}$ state factors through $(\mathcal{O}(B, X), \sigma)$, and we have:

Proposition 8.2. *The system $(\mathcal{O}(B, X), \sigma)$ has a unique $\text{KMS}_{\log 2}$ state, which is given on the nucleus $\mathcal{N} = \{e, a, b, a^{-1}, b^{-1}, ab^{-1}, ba^{-1}\}$ by*

$$\phi(u_g) = \begin{cases} 1 & \text{for } g = e \\ \frac{1}{2} & \text{for } g = b, b^{-1} \\ 0 & \text{for } g = a, a^{-1}, ab^{-1}, ba^{-1}. \end{cases}$$


Proof. We know from Proposition 2.5 that (B, X) is contracting with nucleus \mathcal{N} , so existence and uniqueness of ϕ follow from Theorem 7.3. In Figure 2, there are no stationary paths from $g \in \{a, a^{-1}, ab^{-1}, ba^{-1}\}$ to e , so for such g we have $F_g^k = \emptyset$ for all k and $\phi(u_g) = c_g = 0$. For $g \in \{b, b^{-1}\}$, the only stationary paths go straight from g to e , and there are 2^{k-1} of them; thus $|X|^{-k}|F_g^k| = 2^{-k}2^{k-1} = \frac{1}{2}$, and $\phi(u_g) = c_g = \frac{1}{2}$. \square

Corollary 7.7) implies that these computations of the $\text{KMS}_{\log 2}$ state on the nucleus suffice to determine the state. If we want to know other values of the state, we can use the strategy outlined in §8.2. We illustrate this strategy by calculating $\phi(s_v u_{aba} s_w^*)$.

Example 8.3. By Theorem 7.3 (2), $\phi(s_v u_{aba} s_w^*)$ is either 0 or $2^{-|v|}\phi(u_{aba})$. To compute $\phi(u_{aba})$, we draw the portion of the Moore diagram emanating from aba with a view to finding F_{aba}^k . From the defining relations, we calculate

$$\begin{aligned} aba \cdot x &= x & (aba)|_x &= (ab)|_{a \cdot x} a|_x = (ab)|_y b = a|_y e b = b \\ aba \cdot y &= y & (aba)|_y &= (ab)|_{a \cdot y} a|_x = (ab)|_x e = a|_{b \cdot x} a = a|_x a = ba. \end{aligned}$$

We then note that $ba \cdot x = b \cdot y = y$, which forces $ba \cdot y = x$, and hence there are no stationary paths going from aba to e through ba . Now we can delete any edges in the Moore diagram for the nucleus with unequal labels, and find that all the stationary paths from aba to e lie in the diagram



We deduce that $|F_{aba}^k| = 2^{k-2}$ for $k \geq 2$, and hence

$$\phi(u_{aba}) = c_{aba} = \lim_{k \rightarrow \infty} 2^{-k}|F_{aba}^k| = \frac{1}{4}.$$

Thus Theorem 7.3 (2) gives


$$\phi(s_v u_{aba} s_w^*) = \begin{cases} 2^{-|v|-2} & \text{if } v = w \\ 0 & \text{otherwise.} \end{cases}$$

8.4. The Grigorchuk group.

Proposition 8.4. *Let (G, X) be the self-similar action of the Grigorchuk group from §2.4. Then $(\mathcal{O}(G, X), \sigma)$ has a unique $\text{KMS}_{\log 2}$ state ϕ which is given on the nucleus $\mathcal{N} = \{e, a, b, c, d\}$ by*

$$\phi(u_g) = \begin{cases} 1 & \text{for } g = e \\ 0 & \text{for } g = a \\ 1/7 & \text{for } g = b \\ 2/7 & \text{for } g = c \\ 4/7 & \text{for } g = d. \end{cases}$$

Proof. We know from Proposition 2.7 that the Grigorchuk action is contracting with nucleus \mathcal{N} , and $|X| = 2$, so Theorem 7.3 (3) implies that there is a unique $\text{KMS}_{\log 2}$ state ϕ . A look at the Moore diagram shows that there are no stationary paths starting at a , and hence there are no stationary paths going to e through a . Thus it suffices to count paths to e in the following diagram.



In particular, $F_a^k = \emptyset$ for all k , and $\phi(u_a) = c_a = 0$. From d , there are 2^{k-1} paths of length k which go straight to e , 2^{k-4} which first go round the cycle once, and

$$|F_d^k| = 2^{k-1} + 2^{k-4} + \dots + 2^{k-(3j+1)} \quad \text{where } 3j + 1 \leq k \leq 3j + 3.$$

Summing the geometric series gives

$$|F_d^k| = 2^{k-(3j+1)} \left(\frac{(2^3)^{(j+1)} - 1}{2^3 - 1} \right) = \frac{2^{k+2} - 2^{k-(3j+1)}}{7} \quad \text{where } 3j + 1 \leq k \leq 3j + 3.$$

Thus

$$|X|^{-k} |F_d^k| = 2^{-k} |F_d^k| = \frac{4 - 2^{-(3j+1)}}{7} \quad \text{where } 3j + 1 \leq k \leq 3j + 3,$$

and $\phi(u_d) = c_d = \lim_{k \rightarrow \infty} |X|^{-k} |F_d^k| = \frac{4}{7}$. There are similar formulas for c and b :

$$|F_c^k| = |F_d^{k-1}| = \frac{2^{k+1} - 2^{k-(3j+2)}}{7} \quad \text{where } 3j + 2 \leq k \leq 3j + 4, \text{ and} \quad (8.5)$$


$$|F_b^k| = |F_d^{k-2}| = \frac{2^k - 2^{k-(3j+3)}}{7} \quad \text{where } 3j + 3 \leq k \leq 3j + 5,$$

and these formulas imply that $\phi(u_c) = c_c = \frac{2}{7}$ and $\phi(u_b) = c_b = \frac{1}{7}$. \square

Example 8.5. We calculate the value of the state ϕ in Proposition 8.4 on the generator u_{cadac} . We need the part of the Moore diagram emanating from $cadac$ with stationary edges. We calculate, using either the defining relations (2.8) or the same information encoded in the Moore diagram of Figure 3:

$$\begin{aligned} cadac \cdot x &= x & (cadac)|_x &= (cada)|_x a = (cad)|_y a = (ca)|_y ba = c|_x ba = aba \\ cadac \cdot y &= y & (cadac)|_y &= (cada)|_y d = (cad)|_x d = (ca)|_x d = c|_y d = d^2 = e \\ aba \cdot x &= x & (aba)|_x &= (ab)|_y = a|_y c = c \\ aba \cdot y &= y & (aba)|_y &= (ab)|_x = a|_x a = a. \end{aligned}$$

This gets us into the nucleus. Now adding the stationary edges from the Moore diagram of the nucleus gives a diagram which contains all the stationary paths from $cadac$ to e :



We need to count the paths from cadac to e in this diagram. They go either straight to e , or straight to c . Using the formula in (8.5) for F_c^1 , we have

$$|F_{\text{cadac}}^k| = 2^{k-1} + |F_c^{k-2}| = 2^{k-1} + \frac{2^{k-1} - 2^{k-(3j+4)}}{7} \quad \text{where } 3j + 4 \leq k \leq 3j + 6,$$

and hence

$$|X|^{-k} |F_{\text{cadac}}^k| = 2^{-1} + \frac{2^{-1} - 2^{-(3j+4)}}{7} \quad \text{where } 3j + 4 \leq k \leq 3j + 6.$$

Thus $\phi(u_{\text{cadac}}) = c_{\text{cadac}} = \lim_{k \rightarrow \infty} |X|^{-k} |F_{\text{cadac}}^k| = \frac{4}{7}$.

REFERENCES

- [1] L. Bartholdi and B. Virág, *Amenability via random walks*, Duke Math. J. **130** (2005), 39–56.
- [2] J.-B. Bost and A. Connes, *Hecke algebras, type III factors and phase transitions with spontaneous symmetry breaking in number theory*, Selecta Math. (N.S.) **1** (1995), 411–457.
- [3] O. Bratteli and D.W. Robinson, *Operator Algebras and Quantum Statistical Mechanics II*, Second Edition, Springer-Verlag, Berlin, 1997.
- [4] N. Brownlowe and I. Raeburn, *Exel’s crossed product and relative Cuntz-Pimsner algebras*, Math. Proc. Cambridge Philos. Soc. **141** (2006), 497–508.
- [5] J. Cuntz, C. Deninger and M. Laca, *C*-algebras of Toeplitz type associated with algebraic number fields*, Math. Ann., available online.
- [6] R. Exel, *A new look at the crossed-product of a C*-algebra by an endomorphism*, Ergodic Theory Dynam. Systems **23** (2003), 1–18.
- [7] R. Exel, A. an Huef and I. Raeburn, *Purely infinite simple C*-algebras associated to integer dilation matrices*, Indiana Univ. Math. J. **60** (2011), 1033–1058.
- [8] R. Exel and M. Laca, *Partial dynamical systems and the KMS condition*, Comm. Math. Phys. **232** (2003), 223–277.
- [9] N.J. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. **48** (1999), 155–181.
- [10] R.I. Grigorchuk and A. Żuk, *On a torsion-free weakly branch group defined by a three state automaton*, Internat. J. Algebra Comput. **12** (2002), 223–246.
- [11] D. Harari and E. Leichtnam, *Extension du phénomène de brisure spontanée de symétrie de Bost-Connes au cas de corps globaux quelconques*, Selecta Math. (N.S.) **3** (1997), 205–243.
- [12] A. an Huef, M. Laca, I. Raeburn and A. Sims, *KMS states on the C*-algebras of finite graphs*, arXiv:1205.2194.
- [13] M. Laca, *Semigroups of *-endomorphisms, Dirichlet series, and phase transitions*, J. Funct. Anal. **152** (1998), 330–378.
- [14] M. Laca and S. Neshveyev, *KMS states of quasi-free dynamics on Pimsner algebras*, J. Funct. Anal. **211** (2004), 457–482.
- [15] M. Laca and S. Neshveyev, *Type III₁ equilibrium states of the Toeplitz algebra of the affine semigroup over the natural numbers*, J. Funct. Anal. **261** (2011), 169–187.
- [16] M. Laca and I. Raeburn, *Phase transition on the Toeplitz algebra of the affine semigroup over the natural numbers*, Adv. Math. **225** (2010), 643–688.
- [17] M. Laca, I. Raeburn, and J. Ramagge, *Phase transition on Exel crossed products associated to dilation matrices*, J. Funct. Anal. **261** (2011), 3633–3664.
- [18] N.S. Larsen and I. Raeburn, *Projective multi-resolution analyses arising from direct limits of Hilbert modules*, Math. Scand. **100** (2007), 317–360.
- [19] V. Nekrashevych, *Cuntz-Pimsner algebras of group actions*, J. Operator Theory **52** (2004), 223–249.
- [20] V. Nekrashevych, *Self-Similar Groups*, Math. Surveys and Monographs vol. 117, Amer. Math. Soc., Providence, 2005.
- [21] V. Nekrashevych, *C*-algebras and self-similar groups*, J. reine angew. Math. **630** (2009), 59–123.
- [22] J.A. Packer and M.A. Rieffel, *Wavelet filter functions, the matrix completion problem, and projective modules over C(Tⁿ)*, J. Fourier Anal. Appl. **9** (2003), 101–116.
- [23] G.K. Pedersen, *C*-Algebras and Their Automorphism Groups*, London Math. Soc. Monographs, vol. 14, Academic Press, London, 1979.
- [24] M.V. Pimsner, *A class of C*-algebras generalizing both Cuntz-Krieger algebras and crossed products by Z*, Fields Institute Commun. **12** (1997), 189–212.

- [25] J.-F. Planchat, *Fundamental C^* -algebras associated to automata groups*, arXiv:1204.1517.
- [26] I. Raeburn and D.P. Williams, *Morita Equivalence and Continuous-Trace C^* -Algebras*, Math. Surveys and Monographs, vol. 60, Amer. Math. Soc., Providence, 1998.

MARCELO LACA, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VICTORIA,
VICTORIA, BC V8W 3P4, CANADA
E-mail address: laca@math.uvic.ca

IAIN RAEBURN, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTAGO, PO BOX
56, DUNEDIN 9054, NEW ZEALAND
E-mail address: iraeburn@maths.otago.ac.nz

JACQUI RAMAGGE, SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, UNIVERSITY OF WOL-
LONGONG, NSW 2522, AUSTRALIA
E-mail address: ramagge@uow.edu.au

MICHAEL F. WHITTAKER, SCHOOL OF MATHEMATICS AND APPLIED STATISTICS, UNIVERSITY OF
WOLLONGONG, NSW 2522, AUSTRALIA
E-mail address: mfwhittaker@gmail.com