



2013

Summary of vertebrate and molluscan assemblages excavated from late-Pleistocene and Holocene deposits at Khao Toh Chong Rockshelter, Krabi, Thailand

Cyler Conrad
University of New Mexico

Hannah G. Van Vlack
San Jose State University

Ben Marwick
University of Washington, bmarwick@uow.edu.au

Cholawit Thongcharoenchaikit
National Science Museum, Thailand

Rasmi Shoocongdej
Silpakorn University

See next page for additional authors

Publication Details

Conrad, C., Van Vlack, H. G., Marwick, B., Thongcharoenchaikit, C., Shoocongdej, R. & Chaisuwan, B. (2013). Summary of vertebrate and molluscan assemblages excavated from late-Pleistocene and Holocene deposits at Khao Toh Chong Rockshelter, Krabi, Thailand. *The Thailand Natural History Museum Journal*, 7 (1), 11-22.

Summary of vertebrate and molluscan assemblages excavated from late-Pleistocene and Holocene deposits at Khao Toh Chong Rockshelter, Krabi, Thailand

Abstract

The zooarchaeological record of a recently excavated rockshelter site in peninsular Thailand is summarized. Detailed identification of mammalian, reptilian, piscine and molluscan taxa indicate a unique foraging pattern of prehistoric humans throughout the late-Pleistocene to Holocene.

Disciplines

Medicine and Health Sciences | Social and Behavioral Sciences

Publication Details

Conrad, C., Van Vlack, H. G., Marwick, B., Thongcharoenchaikit, C., Shoocongdej, R. & Chaisuwan, B. (2013). Summary of vertebrate and molluscan assemblages excavated from late-Pleistocene and Holocene deposits at Khao Toh Chong Rockshelter, Krabi, Thailand. *The Thailand Natural History Museum Journal*, 7 (1), 11-22.

Authors

Cyler Conrad, Hannah G. Van Vlack, Ben Marwick, Cholawit Thongcharoenchaikit, Rasmi Shoocongdej, and Boonyarit Chaisuwan

Summary of Vertebrate and Molluscan Assemblages Excavated from Late-Pleistocene and Holocene Deposits at Khao Toh Chong Rockshelter, Krabi, Thailand

Cyler Conrad^{*1}, Hannah G. Van Vlack², Ben Marwick³, Cholawit
Thongcharoenchaikit⁴, Rasmi Shoocongdej⁵ and Boonyarit Chaisuwan⁶

¹University of New Mexico, Department of Anthropology, MSC01-1040, Anthropology 1,
Albuquerque, New Mexico, USA

²San Jose State University, Department of Anthropology, Clark Hall Suite 469, One
Washington Square, San Jose, California, USA

³University of Washington, Department of Anthropology, Box 353100, Seattle, Washington,
USA

⁴National Science Museum, Technopolis, Khlong 5, Khlong Luang, Pathum Thani, Thailand

⁵Silpakorn University, Department of Archaeology, Naphrathard Road, Bangkok, Thailand

⁶15th Regional Office of Fine Arts Department, Srisoonthon Road, Phuket, Thailand

ABSTRACT: The zooarchaeological record of a recently excavated rockshelter site in peninsular Thailand is summarized. Detailed identification of mammalian, reptilian, piscine and molluscan taxa indicate a unique foraging pattern of prehistoric humans throughout the late-Pleistocene to Holocene.

KEY WORDS: vertebrate and molluscan assemblages, late-Pleistocene and Holocene deposits, Khao Toh Chong, Krabi, Thailand.

INTRODUCTION

Khao Toh Chong Rockshelter is a site which was excavated June through July 2011 outside of Krabi, Thailand (Fig. 1). Khao Toh Chong (KTC) is significant to Southeast Asian archaeology because of its similarities and proximity to Lang Rongrien, a site with a long but sparse archaeological sequence (Anderson, 1990; Mudar and Anderson, 2007). KTC has potential to address questions of human adaptation raised by previous work in the region due to its high-resolution

archaeological deposit. The archaeological and faunal records indicate that the excavated deposit spans from the late-Pleistocene until recent times (Fig. 3; see Marwick *et al.*, 2013 for more details). During excavations the deposit was observed to be relatively undisturbed with intact stratigraphy. Faunal remains indicate prehistoric human exploitation of freshwater dwelling shellfish and turtles or tortoises, as well as a variety of small and large sized mammals. Mollusk discard rates appear to peak in the early Holocene, coincident with the increased

*Corresponding author.
E-mail: cylerc@unm.edu

monsoonal precipitation. Vertebrate remains show the reliance on artiodactyla during the terminal Pleistocene, followed by the exploitation of reptiles. We attribute this shift to adaptation of human foraging and hunting behaviours, to local environmental conditions.

MATERIALS AND METHODS

Two trenches, A and B, were excavated in four one by one meter units (Fig. 2). In Trench A, the southernmost trench, excavations reached a depth of 1.1 meters below the surface (mbs) (Fig. 3). In trench B, the northernmost trench, excavations were obstructed by bedrock in the northwest and southwest units. Subsequently, excavation depths in trench B extended to approximately 2.0 meters in the northeast and southeast units. A 0.03 m screen was used to sieve the upper strata, while a 0.10 m screen was used to sieve lower strata sediment during the excavation because the high clay content made it difficult for the deposit to pass through the smaller mesh.

Zooarchaeological analyses were performed on all faunal material excavated from the southwest unit of Trench A. Mammalian, reptilian, and piscean taxa were identified using standard zooarchaeological reference manuals (Lekagul and McNeely, 1977; Auetrakulvit, 2004; France,


2009 and Ankel-Simons, 2000). Molluscan taxa were identified using the comparative collections of the National Science Museum of Thailand and additional mollusk references manuals (Swennen *et al.*, 2001; Jiwaluk *et al.*, 2007 and Sucharit and Panha, 2008). The identification processes were hampered by the lack of comparative skeletal material for mammalian, reptilian and piscean taxa. As such, identification of the former was generally broad based in order to limit identification error and provide a basis for future analysis. Avian taxa were absent from this unit of the KTC assemblage. Number of identified specimen (NISP) counts were used to record the abundance of faunal taxa per stratigraphic unit (Lyman, 2008).

A digital palm scale was used to record the mass of identified elements to the nearest 0.1 gram. Standard inter-landmark dimensions were taken in millimeters, when applicable. In addition, identification of cut marks and burning were noted for both mollusk and vertebrate remains. Photographs of complete mollusk specimens were taken using a Canon EOS60D digital camera (Figs. 4 and 5). All archaeofaunal specimens are deposited at the Department of Archaeology, Silpakorn University, Bangkok, Thailand.



Figure 1. Approximate location of Lang Rongrien Rockshelter (1) and Khao Toh Chong Rockshelter (2) outside of Krabi, southern Thailand.

Figure 4. Shells from unit ASW- A. *Nerita balteata* B. *Neritina arata* C. *Planorbis jacobsoni* D. *Cyclophorus suliformis*



C. Conrad 2011

Figure 2. KTC trench location and site morphology (Marwick *et al.*, 2013).




Figure 3. KTC stratigraphy, radiocarbon dates and sample origin in Trench A (Marwick *et al.*, 2013).



Figure 4. Shells from unit ASW- A. *Nerita balteata* B. *Neoradina prasongi* C. *Plectopylis degerbolae* D. *Cyclophorus malaynus*.

Table 2. Mollusk NISP counts per stratigraphic context.

NISP Taxa per Stratigraphic Context	2	3	4	5	6	7H	8	7L	NISPΣ
Gastropoda									
Neritidae gen. indet.	-	-	1	-	-	-	-	-	1
<i>Nerita balteata</i>	-	-	-	-	2	-	-	-	2
<i>Cyclophorus</i> sp.	-	-	-	-	-	7	-	-	7
<i>Cyclophorus</i> cf. <i>saturnus</i>	-	-	-	-	-	-	-	1	1
<i>Cyclophorus malayanus</i>	-	-	-	-	9	1	-	1	11
Cyclophoridae gen. indet.	-	2	3	2	1	28	5	30	71
<i>Rhiostoma jalorensis</i>	-	-	2	2	10	9	2	2	27
<i>Rhiostoma</i> sp.	-	-	-	-	-	6	3	2	11
<i>Filopaludina</i> sp.	-	-	2	-	-	-	-	-	2
Viviparidae gen. indet.	-	1	-	-	-	-	-	-	1
<i>Pila</i> sp.	-	-	-	-	-	2	4	-	6
Ampullariidae gen. indet.	-	-	-	-	-	3	-	3	6
<i>Neoradina prasongi</i>	-	8	134	52	82	1584	2215	771	4846
<i>Telescopium telescopium</i>	-	-	3	2	3	4	2	-	14
Muricidae gen. indet.	-	1	4	-	1	-	-	-	6
<i>Plectopylis degerbolae</i>	-	-	1	-	1	2	3	5	12
<i>Amphidromus atricallosus</i>	-	-	1	-	1	1	-	-	3
Bivalvia									
<i>Anadara</i> sp.	-	1	-	-	-	-	-	-	1
Arcidae gen. indet.	-	-	3	-	1	-	-	-	4
<i>Pseudodon</i> sp.	-	-	1	9	-	-	-	-	10
Amblemidae gen. indet.	-	-	-	-	56	72	143	372	643
Corbiculidae gen. indet.	-	-	-	-	-	1	-	-	1




Figure 6. Shifting *Neoradina prasongi*, Amblemidae, and Cyclophoridae abundance throughout the late-Pleistocene and Holocene at KTC.

DISCUSSION

KTC rockshelter has a remarkably undisturbed record of mammalian, reptilian, piscine and molluscan assemblages present in situ. The invertebrate record at the rockshelter provides a unique and detailed description of molluscan taxa and subsistence patterns during periods of occupation. *Neoradina prasongi* molluscs were abundant during the late-Pleistocene, but much less frequent during the mid and late Holocene.

As Brandt (1974) describes, *Neoradina prasongi* molluscs live in fresh-water stream habitats. At KTC, *N. prasongi* shells constitute the bulk of molluscan food waste in the archaeological assemblage. The period of peak discard rates for *N. prasongi* is approximately 13ky cal B.P., suggesting that the most intensive use of the rockshelter occurred during this time. Utilization of this mollusk also indicates that freshwater stream habitats occurred in proximity to the rockshelter during the late-Pleistocene and early Holocene. Mudar and Anderson (2007) have suggested that during the late-Pleistocene a drier and more open environment occurred in the Krabi region, this period was also characterized by increased monsoon seasonality. During the peak mollusk discard period of the late-Pleistocene, a drier and more open environment would have allowed hunter-gatherer groups to utilize large artiodactyls in the grassland-savannah habitat, alongside *N. prasongi* in seasonally abundant

fresh-water stream resources. Occurrence of abundant turtle or tortoise remains at KTC also suggests that fresh-water stream habitats were found in proximity to the site throughout the late-Pleistocene and early-Holocene.

At Lang Rongrien the abundant Testudines elements through time provided evidence that a previously unidentified record of Southeast Asian turtle and tortoise exploitation occurred at the rockshelter (Mudar and Anderson, 2007). This record also appears at KTC with the presence of abundant Testudines specimens throughout the late-Pleistocene and Holocene deposits. Although, at archaeological sites located near KTC and Lang Rongrien, this record was not identified. Moh Khiew Cave and Sakai Cave in southern peninsular Thailand both had very low abundances of Testudines remains during the late-Pleistocene and Holocene (Pookajorn, 1996). Generally, Testudines elements were only identified by presence or absence at these sites. Since Lang Rongrien and KTC both provide evidence of turtle and tortoise exploitation, this represents a unique fresh-water stream foraging pattern. It also suggests that fresh-water stream habitats were abundant during the late-Pleistocene and early Holocene, in the Krabi region. Furthermore, previously identified shifts in monsoon seasonality, and intensity, during this time suggests that precipitation may have increased during the early Holocene in Peninsular Thailand, matching the

faunal record at KTC (Marwick and Gagan, 2011).

The declining exploitation of freshwater *N. prasongi* molluscs into the Holocene may reflect the shift from freshwater to mangrove swamp habitats in this region or a shift in the foraging dynamics of prehistoric groups (Shoocongdej, 2000 and 2010). Rising sea-levels throughout the Holocene would have shifted mangrove environments closer to the rockshelter overtime, which may have influenced the abundance and distribution of locally available resources and freshwater stream environments (Anderson, 1990; Horten *et al.*, 2005; Tjia, 1996, and Sinsakul, 1992). As Barker *et al.* (2005) describes, archaeologists in Southeast Asia have recently tried to identify how effectively late-Pleistocene hunter-gatherers utilized tropical rainforest environments. Initial faunal data from KTC describes a pattern of hunter-gather groups utilizing a diverse range of locally available taxa in the tropical rainforest environment, suggesting that hunter-gatherers at KTC were able to effectively adapt to shifts in local environmental conditions. Additionally, radiocarbon dates suggest that once intensive harvesting of *N. prasongi* ends during the middle-Holocene, the emergence of rice agriculture and farming in mainland Southeast Asia appears (Barker *et al.*, 2005; Castillo, 2011 and White *et al.*, 2004). Thus, declines in mollusk utilization may reflect a pattern of rising sea-levels, shifting environmental conditions,

and occurrence of agriculture during the mid and late-Holocene in peninsular Thailand.

Our data from KTC not only suggests that a subsistence change occurred at the Pleistocene-Holocene transition, but that foragers utilizing the rockshelter displayed a unique pattern of faunal exploitation not previously seen at archaeological sites in Thailand. As Bulbeck (2003) describes, large abundances of shellfish in rockshelter sites tend to date to the middle-Holocene when a transition towards a broad-spectrum diet occurs, not during the terminal Pleistocene. The earlier peak in the molluscan assemblage at KTC suggests that a different pattern of shellfish exploitation occurred here, one that we link to local environmental conditions. This suggests that the diversity of past human subsistence behaviours in this region may have been underestimated and broader diets may have appeared earlier than previously observed. Khao Toh Chong is a valuable archaeological and palaeoecological resource aiding the understanding of hunter-gatherer subsistence behaviours in Southeast Asia during the late-Pleistocene and Holocene.

ACKNOWLEDGEMENTS

Archaeological research at Khao Toh Chong would not have been possible without the help and support of Rasmi Shoocongdej and the Fine Arts Department, Bangkok, Thailand. We are sincerely grateful for the laboratory

space provided to undertake this analysis in the Archaeological Digital Archives Laboratory at Silpakorn University in Bangkok, Thailand. Without mollusk comparative collections from the National Science Museum of Thailand our efforts in this research process would have been severely hampered, and we are grateful to Cholawit Thongcharoenchaikit for providing them. Also, thank you to the 2011 field school participants Seungki Kwak, Jessica Butler, Anna Hopkins, Rachel Vander Houwen, Fitriwati, Kate Lim, Thanh Son, Kim Sreang Em, Kyaw Minn Htin, Borisut Boriphon, Praewchompoo Chunchaurai, Chonchanok Samrit and Supalak Mheetong. Additional on-site identification assistance was provided by Dr. Prasit Auetrakulvi, Associate Professor of Archaeology at Silpakorn University, Bangkok, Thailand and Cholawit Thongcharoenchaikit, archaeologist and paleontologist for the National Science Museum of Thailand, Bangkok. Dr. Marwick was supported by a Luce/ACLS Post-doctoral fellowship (B9471 01 180401 7301) and the University of Washington Office of Global Affairs Herbert H. Gowen International Studies Endowment for the initial excavation of, and analyses for, this archaeological site.

REFERENCES

- Anderson, D.D. 1990. *Lang Rongrien rockshelter: A Pleistocene, early Holocene archaeological site from Krabi, southwestern Thailand*. Philadelphia: University Museum, University of Pennsylvania. 86 pp.
- Ankel-Simons, F. 2000. *Primate anatomy: An introduction*. San Diego: Academic Press. 752 pp.
- Auetrakulvit, P. 2004. *Faunes du pléistocène final à l'holocène de Thaïlande: approche archéozoologique*. (Doctoral dissertation). L'Université de Provence Aix-Marseille I, Marseille.
- Barker, G., T. Reynolds and D. Gilbertson. 2005. The Human Use of Caves in Peninsular and Island Southeast Asia: Research Themes. *Asian Perspectives*. 44 (1): 1-15.
- Bulbeck, F. 2003. Hunter-gatherer occupation of the Malay Peninsula from the Ice Age to the Iron Age. In: J. Mercader (ed.), *Under the canopy: The archaeology of tropical rain forests*. Piscataway: Rutgers University Press.
- Brandt, R.A.M. 1974. The non-marine aquatic Mollusca of Thailand. *Arch. Molluskenkunde*. 105: 1-423.
- Castillo, C. 2011. Rice in Thailand: The Archaeobotanical Contribution. *Rice*. 4(3-4): 114-120.
- France, D.L. 2009. *Human and Nonhuman Bone Identification: A Color Atlas*. Boca Raton: CRC Press. 734 pp.
- Francis, C. 2008. *A Guide to the Mammals of Southeast Asia*.

- Princeton: Princeton University Press. 392 pp.
- Horten, B.P., P.L. Gibbard, G.M. Milne, R.J. Morley and C. Purintavaragul. 2005. Holocene sea levels and palaeoenvironments, Malay-Thai Peninsula, southeast Asia. *The Holocene* 15 (8): 1199-1213.
- Jiwaluk, J., P. Promprasri and O. Nakchinda. 2007. *Fresh-water Bivalves of Thailand*. The Agricultural Cooperation Federation of Thailand Press. Bangkok. (in Thai).
- Lekagul, B. and J.A. McNeely. 1977. *Mammals of Thailand*. Bangkok: Association for the Conservation of Wildlife. 758 pp.
- Lyman, L. 2008. *Quantitative Paleozoology*. Cambridge, Cambridge University Press. 348 pp.
- Marwick, B. and M. Gagan. 2011. Late Pleistocene monsoon variability in northwest Thailand: an oxygen isotope sequence from the bivalve *Margaritanopsis laeosensis* excavated in Mae Hong Son province. *Quaternary Science Reviews*. 30: 3088-3098.
- Marwick, B., R. Shoocongdej, C. Thongcharoenchaikit, B. Chaisuwan, C. Khowkhiew and S. Kwak. 2013. Hierarchies of engagement and understanding: Community engagement during archaeological excavations at Khao Toh Chong rockshelter, Krabi, Thailand. In: S. O'Connor (ed.), *Transcending the Culture-Nature Divide in Cultural Heritage: Views from the Asia-Pacific Region*. Terra Australis, ANU E Press. 203 pp.
- Mudar, K. and D. Anderson. 2007. New Evidence for Southeast Asian Pleistocene Foraging Economies: Faunal Remains from the Early Levels of Lang Rongrien Rockshelter, Krabi, Thailand. *Asian Perspectives*. 46(2): 298-334.
- Pookajorn, S. 1996. Human activities and environmental changes during the late Pleistocene to middle Holocene in southern Thailand and Southeast Asia. In: L. Guy Straus *et al.* (eds.), *Humans at the End of the Ice Age*. New York, Plenum Press, pp. 201-213.
- Shoocongdej, R. 2000. Forager mobility organization in seasonal tropical environments of western Thailand. *World Archaeology*. 32(1): 14-40.
- Shoocongdej, R. 2010. Subsistence-Settlement Organization during the Late Pleistocene-Early Holocene: The Case of Lang Kamnang Cave, Western Thailand. In: B. Bellina *et al.* (eds.), *50 Years of Southeast Asian Archaeology: Essays in Honor of Ian Glover*. River Books, Bangkok, pp. 51-66.
- Sinsakul, S. 1992. Evidence of quaternary sea level changes in the coastal areas of Thailand: a review. *Journal of*

- Southeast Asian Earth Sciences*. 7(1): 23-37.
- Sri-aroon, P., C. Lohachit and M. Harada. 2005. Brackish-water mollusks of Surat Thani Province, southern Thailand. *The Southeast Asian Journal of Tropical Medicine and Public Health*. 36: 180-188.
- Sucharit, C. and S. Panha. 2008. *Land Snails of Khao Nun National Park*. Bangkok Press. Bangkok. (in Thai).
- Swennen, C., R.G. Moolenbeck, N. Ruttanadakul, H. Hobbelink, H. Dekker and S. Hajisamae. 2001. *The molluscs of the southern Gulf of Thailand*. Bangkok: Biodiversity Research and Training Program. 210 pp.
- Tjia, H. D. 1996. Sea-Level Changes in the Tectonically Stable Malay-Thai Peninsula. *Quaternary International: the Journal of the International Union for Quaternary Research*. 31: 95-101.
- White, J. C., D. Penny, L. Kealhofer and B. Maloney. 2004. Vegetation changes from the late Pleistocene through the Holocene from three areas of archaeological significance in Thailand. *Quaternary International*. 113: 111-132.