



PRIFYSGOL
BANGOR
UNIVERSITY

Microbial enzyme activity and stoichiometry signal the effects of agricultural intervention on nutrient cycling in peatlands

Qin, Lei; Freeman, Chris; Jia, Xueying; Zhang, Zhongsheng; Liu, Bo; Zhang, Shaoqing; Jiang, Ming

Ecological Indicators

DOI:

[10.1016/j.ecolind.2020.107242](https://doi.org/10.1016/j.ecolind.2020.107242)

Published: 01/03/2021

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Qin, L., Freeman, C., Jia, X., Zhang, Z., Liu, B., Zhang, S., & Jiang, M. (2021). Microbial enzyme activity and stoichiometry signal the effects of agricultural intervention on nutrient cycling in peatlands. *Ecological Indicators*, 122, [107242]. <https://doi.org/10.1016/j.ecolind.2020.107242>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Microbial enzyme activity and stoichiometry signal the effects of agricultural**
2 **intervention on nutrient cycling in peatlands**

3 Lei Qin¹, Chris Freeman², Xueying Jia³, Zhongsheng Zhang¹, Bo Liu¹, Shaoqing Zhang¹,
4 Ming Jiang^{1,4*}

5

6 ¹Key Laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and
7 Agroecology, Chinese Academy of Sciences, Changchun 130102, China.

8 ²School of Natural Sciences, Bangor University, Bangor LL57 2UW, UK.

9 ³Taiyuan Normal University, Taiyuan 030619, China

10 ⁴Joint Key Lab of Changbaishan Wetland and Ecology, Jilin Province, Changchun 130102,
11 China

12

13

14

15

16

17 *Corresponding author: Ming Jiang (Email: jiangm@iga.ac.cn; Address: Key Laboratory of
18 Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology,
19 Chinese Academy of Sciences, Changchun 130102, China)

20

21

22

23

24

25

26

27

28

29 **Abstract:**

30 Fertilization in agricultural peatlands accelerates nutrient cycling and creates a potential
31 risk to nearby natural peatlands. Here, using undisturbed peatlands as reference, we studied
32 soil carbon (C), nitrogen (N), phosphorus (P) and the key enzymes for nutrient cycling at 0-50
33 cm soil depth in agricultural, nearby disturbed peatlands in a temperate fen in Northeast China.
34 Agricultural intervention significantly increased total P in agricultural and disturbed peatlands,
35 and decreased soil organic carbon content and total N in surface soil of agricultural peatlands,
36 however total N significantly accumulated at 20-30 cm soil both in agricultural and disturbed
37 peatlands ($p < 0.05$). Both N-acetyl- β -glucosaminidase and phosphatase significantly declined
38 in agricultural peatlands, while only phosphatase decreased in disturbed peatlands ($p < 0.05$),
39 and linear regression models showed strong effects of changes of soil nutrient levels on
40 enzyme activities. The ratios of β -D-glucosidase to N-acetyl- β -glucosaminidase and
41 phosphatase markedly increased in agricultural peatlands and showed higher ratios in deeper
42 soil of disturbed peatlands, suggesting relatively higher microbial demand for carbon.
43 Nonmetric multidimensional scaling analysis showed that variations of enzyme activity and
44 stoichiometry can be used to reveal agricultural disturbance, and further redundancy analysis
45 identified that total P and SOC explained 38.3% and 8.3% of the variance. Overall, our
46 findings show that microbial enzymatic activity and stoichiometry can be effective and
47 sensitive indicators of agricultural intervention and nutrient changes in peatlands, which
48 implies that they can be used in monitoring of future fertilization management strategies
49 aimed at fostering more sustainable agriculture.

50 **Key words:** fertilization; peatlands; management strategies; nutrient input

51 **Introduction**

52 Peatlands represent a significant atmospheric carbon (C) sink, and hold about 30% of
53 global soil C, despite covering just 3% of the global terrestrial surface (Gorham, 1991). Plant
54 productivity in these ecosystems exceeds decay leading to accumulation over hundreds to
55 thousands of years in the form of increased peat depth and carbon storage (Yu et al., 2010).
56 Peatlands are defined as soils with a high carbon content ($>30\%$) and an organic horizon

57 larger than 30 cm (Rydin and Jeglum, 2006), and in many well developed peatlands, peat
58 depth can exceed two meters (Yu et al., 2010). In recent decades, the need for food and
59 energy, has led to up to 50.9 Mha peatlands being converted to agriculture, grasslands and
60 forestry for food and energy supply (Leifeld and Menichetti, 2018), triggering vast carbon
61 losses (Saurich et al., 2019).

62 There are about 25 Mha of agricultural peatlands worldwide (Tubiello et al., 2016),
63 representing ~50% of drained peatlands (50.9 Mha) (Leifeld and Menichetti, 2018). Oxygen
64 recovery following anoxic conditions directly promotes microbial metabolic process, turning
65 peatlands from a carbon sink into a hotspot of carbon mineralization (Eickenscheidt et al.,
66 2015; Leifeld and Menichetti, 2018). In addition to oxygen, fertilizer application also
67 contributes to the acceleration of microbial respiration (Eickenscheidt et al., 2015; Saurich et
68 al., 2019). Thus, these all contribute to carbon losses. Bader et al. (2017). reported that
69 organic matter decomposition in agricultural utilized peatlands occurs at a higher rate than
70 either grasslands or peatlands used for forestry.

71 The additional inorganic nutrients associated with fertilization can easily leach into
72 deeper peat soil (Kogel-Knabner et al., 2010), compounding these effects. Moreover, some
73 studies also show that fertilization increases the nutrient burden of surface and groundwater
74 (Koerselmann et al., 1990; Steinmuller et al., 2016; Berger et al., 2017), potentially leading to
75 eutrophication of nearby pristine peatlands (Wright and Reddy, 2001; Prenger and Reddy,
76 2004; Steinmuller et al., 2016). Clearly, fertilization is far more detrimental to peatlands than
77 is often assumed, yet few studies have evaluated agricultural fertilization disturbance in
78 peatlands.

79 Microbial enzyme activities are widely recognized as sensitive indicators of changes in
80 soil function under agricultural management (Lagomarsino et al., 2009; Zagal et al., 2009;
81 Pajares et al., 2009). Soil microorganisms produce extracellular enzymes to acquire nutrients
82 to satisfy the demand for energy for growth, thereby influencing carbon and nutrient cycling
83 (Sinsabaugh and Moorhead, 1994; Luo et al., 2017). Microorganisms also change nutrient


84 acquisition strategies during fertilization (Sinsabaugh and Moorhead, 1994). Although soil
85 physico-chemical properties can reflect the agricultural intervention, short-term changes are
86 usually not easy to detect (Lagomarsino et al., 2009) and fertilization-induced plant
87 composition and biomass changes also influence soil physico-chemical properties (Keller et
88 al., 2006). Peatlands are nutrient-poor systems (Bragazza et al., 2006), and inorganic N and P
89 fertilizer inputs are well known to change microbial enzyme activity and stoichiometry
90 (Pinsonneault et al., 2016; Song et al., 2019). Previous studies have found that agricultural N
91 and P input changed variation in the activities of P and N hydrolase in natural peatlands
92 (Wright and Reddy, 2001; Prenger and Reddy, 2004). While modern agriculture now attempts
93 to create an environment-friendly strategy for sustainable management (Zhang et al., 2012),
94 for agricultural peatlands, maintaining food yields and decreasing fertilizer input in deeper
95 soil itself and its nearby natural peatlands are optimum. Therefore, a full evaluation of the
96 effects of agricultural intervention on microbial enzyme activity and stoichiometry is needed
97 to improve fertilization management in its widest context.

98 In Northeast China, peatlands have been increasingly cultivated in order to increase food
99 supply since the 1950s, and as cultivated peatlands continue to experience frequent flooding,
100 most were cultivated for rice production. In this study, we investigate the effects of
101 fertilization on both agricultural and nearby natural peatlands in terms of soil carbon (C),
102 nitrogen (N), phosphorus (P) and the key enzymes for C, N and P nutrient cycling. We
103 anticipated (1) that microbial enzymatic activities would be strongly correlated with changes
104 in soil nutrient levels under agricultural intervention and (2) that the variations in microbial
105 enzymatic activity and stoichiometry would exhibit greater similarities in agricultural
106 peatlands than undisturbed peatlands. The information obtained from our study would be
107 important for monitoring nutrient flows from agricultural peatlands and their effects on soil
108 nutrient cycling, potentially greatly improve future strategies for fertilization management.

109 **2 Materials and methods**

110 **2.1 Study site**

111 This study was conducted at the Jinchuan Peatlands of Changbai Mountain, Northeastern
112 China. The annual average temperature and precipitation are 4.1 °C and 704.2 mm,
113 respectively. The peat is 4 to 6 m deep in this area and typical plant include *Carex chmidtii*,
114 *Etulao valifolia*, *Phragmites australis* and *Thelypteris palustris*. Peatland reclamation began
115 1960's mainly for paddy creation. In general, N fertilizer (urea) is applied mid-May, and
116 throughout June for rice growth, totalling 260 kg N ha⁻¹ year⁻¹. In contrast, P fertilizer is
117 applied in mid-May only, amounting to 70 kg P ha⁻¹ year⁻¹. These agricultural peatlands have
118 been found to influence hydrology and soil organic carbon accumulation of nearby natural
119 peatlands (Zhang et al., 2016; Wang et al., 2017).



120
121 **Figure 1** Sampling sites in Jinchuan Peatlands, a temperate fen in Northeastern China

122 2.2 Sample collection

123 We selected agricultural peatlands (paddy fields), peatlands disturbed by their close
124 proximity to these agricultural peatlands (disturbed peatlands) and undisturbed peatlands with
125 similar plant composition (Figure 1) as described by Zhang et al. (2016). We established five
126 random sites in each type and distance between each site was at least 20 meters. Five soil
127 cores were collected from each site using a core sampler at 0-10 cm, 10-20 cm, 20-30 cm, 30-
128 40 cm and 40-50 cm soil depths, then each sample from a given depth mixed to make a
129 composite sample for that depth, totally 75 samples in all. Each sample was divided into two

130 parts, one was used for analysis of soil properties and the other for measuring microbial
131 enzyme activity and stored at 4 °C.

132 **2.3 Sample analysis**

133 Soil organic carbon (SOC) was determined by the dichromate oxidation method. Soil
134 total nitrogen (TN) was analyzed using the Kjeldahl method. And soil total phosphorus (TP)
135 was measured by the Mo-Sb colorimetric method.

136 The determination of microbial enzyme activity was performed as described by Saiya-
137 Cork et al. (2008), which began within one week of sample collection. We selected of β -D-
138 glucosidase (BDG), N-acetyl- β -glucosaminidase (NAG) and phosphatase (PHO) as indicators
139 for C-, N- and P-cycling, respectively (Luo et al., 2017). Briefly, 1.0 g soil was homogenized
140 in 125 ml of acetate buffer (50 mM, pH 5.0) in a blender for 1 min. We conducted assays
141 using 96-well microtiter plates, with eight replicate wells per sample per assay. The analysis
142 included eight replicate wells for each blank (50 μ l of acetate buffer plus 200 μ l of sample
143 suspension), a negative control (50 μ l substrate solution plus 200 μ l of acetate buffer), and a
144 quench standard (50 μ l of standard 10 mM 4 methylumbelliferone plus 200 μ l sample
145 suspension). The microplates were incubated in the dark at 20 °C for 4 h. To stop the reaction,
146 a 10 μ l aliquot of 1 M NaOH was added to each well. Fluorescence was measured using a
147 microplate fluorometer with 365 nm excitation and 450 nm emission filters (Synergy H4
148 BioTek, USA). After correcting for negative controls and quenching, activities were
149 expressed in units of $\text{nmol h}^{-1} \text{g}^{-1}$ dry soil.

150 **2.4 Statistical analysis**

151 One-way ANOVA was used to determine statistical differences in soil nutrient levels and
152 microbial enzyme activity. Significant differences between means were established by
153 Duncan test at $p < 0.05$. The relationship between soil nutrient and microbial enzyme activity
154 were assessed using linear regression model within agricultural, disturbed and undisturbed
155 peatlands. Microbial enzyme activity was log-transformed to meet the assumptions of
156 homoscedasticity. These statistical analyses were performed by SPSS 23.0.

157 We used Redundancy Analysis (RDA) to know the contribution of environmental
 158 variables (SOC, TN, TP and their stoichiometries) to variation of microbial enzyme activity
 159 and stoichiometry. Variables were log-transformed and centered to equalize the weight of
 160 variables with ranges of different orders of magnitude. Interactive forward selection
 161 procedures with unrestricted permutation tests (499 permutations) were used to determine the
 162 significant environmental variables to be included in final models. Further, multivariate
 163 analysis was performed using two-dimensional nonmetric multidimensional scaling (NMDS)
 164 using a Bray–Curtis dissimilarity matrix to calculate the similarities of microbial enzyme
 165 activity and stoichiometry among different sites. These statistical analyses were performed
 166 using Canoco 5.0 (Microcomputer Power, Ithaca, NY, USA).

167 **3 Results**

168 **3.1 The effect of agriculture on soil C, N and P**

169 Compared with undisturbed peatlands, agricultural peatlands SOC content decreased by
 170 29.7 %, 11.8 % and 2.3 % at 0-10 cm, 10-20 cm, and 20-30 cm soil layers ($p < 0.05$, Table 1).
 171 TN content decreased by 22.4% at 0-10 cm but increased by 6.3 % at 20-30 cm soil depth
 172 ($p < 0.05$, Table 1). TP content decreased with depth, with the highest 51.7 % at 0-10 cm and
 173 lowest 32.7 % in the 40-50 cm soil layer ($p < 0.05$, Table 1). In disturbed peatlands, SOC did
 174 not show any differences down the profile, TN increased by 7.1 % and 8.3 % at 10-20 cm and
 175 20-30 cm ($p < 0.05$, Table 1), respectively. TP increased by 15.9 %, 23.2 %, 26.0 %, and
 176 16.6 % at 0-10 cm, 10-20 cm, 20-30 cm and 30-40 cm, respectively ($p < 0.05$, Table 1).

177 In agricultural peatlands, SOC:TN at 10-30 cm, and both SOC:TN and TN:TP at 0-50
 178 cm were significantly lower than those in undisturbed peatlands ($p < 0.05$, Table 1). And in
 179 disturbed peatlands, SOC:TN decreased at 10-20 cm, and both SOC:TP and TN:TP at 10-40
 180 cm significantly decreased compared with undisturbed peatlands ($p < 0.05$, Table 1).

181

182 Table 1 The effects of agricultural intervention on soil properties

Depth (cm)	SOC (g kg ⁻¹)	TN (g kg ⁻¹)	TP (g kg ⁻¹)	SOC:TN	SOC:TP	TN:TP
---------------	------------------------------	-----------------------------	-----------------------------	--------	--------	-------

Agricultural Peatlands	0-10	296.76b (30.35)	15.07 b (1.32)	1.12 a (0.05)	19.62 a (0.85)	269.11 b (34.74)	13.56 b (1.38)
	10-20	386.86b (9.01)	19.20 b (0.48)	1.12 a (0.04)	20.16 c (0.26)	348.09 c (12.40)	17.27 c (0.57)
	20-30	423.09b (6.2)	20.80 a (0.27)	1.04 a (0.05)	20.80 b (0.56)	422.30 c (24.48)	20.24 c (0.80)
	30-40	437.54 a (7.93)	20.77 a (0.54)	0.97 a (0.03)	21.13 a (0.70)	453.74 c (19.75)	21.51 c (0.86)
	40-50	431.76b (8.29)	20.21 a (0.40)	0.96 a (0.04)	21.39 a (0.47)	450.18 b (23.26)	21.02 b (0.82)
Disturbed Peatlands	0-10	426.47 a (7.10)	20.14 a (0.76)	0.86 b (0.01)	21.28 a (0.78)	497.68 a (15.93)	23.50 a (1.06)
	10-20	4443.79 a (4.01)	20.66 a (0.020)	0.90 b (0.02)	21.48 b (0.14)	493.76 b (13.74)	22.99 b (0.64)
	20-30	455.08 a (3.89)	21.19 a (0.38)	0.83 b (0.01)	21.51 ab (0.53)	549.01 b (5.68)	25.57 b (0.55)
	30-40	451.28 a (8.36)	19.55 a (0.83)	0.78b (0.01)	23.18 a (0.68)	580.23 b (11.06)	25.12 b (0.95)
	40-50	453.93 a (4.49)	20.68 a (0.77)	0.79 b (0.02)	22.06 a (0.77)	574.88 a (10.89)	26.14 a (0.68)
Undisturbed Peatlands	0-10	422.10 a (6.05)	19.41 a (0.35)	0.74 c (0.01)	21.79 a (0.61)	588.52 a (23.92)	26.25 a (0.65)
	10-20	438.44 a (3.17)	19.30 b (0.25)	0.73 c (0.02)	22.74 a (0.46)	600.94 a (17.70)	26.43 a (0.53)
	20-30	442.44 ab (5.55)	19.57 b (0.30)	0.66 c (0.01)	22.63 a (0.20)	672.57 a (6.48)	29.74 a (0.48)
	30-40	439.89 a (5.55)	18.87 a (0.52)	0.67 c (0.02)	22.27 a (0.81)	660.98 a (22.58)	28.47 a (0.95)
	40-50	444.63 ab (6.67)	19.22 a (0.45)	0.73 b (0.02)	23.21a (0.84)	613.35 a (24.51)	26.44 a (0.57)

183 Notes: Different letters indicate significant differences at the same depth. Soil organic carbon
184 (SOC); Total nitrogen (TN); Total phosphorus (TP).

185 3.2 The effect of agricultural intervention on microbial enzymatic activities

186 In undisturbed peatlands, BDG and PHO activity showed high similarity at 0-50 cm
187 depth (Figure 2a & c). Compared with undisturbed peatlands, in agricultural peatlands, BDG
188 activity did not show any differences, although NAG and PHO activity significantly
189 decreased across the depths (Figure 2b). In disturbed peatlands, BDG activity significantly
190 increased at 0-10 cm and 30-40 cm, NAG activity significantly increased at 0-10 cm and
191 decreased at 20-30 cm and 40-50 cm, and PHO activity significantly declined at all the depths.

192 In agricultural and disturbed peatlands, both BDG:NAG and BDG:PHO were
193 significantly higher at all the depth than those in undisturbed peatlands ($p < 0.05$, Figure 3a &

194 b). Overall, they also showed higher ratios in agricultural peatlands than disturbed peatlands
195 (Figure 3a & b).

196


197 **Figure 2** The effects of agricultural intervention on microbial enzyme activities. (a) β -D-
198 glucosidase (BDG); (b) N-acetyl- β -glucosaminidase (NAG); (c) phosphatase (PHO). Different
199 letters indicated significant differences at same depth.

200

201 **Figure 3** The effects of agricultural intervention on microbial enzyme stoichiometry. (a) ratio of
202 β -D-glucosidase to N-acetyl- β -glucosaminidase (BDG:NAG); (b) ratio of β -D-glucosidase to
203 phosphatase (BDG:PHO); (c) ratio of N-acetyl- β -glucosaminidase to phosphatase (NAG:PHO).
204 Different letters indicated significant differences at the same depth.

205 **3.3 Correlation between soil nutrients and microbial enzyme activities**

206 Linear regression model showed that SOC significantly increased BDG, NAG and PHO
207 activity, respectively ($p < 0.05$, Figure 4adg). TN was positively correlated with BDG and
208 NAG activity, respectively ($p < 0.05$, Figure 4b&e). And TP significantly decreased NAG and
209 PHO activity, respectively ($p < 0.01$, Figure 4f&i).



210

211 **Figure 4** The relationship between soil nutrient and microbial enzyme activity. β -D-glucosidase
 212 (BDG); N-acetyl- β -glucosaminidase (NAG); phosphatase (PHO); Soil organic carbon (SOC); total
 213 nitrogen (TN); total phosphorus (TP).

214 **3.4 Influences of agricultural intervention on microbial enzyme activity and**
 215 **stoichiometry**


216 Clusters in enzymatic activity and stoichiometry in agricultural, disturbed and
 217 undisturbed peatlands can be seen in the NMDS ordination graph (Figure 5). Variations in
 218 agricultural and disturbed peatlands showed a degree of similarity, which was absent in that
 219 of undisturbed peatlands.

220 The SOC, TN, TP and their stoichiometries explain 50.8% of the variation of microbial
 221 enzymatic activity and their stoichiometry based on ordination analysis using redundancy

222 analysis (Table 2 & Figure 6). And TP and SOC were determined to be factors significantly
 223 explaining 38.3% and 8%, respectively (Table 2 & Figure 6).

224

225 **Figure 5** NMDS analysis of the composition of microbial enzyme activity and stoichiometry.
 226 Squares represent agricultural peatlands, circles represent disturbed peatlands, triangles represent
 227 undisturbed peatlands. Yellow, red, green, blue, and black represent 0-10 cm, 10-20 cm, 20-30 cm,
 228 30-40 cm and 40-50 cm, respectively. Resemblance distance measure: Bray-Curtis similarity.



229

230 **Figure 6** Redundancy analysis ordination plot of enzymatic variables constrained by
 231 environmental variables. Enzymatic data were log-transformed and centered to normalize weights
 232 of data due to differences in orders of magnitude and ranges.

233 **Table 2** Results of redundancy analysis model of enzymatic variation using environmental
 234 variables, determined by forward selection procedure with unrestricted permutation tests
 235 the contribution of variables. Soil organic carbon (SOC); Total nitrogen (TN); Total phosphorus
 236 (TP).

Variables	Explain(%)	Contribution(%)	Pseudo F	p
TP	38.3	75.5	45.3	0.002
SOC	8.0	15.8	10.7	0.002

TN	0.8	1.5	1.1	0.358
SOC:TN	1.7	3.3	2.3	0.354
SOC:TP	1.2	2.3	1.6	0.358
TN:TP	0.8	1.6	1.1	0.354
Total	50.8	100		

237 **4 Discussion**

238 **4.1 Soil C, N and P changes associated with contrasting levels of agricultural influence**

239 Agricultural activities have been found to influence hydrology in peatlands, which
240 increases surface peat aerobic decomposition in agricultural and disturbed peatlands (Zhang et
241 al., 2016; Wang et al., 2017). Peatlands are widely acknowledged as N or P limited ecosystem
242 (Hill et al., 2012), and as such, any increase in nutrient availability potentially stimulates
243 organic matter decomposition, and causes fertilization-induced plant biomass changes (Keller
244 et al., 2006), which further influences SOC. These could best explain the large soil organic
245 carbon loss in surface soil of agricultural peatlands and weak variance in disturbed peatlands.

246 Inorganic fertilization would be expected to cause nutrients to leach deeper into a given
247 soil or from agricultural peatlands to nearby natural peatlands, changing soil nutrient levels
248 therein (Kogel-Knabner et al., 2010; Steinmuller et al., 2016). In our study we found that in
249 agricultural peatlands, TN was 22.4% lower in surface 0-10 cm soil than in less disturbed
250 soils suggesting a higher degree of N mobility, and losses potentially through leaching. In the
251 nearby peatlands, TN remained unchanged at the surface 0-10cm but increased 7.3-8.3 % at
252 10-30 cm depth (Table 1). Total P in agricultural peatlands was higher than in disturbed
253 peatlands, especially in the surface soil (Table 1). Previous studies have shown N or P
254 fertilizer uptake by crops and harvest can reduce nutrient abundance in surface soils (Cao et
255 al., 1984), however, plant-assimilated N and P would return to the soil through plant
256 decomposition in natural peatlands. It is likely that this is observed because reactive nitrogen
257 is easily leached (Kogel-Knabner et al., 2010) and liable to oxidation through nitrification,
258 denitrification and feammox (Yang et al., 2012; Shi et al., 2017), all contributing to N loss in
259 these upper layers. In contrast, P is easily absorbed by soil minerals (Zhao et al., 2018), and

260 previous studies show that surface soils in this area contain an abundance of minerals (Qin et
261 al., 2020), which would help to retain P surface soil of these agricultural peatlands. However,
262 many factors influence soil nutrient levels, and overall there is little doubt that the observed
263 increased N and P levels indicate that fertilization affects soil nutrient levels in agricultural
264 peatlands and then further impacts neighboring peatlands. Soil C:N has been suggested as an
265 indicator for nutrient transformation (Spohn et al., 2013; Hu et al., 2019), in our study, there
266 were no differences in C:N at 30-50 cm in agricultural or disturbed peatlands compared with
267 undisturbed peatlands, providing reassurance that there are limits to the depths affected by
268 fertilizer application. This is most likely due to the low hydraulic conductivity at depth in
269 peatlands.

270 **4.2 The response of microbial enzymatic activity to agricultural intervention**

271 Compared soil physico-chemical properties, microbial enzymatic activities may provide
272 an earlier warning of the effects of agricultural intervention by reflecting changes in C, N and
273 P levels as they begin to occur (Gil-Sotres et al., 2005; Lagomarsino et al., 2009; Burns et al.,
274 2013). In our study, both NAG and PHO activity significantly decreased down the profile in
275 agricultural peatlands, a trend also seen at least for PHO activity, in the more disturbed
276 peatlands. Ratios of BDG to NAG and BDG to PHO significantly increased down the profile
277 in agricultural peatlands and deeper soil in disturbed peatlands, suggesting greater microbial
278 demand for carbon (Luo et al., 2017). Further regression analysis showed that SOC positively
279 influence BDG, NAG and PHO activity, while total N positively influenced BDG and PHO
280 activity, however, total P negatively influenced BDG and PHO activity ($p < 0.05$, Figure 4b &
281 e). According to the resource allocation models, increased inorganic N or P availability could
282 decrease microbial N and P acquiring enzyme activities leading to increases in of microbial C
283 acquisition (Sinsabaugh and Moorheas, 1994; Pinsonneault et al., 2016). C and N are essential
284 substrates and nutrients for microorganisms to be able to synthesize enzymes and support
285 productivity (Olander and Vitousek, 2000).

286 However, as proteins, phosphatases have relatively high N concentrations (between 8%
287 and 32%), may represent a significant investment of N (Treseder and Vitousek, 2001), and

288 high P may also decrease microbial N demand. Furthermore, microbial growth and function
289 could also be inhibited in presence of excessive P (Conrad et al., 2000; Li et al., 2017). Based
290 on the degree of correlation, P strongly inhibited NAG activity despite its promotion by SOC
291 and total N (Figure 4).

292 Clearly the microbial enzyme activities and their respective stoichiometries sensitively
293 reflect the agricultural intervention based on NMDS. Moreover, RDA analysis confirmed that
294 total P and SOC were determining factors that significantly explain 38.3% and 8% of the
295 variation, respectively. These observations confirm that microbial enzymatic activity and
296 stoichiometry are strongly influenced by changes of nutrient levels and agricultural
297 intervention (Lagomarsino et al., 2009; Jian et al., 2016).

298 **Implications for managements in agricultural peatlands**

299 N fertilizer consumption has increased by 18% over a period of just 20 years as part of
300 global efforts to increase crop yields (Allen and Beatty, 2011). Generally, N fertilizers are
301 over-applied, at rates far exceeding the maximum demand of the crop (Allen and Beatty,
302 2011). Not only is this a waste of resources but it also results in nitrogen pollution of the
303 atmosphere, of rivers and of the oceans. Nutrient inputs also destabilize carbon stores by
304 increasing organic matter decomposition, and thereby indirectly decreases soil N retention
305 capacity (Zhu and Wang et al., 2011). This can mislead farmers into believing that crops need
306 even more N fertilization with long-term tillage, further exacerbating N fertilization rates. P is
307 easily absorbed by mineral (Emsens et al., 2017), leaching and runoff with water is the main P
308 loss pathways, from this point, P fertilization seems much abundant than N fertilizer in such
309 agricultural system, it is necessary to reduce the quantity of P fertilizer.

310 Modern agricultural practice now sees integrated nutrient management as the most
311 sustainable strategy for increasing food production as this decreases chemical fertilizer
312 consumption (Zhang et al., 2012). Many studies have focused on crop nutrient uptake,
313 nutrient supply in root zone, and fertilization loss (Zhang et al., 2012; Yousaf et al., 2016).
314 However, the environment peat occurs high water table levels and soils with high hydraulic
315 conductivity near the surface that favor lateral water movement. Agricultural activity in such

316 peatlands can as a consequence impart wider effects on nearby natural peatlands as a result of
317 the connectivity through surface and groundwater flows. Thus, monitoring and evaluation is
318 important for maintaining natural peatland system stability and ecological services. Our study,
319 suggests that soil microbial enzymatic activity and stoichiometry could provide an effective
320 early warning indicator for risks from agricultural peatlands to their adjacent natural systems.
321 Moreover, coupled analyses of soil properties and microbial enzyme activities could provide a
322 detailed insight into soil carbon and nutrient cycling, which can ensure that the risks from
323 fertilizer management in agricultural peatlands towards their adjacent natural ecosystems is
324 minimized.

325 **Conclusions:**

326 Compared with undisturbed peatlands, soil properties in agricultural and disturbed
327 peatlands showed significant impacts from disturbance that were strongly correlated with
328 changes in microbial enzymatic activities. When coupled with consequent changes in
329 microbial enzymatic stoichiometry and changes detected soil nutrient levels, far more
330 sensitive indicators of ecological changes are achievable than by measuring soil properties
331 alone. Variations in microbial enzymatic activity and stoichiometry proved to be highly
332 responsive to agricultural intervention. Thus, such measures are proposed as valuable
333 indicators of agricultural intervention that could be of great value in monitoring the success of
334 future fertilization strategies aimed at a more sustainable approach to agriculture.

335 **Acknowledgments**

336 This work was supported by the National Natural Science Foundation of China
337 (41771120, U19A2042) and Jilin Provincial Science and Technology Department
338 (20190201308JC, 20200201195JC), and the Youth Innovation Promotion Association CAS
339 378 (2019234).

340

341 **References:**

342 Allen, G., Beatty, P.H., 2011. Fertilizing nature: A tragedy of excess in the commons. *Plos*
343 *Biol.* 9, e1001124.

344 Bader, C., Mueller, M., Schulin, R., Leifeld, J., 2018. Peat decomposability in managed
345 organic soils in relation to land use, organic matter composition and temperature.
346 *Biogeosciences*, 15, 703-719.

347 Berger, S., Gebauer, G., Blodau, C., Knorr, K.H., 2017. Peatlands in a eutrophic world -
348 Assessing the state of a poor fen-bog transition in southern Ontario, Canada, after long
349 term nutrient input and altered hydrological conditions. *Soil Biol. Biochem.*, 114, 131-
350 144.

351 Bragazza, L., Freeman, C., Jones, T.G., et al., 2006. Atmospheric nitrogen deposition
352 promotes carbon loss from peat bogs. *PNAS* 103, 19386-19389.

353 Burns, R.G., DeForest, J.L., Marxsen, J., Sinsabaugh, R.L., et al., 2013. Soil enzymes in a
354 changing environment: Current knowledge and future directions. *Soil Biol. Biochem.* 58,
355 216-234.

356 Cao, Z.H., DeDatta, S.K., Fillery, I.R.P., 1984. ¹⁵Nitrogen balance and residual effects of
357 urea-N in wetland rice fields as affected by deep placement techniques. *Soil Sci. Soc.*
358 *Am. J.* 48, 203-208.

359 Conrad, R., Klose, M., Claus, P., 2000. Phosphate inhibits acetotrophic methanogenesis on
360 rice roots. *Appl. Environ. Microb.* 66, 828-831

361 Eickenscheidt, T., Heinichen, J., Droesler, M., 2015. The greenhouse gas balance of a drained
362 fen peatland is mainly controlled by land-use rather than soil organic carbon content.
363 *Biogeosciences*, 12, 5161-5184.

364 Emsens, W.J., Aggenbach, C.J.S, Smolders, A.J.P, Zak, D., van Diggelen, R., 2017.
365 Restoration of endangered fen communities: the ambiguity of iron–phosphorus binding
366 and phosphorus limitation. *J. Appl. Ecol.* 54, 1755-1764.

367 Gil-Sotres, F., Trasar-Cepeda, C., Leiros, M.C., Seoane, S., 2005. Different approaches to
368 evaluating soil quality using biochemical properties. *Soil Biol. Biochem.* 37, 877-887.

369 Gorham, E., 1991. Northern peatlands: role in the carbon cycle and probable responses to
370 climatic warming. *Ecol. Appl.* 1, 182-195.

371 Hill, B. H. et al. 2014. Ecoenzymatic stoichiometry and microbial processing of organic
372 matter in northern bogs and fens reveals a common P-limitation between peatland types.
373 *Biogeochemistry* 120, 203-224.

374 Hu, C., Li, F., Xie, Y.H., Deng, Z.M., Hou, Z.Y., Li, X., 2019. Spatial distribution and
375 stoichiometry of soil carbon, nitrogen and phosphorus along an elevation gradient in a
376 wetland in China. *Eur. J. Soil Sci.* 70, 1128-1140.

377 Jian, S.Y., Li, J.W., Chen, J., Wang, G.s., Mayes, M.A., Dzantor, K., Hui, D.F., Leo, Y.Q.,
378 2016. Soil extracellular enzyme activities, soil carbon and nitrogen storage under
379 nitrogen fertilization: A meta-analysis. *Soil Biol. Biochem.* 10, 32-43.

380 Keller, J.K., Bauers, A.K., Bridgham, S.D., Kellogg, L.E., Lversen, C.M. 2006. Nutrient
381 control of microbial carbon cycling along an ombrotrophic-minerotrophic peatland
382 gradient. *J. Geophys. Res-Bioge.* G03006.

383 Kogel-Knabner, I., Amelung, W., Cao, Z.Z., Fiedler, S., Frenzel, P., Jahn, R., Kalbitz, K.,
384 Kolbl, A., Schloter, M., 2010. Biogeochemistry of paddy soils. *Geoderma* 157, 1-14.

385 Koerselmann, W., Bakker, S.A., Blorn, M., 1990. Nitrogen phosphorous and potassium
386 budgets for two small fens surrounded by heavily fertilized pastures. *J. Ecol.* 78,
387 428e442.

388 Lagomarsino, A., Moscatelli, M.C., Di Tizio, A., Mancinelli, R., Grego, S., Marinari, S., 2009.
389 Soil biochemical indicators as a tool to assess the short-term impact of agricultural
390 management on changes in organic C in a Mediterranean environment. *Ecol. Indic.* 9,
391 518-527.

392 Leifeld, J. Menichetti, L., 2018. The underappreciated potential of peatlands in global climate
393 change mitigation strategies. *Nat. Commun.* 9, 1071.

394 Li, L.N., Qu, Z., Jia, R., Wang, B.L., Wang, Y.Y., Qu, D., 2017. Excessive input of
395 phosphorus significantly affects microbial Fe(III) reduction in flooded paddy soils by
396 changing the abundances and community structures of Clostridium and Geobacteracea.
397 *Sci. Total Environ.* 607, 982-991.

398 Li, T., Bu, Z., Liu, W., Zhang, M., Peng, C., Zhu, Q., Shi, S., Wang, M., 2019. Weakening of
399 the 'enzymatic latch' mechanism following long-term fertilization in a minerotrophic
400 peatland. *Soil Biol. Biochem.* 136, 107528.

401 Luo, L., Meng, H., Gu, J., 2017. Microbial extracellular enzymes in biogeochemical cycling
402 in ecosystems. *J. Environ. Manage.* 197, 539-549.

403 Olander, L.P., Vitousek, P.M., 2000. Regulation of soil phosphatase and chitinase activity by
404 N and P availability. *Biogeochemistry*, 49, 175-190.

405 Pajares, S., Gallardo, J.F., Masciandaro, G., et al., 2009. Biochemical indicators of carbon
406 dynamic in an Acrisol cultivated under different management practices in the central
407 Mexican highlands. *Soil Till. Res.* 105, 156-163.

408 Pinsonneault, A.J., Moore, T.R., Roulet, N.T., 2016. Effects of long-term fertilization on peat
409 stoichiometry and associated microbial enzyme activity in an ombrotrophic bog.
410 *Biogeochemistry*, 129, 149-164.

411 Prenger, J.P., Reddy, K.R., 2004. Microbial enzyme activities in a freshwater marsh after
412 cessation of nutrient loading. *Soil Sci. Soc. Am. J.* 68, 1796-1804.

413 Qin L., Wei T., Liang Y., Freeman C., Ming, J., 2020. Nitrogen availability influences
414 microbial reduction of ferrihydrite-organic carbon with substantial implications for
415 exports of iron and carbon from peatlands. *Appl. Soil Ecol.* 153, 103637.

416 Rydin, H., Jeglum, J.K., 2006. *The biology of peatlands*, Oxford University Press Inc., New
417 York.

418 Saiya-Cork, K.R., Sinsabaugh, R.L., Zak, D.R., 2002. The effects of long term nitrogen
419 deposition on extracellular enzyme activity in an *Acer saccharum* forest soil. *Soil Biol.*
420 *Biochem.* 34, 1309-1315.

421 Saurich, A., Tiemeyer, B., Dettmann, U., Don, A., 2019. How do sand addition, soil moisture
422 and nutrient status influence greenhouse gas fluxes from drained organic soils? *Soil Biol.*
423 *Biochem.* 135, 71-84.

424 Shi, Y., Wang, Z.Q., He, C.G., Zhang, X.Y., Sheng, L.X., Ren, X.D., 2017. Using ¹³C
425 isotopes to explore denitrification-dependent anaerobic methane oxidation in a paddy-
426 peatland. *Sci. Rep.* 7, 40848.

427 Song, Y.Y., Song, C.C., Ren, J.S., Zhang, X.H., Jiang, L., 2019. Nitrogen input increases
428 *Deyeuxia angustifolia* litter decomposition and enzyme activities in a marshland
429 ecosystem in Sanjiang Plain, Northeast China. *Wetlands* 39, 549-557.

430 Spohn, M., Babka, B., Giani, L., 2013. Changes in soil organic matter quality during sea-
431 influenced marsh soil development at the North Sea coast. *Catena*, 107, 110–117.

432 Steinmuller, H.E., Graham, S.A., White, J.R., McKee, M., Mendelsohn, I.A., 2016. A
433 decadal-scale nutrient loading study in a coastal wetland: Impaction soil microbial
434 processes. *Ecol. Eng.* 97, 58-63.

435 Treseder, K.K., Vitousek, P.M., 2001. Effects of soil nutrient availability on investment in
436 acquisition of N and P in Hawaiian rain forests. *Ecology* 82, 946-954.

437 Tubiello, F.N., Biancalani, R., Salvatore, M., Rossi, S., Conchedda, G., 2016. A worldwide
438 assessment of greenhouse gas emissions from drained organic soils. *Sustainability*, 8,
439 371.

440 Wang, Z.C., Liu, S.S., Huang, C., Liu, Y.Y., Bu, Z.J., 2017. Impact of land use change on
441 profile distributions of organic carbon fractions in peat and mineral soils in Northeast
442 China. *Catena*, 152, 1-8.

443 Wright, A.L., Reddy, K.R., 2001. Phosphorus loading effects on extracellular enzyme activity
444 in Everglades wetland soils. *Soil Sci. Soc. Am. J.* 65, 588-595.

445 Yang, W.H, Weber, K.A, Silver, W.L., 2012. Nitrogen loss from soil through anaerobic
446 ammonium oxidation coupled to iron reduction. *Nat. Geosci.* 5, 538–541.

447 Yousaf, M., Li, X., Zhang, Z., et al., 2016. Nitrogen fertilization management for enhancing
448 crop productivity and nitrogen use efficiency in a rice-Oilseed rape rotation system in
449 China. *Front. Plant Sci.* 7, 1946.

450 Yu, Z. C., J. Loisel, D. P. Brosseau, D. W. Beilman, and S. J. Hunt. 2010. Global Peatland
451 Dynamics since the Last Glacial Maximum. *Geophys. Res. Lett.* 37, L13402.

452 Zagal, E., Munoz, G., Quiroz, M., et al.,2009. Sensitivity of early indicators for evaluating
453 quality changes in soil organic matter. *Geoderma*, 2009,151:191-198.

454 Zhang, T., Shi, S., Zhang, W., Wu, Y., Yang, M., Wang, P. 2016. Environmental factors and
455 dissolved organic carbon content in a Jinchuan peatland. *Acta Ecol. Sinica* 36,160-165.

456 Zhang, F., et al., 2012. integrated nutrient management for food security and environmental
457 quality in China. *Adv. Agron.* 116, 1-40.

458 Zhao, Z.J., Jin, R., Fang, D., Wang, H., Dong, Y., Xu, R.K., Jiang, J., 2018. Paddy cultivation
459 significantly alters the forms and contents of Fe oxides in an Oxisol and increases
460 phosphate mobility. *Soil Till. Res.* 184, 176-180.

461 Zhu, W.X., Wang, W.W., 2011. Does soil organic matter variation affect the retention of
462 $^{15}\text{NH}_4^+$ and $^{15}\text{NO}_3^-$ in forest ecosystems? *Forest Ecol. Manag.* 261, 675-682.

463

464

465

466

467

468 **Figures & Tables Captions:**

469 **Figure 1** Sampling sites in Jinchuan Peatlands, a temperate fen in Northeastern China.

470 **Figure 2** The effects of agricultural intervention on microbial enzyme activities. (a) β -D-
471 glucosidase (BDG); (b) N-acetyl- β -glucosaminidase (NAG); (c) phosphatase (PHO). Different
472 letters indicated significant differences at same depth.

473 **Figure 3** The effects of agricultural intervention on microbial enzyme stoichiometry. (a) ratio of
474 β -D-glucosidase to N-acetyl- β -glucosaminidase (BDG:NAG); (b) ratio of β -D-glucosidase to
475 phosphatase (BDG:PHO); (c) ratio of N-acetyl- β -glucosaminidase to phosphatase (NAG:PHO).
476 Different letters indicated significant differences at same depth.

477 **Figure 4** The relationship between soil nutrient and microbial enzyme activity. β -D-glucosidase
478 (BDG); N-acetyl- β -glucosaminidase (NAG); phosphatase (PHO); Soil organic carbon (SOC); total
479 nitrogen (TN); total phosphorus (TP).

480 **Figure 5** NMDS analysis of the composition of microbial enzyme activity and stoichiometry.
481 Squares represent agricultural peatlands, circles represent disturbed peatlands, triangles represent
482 undisturbed peatlands. Yellow, red, green, blue, and black represent 0-10 cm, 10-20 cm, 20-30 cm,
483 30-40 cm and 40-50 cm, respectively. Resemblance distance measure: Bray-Curtis.

484 **Figure 6** Redundancy analysis ordination plot of enzymatic variables constrained by
485 environmental variables. Enzymatic data were log-transformed and centered to normalize
486 weights of data due to differences in orders of magnitude and ranges.

487 **Table1** The effects of agriculture intervention on soil properties. Different letters indicate
488 significant differences at same depth. Soil organic carbon (SOC); Total nitrogen (TN); Total
489 phosphorus (TP).


490 **Table 2** Results of redundancy analysis model of enzymatic variation using environmental
491 variables, determined by forward selection procedure with unrestricted permutation tests
492 the contribution of variables.

493

494


495

496 **Figure 1**



499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514


515 **Figure 2**



516
 517
 518

519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535

Figure 3



536
 537
 538

539

540

541

542

543

544

545

546

547

548

549

550

551


552

553

554

555

556 **Figure 4**



● Agricultural Peatlands ● Disturbed Peatlands ● Undisturbed Peatlands

557

558

559

560

561

562


563

564

565

566

567 **Figure 5**



568

569

similarity.

570

571

572

573

574

575

576

577

578


579

580

581

582

583 **Figure 6**



584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600 **Table1**

601

Table1 The effects of agricultural intervention on soil properties

	Depth (cm)	SOC (g kg ⁻¹)	TN (g kg ⁻¹)	TP (g kg ⁻¹)	SOC:TN	SOC:TP	TN:TP
Agricultural Peatlands	0-10	296.76b (30.35)	15.07 b (1.32)	1.12 a (0.05)	19.62 a (0.85)	269.11 b (34.74)	13.56 b (1.38)
	10-20	386.86b (9.01)	19.20 b (0.48)	1.12 a (0.04)	20.16 c (0.26)	348.09 c (12.40)	17.27 c (0.57)
	20-30	423.09b (6.2)	20.80 a (0.27)	1.04 a (0.05)	20.80 b (0.56)	422.30 c (24.48)	20.24 c (0.80)
	30-40	437.54 a (7.93)	20.77 a (0.54)	0.97 a (0.03)	21.13 a (0.70)	453.74 c (19.75)	21.51 c (0.86)
	40-50	431.76b (8.29)	20.21 a (0.40)	0.96 a (0.04)	21.39 a (0.47)	450.18 b (23.26)	21.02 b (0.82)
Disturbed Peatlands	0-10	426.47 a (7.10)	20.14 a (0.76)	0.86 b (0.01)	21.28 a (0.78)	497.68 a (15.93)	23.50 a (1.06)
	10-20	4443.79 a (4.01)	20.66 a (0.0.20)	0.90 b (0.02)	21.48 b (0.14)	493.76 b (13.74)	22.99 b (0.64)
	20-30	455.08 a (3.89)	21.19 a (0.38)	0.83 b (0.01)	21.51 ab (0.53)	549.01 b (5.68)	25.57 b (0.55)
	30-40	451.28 a (8.36)	19.55 a (0.83)	0.78b (0.01)	23.18 a (0.68)	580.23 b (11.06)	25.12 b (0.95)
	40-50	453.93 a (4.49)	20.68 a (0.77)	0.79 b (0.02)	22.06 a (0.77)	574.88 a (10.89)	26.14 a (0.68)
Undisturbed Peatlands	0-10	422.10 a (6.05)	19.41 a (0.35)	0.74 c (0.01)	21.79 a (0.61)	588.52 a (23.92)	26.25 a (0.65)
	10-20	438.44 a (3.17)	19.30 b (0.25)	0.73 c (0.02)	22.74 a (0.46)	600.94 a (17.70)	26.43 a (0.53)
	20-30	442.44 ab (5.55)	19.57 b (0.30)	0.66 c (0.01)	22.63 a (0.20)	672.57 a (6.48)	29.74 a (0.48)
	30-40	439.89 a (5.55)	18.87 a (0.52)	0.67 c (0.02)	22.27 a (0.81)	660.98 a (22.58)	28.47 a (0.95)
	40-50	444.63 ab (6.67)	19.22 a (0.45)	0.73 b (0.02)	23.21a (0.84)	613.35 a (24.51)	26.44 a (0.57)

602 Notes: Different letters indicate significant differences at the same depth. Soil organic carbon

603 (SOC); Total nitrogen (TN); Total phosphorus (TP).

604

605

606

607

608

609

610 **Table 2** Results of redundancy analysis model of enzymatic variation using environmental
611 variables, determined by forward selection procedure with unrestricted permutation tests
612 the contribution of variables. Soil organic carbon (SOC); Total nitrogen (TN); Total phosphorus
613 (TP).

Variables	Explain(%)	Contribution(%)	Pseudo F	p
TP	38.3	75.5	45.3	0.002
SOC	8.0	15.8	10.7	0.002
TN	0.8	1.5	1.1	0.358
SOC:TN	1.7	3.3	2.3	0.354
SOC:TP	1.2	2.3	1.6	0.358
TN:TP	0.8	1.6	1.1	0.354
Total	50.8	100		

614