



ORIGINAL PAPER

Recreational angling as a pathway for invasive non-native species spread: awareness of biosecurity and the risk of long distance movement into Great Britain

E. R. C. Smith  · H. Bennion · C. D. Sayer · D. C. Aldridge · M. OwenReceived: 28 August 2018 / Accepted: 3 December 2019 / Published online: 9 January 2020
© The Author(s) 2020

Abstract Identifying and establishing the relative importance of different anthropogenic pathways of invasive non-native species (INNS) introduction is critical for effective management of their establishment and spread in the long-term. Angling has been identified as one of these pathways. An online survey of 680 British anglers was conducted to establish patterns of movement by British anglers abroad, and to establish their awareness and use of biosecurity practices. The survey revealed that 44% of British anglers travelled abroad for fishing, visiting 72 different countries. France was the most frequently visited country, accounting for one-third of all trips abroad. The estimated time taken to travel from Western Europe into Great Britain (GB) is within the time frame that INNS have been shown to survive on damp angling equipment. Without biosecurity, it is therefore highly likely that INNS could be unintentionally transported into GB on damp angling gear.

Since the launch of the Check, Clean Dry biosecurity campaign in GB in 2011, the number of anglers cleaning their equipment after every trip has increased by 15%, and 80% of anglers now undertake some form of biosecurity. However, a significant proportion of the angling population is still not implementing sufficient, or the correct biosecurity measures to minimize the risk of INNS dispersal on damp angling equipment. With the increase in movement of anglers abroad for fishing, further work is required to establish the potential for INNS introduction through this pathway.

Keywords Angling · Biosecurity · Awareness · Invasive species · Human pathways

Introduction

Introduction of non-native species by human-mediated jump dispersal is well documented and encapsulates a variety of activity, from the unintentional harbouring of non-native species within shipping cargo (Suarez et al. 2001) to intentionally introducing species for economic purposes such as aquaculture in the case of the Signal crayfish (*Pacifastacus leniusculus*) (Holdich et al. 2004). Although many anthropogenic jump dispersal mechanisms or ‘pathways’ have been identified (Hulme 2009), the relative

E. R. C. Smith (✉) · H. Bennion · C. D. Sayer
Environmental Change Research Centre, University
College London, Pearson Building Gower Street, London,
UK
e-mail: emily.r.smith@ucl.ac.uk

D. C. Aldridge
Department of Zoology, University of Cambridge,
Cambridge, UK

M. Owen
Angling Trust, Eastwood House, 6 Rainbow Street,
Leominster, UK

importance of each pathway is unknown. Related to this, is the increasing recognition that, for many invasive non-native species (INNS) the most cost-effective approach to minimising their environmental and socio-economic impacts is prevention of initial establishment in the first place (Leung et al. 2002; Finnoff et al. 2007; Caplat and Coutts 2011; Brundu 2015). Once an INNS is introduced, unless it is detected early and rapid eradication is undertaken, it often becomes highly expensive, and in some cases impossible to completely eradicate (Mack et al. 2000; Kolar and Lodge 2001; Wittenberg and Cock 2001; Simberloff et al. 2013). Recognising the long-term economic and environmental benefits of preventing further INNS invasions, prevention has been placed at the forefront of the EU Regulation of Invasive Alien Species (1143/2014) (Beninde et al. 2014). Following the introduction of this regulation it is now an obligation for EU Member States to investigate and prioritise potential pathways of human INNS introduction (Trouwborst 2015). An INNS pathway refers to a suite of processes or human activities, that result in the intentional or unintentional movement of an INNS from its natural range, either past or present, into a new environment (Genovesi and Shine 2004; Pyšek et al. 2011). Vectors are distinguished as the physical means or agent such as a ship, vehicle wheels or angling net, via which INNS are moved outside their native range. Through the creation of pathway action plans (PAPs), resources can be allocated to target the most significant pathways, or a particular aspect of a vector identified as the weakest link or greatest biosecurity threat. Managing pathways of human introduction represents a more effective approach than individual INNS management as it reduces the risks of all non-native species using that pathway. This is particularly important as the dispersal mechanisms of many non-native species remain uncertain, and due to time lags it is hard to predict which non-native species may, or may not become invasive in the future (Essl et al. 2015).

Recreational angling has been identified in the EU Regulation and the convention on biological diversity (CBD) as a potential human pathway of INNS introduction (Hulme 2009; Harrower et al. 2018). Used traditionally for the provision of food, angling has also evolved into a popular catch-and-release sport in Western countries, with a rod and line used to catch a variety of fish species (Von Brandt 1964; Pitcher and

Hollingworth 2002). Grouped together with aquaculture and other leisure activities, angling has been reported to account for more than 40% of aquatic INNS invasions in Europe (DAISIE 2009). Angling is a highly popular activity, with an estimated 11.7% and 4.8–6.5% of the population in the United States and Europe participating in fishing every year (Hickley 2018). Around 9% of the population in England and Wales aged 12 years or older took part in angling in 2009–2010, equating to around 4.2 million people (Simpson and Mawle 2010; Sports England 2011). However, despite the link between angling and non-native species being reported for many years (Maitland 1987; Winfield et al. 1996; William and Moss 2001; Zięba et al. 2010) the relative importance of angling as a pathway and vector for non-native species dispersal is still relatively unknown. A few studies have been undertaken to investigate the role of angling in the secondary dispersal of INNS between water bodies (Gates et al. 2009; Anderson et al. 2014), and others have reported the potential for INNS introduction and spread from the use of live bait by anglers (Keller et al. 2007; Kilian et al. 2012; Drake and Mandrak 2014; Cerri et al. 2017). In North America, higher numbers of non-native species have also been found to coincide with areas of greater recreational fishing demand (Davis and Darling 2017). However, there have been limited, if any, studies undertaken to investigate the potential for long-distance jump dispersal of INNS between continents/countries on damp angling equipment. This is despite a recent increase in the number of tourists travelling abroad for recreational activities including angling (Hulme 2015).

Many INNS can survive for a few days (Stebbing et al. 2011; Bacela-Spychalska et al. 2013) and in some cases up to two weeks in damp angling equipment and clothing (Fielding 2011; Anderson et al. 2015). In 2011 around 64% of British anglers stated that they fished in more than one catchment per fortnight (Anderson et al. 2014). The high frequency of anglers returning from fishing within the time frame of INNS persistence in damp equipment suggests that angling gear could act as vector for the spread of INNS between waterbodies. Thus, mechanisms need to be implemented to ensure any invasive species present on equipment are removed or killed before re-use. Recognising this, the biosecurity campaign check, clean, dry (CCD) was launched in Great Britain by Defra in 2011. Biosecurity refers to the undertaking of

a set of measures which individually, or collectively, contribute to a reduction in the risk of spreading INNS, including plants, animals and microbes (Dobson et al. 2013; Shannon et al. 2018). The aim of the CCD campaign is to provide simple biosecurity guidance to recreational water users in order to increase awareness of INNS and in turn to minimise their spread. There are further measures that complement the CCD including strategic planning to ensure sites without INNS are visited prior to sites with known INNS populations, and/or rotating different sets of equipment between sites (Dunn and Hatcher 2015). By preventing the spread of INNS in the first place, it may save substantial environmental and economic costs in the long-term due to damage to the environment, and expenses to remove INNS.

Public engagement and compliance will be essential for the success of this biosecurity campaign (Bremner and Park 2007; Garcia-Llorente et al. 2008; Gozlan et al. 2013). People are often the weakest link in the control of INNS species (Cliff and Campbell 2012) and it can take time for individuals to adopt biosecurity measures as a new social norm (Rogers 2003; Prinbeck et al. 2011; Sutcliffe et al. 2018). Consequently, monitoring the uptake of biosecurity by recreational users is essential to assess the success of the campaign and to identify future priorities. However, except for a baseline study conducted during the first year of the CCD launch (Anderson et al. 2014), changes in the biosecurity behaviour of recreational water users including British anglers is unknown. This study explores changes in angling biosecurity behaviour since the launch of the CCD campaign, and assesses the risk of recreational angling activity unintentionally introducing, or spreading, non-native species into Great Britain (GB) from abroad on damp angling equipment (boots, nets). We focus on the dispersal of INNS species potentially transmitted in angling equipment such as macrophytes and macroinvertebrates. Although parasites and diseases such as the Salmon louse (*Gyrodactylus salaris*) are not explicitly investigated, there is also potential for dispersal of these in contaminated angling equipment (Peeler et al. 2004).

Methodology

A structured online questionnaire survey was conducted between the 8th of July and 31st of October

2015. The survey was produced using the online software, SurveyMonkey. The use of the internet for data collection is accepted as an effective approach to data collection, providing access to a geographically dispersed population, and a sampling size not always achievable using an interview-based approach (Couper et al. 2007; Couper and Miller 2008). The questionnaire was publicised to anglers by Angling Trust social media (Facebook and Twitter) and also circulated via email to their members. The Angling Trust is an organisation that represents all game, coarse and sea anglers in England and Wales on environmental and angling issues. As a result, there is potential for a high response from anglers that have an interest in the natural environment as they are more likely to engage with Angling Trust ideas. To account for this, the questionnaire was also circulated to angling clubs, relevant angling magazines, and promoted at three GB angling events. This included two regional angling forums which brought together angling clubs in the southwest and southeast of England, and the Country Land and Business Association (CLA) game fair in northern England. The CLA is a membership organisation for owners of land, property and business in England and Wales, and the fair is well attended by members and the general public. The different events are attended by different angling club representatives and provided an opportunity to promote the questionnaire across a reasonable geographic coverage, whilst minimising bias in responses from particular regions. All of the events were attended in July 2015. Hard copies of the questionnaires were also made available to minimise potential for selection bias by excluding anglers that do not use the Internet. Despite attempts to reduce potential bias through promotion of the questionnaire at other angling events, it should be recognised that data derived from this survey are assumed to represent the maximum percentage of anglers currently conducting biosecurity in GB.

Questionnaire survey design

This study focused on quantifying the potential for recreational angling to facilitate jump dispersal of NNS from Europe to GB by investigating the frequency at which anglers travelled to different countries and undertook biosecurity after a fishing trip. Given this overall aim, a closed-format

questionnaire was deemed the most appropriate approach. Questions that required more extensive individual responses such as names of fishing sites had a ‘free-text’ option included. Interviews and group discussions would have provided a greater insight into why individuals behave in particular ways and how this is influenced by different factors (Longhurst 2010). However, interviews and group discussions would not have reached the high volume of respondents required in this study. Using a web-based approach enabled access to greater numbers of anglers across a larger geographical area within GB (Schmidt 1997).

The questionnaire was organised into marked sections applying filter questions to avoid asking irrelevant questions to the respondents. For example, after asking an individual whether they went fishing abroad, if a respondent answered ‘no’ the questionnaire would automatically skip to the next relevant section. This ensured that the questionnaire was as easy to follow and fill in as possible, thus maximising the number of respondents that completed the questionnaire.

The questionnaire was phrased to allow comparison against the baseline angling awareness survey undertaken by Anderson et al. (2014) in 2011. The first section focused on frequency and patterns of movement of anglers within GB and abroad. Answers were generally quantitative, employing statements such as fishing once a week, every two weeks rather than more generic ‘often’, ‘sometimes’ statements thereby providing a more accurate representation of their activity (Angelsen and Lund 2011). The second section explored the use of different equipment such as nets, slings, waders, and the frequency with which equipment was cleaned and dried. The CCD campaign, as launched in 2011 has been used to promote awareness of INNS and simple biosecurity guidance that can be undertaken by the general public and practitioners in the field to reduce the risk of spreading INNS. It is focused on three main elements: ‘Check’—examining equipment, boats and clothing and removing any fragments of plants, mud or other material, ‘Clean’—thoroughly washing equipment and clothing in hot water or disinfectant, and ‘Dry’—leaving equipment and clothing to dry in the sunlight for at least two days. As these are the key messages promoted by the campaign, these were used to phrase questions around biosecurity procedures conducted by anglers. The final

section of the questionnaire included questions on angler awareness of the CCD campaign and INNS. It is recognised that, by using the terminology ‘INNS’, the questionnaire overlooks non-native species, which after a lag phase, have the potential to become invasive at a later stage (Crooks et al. 1999). However, the focus on the study was to ascertain anglers awareness of INNS. Thus, although biosecurity measures undertaken by anglers are likely to minimise introduction of all non-native species being spread by this vector, to ensure clarity in the questionnaire only the term INNS was used. This section was placed at the end of the survey to minimise the risk of conditioning the respondents’ answers surrounding their cleaning and drying behaviour in the earlier section of the questionnaire. This survey complied with University College London (UCL) guidelines on ethical conduct. Respondents were asked for their age, gender and the first 3–4 digits of their postcode. This information would not enable any respondent to be identified. All data were collected and stored anonymously.

A pilot study was undertaken to pre-test the survey before publishing it online. This ensured that questions were interpreted correctly and that sufficient answer options were available for the closed questions (Gaddis 1998). Ten anglers were asked to undertake the online survey. Following the pilot, minor modifications were made to the final questionnaire to improve question clarity and to include additional tick box options in certain questions such as additional angling equipment. The final questionnaire is available in “[Appendix](#)”.

Data analysis

Differences in biosecurity behaviour between different types of freshwater anglers were analysed. Anglers that fished mainly for Common carp (*Cyprinus carpus*) were treated as a separate group from general coarse anglers who target other freshwater species such as Bream (*Abramas* spp.), Roach (*Rutilus* spp.) and Tench (*Tinca* spp.) Many anglers undertake sea fishing alongside freshwater fishing. However, due to differences in the environmental tolerances of freshwater and marine INNS, particularly in relation to salinity, anglers that only undertook sea fishing were removed from the analysis. This accounted for three respondents only. Subsequently, five different types of anglers were derived: game, competition, lure, coarse-

other and coarse-carp. Match anglers are those that fish in competitions in contrast with the other groups that fish simply for pleasure. Demographic information obtained for the 2015 GB Environment Agency (EA) rod licence data was used to test the representativeness of the sample compared to the overall GB angling population.

Risk categories were ascertained for each respondent based on the CCD campaign. Four categories of risk were assigned: 'Low', 'Minor', 'Moderate' and 'Major' (Table 1). Anglers categorised as 'Low' risk, cleaned and dried their equipment after every trip. The category 'Low' risk was chosen rather than 'No' risk as there is always a small risk that an INNS could be unintentionally transmitted. Anglers classified as 'minor' risk, cleaned and/or dried their equipment after every 2–5 trips, 'moderate' every 6–10 or 11+ trips, and 'major' risk did not clean and/or dry their angling equipment at all. For further clarification, respondents were classified according to their most infrequent cleaning or drying activity. For example, an angler that cleaned their equipment every 6–10 trips, and dried their equipment every time was placed in the moderate risk category. A limitation of this approach is that it assumes equal importance of cleaning and drying in minimising the risk of invasive species being spread. However, some studies suggest that cleaning equipment using hot water is more effective than drying for rapid decontamination of equipment, causing 99% mortality within an hour, compared to drying that took several days (Anderson et al. 2014). For the initial risk analysis, it was also assumed that respondents were cleaning and drying their equipment in accordance with the Check, Clean, Dry campaigns, using hot water at 45 °C (Anderson et al. 2015) and drying their equipment until it was completely dry. This assumption was reviewed in the analysis.

To assess temporal changes in the biosecurity activity of anglers, only anglers that fished at least once a fortnight were included to reflect the approach used in the 2011 baseline data collection. Consequently, for this part of the analysis only 79% (anglers that fished once a fortnight) of the 680 responses were used.

The first 3–4 digits of the respondent's postcode were converted into longitude and latitude data using Doogal (<http://www.doogal.co.uk/BatchGeocoding.php>). These data were then superimposed onto a map of GB in ArcMap (version 10.3.1) to assess the geographic distribution of the sample angler population, and to identify any spatial patterns in the distribution of anglers of different risk in GB.

Kolmogorov–Smirnov tests were undertaken in SPSS 24 to determine the representativeness of the sample questionnaire in relation to the entire British freshwater angling population. Age and gender demographic data were compared against environment agency (EA) rod licence data for 2015 following similar comparisons conducted by Anderson et al. (2014) and White et al. (2005). Rod licence was used as any angler wishing to fish in freshwater bodies in GB requires a licence. Chi squared tests were employed to determine relationships between the risk of types of anglers, their risk categories and awareness of the CCD. As there were less than five anglers who stated that they mainly lure fish, these were removed from this aspect of the analysis to meet the assumption of the Chi squared test. Both tests had over 500 sets of observations indicating robust p-values (Jaeger 2008). Post-hoc Cramer tests were applied to the risk and biosecurity awareness Chi squared tests to assess the significance and size of the effect.

Table 1 Categorisation of anglers' risk based on their cleaning and drying frequency

Risk category	Cleaning and drying frequency	Example
Low	Every trip	Individual cleans and dries after every angling trip
Minor	Undertake both every 2–5 trips	Angler may clean his/her equipment every trip but only dries it every 2–5 trips or vice versa
Moderate	Every 6–10 trips for both cleaning and drying	Angler may only clean his/her equipment every 6–10 trips, but dries every 2–5 or vice versa
Major	Does not undertake both parts of the biosecurity process (clean, dry).	Angler cleans his/her equipment after a trip but does not dry it

Results

Data representativeness

Six-hundred and eighty questionnaires were collected (Fig. 1). This included 637 from the online survey and

43 from hard-copy questionnaires. Respondents represented all of the different types of angling. Respondents represented all of the different groups of angling. Coarse (excluding carp) and game anglers were the most popular types of anglers accounting for 46% and 28% of respondents respectively. 98% of the




Fig. 1 Spatial distribution of anglers that responded to the questionnaire. Anglers that fish abroad are shown in black whilst anglers that only fish in the UK are shown in grey. Locations were identified using the first 3–4 digits of respondents postcode

Table 2 Frequency of fishing trips of British anglers within the UK (%), by fishing type. The group coarse carp refers to anglers that primarily fish for common carp *Cyprinus carpio*

and is treated as a separate group from anglers that fish primarily for other fish species such as roach, tench, bream and rudd (Coarse excluding carp)

	Frequency of fishing per angler type (%)							
	More than once a week	Once a week	Fortnightly	Every 3 weeks	Once a month	Once every 2 months	Once every 3 months	Less than once every 3 months
All	32.1	29.1	17.3	7.1	7.5	2.0	1.4	3.4
Coarse carp	29.7	35.2	17.6	3.3	8.8	2.2	1.1	2.2
Coarse (excluding common carp <i>Cyprinus carpio</i>)	31.6	30.5	16.2	8.1	7.7	2.2	0.7	2.9
Lure	18.8	43.8	37.5	0.0	0.0	0.0	0.0	0.0
Game	31.7	22.6	18.3	9.1	7.3	2.4	3.0	5.5
Competition	54.5	33.3	9.1	0.0	3.0	0.0	0.0	0.0

respondents were male, with the greatest proportion of respondents were aged 65+ (34%) and 55–64 (29%). No significant difference was detected between the demographic ratios of the two groups (K–S Test, $D = 0.13$, $p > 0.05$). The majority of respondents lived in England (Fig. 1). No respondents came from the Republic of Ireland. Motor vehicles were the primary mode of transport for 95% of respondents visiting angling waters in Britain.

Seventy-nine percent of all respondents fished at least once a fortnight, and 61% fished at least once a week (Table 2). Lure and competition anglers fished most frequently, with 100% and 97% of anglers fishing once a fortnight respectively. Game anglers fished the least often, with 72.6% of this group fishing once a fortnight. There was no significant difference between the frequency of fishing trips and type of angler ($n = 576$, $df = 4$, $p = 0.138$).

Fishing abroad

Three hundred of the respondents (44%) used their fishing equipment abroad (Fig. 1), visiting over 70 different countries (Table 3) on six continents. Some 82% of anglers fishing abroad visited at least one European country, with 22 of the current 28 EU Member States listed as a fishing destination. 177 (59%) of British anglers fishing abroad only visited water bodies and fisheries in Europe. Countries in Western Europe were the most popular angler destination, with France and Ireland the most frequently visited countries accounting for 33.3% and 27% of

trips abroad respectively (Fig. 2). The USA and Canada were the most frequently visited countries outside of Europe (17.3% and 10.7% abroad trips, respectively). A total of 49 (16.3%) anglers fishing abroad exclusively visited sites outside of Europe.

Cars and vans were the primary mode of transport for some 43% of the anglers fishing abroad. Airplane travel represented the second most popular mode of transport for anglers fishing abroad, accounting for 34.7% of travel. For British anglers that fished exclusively in Western Europe (Scandinavia, the Netherlands, France, Spain, Ireland, Iceland and Portugal) some 64.7% used motor vehicles as their primary mode of transport. 18.4% and 16.2% of anglers also used airplanes and ferries to travel to these Western European countries. 69.4% of anglers fishing exclusively in France and The Netherlands travelled primarily by car or van.

Angler risk

Some 46% and 45% of anglers that fished at least once a week or fortnightly, respectively, were categorised as low risk, cleaning and drying their equipment after every trip (Table 4). Minor and moderate risk accounted for 23.5% and 9.7% of anglers, respectively. In total, 80% of anglers were conducting some form of biosecurity occasionally after a fishing trip. Major risk anglers that were not cleaning and/or drying their equipment after every trip accounted for 19.5% of anglers. Some 50.4% of anglers fishing less than once per fortnight were considered low risk.

Table 3 Frequency of travel of British anglers to different countries for fishing as proportion of the total number (n = 680) of respondents and a percentage of anglers fishing abroad (total anglers travelling abroad n = 300) (%). Islands placed within brackets were grouped together to represent a single country

Country	Total number of respondents	Percentage of total anglers	Percentage of anglers traveling abroad
Europe			
France	100	14.7	33.3
Ireland	81	11.9	27.0
Spain	44	6.5	14.7
Netherlands, Norway	24	3.5	8.0
Germany	14	2.1	4.7
Iceland	12	1.8	4.0
Italy	11	1.6	3.7
Denmark	10	1.5	3.3
Cyprus, Greece, Portugal, Sweden	8	1.2	2.7
Belgium, Turkey	5	0.7	1.7
Slovenia	4	0.6	1.3
Poland, Romania	3	0.4	1.0
Czech Republic, Hungary, Malta, Bosnia and Herzegovina	2	0.3	0.7
Finland, Luxembourg, Slovakia, Switzerland	1	0.2	0.3
North and South America			
USA	52	7.6	17.3
Canada	32	4.7	10.7
Cuba	12	1.8	4.0
(Canary Islands, Tenerife, Lanzarote, Grand Union), (Trinidad and Tobago)	7	1.0	2.3
Argentina, Cyprus, Thailand	6	0.9	2.0
Antigua, (West Indies, Caribbean, British Virgin Islands, Barbados)	5	0.7	1.7
Alaska, Brazil	4	0.6	1.3
Mexico	3	0.4	1.0
Cayman, Chile,	2	0.3	0.7
Guyana, Peru, Suriname, Venezuela, Jamaica	1	0.2	0.3
Russia			
Russia	8	1.2	2.7
Kazakhstan	1	0.2	0.3
Africa			
South Africa	7	1.0	2.3
Seychelles	4	0.6	1.3
Belize, Kenya	3	0.4	1.0
Egypt, Gambia, Mauritius,	2	0.3	0.7
Guyana, Morocco, Myanmar, Nepal, Oman, Peru, Uganda, Zambia	1	0.2	0.3
Asia			
India, Myanmar, Nepal, Oman, Outer Mongolia, Philippines, Singapore,	1	0.2	0.3
Australasia			
New Zealand	14	2.1	4.7
Australia	9	1.3	3.0
Tasmania	1	0.2	0.3




Fig. 2 Movement of British anglers to different fishing destinations in Europe. Values are given as a percentage of the number of British anglers travelling abroad. Colours were assigned from a gradient of yellow (low), orange (medium) and

red (high) to represent the percentage of British anglers visiting each European country. Countries which were not visited by any British anglers are shown in grey. The individual numbers are available in Table 3

Table 4 Risk categorisation of anglers fishing at least once a week or once a fortnight (%)

	Anglers fishing once a week	Anglers fishing once a fortnight	Anglers fishing less than once a fortnight
Low	46.1	44.8	50.4
Minor	23.6	23.7	21.0
Moderate	11.8	12.0	7.6
Major	18.5	19.5	21.0

There was no spatial pattern in the distribution of anglers of different biosecurity risk within GB (Fig. 3).

The biosecurity risk of anglers fishing at least once a fortnight was investigated and a similar percentage for the angler risk was identified. Over 40% of anglers fishing at least once a fortnight were low risk (Table 4). Twenty percent of anglers that fished at

least once a fortnight were classified as major risk. 17% of anglers fishing once a fortnight never cleaned or dried their equipment after fishing.

Except for competition anglers, 40% of anglers represented by each angler type were categorised as low risk. The carp and game angler categories had the greatest proportion of low risk anglers at 55% and 52.2%, respectively. Carp anglers had the lowest




Fig. 3 Geographic distribution of anglers of different risk throughout Britain. Locations were identified using the first 3–4 digits of their postcode

percentage of high risk anglers, with 12.5% compared to over 20% for coarse, game and competition (match) anglers. However, these differences were not significant ($n = 525$, $df = 3$ $p = 0.105$).

Some 46% of anglers had heard of the CCD campaign. Anglers that had heard of CCD were more likely to undertake biosecurity after every trip (Fig. 4).

One-quarter of anglers that had heard of the campaign cleaned and dried their equipment after every trip. 17.6% of anglers that had not heard of the campaign were classified as a moderate or major biosecurity risk. 12.3% of anglers that had heard of the campaign fell into these two categories. Differences in the risk of anglers based on their awareness of the CCD




Fig. 4 Awareness of British anglers of the Check, Clean Dry biosecurity campaign and their risk category according to the frequency they clean and dry their equipment

campaign were significant ($X^2 = 9.017$, $n = 528$, $df = 3$, $p = 0.03$). A post hoc Cramer’s test of a significant Chi squared test revealed a weak (0.131), significant relationship between the awareness of anglers of the CCD campaign and their risk category ($p = 0.03$).

Of the anglers that undertook biosecurity, 33% cleaned their equipment using hot water. Over 40% used cold water, and 10.8% washed their equipment at

a water bank (Fig. 5). For 37% of anglers cold water was the sole method used to clean their equipment, without any application of detergent or disinfectant. The use of cold water as the only cleaning approach also accounted for 31% of anglers in the low risk category. Some 16.2% of anglers did not conduct any cleaning.

Temporal changes in angler biosecurity behaviour

The proportion of anglers cleaning and drying their equipment after every trip rose from 21% in 2011 to 35.5% in 2015 (Fig. 6). Cleaning frequency also rose over this period from 22 to 37.8%. In contrast, drying frequency fell from 80 to 52.8%. Coinciding with an increase in low risk anglers, the percentage of high-risk anglers not undertaking any biosecurity rose from 11.9% in 2011 to 19.5% in 2015. Restricting analysis to anglers fishing fortnightly and going abroad on fishing trips, the proportion of high-risk anglers increased from 18% to almost 31.8%.




Fig. 5 Methods used by British anglers to clean their equipment after a fishing trip. Some anglers used multiple methods, as a result, the sum of percentages is greater than 100%. ‘Dip’

refers to disinfection through equipment by submersion in a container containing disinfectant provided by the fishery




Fig. 6 Temporal change in the biosecurity of anglers fishing at least once per fortnight since the launch of the Check, Clean Dry campaign in March 2011. Baseline 2011 data was sourced from Anderson et al. (2014)

Table 5 Estimated duration in hours of ferry journeys between the UK and the Netherlands, Belgium France and Ireland. (Source: Brittany Ferries and P&O Ferries <http://www.poferries.com/en/portal> Accessed 02/06/2016)

Ferry routes between Europe, Ireland and GB	Estimated duration (h)	Frequency of ferries (number per day)	Number of cars per ferry
Dover–Calais	1.50	23	520–1059
Hull–Rotterdam	12.00	1	250–850
Hull to Zeebrugge	13.25	1	250–850
Poole to Cherbourg	4.50	1	590
Portsmouth to Caen	6.00	4	600–800
Portsmouth to Cherbourg	3.00	2	235
Portsmouth to Le Havre	3.45	1	160–200
Portsmouth to St Malo	8.00	1–2	580
Plymouth to Roscoff	5.00	5	470
Cairnryan to Belfast	2.25	6	660
Cairnryan to Larne	2.00	7	316–375
Fishguard to Rosslare	3.25	2	564
Liverpool to Belfast	8.00	2	85
Liverpool to Dublin	7.50	3	80–125

Discussion

Angling as a pathway for the unintentional introduction of INNS from Europe

Responding to the obligation for GB to investigate potential human pathways and vectors of INNS introduction, this study represents the first known study assessing the potential for anglers to act as unintentional vectors for the spread of invasive species between countries in Europe. Over 40% of anglers used their equipment abroad for fishing. With 4 million estimated anglers in GB (EA 2004) this extrapolates to around 1.76 million GB anglers potentially travelling abroad with their angling equipment, often to two countries or more. This includes potentially 588,000 travelling to France for fishing, and 847,100 travelling to a country in Western Europe including The Netherlands and Norway. Horizon scanning studies indicate there are at least 16 freshwater invasive species present within Western Europe that are of medium or high-risk of entering GB (Roy et al. 2014; Gallardo et al. 2016), including at least 10 aquatic Ponto-Caspian INNS (Gallardo and Aldridge 2013a). In addition to invasive species, invasive parasites and pathogens such as the ecto-parasite Salmon louse (*Gyrodactylus salaris*) also represent a major biosecurity concern to British waters. *Gyrodactylus salaris* has had devastating impacts on salmon populations in invaded Norwegian rivers and if introduced to GB is likely to have similar negative impacts on GB salmon populations (Peeler et al. 2004). Given the bioclimatic similarities between Western Europe and GB (Gallardo and Aldridge 2013b), it is anticipated that any INNS establishing in these regimes have a high likelihood of being able to survive and spread within GB (Gallardo and Aldridge 2013b, 2015). Consequently, Western Europe represents a substantial source for new invasive species that could be introduced by recreational pathways such as angling.

In addition to the establishment of new INNS there is also the risk of introducing new genetic and phenotypic strands of INNS already established in GB. Some INNS are limited in their current distribution due to genetic or fitness bottlenecks, meaning they are not adequately suited to the environment they have invaded (Crooks et al. 1999). The introduction of new phenotypic variants from different source regions

could release the INNS from these environmental restrictions and facilitate expansions in their distribution, thereby increasing impacts on invaded habitats (Lavergne and Molofsky 2007; Forsman 2014). In GB, some invasive species with limited distribution such as Floating water primrose (*Ludwigia grandiflora*) have been targeted for eradication. The introduction of new phenotypic strands or populations could therefore undermine efforts to control or eradicate these INNS.

With over 40% of British anglers primarily travelling to European fishing sites by motor vehicle, there is a substantial risk of invasive species being transported back into GB on damp angling equipment. Current estimates of the desiccation tolerance of INNS indicate that some are capable of surviving for up to 15 days on damp angling equipment, with this including invasive species already established in GB such as Killer shrimp (*Dikerogammarus villosus*) and Zebra mussel (*Dreissena polymorpha*) (Fielding 2011; Anderson et al. 2014). The ability of INNS species to survive the return journey on damp equipment in motor vehicles needs to be further tested but results from current desiccation studies on INNS, coupled with the short travel time (2–14 h to return from Western Europe to GB) (Table 5) suggests potential for a number of high-risk INNS to be unintentionally transported back from Europe to GB via this conduit. Except for a few studies on individual lakes (Bacela-Spychalska et al. 2013), the presence of INNS in European fishing lakes is little known. The determination of new INNS of high risk of being introduced in GB could potentially provide an alternative or complementary approach to horizon scanning.

Awareness and implementation of biosecurity

It should be recognised that self-report style questionnaires are vulnerable to social desirability response bias, with participants potentially stating answers that they believe to be socially acceptable, or desirable by the researcher (Randall and Fernandes 1991; Lajunen and Summala 2003). This cannot be factored out of any questionnaire (Brace 2008). As a result, it is possible that some respondents may overestimate how often they clean and dry their equipment in order to satisfy the surveyor (Cliff and Campbell 2012). Therefore, although the demographic analysis indicated this study was representative of British angler population holding a rod licence in 2015, the findings

of this questionnaire should be interpreted with caution. Furthermore, the opt-in nature of this questionnaire means there is potential for a greater response from individuals that are aware and care about conservation issues, or who represent more affluent members of the angling community due to the recruitment of responses via the Internet and at the game fair event (White et al. 2005). These individuals are therefore more likely to have excess income to spend on fishing trips abroad. The percentages presented here should therefore be seen as representing a maximum estimate for anglers fishing abroad and undertaking biosecurity. Taking these factors into account, despite the potential respondent errors, the marked increase in biosecurity implementation since 2011 can undoubtedly be attributed to greater uptake of biosecurity. Therefore, there is evidence that anglers are becoming more aware of the risk of invasive species, resulting in the implementation of measures aimed at reducing the risk of dispersing species between water bodies.

Despite the substantial increase in the number of anglers undertaking biosecurity in our study, only 48% of anglers claimed to be aware of the Check Clean Dry campaign. This compares to New Zealand where 80% of recreational users are aware of an equivalent initiative (Anderson 2015). Initiated in 2004, the New Zealand campaign represents a long-established initiative, promoted through a national campaign, and implemented through regional biosecurity plans. Greater levels of awareness may therefore be partially due to the longer exposure of water users to the campaign. However, differing levels in awareness of the campaign, may also be partially attributed to the communication channels through which individuals are hearing about the campaign. Whilst 54% of water users in the regional area of Bay of Plenty, New Zealand had heard of the campaign through signage at boat ramps (Anderson 2015), the majority of British anglers were made aware of the CCD through angling magazines or environmental organisations. Consequently, although British anglers were being informed of the importance of biosecurity, this may not be explicitly tied to the Check, Clean Dry campaign, with this reflected by a weak, but significant association recorded between anglers' awareness of the campaign and their likelihood of frequently undertaking biosecurity. Therefore, it is suggested that practitioners should exercise caution in using awareness of the

Check Clean Dry campaign as the sole predictor of biosecurity uptake by the public in GB. Instead, a combination of factors, including measures of action after leaving the water should be used to monitor uptake of biosecurity procedures.

There has been a marked increase in the total proportion of anglers undertaking some form of biosecurity, in terms of either cleaning or drying their equipment occasionally after a fishing trip. However, over the same time period there has also been a 7% increase in the number of anglers who are not undertaking any biosecurity. INNS are highly adaptable species, capable of regenerating and spreading from a single plant node, asexual invertebrate or egg-bearing macroinvertebrate (Havel and Shurin 2004; Hussner 2009; Okada et al. 2009; Pigneur et al. 2011; Bruckerhoff et al. 2015; Riccardi 2015). Consequently, the unintentional introduction of a single viable plant fragment or live INNS specimen is all that is required to enable a new INNS population to establish. Further work is therefore required to engage with anglers that are still not conducting adequate biosecurity measures. This includes identifying the factors that are currently preventing anglers from undertaking biosecurity. Anglers stated that the availability of a cleaning station and the visual cleanliness of the equipment were some of the main reasons affecting whether an angler cleaned their equipment after use, with the financial cost of undertaking biosecurity and the availability of information being less important. These factors have also been reported as some of the main reasons inhibiting biosecurity for canoeists and boaters (Anderson et al. 2014; De Ventura et al. 2017). Going forward, the importance of routinely cleaning equipment needs to be reiterated, and more resources need to be assigned to ensure easy access to cleaning facilities at the angling waters. In addition to promotion of the CCD campaign, greater clarification is still required on the appropriate methods for cleaning equipment. The use of hot water is increasingly considered to be one of the most efficient, environmentally friendly and cost-effective methods for cleaning equipment and clothing (Beyer et al. 2010; Perepelizin and Boltovskoy 2011; Stebbing et al. 2011; Anderson et al. 2015; Sebire et al. 2018). Disinfectants such as Virkon[®] Aquatic and Virasure[®] have also been proposed as effective approaches to decontaminate equipment and small watercraft (Coughlan et al. 2018; Cuthbert et al. 2018). However,

although the percentage of anglers cleaning their equipment has risen since the launch of the CCD guidance, 50% of anglers are using cold water. For 'low' risk anglers cleaning their equipment after every trip, cold water cleaning accounted for the only cleaning method for 31% of the category. These findings indicate that although anglers are undertaking cleaning approaches, their 'cleaning' method may not be effective in killing any attached INNS. It is therefore essential that promoters of the CCD campaign provide clearer messaging regarding effective cleaning.

Conclusions

Following the launch of the EU Regulation (1143/2014) in 2015, EU Member States are obliged to investigate potential anthropogenic pathways of INNS introduction and create pathway action plans (PAPs) for INNS pathways identified as being a risk (Caffrey et al. 2014; Beninde et al. 2014). This study represents the first attempt at quantifying the importance of angling as an international pathway, providing estimates of the volume of British anglers travelling to Europe for recreational fishing as well as valuable insights into changes in anglers' behaviour since the launch of the invasive species-specific CCD campaign. Although this study has focused on angling within GB, it needs to be recognised that the angling pathway is potentially a global one. With limited biogeographic boundaries between many countries in continental Europe (Rahel and Olden 2008), the potential two-way cross-border movement of INNS by anglers could be significant for many countries. As a result, British anglers travelling abroad could also unintentionally introduce new populations of INNS into water bodies in the destination country. The findings of this study are therefore highly relevant to any country that receives a high volume of British anglers including Ireland and France. This is clearly exemplified by the recent outbreak of Crayfish plague (*Aphanomyces astaci*) in the Republic of Ireland. Considered a last refuge for many native European freshwater species, Ireland is an Ark site for White clawed crayfish (*Austropotamobius pallipes*). Until recently there were no reported occurrences of the invasive Signal crayfish (*P. leniusculus*) or the crayfish plague that *P. leniusculus* carries. However, in 2017, the presence of the plague was confirmed in the River Suir,

County Tipperary, Republic of Ireland, and at time of writing had spread into four different catchments. No signal crayfish have been found so the source of the plague is unknown. There have been some suggestions that it may have been introduced on damp equipment (kayaks, nets, pleasure boats, waders). However, as there are many different users of these catchments, the original source of the introduction cannot be verified. Further research into the ability of pathogens to survive on equipment, and investigations into the presence of invasive species in private fisheries, sailing clubs or other water bodies will help to disentangle the potential sources of different groups of species or pathogens by each pathway.

Since the launch of the CCD campaign in 2011, the percentage of anglers undertaking biosecurity after every trip has almost doubled. Although changes to other recreational water users are unknown, this suggests that the campaign has been successful in increasing awareness of invasive species and encouraging the public to undertake biosecurity measures. The observed success of the CCD campaign as reported in this study, can be used to inform the angling PAP promoting the use of biosecurity as an invasive management tool. These plans are pathway-specific and outline the main policy and management approaches available for the various stakeholders involved. In addition to this, the findings of this study are also applicable to other freshwater pathways where biosecurity is being used as a management technique. This includes the use of recreational boat and kayak activity. Exchanges of best practice between different countries and recreational users could therefore be highly effective in reducing the risk of spread of invasive species.

Further work is required to determine what, if any, invasive species are present in European fishing lakes, and to assess the ability of INNS to survive car trips from Europe back to GB. The findings of this work indicate that angling could be an important pathway for the movement of aquatic INNS, particularly from Western Europe into GB.

Acknowledgements The research was supported by a studentship from the UK Natural Environment Research Council (Grant No. NE/L002485/1) as part of the London NERC Doctoral Training Partnership, and a CASE partnership with the Angling Trust. The author would like to thank the Angling Trust for their advice and support in the development of the survey.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your

intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix

Invasive Non-native Species Survey

Demographic information

1. Age

Under 18		45-54	
18-24		45-54	
25-34		65+	
34-44			

2. Sex

Male	
Female	

3. Please enter the first three or four digits of your postcode (This will enable us to estimate how far different water users travel to take part in their activities. Using the first 3-4 digits will not reveal your home location to us, only the general area that you live in.)

Postcode:

4. Which type of angling do you do? (Choose all that are appropriate)

Game	
Coarse - carp	
Coarse - lure	
Coarse - other	
Competition	
Sea	

5. Which type of angling do you do the most often? (Please only choose one from the selection)

Game	
Coarse - carp	
Coarse - lure	
Coarse - other	
Competition	
Sea	

Threats to UK waterbodies

6. What do you think are the biggest threats to species in UK rivers and lakes?

Please rate the following from 1 to 5 (1 = smallest threat 5 = greatest threat. Only use each number once).

	1	2	3	4	5
Predation					
Pollution					
Invasive non-native species					
Climate change					
Decrease in the quality of habitat					

Movement patterns

7. How frequently do you do angling?

More than once a week		Once every month	
Once a week		Once every 2 months	
Once every 2 weeks		Once every 3 months	
Once every 3 weeks		Less than once every 3 months	

8. On average, how long do you spend at the site when you fish?

Under 2 hours		8-10hours	
2-4hours		10-12hours	
4-6hours		Over 12 hours	
6-8hours			

9. Please list the 3 UK angling venues where you last went fishing in the UK

Venue name (including nearest town, county)	
Venue name (including nearest town, county)	
Venue name (including nearest town, county)	

10. Please list the 3 UK angling venues that you go to the **most frequently**

Venue name (including nearest town, county)	
Venue name (including nearest town, county)	
Venue name (including nearest town, county)	

11. How do you travel to these UK venues? (Choose one or more as relevant)

Motor vehicle e.g. car/van		Train	
Motorbike		Bus	
Cycle		Tube	
Walk		Other (please state)	

12. What form of transport do you use the most often to travel to the UK angling venues?

Motor vehicle e.g. car/van		Train	
Motorbike		Bus	
Cycle		Tube	
Walk		Other (please state)	

Fishing in Europe

13. Have you ever used your own angling equipment outside of the UK?

Yes	
No	

If no, please skip to question 19.

14. If yes, which countries do you go to for fishing?

15. Can you list the 3 angling venues that you most **recently** went fishing in **Europe** (outside of the UK)

Venue name (including nearest town, county)	
Venue name (including nearest town, county)	
Venue name (including nearest town, county)	

16. Can you list 3 angling venues you **most frequently visit in Europe** (outside of the UK)

Venue name (including nearest town, country)	
Venue name (including nearest town, country)	
Venue name (including nearest town, country)	

17. If yes, how do you travel to these countries?

Motor vehicle e.g. car/van		Ferry	
Motorbike		Bus	
Airplane		With a fishing company	
Eurostar		Other (please state)	

18. Which mode of transport do you use the most often to get to the European venues?
Please only select one

Motor vehicle e.g. car/van		Ferry	
Motorbike		Bus	
Airplane		With a fishing company	
Eurostar		Other (please state)	

Equipment use

19. Which of the following items of equipment do you use? (Select all that are appropriate)

Neoprene waders/ wellies		Bass bags	
Felt waders/ wellies		Bait boat	
Pike tube		Weigh sling	
Landing net		Fly fishing belly boats	
Keep net		Row boat	
Carp/pike/ catfish cradle		Other (please state)	
Carp sack			

20. Where do you store your equipment between trips?

Indoors	
Shed or garage	
Outdoors	
Other (please state)	

21. If you use waders, how many hours do you typically keep them in the water for?

Less than one hour		7-8 hours	
1-2 hours		8-9 hours	
2-3 hours		9-10 hours	
3-4 hours		10-11 hours	
4-5 hours		11-12 hours	
5-6 hours		Over 12 hours	
6-7 hours		Do not use waders	

22. If you use a keep net, how many hours do you typically keep it in the water for?

Less than one hour		7-8 hours	
1-2 hours		8-9 hours	
2-3 hours		9-10 hours	
3-4 hours		10-11 hours	
4-5 hours		11-12 hours	
5-6 hours		Over 12 hours	
6-7 hours		Do not use a keep net	

23. Do you ever enter the water whilst you fish?

Yes	
No	

24. Do you ever clean your equipment between trips?

Yes	
No	

If no, then please move to question 28

25. If yes, then how frequently?

After every trip	
After 2-5 trips	
After 6-10 trips	
After 11+ trips	

26. What do you use to clean your equipment?

Hot water	
Cold water	
Detergent	
Disinfectant	
Other (please state)	

27. How important are the following factors when deciding whether to clean your equipment after a trip?

Please rate the following from 1 to 5 91= not at all important, 5 extremely important. Only use each number once)

	1	2	3	4	5
The availability of a hose/ cleaning station					
The cost of cleaning your equipment					
The time it takes to clean your equipment					
The availability of information about what to do					
How clean your equipment looks at the end of your trip					

28. Do you ever dry your equipment for over 24 hours between each trip? e.g. hang on washing line or in airing cupboard?

Yes	
No	

If no, then please skip to question 30

29. If you do dry your equipment, how frequently?

After every trip	
After 2-5 trips	
After 6-10 trips	
After 11+ trips	

Bait

30. Do you ever use live bait ?

Yes	
No	

If no, please move to question 34

31. What type of bait do you use?

Fish	
Maggots	
Worms	
Other (please state)	

32. Where do you source your bait from?

Angling shop	
Collect from the wild	
Other (please state)	

33. What do you do with your bait at the end of your angling trip? (Choose all that apply)

Take it home and keep for next trip	
Take it home to dispose of it	
Leave at the water body	
Give to other anglers nearby	
Other (please state)	

The Check, Clean Dry Campaign

34. Have you heard of the Check, Clean Dry campaign in the UK?

Yes	
No	

If no, please move onto question 36

35. If yes, where did you hear about it?

Invasive Non-native Species

36. Have you heard of invasive species?

Yes	
No	

If no, then please move onto question 38

37. Please list any invasive species that you can name

Conservation Organisations

38. Are you part of any conservation organisations?

Yes	
No	

39. If yes, which organisations

40. Are you a member of any angling clubs or syndicates?

Yes	
No	

41. Can you please list these clubs/syndicates

You have now reached the end of the questionnaire. Thank-you for participating.

If you have any comments or questions about anything you have come across in this survey, please contact Emily Smith at Emily.smith@anglingtrust.net

References

- Anderson LG (2015) Managing aquatic non-native species: the role of biosecurity. Ph.D. Dissertation, University of Leeds, Leeds
- Anderson LG, White PCL, Stebbing PD, Stentiford GD, Dunn AM (2014) Biosecurity and vector behaviour: evaluating the potential threat posed by anglers and canoeists as pathways for the spread of invasive non-native species and pathogens. *PLoS ONE* 9:1–10
- Anderson LG, Dunn AM, Rosewarne PJ, Stebbing PD (2015) Invaders in hot water: a simple decontamination method to prevent the accidental spread of aquatic invasive non-native species. *Biol Invasions* 17:2287–2297
- Angelsen A, Lund JF (2011) Designing the household questionnaire. In: Angelsen A, Larsen HO, Lund JF, Smith-Hull C, Wunder S (eds) *Measuring livelihoods and environmental dependence: methods for research and fieldwork*. Earthscan, London, pp 107–126
- Bacela-Spychalska K, Grabowski M, Rewicz T, Konopacka A, Wattier R (2013) The killer shrimp *Dikerogammarus villosus* (Crustacea, Amphipoda) invading alpine lakes: overland transport by recreational boats and scuba-diving gear as potential entry vectors? *Aquat Conserv* 23:606–618
- Beninde J, Fischer ML, Hochkirch A, Zink A (2014) Ambitious advances of the European Union in the legislation of invasive alien species. *Conserv Lett* 49:1–17
- Beyer J, Moy P, Stasio B (2010) Acute upper thermal limits of three aquatic invasive invertebrates: hot water treatment to prevent upstream transport of invasive species. *Environ Manag* 47:67–76
- Brace I (2008) *Questionnaire design: how to plan, structure and write survey material for effective market research*. Kogan Page Publishers, London
- Bremner A, Park K (2007) Public attitudes to the management of invasive non-native species in Scotland. *Biol Conserv* 139:306–314
- Bruckerhoff L, Havel J, Knight S (2015) Survival of invasive aquatic plants after air exposure and implications for dispersal by recreational boats. *Hydrobiologia* 746:113–121
- Brundu G (2015) Plant invaders in European and Mediterranean inland waters: profiles, distribution, and threats. *Hydrobiologia* 1:61–79
- Caffrey JM, Baars J-R, Barbour JH, Boets P, Boon P, Davenport K, Dick JTA, Early J, Edsman L, Gallagher C, Gross J, Heiniman P, Horrill C, Hudin S, Hulme PE, Hynes S, MacIsaac HJ, McLoone P, Millane M, Moen TL, Newman J, O’Conchuir R, O’Farrell M, O’Flynn C, Oidtmann B, Renals T, Riccardi A, Roy H, Shaw R, Weyl O, Williams F, Lucy FE (2014) Tackling invasive alien species in Europe: the top 20 issues. *Manag Biol Invasions* 5:1–20. <https://doi.org/10.3391/mbi.2014.5.1.01>
- Caplat P, Coutts SR (2011) Integrating ecological knowledge, public perception and urgency of action into invasive species management. *Environ Manag* 48:878
- Cerri J, Ciappelli A, Lenuzza A, Nocita A, Zaccaroni M (2017) The randomised response technique: a valuable approach to monitor pathways of aquatic biological invasions. *Fish Manag Ecol* 24:504–511
- Cliff N, Campbell ML (2012) Perception as a tool to inform aquatic biosecurity risk assessments. *Aquat Invasions* 7:387–404
- Coughlan NE, Cuthbert RN, Kelly TC, Jansen MAK (2018) Parched plants: viability of invasive aquatic macrophytes following exposure to various desiccation regimes. *Aquat Bot* 150:9–15
- Couper MP, Miller PV (2008) Web survey methods introduction. *Public Opin Q* 72:831–835

- Couper MP, Kapteyn A, Schonlau M, Winter J (2007) Non-coverage and nonresponse in an internet survey. *Soc Sci Res* 36:131–148
- Crooks JA, Soulé ME, Sandlund OT (1999) Lag times in population explosions of invasive species: causes and implications. *Invasive Species Biodivers Manag* 103:125
- Cuthbert RN, Crane K, Caffrey JM, MacIssac HJ (2018) A dib or a dab: assessing the efficacy of Virasure® Aquatic disinfectant to reduce secondary spread of the invasive curly waterweed *Lagarosiphon major*. *Manag Biol Invasions* 9:3. <https://doi.org/10.3391/mbi.2018.9.3.08>
- Davis AJS, Darling JA (2017) Recreational freshwater fishing drives non-native aquatic species richness patterns at a continental scale. *Divers Distrib* 23:692–702
- De Ventura L, Weissert N, Tobias R, Kopp K, Jokela J (2017) Identifying target factors for interventions to increase boat cleaning in order to prevent spread of invasive species. *Manag Biol Invasions* 8:71–84
- Dobson A, Barker K, Taylor SL (2013) *Biosecurity: the socio-politics of invasive species and infectious diseases*. Routledge, London
- Drake DAR, Mandrak NE (2014) Ecological risk of live bait fisheries: a new angle on selective fishing. *Fisheries* 39:201–211
- Dunn AM, Hatcher MJ (2015) Parasites and biological invasions: parallels, interactions, and control. *Trends Parasitol* 31:189–199
- EA (2004) *Our nations' fisheries: the migratory and freshwater fisheries of England and Wales—a snapshot*. Bristol, Environment Agency
- Essl F, Bacher S, Blackburn TM, Booy O, Brundu G, Brunel S, Cardoso AC, Eschen R, Gallardo B, Galil B, Garcia-Berthou E (2015) Crossing frontiers in tackling pathways of biological invasions. *Bioscience* 65:769–782. <https://doi.org/10.1093/biosci/biv082>
- Fielding N (2011) *Dikergammarus villosus*: preliminary trials on resistance to control measures. *Freshw Biol Assoc Newslett* 54
- Finnoff D, Shogren JF, Leung B, Lodge D (2007) Take a risk: preferring prevention over control of biological invaders. *Ecol Econ* 62:216–222
- Forsman A (2014) Effects of genotypic and phenotypic variation on establishment are important for conservation, invasion, and infection biology. *PNAS* 111:302–307
- Gaddis SE (1998) How to design online surveys. *Train Dev J* 52:67–71
- Gallardo B, Aldridge DC (2013a) Priority setting for invasive species management of Ponto-Caspian invasive species into Great Britain. *Ecol Appl* 23:352–364
- Gallardo B, Aldridge DC (2013b) The dirty dozen: socio-economic factors amplify the invasion potential of 12 high risk aquatic invasive species in Great Britain and Ireland. *J Appl Ecol* 50:757–766
- Gallardo B, Aldridge DC (2015) Is Great Britain heading for a Ponto-Caspian invasional meltdown. *J Appl Ecol* 23:352–364
- Gallardo B, Clavero M, Sánchez MI, Vilà M (2016) Global ecological impacts of invasive species in aquatic ecosystems. *Glob Chang Biol* 22:151–163
- García-Llorente M, Martín-López B, González JA, Alcorlo P, Montes C (2008) Social perceptions of the impacts and benefits of invasive alien species: implications for management. *Biol Conserv* 141:2969–2983
- Gates KK, Guy CS, Zale AV, Horton TB (2009) Angler awareness of aquatic nuisance species and potential transport mechanisms. *Fish Manag Ecol* 16:448–456
- Genovesi P, Shine C (2004) European strategy on invasive alien Species: convention on the conservation of european wildlife and habitats (Bern Convention). CoC, Strasbourg. <https://www.cbd.int/doc/external/cop-09/bern-01-en.pdf>. Accessed 29 July 2018
- Gozlan RE, Burnard D, Andreou D, Britton JR (2013) Understanding the threats posed by non-native species: public vs. conservation managers. *PLoS ONE* 8:e53200. <https://doi.org/10.1371/journal.pone.0053200>
- Harrower CA, Scalera R, Pagad S, Schonrogge K, Roy HE (2018) Guidance for the interpretation of CBD categories on introduction pathways. IUCN, Gland
- Havel JE, Shurin JB (2004) Mechanisms, effects, and scales of dispersal in freshwater zooplankton. *Limnol Oceanogr* 49:1229–1238
- Hickley P (2018) Recreational fisheries—social, economic and management. In: Hickley P, Tompkins H (eds) *Recreational fisheries: social, economic and management aspects*. Wiley, Oxford
- Holdich DM, Sibley P, Peay S (2004) The White-clawed crayfish—a decade on. *Br Wildl* 15:153–164
- Hulme PE (2009) Trade, transport and trouble: managing invasive species pathways in an era of globalization. *J Appl Ecol* 46:10–18
- Hulme PE (2015) Invasion pathways at a crossroad: policy and research challenges for managing alien species introductions. *J Appl Ecol* 52:1418–1424
- Hussner A (2009) Growth and photosynthesis of four invasive aquatic plant species in Europe. *Weed Res* 49:506–515
- Jaeger TF (2008) Categorical data analysis: away from ANOVAs (transformation or not) and towards Logit Mixed Models. *J Mem Lang* 59:434–446
- Keller RP, Cox AN, Van Loon C, Lodge DM, Herborg LM, Rothlisberger J (2007) From bait shops to the forest floor: earthworm use and disposal by anglers. *Am Midl Nat* 158:321–328
- Kilian JV, Klauda RJ, Widman S, Kashiwagi M, Bourquin R, Weglein D, Schuster J (2012) An assessment of a bait industry and angler behaviour as a vector of invasive species. *Biol Invasions* 14:1469–1481
- Kolar CS, Lodge DM (2001) Progress in invasion biology: predicting invaders. *Trends Ecol Evol* 16:199–204
- Lajunen T, Summala H (2003) Can we trust self-reports of driving? Effects of impression management on driver behaviour questionnaire responses. *Transp Res Part F* 6:97–107
- Lavergne S, Molofsky J (2007) Increased genetic variation and evolutionary potential drive the success of invasive grass. *PNAS* 104:3883–3888
- Leung B, Lodge DM, Finnoff D, Shogren JF, Lewis MA, Lamberti G (2002) An ounce of prevention or a pound of cure: bioeconomic risk analysis of invasive species. *Proc R Soc Lond (Biol)* 269:2407–2413
- Longhurst R (2010) Semi-structured interviews and focus groups. In: Clifford N, French S, Valentine G (eds) *Key methods in geography*. SAGE, London, pp 103–115

- Mack R, Simberloff D, Lonsdale WM, Evans H, Clout M, Bazzaz FA (2000) Biotic invasions: causes, epidemiology, global consequences, and control. *Ecol Appl* 10:689–710
- Maitland PS (1987) Fish introductions and translocations—their impact in the British Isles. In: Maitland PS, Turner AK (eds) *Angling and wildlife in fresh waters*. ITE Symposium, pp 57–65
- Okada M, Grewell BJ, Jasieniuk M (2009) Clonal spread of invasive *Ludwigia hexapetala* and *L. grandiflora* in freshwater wetlands of California. *Aquat Bot* 91:123–129
- Peeler EJ, Gardiner RR, Thrush MA (2004) Qualitative risk assessment of routes of transmission of the exotic fish parasite *Gyrodactylus salaris* between river catchments in England and Wales. *Prev Vet Med* 64:175–189
- Perepelizin PV, Boltovskoy D (2011) Hot water treatment (chronic upper lethal temperature) mitigates biofouling by the invasive asian mussel *Limnoperna fortunei* in industrial installations. *Environ Sci Technol* 45:7868–7873
- Pigneur LM, Marescaux J, Roland K, Etoundi E, Descy JP, Van Doninck K (2011) Phylogeny and androgenesis in the invasive Corbicula clams (*Bivalvia*, *Corbiculidae*) in Western Europe. *BMC Evol Biol* 11:147–163
- Pitcher TJ, Hollingworth C (2002) *Recreational fisheries: ecological, economic and social evaluation*. Blackwell Science, London
- Prinbeck G, Lach D, Chan S (2011) Exploring stakeholders' attitudes and beliefs regarding behaviors that prevent the spread of invasive species. *Environ Educ Res* 17:341–352
- Pyšek P, Jarošík V, Pergl J (2011) Alien plants introduced by different pathways differ in invasion success: unintentional introductions as a threat to natural areas. *PLoS ONE* 6:1–11
- Rahel FJ, Olden JD (2008) Assessing the effects of climate change on aquatic invasive species. *Conserv Biol* 22:521–533
- Randall DM, Fernandes MF (1991) The social desirability response bias in ethics research. *J Bus Ethics* 10:805–817
- Ricciardi A (2015) Ecology of invasive alien invertebrates. In: Thorp JH, Rogers DC (eds) *Ecology and general biology: freshwater invertebrates*. Academic Press, Cambridge, pp 83–91
- Rogers EM (2003) *Diffusion of innovations*, 5th edn. Free Press, New York
- Roy HE, Peyton J, Aldridge DC, Bantock T, Blackburn TM, Britton R, Clark P, Cook E, Dehnen-Schmutz K, Dines T, Dobson M, Edwards F, Harrower C, Harvey MC, Minchin D, Noble DG, Parrott D, Pocock MJO, Preston CD, Roy S, Salisbury A, Schönrogge K, Sewell J, Shaw RH, Stebbing P, Stewart AJA, Walker KJ (2014) Horizon scanning for invasive alien species with the potential to threaten biodiversity in Great Britain. *Glob Change Biol* 20:3859–3871
- Schmidt WC (1997) World-Wide Web survey research: benefits, potential problems, and solutions. *Behav Res Methods Instrum Comp* 29:274–279
- Sebire M, Rimmer G, Hicks R, Parker SJ, Stebbing PD (2018) A preliminary investigation into biosecurity treatments to manage the invasive killer shrimp (*Dikergammarus villosus*). *Manag Biol Invasions* 9:101–113
- Shannon C, Quinn CH, Stebbing PD, Hassal C, Dunn A (2018) The application of hot water to reduce the introduction and spread of aquatic invasive alien species. *Manag Biol Invasions* 9:417–423
- Simberloff D, Martin J-L, Genovesi P, Maris V, Wardle DA, Aronson J, Courchamp F, Galil B, Garcia-Berthou E, Pascal M, Pysek P, Sousa R, Tabacchi E, Vila M (2013) Impacts of biological invasions: what's what and the way forward. *Trends Ecol Evol* 28:58–66
- Simpson D, Mawle GW (2010) *Public attitudes to angling 2010*. Bristol, Environment Agency
- Sports England (2011) *Fishing for answers: final report of the social and community benefits of angling project*. Substance, Bristol
- Stebbing PB, Sebire M, Lyons B (2011) Evaluation of a number of treatments to be used as biosecurity measures in controlling the spread of the invasive killer shrimp (*Dikergammarus villosus*)—final report. Lowestoft, Cefas
- Suarez AV, Holway DA, Case TJ (2001) Patterns of spread in biological invasions dominated by long-distance jump dispersal: insights from Argentine ants. *PNAS* 98:1095–1100
- Sutcliffe C, Quinn CH, Shannon C, Glover A, Dunn AM (2018) Exploring the attitudes to and uptake of biosecurity practices for invasive non-native species: views amongst stakeholder organisations working in UK natural environments. *Biol Invasions* 20:399–411
- Trouwborst A (2015) The Bern convention and EU regulation 1143/2014 on the prevention and management of the introduction and spread of invasive alien species. https://papers.ssrn.com/sol3/papers.cfm?abstract_id=2623496. Accessed 26 July 2018
- von Brandt A (1964) *Modern fishing of the world 2*. Fishing News Books Ltd, London
- White PCL, Jennings NV, Renwick AR, Barker NHL (2005) Questionnaires in ecology: a review of past use and recommendations for best practice. *J Appl Ecol* 42:421–430
- Williams AE, Moss B (2001) *Angling and conservation at sites of special scientific interest in England: economics, attitudes and impacts*. *Aquat Conserv* 11:357–372
- Winfield IJ, Adams CE, Fletcher JM (1996) Recent introductions of the ruffe (*Gymnocephalus cernuus*) to three United Kingdom lakes containing Coregonus species. *Ann Zool Fenn* 33:459–466
- Wittenberg R, Cock MJW (2001) *Invasive alien species: a toolkit of best prevention and management practices*. CABI, Wallingford
- Zięba G, Copp GH, Davies GD, Stebbing PD, Wesley KJ, Britton JR (2010) Recent releases and dispersal of non-native fishes in England and Wales, with emphasis on sunbleak *Leucaspis delineatus*. *Aquat Invasions* 5:155–161

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.