



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

A Study on Ningxia Desert Grassland (*Stipa breviflora*) Utilization

Kechang Li

Ningxia Grassland Station, China

Xin Wu

Ningxia Grassland Station, China

Jinhua Wang

Ningxia Grassland Station, China

Zhao Yu

Ningxia Grassland Station, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/9-4/5>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

A study on Ningxia desert grassland (*stipa breviflora*) utilization

Li Kechang , Wu Xin , Wang Jinhua , Yu Zhao

The Grassland Management and Service Station ,Ningxia Autonomous Region . 750002 ,Yinchuan

Key words : stipa breviflora ,desert grassland ,grazing density ,grazing regime ,control measures

Abstracts We studied the effect of grazing density ,grazing regime and related management measures on the desert grassland dominated by Stipa breviflora in the middle arid and semi-arid areas of Ningxia .

Study methods and contents The Sidunzi village in Yanchi county of Ningxia was selected as the trial area and the dominant plant specie was Stipa breviflora .The experimental area was 400hm² ,and the demonstration area was 800hm² . The selected animal was the Ningxia tan sheep .

Experimental results and findings

Grazing density The experimental results showed a strong negative relationship between the unit sheep weight and the grazing density . The regression equation was : $G_a = 18.48 - 7.01G$ ($r = -0.97^{**}$) ; A strong positive relationship existed between the unit grassland area (1hm²) weight and the grazing density ,The regression equation was : $G_h = 0.31 + 17.60G - 6.60G^2$ ($r = 0.87^*$) . It showed that the grazing density suitable for this grassland should be around 0.750 sheep/hm² (see chart 1) . When the grazing density exceeded 0.750 sheep/hm² ,ewes exhibited barrenness ,lower lambing rate and delayed pregnancy . A close relationship existed between barrenness or lambing rate and the grazing density ,respectively : $Y = -18.10 + 30.98X$ ($r = 0.97^{**}$) , $Y = 136.50 - 65.68X$ ($r = -0.95^*$) .




chart 1 Relations between the tan sheep unit weight increased and hectare weight increased under the different grazing densities

Grazing regime Assume 0.75 sheep/hm² as the grazing density and take the free grazing as a reference ,several rotational grazing experiments were begun in 2 to 6 blocks to test this grazing regime . The experimental results indicated that the biomass of the plant communities ,coverage and density in six rotation grazing blocks and those in four rotation grazing blocks were remarkably higher than the ones with free grazing ($P < 0.05$) . The trend of density and coverage of the palatable forage was as follows : grazing rotated in 6 blocks > grazing rotated in 4 blocks > grazing rotated in 2 blocks > free grazing . During the growing period the stocking capacities in six rotation graze blocks ,four rotation graze blocks and two rotation graze blocks separately were higher than the one in the free grazing area by 21.5% ,15.8% and 8% ,respectively . The weights of the tan sheep increased during the grazing period . This result showed that the density of 0.75 sheep /hm² was good for this grassland .

Management measures (1) Develop contracts with households or joint households ,(2) Build fences for all contracted grasslands which would not be larger than 200hm² ;(3) Define the grazing capacity for each grassland in accordance with the grazing density .(4) Combine several households into a joint grazing team who would be issued a Grazing Permit and implement rotational grazing in four to six rotated grazing blocks .

References

Mr . Li Kechang & Mr . Wu Yuanqing Study on Ningxia grassland science ,Ningxia People's Press .
Mr . Ma Ruiwen ,Scientific grassland using research ,Ningxia People's Press .