

T cell-mediated oral tolerance is intact in germ-free mice

K. L. W. Walton,^{*†} J. A. Galanko,[‡]
R. Balfour Sartor^{*‡§} and N. C. Fisher^{*‡¶}
^{*}Department of Medicine, [†]SPIRE Program,
[‡]Center for Gastrointestinal Biology and Disease,
and [§]Department of Microbiology and
Immunology, University of North Carolina,
Chapel Hill, NC, USA

Summary

Commensal enteric bacteria stimulate innate immune cells and increase numbers of lamina propria and mesenteric lymph node (MLN) T and B lymphocytes. However, the influence of luminal bacteria on acquired immune function is not understood fully. We investigated the effects of intestinal bacterial colonization on T cell tolerogenic responses to oral antigen compared to systemic immunization. Lymphocytes specific for ovalbumin–T cell receptor (OVA–TCR Tg⁺) were transplanted into germ-free (GF) or specific pathogen-free (SPF) BALB/c mice. Recipient mice were fed OVA or immunized subcutaneously with OVA peptide (323–339) in complete Freund's adjuvant (CFA). Although the efficiency of transfer was less in GF recipients, similar proportions of cells from draining peripheral lymph node (LN) or MLN were proliferating 3–4 days later *in vivo* in GF and SPF mice. In separate experiments, mice were fed tolerogenic doses of OVA and then challenged with an immunogenic dose of OVA 4 days later. Ten days after immunization, lymphocytes were restimulated with OVA *in vitro* to assess antigen-specific proliferative responses. At both high and low doses of OVA, cells from both SPF and GF mice fed OVA prior to immunization had decreased proliferation compared to cells from control SPF or GF mice. In addition, secretion of interferon (IFN)- γ and interleukin (IL)-10 by OVA–TCR Tg⁺ lymphocytes was reduced in both SPF and GF mice fed OVA compared to control SPF or GF mice. Unlike previous reports indicating defective humoral responses to oral antigen in GF mice, our results indicate that commensal enteric bacteria do not enhance the induction of acquired, antigen-specific T cell tolerance to oral OVA.

Keywords: commensal bacteria, germ-free mice, mucosal immunity, oral tolerance

Accepted for publication 14 December 2005
Correspondence: Kristen Walton, Department of
Medicine, 7317 MBRB, CB#7032, University of
North Carolina, Chapel Hill, NC 27599–7032,
USA.
E-mail: kristen_williams@med.unc.edu
[¶]Current address: Nancy C. Fisher, BD Technol-
ogies, Research Triangle Park, NC, USA.

Introduction

The intestinal mucosa is a major interface between an organism and its environment. Resident immune cells encounter continuously a wide variety of antigens, derived from food, commensal enteric flora or ingested pathogens. Thus, the intestinal immune system faces the challenge of maintaining a state of tolerance to self-antigens and non-pathogenic non-self-antigens, while remaining poised for a rapid, protective response to pathogens. Food antigens encountered in the gastrointestinal tract elicit typically a state of antigen-specific systemic hyporesponsiveness, called oral tolerance. Oral tolerance can occur through clonal deletion [1], anergy [2] or the generation of regulatory T cells [3,4], depending on fac-

tors such as antigen dose. Co-stimulatory molecules such as CD86 [5] and CTLA-4 [6], but not CD40 [7,8], are important in the induction of oral tolerance. In addition, regulatory cells producing transforming growth factor (TGF)- β [3,4] have been implicated in oral tolerance. There are mixed reports in the literature regarding the importance of interleukin (IL)-10, another cytokine considered widely to be immunoregulatory, in oral tolerance to soluble protein antigens [3,6,9,10]. Induction of a potentially regulatory CD4⁺/CD25⁺ cell population has also been demonstrated in low-dose oral tolerance models [10,11].

Alterations in the pathway leading to oral tolerance can cause serious consequences, including food hypersensitivity and pathogenic immune responses to the normal intestinal

microbiota. Indeed, loss of tolerance to commensal enteric bacteria has been demonstrated in inflammatory bowel diseases (IBD) [12–14]. In addition, many rodent models of IBD do not develop disease in a germ-free (GF) setting, but develop colitis rapidly when colonized with specific pathogen-free (SPF) bacteria [15–17] or selected enteric bacterial species [18], and exhibit Th1 responses to commensal enteric bacterial antigens [17,18]. Interestingly, a recent study reported reductions in oral tolerance to keyhole limpet haemocyanin, a soluble protein, in patients with IBD [19]. However, the mechanisms underlying tolerance to the commensal microbiota and the loss of tolerance in disease are not understood.

Commensal enteric bacteria also play an integral role in the development of the mucosal immune system. Significant alterations in gut-associated lymphoid tissues (GALT) occur in GF animals, including reduced numbers of intra-epithelial lymphocytes (IEL) [20], reduced numbers and size of Peyer's patches (PP) [21,22] and reduced secretion of IgA [23]. These alterations are normalized rapidly upon colonization with non-pathogenic commensal bacteria [20–22]. Bacterial antigen-specific T cells may participate in regulating gut immune responses. For example, bacterial antigen-specific T regulatory cells that secrete IL-10 in response to caecal bacterial lysates can suppress pathogenic Th1 cell proliferation and prevent colitis in a co-transfer model [24].

The importance of commensal bacteria in GALT development and gut immune responses supports a hypothesis that oral tolerance may be altered in GF mice. Early studies using sheep red blood cells as the antigenic stimulus showed that GF mice had defective antibody responses to oral antigen, which was restored by co-administration of lipopolysaccharide (LPS) [25]. Th1-dependent but not Th2-dependent antibody responses were reduced in GF mice after feeding with ovalbumin (OVA), while both types of responses were reduced in SPF mice [26]. The authors suggested that this was due to decreased numbers of T cells in GF PP [27]. In contrast, other studies have shown suppression of OVA-specific antibodies in the serum of OVA-fed GF mice to a similar degree to the suppression observed in conventionally housed mice, although the duration of oral tolerance was decreased [28,29]. Additionally, serum from OVA-fed GF BALB/c mice suppressed OVA-specific delayed-type hypersensitivity responses in recipient mice [30]. Differences among these studies may be due to a variety of factors such as antigens used, doses and feeding regimens and methods of assessing tolerance.

In the current studies, we utilized the adoptive transfer of OVA-specific-T cell receptor (OVA-TCR) Tg⁺ lymphocytes from DO11.10 transgenic mice into syngeneic GF and SPF mice to assess the impact of the absence of intestinal bacterial on oral tolerance at the level of T cell activation and sensitization. Because different doses of antigen may induce oral tolerance by different cellular mechanisms [1–4,10,29], we

used both high- and low-dose feeding regimens. We provide evidence that commensal enteric bacteria are not required for the induction of tolerance to oral antigen in adoptively transferred OVA-TCR Tg⁺ T cells.

Materials and methods

Mice

DO11.10 (OVA-TCR Tg⁺) breeder mice on a BALB/c background were originally obtained from D. Y. Loh (Washington University School of Medicine, St. Louis, MO, USA) [31]. SPF BALB/c mice were obtained from Harlan Sprague-Dawley (Indianapolis, IN, USA). BALB/c and DO11.10 mice were maintained under SPF conditions at the University of North Carolina animal facility. GF BALB/c mice were maintained in flexible plastic isolators at the Gnotobiotics Animal Facility of the Center for Gastrointestinal Biology and Disease, located at the Laboratory Animal Resources facility of the North Carolina State University College of Veterinary Medicine. The GF status of the mice was monitored by bi-weekly aerobic and anaerobic culture and Gram stain of stools from each isolator. In addition, stools were cultured from mice at the end of each experiment to confirm that the experimental mice remained sterile. Mice were used at 6–16 weeks of age. All experimental procedures were approved by the University of North Carolina and the North Carolina State University Animal Use and Care Committees.

Adoptive transfer and 5,6-carboxy-succinimidyl-fluorescein-ester (CFSE) labelling of donor cells

Peripheral (axillary and inguinal) and mesenteric lymph nodes from donor DO11.10 Tg⁺ mice were dissected into sterile phosphate-buffered saline (PBS) containing 0.1% bovine serum albumin (BSA) and 1% gentamicin to help ensure sterility. Single-cell suspensions were made by pressing the tissues through nylon mesh. Lymphocytes were washed in sterile PBS and injected intraperitoneally (i.p.) into recipient SPF and GF BALB/c mice. Intraperitoneal rather than intravenous injections were used due to technical issues related to handling GF mice in isolators. To track the proliferation of donor cells *in vivo*, donor lymphocytes were labelled with CFSE (Molecular Probes, Eugene, OR, USA), as described previously [32]. Briefly, unseparated DO11.10 lymphocytes were resuspended in sterile PBS containing 0.1% BSA and 1% gentamicin at 10⁷ cells/ml. CFSE was added to a final concentration of 10 μM and the cells were incubated at 37°C for 10 min. Labelled cells were washed once with sterile PBS containing 0.1% BSA and 1% gentamicin, washed once with sterile PBS and resuspended in PBS for intraperitoneal injection. Lymphocytes (3–8 × 10⁶) were injected into each mouse, depending on the experiment. Higher numbers of cells were used in later experiments after determining that OVA-TCR Tg⁺ cell

transplantation was typically very poor in GF mice given lower cell numbers.

Feeding and immunizations

For short-term experiments, 1 day after adoptive transfer of CFSE-labelled OVA-TCR Tg⁺ cells, mice were fed 100 mg OVA in 250 µl PBS by oral gavage, immunized subcutaneously (s.c.) with OVA p323–339 in complete Freund's adjuvant (CFA) or left untreated. Mice were killed 3 days after OVA treatment. For long-term experiments, mice were divided into fed and control groups. For high-dose experiments, 1 day after adoptive transfer of OVA-TCR Tg⁺ cells, the fed groups of GF and SPF mice were given 100 mg OVA in 250 µl PBS by oral gavage, and control mice were given no OVA. For low-dose experiments, the mice were given either sterile drinking water containing 1 mg/ml or 0.25 mg/ml OVA or sterile OVA-free drinking water. Mice were thus given OVA continuously for 5 days. In both high- and low-dose experiments, all mice were immunized s.c. with OVA peptide 323–339 in CFA 4 days after the last antigen encounter. In a separate control experiment, four GF and four SPF BALB/c mice that were not injected with OVA-TCR Tg⁺ cells were immunized s.c. with OVA p323–339 in CFA to test for reactivity of endogenous T cells to the peptide. Mice were necropsied 10 days after immunization.

Monoclonal antibodies (mAbs) and fluorescence activated cell sorter (FACS) analysis

The clonotypic anti-transgene monoclonal antibody, KJ1-26 [33] was prepared from hybridoma cells. IgG in hybridoma culture supernatants was purified by protein G chromatography (Pharmacia, Uppsala, Sweden) and was conjugated to FITC or biotin. Anti-CD4-phycoerythrin (PE) (CT-CD4) was purchased from Caltag (San Francisco, CA, USA). Anti-CD25-PE (PC61), anti-CD45RB-PE (16 A) and streptavidin-PerCP were purchased from BD Pharmingen (San Diego, CA, USA). Cell surface molecules were detected by staining 10⁶ viable cells with the antibody conjugates indicated for 20 min at 4°C in PBS containing 0.1% BSA and 0.01% sodium azide. Stained cells were washed and, as appropriate, cells were incubated with streptavidin-PerCP for an additional 5 min and then washed. Data were acquired on a FACScan (Becton Dickinson, Mountain View, CA, USA) and analysed with Summit software from Cytomation, Inc. (Ft Collins, CO, USA). A total of 100 000 size-gated lymphocytes were counted routinely.

Proliferation assay

Lymphocytes from the draining lymph node (LN) (axillary) of fed and control mice were cultured at 2 × 10⁵ cells/well in 200 µl complete medium with 10% fetal calf serum (FCS)

(HyClone, Logan, UT, USA), 50 µM 2-mercaptoethanol, 10 mg/ml folate, 2 mM glutamine, 1 mM pyruvate, 0.5% gentamicin and 0.5% penicillin–streptomycin and the concentrations indicated of OVA p323–339. An aliquot of cells from each mouse was subjected to FACS analysis to determine the number of OVA-TCR Tg⁺ cells that were present in each well. Mice with undetectable or very poor transplantation (less than 0.25% OVA-TCR Tg⁺ cells in the total lymphocyte population) were eliminated from further analysis. Cells were cultured for 3 days, with 1 µCi of [³H]-thymidine added for the last 18 h of culture. Cells were harvested onto glass fibre filters and the incorporated radioactivity was counted. The degree of transplantation of OVA-TCR Tg⁺ cells varied from animal to animal as well as across different experiments, and was typically lower in GF recipients (see Results section). To normalize the proliferation data, the counts per minute (cpm) per well were normalized to the number of input OVA-TCR Tg⁺ cells. The percentage of input transgenic cells in the lymphocytes harvested from each mouse was determined by FACS analysis on a separate aliquot of cells on the day of plating. The absolute number of OVA-TCR Tg⁺ cells was calculated by [(% OVA-TCR Tg⁺ cells/100)/2 × 10⁵ total cells plated]. Counts per minute above control per Tg⁺ cell was determined by the following equation: $\text{cpm}/\text{input Tg}^+ \text{ cell} = (\text{cpm}_{\text{stimulated}} - \text{cpm}_{\text{unstimulated}}) / \# \text{ input OVA-TCR Tg}^+ \text{ cells}$.

Analysis of cytokine secretion

Lymphocytes from the draining LN of fed and control SPF and GF mice were cultured in the presence or absence of 1 µM OVA p323–339 as described above for proliferation assays. Culture supernatants were collected after 72 h and frozen at –20 degrees until analysis. The concentrations of interferon (IFN)-γ and interleukin (IL)-10 in the supernatants were determined by a sandwich enzyme-linked immunosorbent assay (ELISA) (BD Pharmingen, San Jose, CA, USA), according to the manufacturer's instructions. The detection limits for both IFN-γ and IL-10 were 15 pg/ml.

Statistics

Student's *t*-test was used to compare differences between two means. For analysis of T cell proliferation experiments, a randomization test was performed. High- and low-dose experiments were analysed as separate groups. For each combination of control *versus* fed, GF or SPF and *in vitro* stimulation with OVA (0.01, 0.10 and 1.0 µM) a randomization test was performed by enumerating all possible sums of within-experiment ranks for the fed group and comparing it to the actual sum of within-experiment ranks for the fed group found in the data. In all cases, *P*-values < 0.05 were considered statistically significant.

Results

Transplantation of OVA-TCR Tg⁺ T cells is less efficient in GF mice

To assess antigen-induced proliferation and tolerance induction, pooled peripheral lymph node (PLN) and mesenteric lymph node (MLN) lymphocytes from OVA-TCR Tg⁺ DO11.10 donor mice were transferred adoptively by intraperitoneal injection into GF and SPF mice. The efficiency of cell transfer was consistently less in GF than in SPF recipients. In a typical experiment, GF BALB/c recipient mice had $1.10 \pm 0.10\%$ CD4⁺/KJ1-26⁺ cells in the PLN and $1.50 \pm 0.05\%$ CD4⁺/KJ1-26⁺ cells in the MLN 4 days after cell transfer, compared to $1.46 \pm 0.09\%$ in the PLN and $2.04 \pm 0.01\%$ in the MLN for SPF recipient mice given identical numbers of the same donor cells ($P < 0.05$ for GF *versus* SPF MLN, and $P = 0.09$ for PLN by Student's *t*-test, $n = 2-3$ for each tissue). Transplantation was not detectable in 20–30% of GF mice consistently across all experiments, regardless of whether or not the mice were exposed to antigen subsequent to adoptive transfer of OVA-TCR Tg⁺ cells. Mice with less than 0.25% CD4⁺/KJ1-26⁺ cells were dropped from all subsequent analyses. It has been reported that GF mice lack expression of secondary lymphoid chemokine (SLC) in PP [27], so it is possible that naive DO11.10-derived lymphocytes may not traffic appropriately to PP or MLN in GF mice. However, CD4⁺/KJ1-26⁺ cells were present typically in roughly similar proportions in the MLN compared to PLN of GF mice. In one experiment, the proportions of CD4⁺/KJ1-26⁺ cells were also assessed in the spleen and blood of recipient GF mice. GF mice with undetectable transplantation of CD4⁺/KJ1-26⁺ cells in the PLN and MLN also had undetectable transplantation in the spleen and blood (data not shown). This indicates that the reduced transplantation observed in the PLN and MLN of GF recipient mice was not likely to be caused by accumulation of adoptively transferred lymphocytes in the spleen or the blood.

OVA-TCR Tg⁺ T cells are activated by oral and subcutaneous OVA in GF mice

CFSE-labelled DO11.10 lymphocytes were transferred to GF and SPF mice, which were subsequently fed either 100 mg OVA, immunized s.c. with the antigenic OVA peptide 323–339 (OVA p323–339) emulsified in CFA or left untreated. Cells from the draining axillary PLN (immunized mice) or MLN (fed mice) were assessed by flow cytometry 3 days later. As shown in Fig. 1, CFSE⁺/KJ1-26⁺ lymphocytes in the PLN or MLN of GF and SPF recipients did not undergo a significant degree of proliferation in mice that were not exposed to OVA (control mice). Fewer CFSE⁺ cells were detected in the PLN and MLN of GF *versus* SPF recipients, as described above. However, in both GF and SPF mice immunized s.c. with OVA in CFA, significant proliferation of KJ1-26⁺ cells




Fig. 1. DO11.10 Tg⁺ cell proliferation in response to antigen in germ-free (GF) or specific pathogen-free (SPF) BALB/c mice. SPF or GF BALB/c mice injected with 5,6-carboxy-succinimidyl-fluorescein-ester (CFSE)-labelled DO11.10 lymphocytes were given no treatment (Control), immunized subcutaneously with ovalbumin (OVA) p323–339 in complete Freund's adjuvant (CFA) (Imm) or fed 100 mg OVA (Fed). Three days later, draining axillary lymph nodes (PLN) were collected from immunized and control mice, and draining mesenteric LN (MLN) were collected from fed and control mice. Proliferation of OVA-specific cells was assessed by decreased CFSE fluorescence. Each histogram shows data from an individual mouse and is representative of four separate experiments with two to four mice per group in each experiment.

was detected in lymphocytes from the draining PLN, based on decreased intensity of CFSE fluorescence (Fig. 1).

Similar T cell reactivities were seen in MLN cells from OVA-fed GF and SPF mice. In agreement with previous reports [7,32,34], MLN cells from fed SPF mice retained more CFSE fluorescence than cells from the draining PLN of immunized SPF mice (Fig. 1), indicating that less proliferation had occurred in the MLN of fed mice. The average number of cell divisions was increased slightly in the MLN of fed GF mice compared to fed SPF mice (GF, 3.4 ± 0.1 *versus* SPF, 2.8 ± 0.3) but this did not achieve statistical significance by




Fig. 2. Proportion of KJ1-26⁺ cells which had divided in germ-free (GF) or specific pathogen-free (SPF) mice. SPF or GF BALB/c were injected with 5,6-carboxy-succinimidyl-fluorescein-ester (CFSE)-labelled DO11.10 and treated as in Fig. 1. The data shown represent the mean \pm s.d. of CD4⁺/KJ1-26⁺ cells that had divided at least once, based on loss of CFSE fluorescence, expressed as a percentage of the total number of CD4⁺/KJ1-26⁺ cells. The bars labelled MLN (mesenteric lymph nodes) show data from MLN cells from fed mice, while the bars labelled PLN (peripheral lymph nodes) are axillary LN cells from immunized mice. No statistically significant differences were detected between GF and SPF mice in either group (GF *versus* SPF MLN, $P = 0.13$), $n = 4$ for each group.

Student's *t*-test ($P = 0.16$). Fewer total cells underwent proliferation in the PLN and MLN of GF mice compared to SPF mice following immunization or feeding, respectively. However, the proportion of cells that had proliferated relative to the total number of CFSE⁺/KJ1-26⁺ cells was again not significantly different in GF and SPF mice in either the fed or the immunized groups (Fig. 2). Taken together, these results indicate that DO11.10 lymphocytes transferred into GF mice are capable of responding to antigen presented by the native antigen-presenting cells in both the PLN and MLN, despite the less efficient transfer of OVA-TCR Tg⁺ cells in GF recipients.

Induction of high-dose oral tolerance to OVA is equal in GF and SPF mice

We next determined whether any defects were apparent in the induction of oral tolerance in GF mice. GF and SPF cell transfer recipient mice were separated into control groups, which received no OVA, and fed groups, which received 100 mg OVA by gavage. Four days after OVA feeding, all the mice were immunized s.c. with OVA p323-339 in CFA. Ten days later, cells from the draining axillary PLN of all mice were restimulated *in vitro* with OVA p323-339. Cells from control SPF and GF mice proliferated strongly in response to *in vitro* restimulation with OVA p323-339 (Fig. 3), showing sensitization of these cells in response to immunization. As expected, cells from SPF mice fed OVA had a significantly




Fig. 3. Oral tolerance is induced in germ-free (GF) mice given a high dose of antigen. Oral tolerance was induced in specific pathogen-free (SPF) and GF BALB/c mice as described in the Methods section. Fed mice were given 100 mg ovalbumin (OVA) by intragastric gavage. Control mice were left untreated. All mice were immunized by subcutaneous injection of OVA p323-339 in complete Freund's adjuvant (CFA); 10 days later, draining axillary LN were collected and cells were restimulated *in vitro* with ovalbumin peptide at the indicated concentrations. Proliferation was assessed by incorporation of methyl-[³H]-thymidine. Data are expressed as counts per minute (cpm) (stimulated - media) per input Tg⁺ cell. Data represent the mean \pm s.e.m. of triplicate wells for two to four mice per group and are representative of three similar experiments. Across all three experiments, proliferation was significantly ($P < 0.05$) inhibited in Fed *versus* Control for SPF at 0.01 and 0.1 μ M OVA p323-339 and for GF at 0.01 μ M OVA p323-339. The *P*-value for GF Fed *versus* Control at 0.1 μ M was 0.10.

diminished proliferation response to 0.01 and 0.1 μ M OVA ($P < 0.05$) (Fig. 3), indicating that these cells had been tolerized by oral OVA. Cells from OVA-fed GF mice showed a similar decrease in proliferative responses (Fig. 3), indicating that oral antigen in GF mice is capable of inducing a similar tolerization of DO11.10 T cells. A statistically significant reduction in DO11.10 cell proliferation in OVA-fed mice relative to control GF mice was not always achieved in a given experiment. This was due largely to the low sample size in individual experiments resulting from exclusion of GF mice with poor T cell transplantation from the analysis. However, analysis across three experiments determined that GF OVA-fed mice showed lower proliferation with 0.01 μ M OVA p323-339 ($P < 0.05$) and a trend towards lower proliferation with 0.1 μ M OVA p323-339 ($P = 0.10$) than the mean proliferation observed with the same concentration of OVA p323-339 in GF control mice.

In a separate control experiment, unmanipulated SPF and GF BALB/c mice were immunized with OVA p323-339 in CFA to test whether endogenous T cell priming was different in GF mice. Only minimal priming of endogenous OVA-TCR Tg⁺ cells was observed in both SPF and GF mice. In GF mice, proliferation of lymphocytes in response to *in vitro* restimulation with 1 μ M OVA p323-339 was 1.8 ± 0.5 -fold over proliferation in unstimulated lymphocytes, compared to 1.7 ± 0.6 -fold stimulation over control levels in SPF mice.

Induction of oral tolerance is intact in GF mice fed low-dose OVA

Tolerance induced by high doses of oral antigen is mediated primarily by clonal deletion or anergy [1,2], while lower doses of soluble antigens induce tolerance mediated primarily by regulatory T cells [3,4,10,11]. We reasoned that, while oral tolerance is intact in GF mice at high doses of antigen, possible differences might occur in the induction of regulatory T cells, which would be reflected by defective induction of oral tolerance by lower doses of antigen. Low-dose OVA feeding regimens vary in the literature but are generally less than 5 mg per day for multiple days, with antigen administration either by gavage or via the drinking water [3–5,9–11]. We chose two low doses, 1 mg/ml or 0.25 mg/ml, given continuously in the drinking water for 5 days. Continuous administration was chosen to minimize the required number of times to open the sterile isolators, in an effort to reduce the likelihood of contamination. GF mice drank slightly higher quantities of OVA-containing water (GF, 4.44 ± 0.50 ml/day *versus* 3.17 ± 0.24 ml/day for SPF, $P < 0.01$). However, both groups drank quantities of OVA (4.4 *versus* 3.2 mg/day and 1.1 *versus* 0.8 mg/day for 5 days for 1 mg/ml and 0.25 mg/ml experiments, respectively) which were well within typical ranges used for low-dose tolerance experiments [3–5,9–11]. These mice consumed very similar doses of OVA to mice from a study which demonstrated intact Th2-dependent antibody responses in GF mice despite oral administration of OVA [26]. Results from a representative experiment using 1 mg/ml OVA for 5 days are shown in Fig. 4. No differences between GF and SPF mice fed OVA were observed in the inhibition of proliferative responses upon *in vitro* restimulation. Similar inhibition was obtained by feeding 0.25 mg/ml OVA for 5 days in both GF and SPF mice (data not shown). When the low-dose experiments were analysed together, proliferation in fed groups was significantly reduced relative to control groups for both GF and SPF at all doses of OVA peptide restimulation ($P < 0.05$ for fed *versus* control mice).

IFN- γ responses are suppressed in SPF and GF mice after oral tolerance induction

Exposure to tolerogenic doses of antigen results typically in suppression of cytokine recall responses, in addition to suppression of proliferative recall responses. Therefore, we next measured whether suppression of cytokine secretion by DO11.10 T cells could be observed in cells from control or OVA-fed SPF and GF mice. Lymphocytes were isolated and restimulated as described for the proliferation experiments, and representative samples from four independent experiments were assessed by ELISA for IFN- γ and IL-10. Samples from one high-dose experiment and three low-dose experiments were analysed, and the pooled data are shown in Fig. 5. IFN- γ secretion was suppressed in OVA-TCR Tg⁺ cells from SPF mice that were fed OVA prior to immunization




Fig. 4. Oral tolerance is induced in germ-free (GF) mice given a low, continuous dose of antigen. Fed mice were given ovalbumin (OVA) continuously for 5 days in the drinking water at 1 mg/ml. Control mice were given normal drinking water. All mice were immunized by subcutaneous injection of OVA p323–339 in CFA; 10 days later, draining axillary LN were collected and restimulated *in vitro* with ovalbumin peptide. Proliferation was assessed by incorporation of methyl-³H-thymidine. Data are expressed as counts per minute (cpm) (stimulated – media) per input Tg⁺ cell. Data represent the mean \pm s.e.m. of triplicate wells for two to four mice per group and are representative of six similar experiments (three experiments using 1 mg/ml OVA and three experiments using 0.25 mg/ml OVA). Across all low-dose experiments, proliferation in Fed groups was statistically less than in Control groups for both GF and specific pathogen-free (SPF) at all doses of OVA p323–339 ($P < 0.05$).

compared to control SPF mice that were not fed prior to immunization (Fig. 5a). IFN- γ secretion was also reduced significantly in OVA-TCR Tg⁺ cells from fed GF mice compared to control GF mice. Interestingly, significantly lower levels of IFN- γ were observed in control GF mice compared to control SPF mice (3.8 ± 1.1 pg/ml per input OVA-TCR Tg⁺ cell for GF controls *versus* 16.6 pg/ml per input OVA-TCR Tg⁺ cell for SPF controls, $P < 0.05$). IL-10 was low but detectable in SPF control mice (Fig. 5b) and was suppressed to levels near or below the assay detection limit in SPF mice that were fed OVA. In contrast, IL-10 secretion was at or very near the assay detection limit in both control and fed GF mice (Fig. 5b). Neither IFN- γ nor IL-10 were detectable in lymphocytes from SPF or GF mice that were not transferred adoptively with OVA-TCR Tg⁺ cells prior to immunization with OVA p323–339 (data not shown).

Discussion

The current studies tested the requirement for commensal intestinal bacteria in the induction of oral tolerance to a soluble protein antigen. We found that T cell oral tolerance responses to high- and low-dose OVA were intact in GF mice, contrary to some [25–27] but not all [28–30] previous reports. These conflicting data may be due to several factors, including the model systems used. Our experiments




Fig. 5. Secretion of cytokines is reduced in ovalbumin-T cell receptor (OVA-TCR) Tg⁺ cells from germ-free (GF) or specific pathogen-free (SPF) mice fed oral antigen. Lymphocytes were isolated from control or OVA-fed SPF and GF mice and restimulated *in vitro* with OVA p323–339. Supernatants were collected after 72 h of culture and were analysed by enzyme-linked immunosorbent assay (ELISA) for interferon (IFN)- γ (a) and interleukin (IL)-10 (b). For each cytokine, the detected concentrations (in pg/ml) were normalized to the number of OVA-TCR Tg⁺ cells in each well to yield pg/ml/OVA-TCR Tg⁺ cell. Data represent the mean \pm s.e.m. for six to eight mice per group and include representative mice from four independent experiments, including both experiments using both high-dose (one experiment) and low-dose (three experiments) feeding regimens. * $P < 0.05$ versus SPF control, # $P < 0.05$ versus GF control.

addressed tolerance in a widely used model, in which monoclonal antigen-specific T cells from SPF DO11-10 mice were transferred adoptively into SPF or GF BALB/c mice. In contrast, prior studies have assessed polyclonal oral tolerance in unmanipulated GF mice based on antibody titres following oral antigen administration.

Reports of oral tolerance limiting antibody responses in GF mice included responses to sheep red blood cells and soluble antigens. Wannemuehler *et al.* [25] showed that GF BALB/c and Swiss mice did not develop tolerance in response to orally administered sheep red blood cells, as determined by splenocyte anti-sheep red blood cell antibody responses. The same group showed in a separate report that LPS-resistant C3H/HeJ mice did not develop humoral oral

tolerance [35]. Another report found that C3H/HeJ mice, as well as TLR-4 defective mice on a C57BL6 \times 129 background, had elevated allergic responses to peanut allergens co-administered orally with cholera toxin [36]. In addition, antibiotic treatment of TLR-4 wild-type mice recapitulated the elevated IgE levels seen in TLR-4-defective mice [36]. IgE and IgG1, but not IgG2, antibody production was suppressed in GF C3H/HeJ mice fed β -lactoglobulin, although the magnitude of suppression was reduced relative to conventionally housed mice [37]. This study also reported a lack of suppression of splenocyte proliferation in GF β -lactoglobulin-fed mice [37]. Sudo *et al.* demonstrated defective oral tolerance in GF BALB/c mice, measured by serum OVA-specific IgE titres in response to challenge with OVA and aluminium hydroxide, which causes an allergic reaction [26]. Interestingly, however, that study also showed suppression of IFN- γ secretion by splenocytes isolated from tolerized GF mice [26], which agrees with the suppression of IFN- γ recall responses observed in our study. Our data show that adoptively transferred OVA-TCR Tg⁺ T cells from an SPF donor mouse are tolerated in a GF setting, demonstrated by suppression of both proliferation and of IFN- γ secretion in OVA-TCR Tg⁺ cells. Collectively, these studies suggest that commensal bacteria influence humoral oral tolerance more strongly than cellular oral tolerance. However, T cell recall responses, as measured by IFN- γ secretion, were decreased in GF mice, although proliferative responses were similar to those observed in SPF mice.

In our low-dose oral tolerance experiments, GF mice drank slightly more OVA protein solution than did SPF mice, but no significant differences in induction of tolerance were observed. It is possible that uptake of OVA following feeding is different in GF *versus* SPF mice. We did not measure directly serum levels of OVA following feeding and so cannot rule out possible differences in uptake of the protein. However, the similar *in vivo* proliferation observed in GF and SPF mice at 3 days after feeding provides evidence that immunologically active OVA is taken up and presented in both SPF and GF mice.

The role of PP in the induction of oral tolerance has been the subject of conflicting reports. Defective oral tolerance in GF BALB/c mice has been linked to decreased T cell numbers in the PP of GF mice [27]. However, there are conflicting reports on the role of PP in induction of oral tolerance [38–40]. Interestingly, the MLN may be a more critical site for oral tolerance induction, as mice lacking both MLN and PP do not show oral tolerance [39,40]. Indeed, fed antigen can be detected in antigen-presenting cells in the MLN as well as PP within a few hours of feeding, and antigen-specific T cell proliferation was induced to a similar extent in the MLN, PLN and spleen of normal mice fed OVA compared to mice made PP-deficient by *in utero* exposure to an anti-IL-7 receptor- α antibody [41]. Subepithelial dendritic cells in areas away from the PP can sample luminal antigens via intra-epithelial dendrites [42]. Our study provides strong evidence

that both early T cell proliferative responses and functional tolerance are not impaired in GF mice, which have relatively underdeveloped PP [21–23]. Taken together with the studies discussed above, these data suggest that PP may not be the sole site of oral tolerance induction.

A decrease in the overall efficiency of transplantation of adoptively transferred DO11.10 cells was noted consistently in GF mice in our studies. Reduced trafficking of these naive DO11.10 cells to mucosal tissues, as might be expected based on previous reports of defective SLC expression in the mucosa of GF mice [27] and T cell transfer in the mucosa of reduced-flora SCID mice [43,44], did not appear to underlie the decrease in transplantation, as the cells did not traffic selectively to other lymphoid compartments such as the PLN, spleen or blood. An important point is that the observed decrease in transplantation efficiency did not affect the functional outcome, as long as T cell transplantation was detectable. DO11.10 cells in GF mice were able to be tolerized upon encounter with oral antigen to a similar extent to those cells injected into SPF mice.

Various populations of regulatory CD4⁺ T cells have been described in oral tolerance, including IL-10-producing Tr1 cells [45], Th3 cells which produce TGF- β [46,47] and CD4⁺/CD25⁺ cells [10,11]. The relative importance of each of these cell populations in oral tolerance is currently unknown. Recent studies have pointed to a role for TGF- β in continuous-feeding oral tolerance models preferentially at lower doses of antigen [3]. Alterations in TGF- β -secreting populations of OVA-TCR Tg⁺ cells in GF mice cannot be ruled out in our studies. We did not assess TGF- β secretion in this study due to the high baseline levels present in FBS added to the culture media. Some [3,48] but not all [6,49] studies have indicated that IL-10 may be a mediator of oral tolerance. In the current studies, IL-10 was suppressed in cultures of cells from SPF tolerized mice and was secreted in very low levels in cultures of cells from control and tolerized GF mice, regardless of the feeding regimen used to induce oral tolerance. However, these T cells were clearly hypoproliferative in response to *in vitro* restimulation with antigen, and thus any possible alterations in cytokine secretion did not affect the induction of T cell hyporesponsiveness. CD25 expression on antigen-specific T cells with regulatory T cell activity has been shown previously in models of oral tolerance [49,50]. In our model system, induction of CD25 was not observed on OVA-specific CD4⁺ cells in SPF or GF mice on day 14 after feeding high or low doses of OVA (data not shown), although we did not test further for regulatory activity or for expression of Foxp3, a transcription factor expressed in regulatory CD25⁺/CD4⁺ T cells [51]. It is possible that native BALB/c regulatory cells suppressed the OVA-TCR Tg⁺ cells. Indeed, the presence of CD4⁺/CD25⁺ cells with suppressive capabilities in the MLN of GF Swiss-Webster mice has been reported recently [52].

While GF mice are free of viable bacteria, the standard GF environment still contains trace amounts of LPS and other

bacteria-derived products from sterilized food or bedding. Bacteria antigens are presented to gut T cells by dendritic cells, and LPS or CpG alone are sufficient to activate dendritic cells to block CD25⁺ T cell-mediated suppression [53]. Thus, T cell responses dependent on bacterial stimulation may be intact in GF mice due to the presence of dietary LPS or CpG. In addition, the OVA-specific T cells used in this study, although naive in terms of OVA exposure, were generated in SPF mice and thus were exposed previously to bacteria. We cannot rule out the possibility that T cells derived from GF mice might have defective oral tolerance. However, to our knowledge there are currently no TCR transgenic mice that have been made GF, and the adoptive transfer of naive OVA-TCR Tg⁺ cells is therefore a useful and valid approach to assess T cell responses to oral antigen. Our findings provide strong evidence that oral antigens are taken up and presented effectively to T cells in GF mice, and that naive T cells are capable of becoming tolerized to oral OVA in the GF environment.

Acknowledgements

The authors gratefully thank Dr Sue Tonkonogy for use of facilities and equipment and Donna Kronstadt for assistance with germ-free mice. The Gnotobiotic and Biostatistics Core facilities of the Center for Gastrointestinal Biology and Disease are supported by NIH grant P30 DK34987. KLWW is supported by the SPIRE program, funded by the MORE division of the National Institutes of General Medical Sciences grant no. GM00678. This work is also funded by NIH grants RO1 DK53347 (RBS) and A1041579 (NCF).

References

- Chen Y, Inobe J, Marks R, Gonnella P, Kuchroo VK, Weiner HL. Peripheral deletion of antigen-reactive T cells in oral tolerance. *Nature* 1995; **376**:177–80.
- Van Houten N, Blake SF. Direct measurement of anergy of antigen-specific T cells following oral tolerance induction. *J Immunol* 1996; **157**:1337–41.
- Faria AM, Maron R, Ficker SM, Slavin AJ, Spahn T, Weiner HL. Oral tolerance induced by continuous feeding: enhanced up-regulation of transforming growth factor-beta/interleukin-10 and suppression of experimental autoimmune encephalomyelitis. *J Autoimmun* 2003; **20**:135–45.
- Weiner HL. Oral tolerance: immune mechanisms and the generation of Th3-type TGF-beta-secreting regulatory cells. *Microbes Infect* 2001; **3**:947–54.
- Liu L, Kuchroo VK, Weiner HL. B7.2 (CD86) but not B7.1 (CD80) costimulation is required for the induction of low dose oral tolerance. *J Immunol* 1999; **163**:2284–90.
- Fowler S, Powrie F. CTLA-4 expression on antigen-specific cells but not IL-10 secretion is required for oral tolerance. *Eur J Immunol* 2002; **32**:2997–3006.
- Sun J, Van Houten N. CD40 stimulation *in vivo* does not inhibit CD4⁺ T cell tolerance to soluble antigens. *Immunol Lett* 2002; **84**:125–32.

- 8 Chung Y, Ko SY, Ko HJ, Kang CY. Split peripheral tolerance: CD40 ligation blocks tolerance induction for CD8 T cells but not for CD4 T cells in response to intestinal antigens. *Eur J Immunol* 2005; **35**:1381–90.
- 9 Marth T, Ring S, Schulte D *et al.* Antigen-induced mucosal T cell activation is followed by Th1 T cell suppression in continuously fed ovalbumin TCR-transgenic mice. *Eur J Immunol* 2000; **30**:3478–86.
- 10 Zhang X, Izikson L, Liu L, Weiner HL. Activation of CD25(+) CD4(+) regulatory T cells by oral antigen administration. *J Immunol* 2001; **167**:4245–53.
- 11 Thorstenson KM, Khoruts A. Generation of anergic and potentially immunoregulatory CD25⁺CD4 T cells *in vivo* after induction of peripheral tolerance with intravenous or oral antigen. *J Immunol* 2001; **167**:188–95.
- 12 Macpherson A, Khoo UY, Forgacs I, Philpott-Howard J, Bjarnason I. Mucosal antibodies in inflammatory bowel disease are directed against intestinal bacteria. *Gut* 1996; **38**:365–75.
- 13 Duchmann R, Kaiser I, Hermann E, Mayet W, Ewe K, Meyer zum Buschenfelde KH. Tolerance exists towards resident intestinal flora but is broken in active inflammatory bowel disease (IBD). *Clin Exp Immunol* 1995; **102**:448–55.
- 14 Duchmann R, May E, Heike M, Knolle P, Neurath M, Meyer zum Buschenfelde KH. T cell specificity and cross reactivity towards enterobacteria, bacteroides, bifidobacterium, and antigens from resident intestinal flora in humans. *Gut* 1999; **44**:812–8.
- 15 Schultz M, Tonkonogy SL, Sellon RK *et al.* IL-2-deficient mice raised under germfree conditions develop delayed mild focal intestinal inflammation. *Am J Physiol* 1999; **276**:G1461–72.
- 16 Sellon RK, Tonkonogy S, Schultz M *et al.* Resident enteric bacteria are necessary for development of spontaneous colitis and immune system activation in interleukin-10-deficient mice. *Infect Immun* 1998; **66**:5224–31.
- 17 Veltkamp C, Tonkonogy SL, De Jong YP *et al.* Continuous stimulation by normal luminal bacteria is essential for the development and perpetuation of colitis in Tg (epsilon26) mice. *Gastroenterology* 2001; **120**:900–13.
- 18 Kim SC, Tonkonogy SL, Albright CA *et al.* Variable phenotypes of enterocolitis in interleukin 10-deficient mice monoassociated with two different commensal bacteria. *Gastroenterology* 2005; **128**:891–906.
- 19 Kraus TA, Toy L, Chan L, Childs J, Mayer L. Failure to induce oral tolerance to a soluble protein in patients with inflammatory bowel disease. *Gastroenterology* 2004; **126**:1771–8.
- 20 Umetsaki Y, Setoyama H, Matsumoto S, Okada Y. Expansion of alpha beta T-cell receptor-bearing intestinal intraepithelial lymphocytes after microbial colonization in germ-free mice and its independence from thymus. *Immunology* 1993; **79**:32–7.
- 21 Shroff KE, Cebra JJ. Development of mucosal humoral immune responses in germ-free (GF) mice. *Adv Exp Med Biol* 1995; **371A**:441–6.
- 22 Pollard M, Sharon N. Responses of the Peyer's patches in germ-free mice to antigenic stimulation. *Infect Immun* 1970; **2**:96–100.
- 23 Talham GL, Jiang HQ, Bos NA, Cebra JJ. Segmented filamentous bacteria are potent stimuli of a physiologically normal state of the murine gut mucosal immune system. *Infect Immun* 1999; **67**:1992–2000.
- 24 Cong Y, Weaver CT, Lazenby A, Elson CO. Bacterial-reactive T regulatory cells inhibit pathogenic immune responses to the enteric flora. *J Immunol* 2002; **169**:6112–9.
- 25 Wannemuehler MJ, Kiyono H, Babb JL, Michalek SM, McGhee JR. Lipopolysaccharide (LPS) regulation of the immune response: LPS converts germfree mice to sensitivity to oral tolerance induction. *J Immunol* 1982; **129**:959–65.
- 26 Sudo N, Sawamura S, Tanaka K, Aiba Y, Kubo C, Koga Y. The requirement of intestinal bacterial flora for the development of an IgE production system fully susceptible to oral tolerance induction. *J Immunol* 1997; **159**:1739–45.
- 27 Maeda Y, Noda S, Tanaka K *et al.* The failure of oral tolerance induction is functionally coupled to the absence of T cells in Peyer's patches under germfree conditions. *Immunobiology* 2001; **204**:442–57.
- 28 Moreau MC, Corthier G. Effect of the gastrointestinal microflora on induction and maintenance of oral tolerance to ovalbumin in C3H/HeJ mice. *Infect Immun* 1988; **56**:2766–8.
- 29 Moreau MC, Gaboriau-Routhiau V. The absence of gut flora, the doses of antigen ingested and aging affect the long-term peripheral tolerance induced by ovalbumin feeding in mice. *Res Immunol* 1996; **147**:49–59.
- 30 Furrir E, Turner MW, Strobel S. Failure of SCID mice to generate an oral tolerogen after a feed of ovalbumin: a role for a functioning gut-associated lymphoid system. *Immunology* 1994; **83**:562–7.
- 31 Murphy KM, Heimberger AB, Loh DY. Induction by antigen of intrathymic apoptosis of CD4⁺CD8⁺TCR^{lo} thymocytes *in vivo*. *Science* 1990; **250**:1720–3.
- 32 Sun J, Dirden-Kramer B, Ito K, Ernst PB, Van Houten N. Antigen-specific T cell activation and proliferation during oral tolerance induction. *J Immunol* 1999; **162**:5868–75.
- 33 Haskins K, Kubo R, White J, Pigeon M, Kappler J, Marrack P. The major histocompatibility complex-restricted antigen receptor on T cells. I. Isolation with a monoclonal antibody. *J Exp Med* 1983; **157**:1149–69.
- 34 Smith KM, Davidson JM, Garside P. T-cell activation occurs simultaneously in local and peripheral lymphoid tissue following oral administration of a range of doses of immunogenic or tolerogenic antigen although tolerized T cells display a defect in cell division. *Immunology* 2002; **106**:144–58.
- 35 Kiyono H, McGhee JR, Wannemuehler MJ, Michalek SM. Lack of oral tolerance in C3H/HeJ mice. *J Exp Med* 1982; **155**:605–10.
- 36 Bashir ME, Louie S, Shi HN, Nagler-Anderson C. Toll-like receptor 4 signaling by intestinal microbes influences susceptibility to food allergy. *J Immunol* 2004; **172**:6978–87.
- 37 Prioult G, Fliss I, Pecquet S. Effect of probiotic bacteria on induction and maintenance of oral tolerance to beta-lactoglobulin in gnotobiotic mice. *Clin Diagn Lab Immunol* 2003; **10**:787–92.
- 38 Fujihashi K, Dohi T, Rennert PD *et al.* Peyer's patches are required for oral tolerance to proteins. *Proc Natl Acad Sci USA* 2001; **98**:3310–5.
- 39 Spahn TW, Fontana A, Faria AM *et al.* Induction of oral tolerance to cellular immune responses in the absence of Peyer's patches. *Eur J Immunol* 2001; **31**:1278–87.
- 40 Spahn TW, Weiner HL, Rennert PD *et al.* Mesenteric lymph nodes are critical for the induction of high-dose oral tolerance in the absence of Peyer's patches. *Eur J Immunol* 2002; **32**:1109–13.
- 41 Kunkel D, Kirchhoff D, Nishikawa S, Radbruch A, Scheffold A. Visualization of peptide presentation following oral application of antigen in normal and Peyer's patches-deficient mice. *Eur J Immunol* 2003; **33**:1292–301.

- 42 Rescigno M, Urbano M, Valzasina B *et al.* Dendritic cells express tight junction proteins and penetrate gut epithelial monolayers to sample bacteria. *Nat Immunol* 2001; **2**:361–7.
- 43 Aranda R, Sydora BC, McAllister PL *et al.* Analysis of intestinal lymphocytes in mouse colitis mediated by transfer of CD4⁺, CD45RB^{high} T cells to SCID recipients. *J Immunol* 1997; **158**:3464–73.
- 44 Camerini V, Sydora BC, Aranda R *et al.* Generation of intestinal mucosal lymphocytes in SCID mice reconstituted with mature, thymus-derived T cells. *J Immunol* 1998; **160**:2608–18.
- 45 Groux H, O'Garra A, Bigler M *et al.* A CD4⁺ T-cell subset inhibits antigen-specific T-cell responses and prevents colitis. *Nature* 1997; **389**:737–42.
- 46 Chen Y, Inobe J, Kuchroo VK, Baron JL, Janeway CA Jr, Weiner HL. Oral tolerance in myelin basic protein T-cell receptor transgenic mice: suppression of autoimmune encephalomyelitis and dose-dependent induction of regulatory cells. *Proc Natl Acad Sci USA* 1996; **93**:388–91.
- 47 Fukaura H, Kent SC, Pietruszewicz MJ, Khoury SJ, Weiner HL, Hafler DA. Induction of circulating myelin basic protein and proteolipid protein-specific transforming growth factor-beta1-secreting Th3 T cells by oral administration of myelin in multiple sclerosis patients. *J Clin Invest* 1996; **98**:70–7.
- 48 Tsuji NM, Mizumachi K, Kurisaki J. Interleukin-10-secreting Peyer's patch cells are responsible for active suppression in low-dose oral tolerance. *Immunology* 2001; **103**:458–64.
- 49 Omata N, Ohshima Y, Yasutomi M, Yamada A, Karasuyama H, Mayumi M. Ovalbumin-specific IgE modulates ovalbumin-specific T-cell response after repetitive oral antigen administration. *J Allergy Clin Immunol* 2005; **115**:822–7.
- 50 Hauet-Broere F, Unger WW, Garssen J, Hoiijer MA, Kraal G, Samsom JN. Functional CD25⁻ and CD25⁺ mucosal regulatory T cells are induced in gut-draining lymphoid tissue within 48 h after oral antigen application. *Eur J Immunol* 2003; **33**:2801–10.
- 51 Sakaguchi S. Naturally arising Foxp3-expressing CD25⁺CD4⁺ regulatory T cells in immunological tolerance to self and non-self. *Nat Immunol* 2005; **6**:345–52.
- 52 Gad M, Pedersen AE, Kristensen NN, Claesson MH. Demonstration of strong enterobacterial reactivity of CD4⁺CD25⁻ T cells from conventional and germ-free mice which is counter-regulated by CD4⁺CD25⁺ T cells. *Eur J Immunol* 2004; **34**:695–704.
- 53 Pasare C, Medzhitov R. Toll pathway-dependent blockade of CD4⁺CD25⁺ T cell-mediated suppression by dendritic cells. *Science* 2003; **299**:1033–6.